

Yves Poumay

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/95916/yves-poumay-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

91
papers

2,680
citations

28
h-index

50
g-index

131
ext. papers

3,002
ext. citations

3.9
avg, IF

4.71
L-index

#	Paper	IF	Citations
91	Preclinical assessment of dual CYP26[A1/B1] inhibitor, DX308, as an improved treatment for keratinization disorders. <i>Skin Health and Disease</i> , 2021 , 1, e22		0
90	Characterization of CYP26B1-Selective Inhibitor, DX314, as a Potential Therapeutic for Keratinization Disorders. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 72-83.e6	4.3	2
89	Experimental Models of Dermatophytosis 2021 , 135-160		1
88	HDAC2 and 7 down-regulation induces senescence in dermal fibroblasts. <i>Aging</i> , 2021 , 13, 17978-18005	5.6	3
87	Deletion of Gene in Human Keratinocytes Demonstrates a Role for TSG-6 to Retain Hyaluronan Inside Epidermis.. <i>JID Innovations</i> , 2021 , 1, 100054		1
86	Skin Disease Models In Vitro and Inflammatory Mechanisms: Predictability for Drug Development. <i>Handbook of Experimental Pharmacology</i> , 2021 , 265, 187-218	3.2	5
85	Histological study of sheep skin transformation during the recreation of historical parchment manufacture. <i>Heritage Science</i> , 2020 , 8,	2.5	3
84	Responses of Reconstructed Human Epidermis to Trichophyton rubrum Infection and Impairment of Infection by the Inhibitor PD169316. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 2080-2089.e6	4.3	7
83	Use of Polarised Light Microscopy to Improve Conservation of Parchment. <i>Studies in Conservation</i> , 2019 , 64, 284-297	0.6	1
82	Methyl-β-cyclodextrin concurs with interleukin (IL)-4, IL-13 and IL-25 to induce alterations reminiscent of atopic dermatitis in reconstructed human epidermis. <i>Experimental Dermatology</i> , 2018 , 27, 435-437	4	16
81	In vitro models of dermatophyte infection to investigate epidermal barrier alterations. <i>Experimental Dermatology</i> , 2018 , 27, 915-922	4	15
80	Non-senescent keratinocytes organize in plasma membrane submicrometric lipid domains enriched in sphingomyelin and involved in re-epithelialization. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2017 , 1862, 958-971	5	5
79	Modeling dermatophytosis in reconstructed human epidermis: A new tool to study infection mechanisms and to test antifungal agents. <i>Medical Mycology</i> , 2017 , 55, 485-494	3.9	15
78	Methyl-β-cyclodextrin treatment combined to incubation with interleukin-4 reproduces major features of atopic dermatitis in a 3D-culture model. <i>Archives of Dermatological Research</i> , 2017 , 309, 63-69	3.3	10
77	Atopic Dermatitis Studies through Models. <i>Frontiers in Medicine</i> , 2017 , 4, 119	4.9	26
76	Hyaluronan Does Not Regulate Human Epidermal Keratinocyte Proliferation and Differentiation. <i>Journal of Biological Chemistry</i> , 2016 , 291, 6347-58	5.4	13
75	TMEM45A Is Dispensable for Epidermal Morphogenesis, Keratinization and Barrier Formation. <i>PLoS ONE</i> , 2016 , 11, e0147069	3.7	6

74	Modelling atopic dermatitis during the morphogenetic process involved in reconstruction of a human epidermis. <i>Current Research in Translational Medicine</i> , 2016 , 64, 179-183	3.7	5
73	The activation of cultured keratinocytes by cholesterol depletion during reconstruction of a human epidermis is reminiscent of monolayer cultures. <i>Archives of Dermatological Research</i> , 2015 , 307, 309-18	3.3	7
72	Microscopy: access to the bigger picture in histology. <i>Nature</i> , 2015 , 519, 291	50.4	
71	Hyaluronidase-1 Is Mainly Functional in the Upper Granular Layer, Close to the Epidermal Barrier. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 3189-3192	4.3	6
70	Epidermal reference genes at the forefront of data interpretation. <i>Experimental Dermatology</i> , 2015 , 24, 738-9	4	
69	Preparation and characterizations of EGDE crosslinked chitosan electrospun membranes. <i>Clinical Hemorheology and Microcirculation</i> , 2015 , 60, 39-50	2.5	11
68	Hyaluronan metabolism in human keratinocytes and atopic dermatitis skin is driven by a balance of hyaluronan synthases 1 and 3. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 2174-2182	4.3	40
67	Knockdown of filaggrin in a three-dimensional reconstructed human epidermis impairs keratinocyte differentiation. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 2938-2946	4.3	80
66	Polysaccharide-coated PCL nanofibers for wound dressing applications. <i>Advanced Healthcare Materials</i> , 2014 , 3, 2032-9	10.1	69
65	High TMEM45A expression is correlated to epidermal keratinization. <i>Experimental Dermatology</i> , 2014 , 23, 339-44	4	23
64	Meaning of relative gene expression in multilayered cultures of epidermal keratinocytes. <i>Experimental Dermatology</i> , 2014 , 23, 754-6	4	3
63	Reconstruction of normal and pathological human epidermis on polycarbonate filter. <i>Methods in Molecular Biology</i> , 2014 , 1195, 191-201	1.4	24
62	In vitro reconstruction of epidermis from primary Darier's disease keratinocytes replicates the histopathological phenotype. <i>Journal of Dermatological Science</i> , 2013 , 71, 138-40	4.3	4
61	Ophiobolin A, a sesterterpenoid fungal phytotoxin, displays higher in vitro growth-inhibitory effects in mammalian than in plant cells and displays in vivo antitumor activity. <i>International Journal of Oncology</i> , 2013 , 43, 575-85	4.4	29
60	TMEM45A is essential for hypoxia-induced chemoresistance in breast and liver cancer cells. <i>BMC Cancer</i> , 2012 , 12, 391	4.8	60
59	Epidermal morphogenesis during progressive in vitro 3D reconstruction at the air-liquid interface. <i>Experimental Dermatology</i> , 2012 , 21, 871-5	4	55
58	Studies of cell signaling in a reconstructed human epidermis exposed to sensitizers: IL-8 synthesis and release depend on EGFR activation. <i>Archives of Dermatological Research</i> , 2012 , 304, 289-303	3.3	14
57	HB-EGF, the growth factor that accelerates keratinocyte migration, but slows proliferation. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 2129-30	4.3	10

56	Development of a chitosan nanofibrillar scaffold for skin repair and regeneration. <i>Biomacromolecules</i> , 2011 , 12, 3194-204	6.9	156
55	Transcriptional profiling after lipid raft disruption in keratinocytes identifies critical mediators of atopic dermatitis pathways. <i>Journal of Investigative Dermatology</i> , 2011 , 131, 46-58	4.3	40
54	In vitro anticancer activity, toxicity and structure-activity relationships of phyllostictine A, a natural oxazatricycloalkenone produced by the fungus <i>Phyllosticta cirsii</i> . <i>Toxicology and Applied Pharmacology</i> , 2011 , 254, 8-17	4.6	13
53	HB-EGF synthesis and release induced by cholesterol depletion of human epidermal keratinocytes is controlled by extracellular ATP and involves both p38 and ERK1/2 signaling pathways. <i>Journal of Cellular Physiology</i> , 2011 , 226, 1651-9	7	17
52	Lipid rafts and the oxidative stress hypothesis. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 1457-9	4.3	5
51	Lack of cadherins <i>Celsr2</i> and <i>Celsr3</i> impairs ependymal ciliogenesis, leading to fatal hydrocephalus. <i>Nature Neuroscience</i> , 2010 , 13, 700-7	25.5	244
50	Inhibition of Akt signaling by exclusion from lipid rafts in normal and transformed epidermal keratinocytes. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 1136-45	4.3	59
49	Study of epidermal differentiation in human keratinocytes cultured in autocrine conditions. <i>Methods in Molecular Biology</i> , 2010 , 585, 71-82	1.4	22
48	p38 MAPK-regulated EGFR internalization takes place in keratinocyte monolayer during stress conditions. <i>Archives of Dermatological Research</i> , 2010 , 302, 229-33	3.3	11
47	Repeated exposures to UVB induce differentiation rather than senescence of human keratinocytes lacking p16(INK-4A). <i>Biogerontology</i> , 2010 , 11, 167-81	4.5	22
46	Proteomic profiling of human keratinocytes undergoing UVB-induced alternative differentiation reveals TRIPartite Motif Protein 29 as a survival factor. <i>PLoS ONE</i> , 2010 , 5, e10462	3.7	24
45	The CYP26 inhibitor R115866 potentiates the effects of all-trans retinoic acid on cultured human epidermal keratinocytes. <i>British Journal of Dermatology</i> , 2009 , 160, 505-13	4	28
44	Candidate housekeeping genes require evaluation before their selection for studies of human epidermal keratinocytes. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 770-3	4.3	44
43	Development of a procedure to simultaneously isolate RNA, DNA, and proteins for [corrected] characterizing cells invading or cultured on chitosan scaffolds. <i>Analytical Biochemistry</i> , 2009 , 393, 145-7	3.1	8
42	Cholesterol and Lipid Rafts as Regulators of Signaling Through the EGF Receptor in Keratinocytes. <i>Open Dermatology Journal</i> , 2009 , 3, 151-158	1.1	2
41	The Reconstructed Human Epidermis Models in Fundamental Research 2009 , 967-976		3
40	Heparin-binding EGF-like growth factor is induced by disruption of lipid rafts and oxidative stress in keratinocytes and participates in the epidermal response to cutaneous wounds. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 717-27	4.3	36
39	Knockdown of PKD1 in normal human epidermal keratinocytes increases mRNA expression of keratin 10 and involucrin: early markers of keratinocyte differentiation. <i>Archives of Dermatological Research</i> , 2008 , 300, 139-45	3.3	18

38	The inhibition of the expression of the small Rho GTPase Rac1 induces differentiation with no effect on cell proliferation in growing human adult keratinocytes. <i>Journal of Cellular Biochemistry</i> , 2008 , 103, 857-64	4-7	11
37	Internalization of EGF receptor following lipid rafts disruption in keratinocytes is delayed and dependent on p38 MAPK activation. <i>Journal of Cellular Physiology</i> , 2008 , 217, 834-45	7	34
36	The dumb ErbB receptor helps healing. <i>Journal of Investigative Dermatology</i> , 2007 , 127, 995-7	4-3	1
35	Functional redundancy of extracellular matrix protein 1 in epidermal differentiation. <i>British Journal of Dermatology</i> , 2007 , 157, 771-5	4	13
34	The small Rho GTPase Rac1 controls normal human dermal fibroblasts proliferation with phosphorylation of the oncoprotein c-myc. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 359, 834-9	3-4	7
33	Modelling the human epidermis in vitro: tools for basic and applied research. <i>Archives of Dermatological Research</i> , 2007 , 298, 361-9	3-3	83
32	Immunogold silver staining associated with epi-fluorescence for cucumber mosaic virus localisation on semi-thin sections of banana tissues. <i>European Journal of Histochemistry</i> , 2007 , 51, 153-8	2-1	8
31	Effects of the CDK-inhibitor CYC202 on p38 MAPK, ERK1/2 and c-Myc activities in papillomavirus type 16 E6- and E7-transformed human keratinocytes. <i>Oncology Reports</i> , 2007 , 18, 999-1005	3-5	4
30	Expression of lysosome-associated membrane protein 1 (Lamp-1) and galectins in human keratinocytes is regulated by differentiation. <i>Archives of Dermatological Research</i> , 2006 , 298, 73-81	3-3	21
29	Effects of the cyclin-dependent kinase inhibitor CYC202 (R-roscovitine) on the physiology of cultured human keratinocytes. <i>Biochemical Pharmacology</i> , 2005 , 70, 824-36	6	18
28	Ultrastructural morphology of the male and female genital tracts of Psoroptes spp. (Acari: Astigmata: Psoroptidae). <i>Experimental and Applied Acarology</i> , 2005 , 36, 305-24	2-1	7
27	Cholesterol depletion upregulates involucrin expression in epidermal keratinocytes through activation of p38. <i>Journal of Investigative Dermatology</i> , 2004 , 123, 564-73	4-3	59
26	Calcium entry into keratinocytes induces exocytosis of lysosomes. <i>Archives of Dermatological Research</i> , 2004 , 296, 30-41	3-3	18
25	A simple reconstructed human epidermis: preparation of the culture model and utilization in in vitro studies. <i>Archives of Dermatological Research</i> , 2004 , 296, 203-11	3-3	130
24	Ultrastructural changes associated with cryopreservation of banana (Musa spp.) highly proliferating meristems. <i>Plant Cell Reports</i> , 2003 , 21, 690-8	5-1	63
23	Analysis of interleukin-1alpha (IL-1alpha) and interleukin-8 (IL-8) expression and release in in vitro reconstructed human epidermis for the prediction of in vivo skin irritation and/or sensitization. <i>Toxicology in Vitro</i> , 2003 , 17, 311-21	3-6	132
22	Cryopreservation for the elimination of cucumber mosaic and banana streak viruses from banana (Musa spp.). <i>Plant Cell Reports</i> , 2002 , 20, 1117-1122	5-1	109
21	The ornithine decarboxylase inhibitor, difluoromethylornithine, inhibits casein kinase II activity, c-Myc expression and normal human keratinocyte proliferation. <i>Archives of Dermatological Research</i> , 2002 , 293, 590-3	3-3	5

20	Ca(2+)/calmodulin-dependent protein kinase (CaM-kinase) inhibitor KN-62 suppresses the activity of mitogen-activated protein kinase (MAPK), c-myc activation and human keratinocyte proliferation. <i>Archives of Dermatological Research</i> , 2002 , 294, 198-202	3.3	11
19	Dual role of protein kinase C on mitogen-activated protein kinase activation and human keratinocyte proliferation. <i>Experimental Dermatology</i> , 2002 , 11, 344-8	4	11
18	Human EGF receptor (HER) family and heregulin members are differentially expressed in epidermal keratinocytes and modulate differentiation. <i>Experimental Cell Research</i> , 2001 , 271, 315-28	4.2	51
17	Differentiation-dependent alternative splicing and expression of the extracellular matrix protein 1 gene in human keratinocytes. <i>Journal of Investigative Dermatology</i> , 2000 , 114, 718-24	4.3	55
16	The tetraspanin CD9 associates with the integrin alpha6beta4 in cultured human epidermal keratinocytes and is involved in cell motility. <i>European Journal of Cell Biology</i> , 2000 , 79, 41-51	6.1	55
15	Differential expression and release of cytokines by an in vitro reconstructed human epidermis following exposure to skin irritant and sensitizing chemicals. <i>Toxicology in Vitro</i> , 1999 , 13, 867-77	3.6	52
14	High-cell-density phorbol ester and retinoic acid upregulate involucrin and downregulate suprabasal keratin 10 in autocrine cultures of human epidermal keratinocytes. <i>Molecular Cell Biology Research Communications: MCBRC: Part B of Biochemical and Biophysical Research Communications</i> , 1999 , 2, 138-44		40
13	Human epidermal keratinocytes upregulate expression of the prolactin receptor after the onset of terminal differentiation, but do not respond to prolactin. <i>Archives of Biochemistry and Biophysics</i> , 1999 , 364, 247-53	4.1	21
12	Science and the Web. <i>Science</i> , 1998 , 280, 1171d-1171	33.3	2
11	Cell density and culture factors regulate keratinocyte commitment to differentiation and expression of suprabasal K1/K10 keratins. <i>Journal of Investigative Dermatology</i> , 1995 , 104, 271-6	4.3	199
10	Basal detachment of the epidermis using dispase: tissue spatial organization and fate of integrin alpha 6 beta 4 and hemidesmosomes. <i>Journal of Investigative Dermatology</i> , 1994 , 102, 111-7	4.3	36
9	Basal cell adhesion to a culture substratum controls the polarized spatial organization of human epidermal keratinocytes into proliferating basal and terminally differentiating suprabasal populations. <i>Epithelial Cell Biology</i> , 1993 , 2, 7-16		2
8	Specific internalization of basal membrane domains containing the integrin alpha 6 beta 4 in dispase-detached cultured human keratinocytes. <i>European Journal of Cell Biology</i> , 1993 , 60, 12-20	6.1	21
7	Basal Cell Adhesion Controls the Polarized Arrangement of Proliferating and Differentiating Keratinocytes in Culture. <i>Journal of Tissue Viability</i> , 1992 , 2, 24-25	3.2	
6	Processing and characterization of the low density lipoprotein receptor in the human colonic carcinoma cell subclone HT29-18: a potential pathway for delivering therapeutic drugs and genes. <i>Bioscience Reports</i> , 1992 , 12, 483-94	4.1	6
5	Influence of a short oxidative stress on the LDL endocytosis by human endothelial cells: an ultrastructural study. <i>Journal of Submicroscopic Cytology and Pathology</i> , 1992 , 24, 61-73		2
4	Utilization of human cultured epidermal keratinocytes: irreversibility of the inhibition of proliferation induced in stored detached cultures. <i>Burns</i> , 1991 , 17, 205-8	2.3	5
3	Endocytosis of low density lipoproteins in human endothelial cells: typical morphological aspects of the high affinity receptor-mediated pathway as revealed by serial sections and acid phosphatase cytochemistry. <i>Journal of Submicroscopic Cytology and Pathology</i> , 1989 , 21, 627-39		

2	Incubation of endothelial cells in a superoxide-generating system: impaired low-density lipoprotein receptor-mediated endocytosis. <i>Journal of Cellular Physiology</i> , 1988 , 136, 289-96	7	13
1	Rapid preparative isolation of concentrated low density lipoproteins and of lipoprotein-deficient serum using vertical rotor gradient ultracentrifugation. <i>Journal of Lipid Research</i> , 1985 , 26, 1476-80	6.3	48