

Aila J Ahola

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/95972/publications.pdf>

Version: 2024-02-01

30
papers

836
citations

567281

15
h-index

501196

28
g-index

31
all docs

31
docs citations

31
times ranked

1445
citing authors

#	ARTICLE	IF	CITATIONS
1	Weight Loss Trajectories in Healthy Weight Coaching: Cohort Study. <i>JMIR Formative Research</i> , 2022, 6, e26374.	1.4	4
2	Symptoms of depression are associated with reduced leisure-time physical activity in adult individuals with type 1 diabetes. <i>Acta Diabetologica</i> , 2021, 58, 1373-1380.	2.5	6
3	Computational modelling of self-reported dietary carbohydrate intake on glucose concentrations in patients undergoing Roux-en-Y gastric bypass versus one-anastomosis gastric bypass. <i>Annals of Medicine</i> , 2021, 53, 1885-1895.	3.8	5
4	Association between symptoms of depression, diabetes complications and vascular risk factors in four European cohorts of individuals with type 1 diabetes – InterDiane Consortium. <i>Diabetes Research and Clinical Practice</i> , 2020, 170, 108495.	2.8	10
5	Perceived Stress and Adherence to the Dietary Recommendations and Blood Glucose Levels in Type 1 Diabetes. <i>Journal of Diabetes Research</i> , 2020, 2020, 1-8.	2.3	7
6	Waist-height ratio and waist are the best estimators of visceral fat in type 1 diabetes. <i>Scientific Reports</i> , 2020, 10, 18575.	3.3	19
7	Effectiveness of a web-based real-life weight management program: Study design, methods, and participants' baseline characteristics. <i>Contemporary Clinical Trials Communications</i> , 2020, 19, 100638.	1.1	6
8	Dietary carbohydrate intake and cardio-metabolic risk factors in type 1 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2019, 155, 107818.	2.8	21
9	Dietary intake in type 1 diabetes at different stages of diabetic kidney disease. <i>Diabetes Research and Clinical Practice</i> , 2019, 155, 107775.	2.8	4
10	Associations of dietary macronutrient and fibre intake with glycaemia in individuals with Type 1 diabetes. <i>Diabetic Medicine</i> , 2019, 36, 1391-1398.	2.3	11
11	Association between depressive symptoms and dietary intake in patients with type 1 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 139, 91-99.	2.8	10
12	Association between habitual coffee consumption and metabolic syndrome in type 1 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018, 28, 470-476.	2.6	21
13	Regression of albuminuria and its association with incident cardiovascular outcomes and mortality in type 1 diabetes: the FinnDiane Study. <i>Diabetologia</i> , 2018, 61, 1203-1211.	6.3	29
14	Association between diet and measures of arterial stiffness in type 1 diabetes – Focus on dietary patterns and macronutrient substitutions. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018, 28, 1166-1172.	2.6	16
15	Adherence to special diets and its association with meeting the nutrient recommendations in individuals with type 1 diabetes. <i>Acta Diabetologica</i> , 2018, 55, 843-851.	2.5	17
16	The serum uric acid concentration is not causally linked to diabetic nephropathy in type 1 diabetes. <i>Kidney International</i> , 2017, 91, 1178-1185.	5.2	40
17	Association between adherence to dietary recommendations and high-sensitivity C-reactive protein level in type 1 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2017, 126, 122-128.	2.8	9
18	Dietary patterns reflecting healthy food choices are associated with lower serum LPS activity. <i>Scientific Reports</i> , 2017, 7, 6511.	3.3	58

#	ARTICLE	IF	CITATIONS
19	Dietary patterns are associated with various vascular health markers and complications in type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1144-1150.	2.3	24
20	Fear of hypoglycaemia and self-management in type 1 diabetes. <i>Journal of Clinical and Translational Endocrinology</i> , 2016, 4, 13-18.	1.4	26
21	Renin-angiotensin-aldosterone blockade is associated with decreased use of antidepressant therapy in patients with type 1 diabetes and diabetic nephropathy. <i>Acta Diabetologica</i> , 2014, 51, 529-533.	2.5	11
22	Patients with type 1 diabetes show signs of vascular dysfunction in response to multiple high-fat meals. <i>Nutrition and Metabolism</i> , 2014, 11, 28.	3.0	17
23	Barriers to self-management of diabetes. <i>Diabetic Medicine</i> , 2013, 30, 413-420.	2.3	206
24	Sense of coherence, food selection and leisure time physical activity in type 1 diabetes. <i>Scandinavian Journal of Public Health</i> , 2012, 40, 621-628.	2.3	38
25	Energy and nutrient intakes and adherence to dietary guidelines among Finnish adults with type 1 diabetes. <i>Annals of Medicine</i> , 2012, 44, 73-81.	3.8	43
26	Purchase of antidepressant agents by patients with type 1 diabetes is associated with increased mortality rates in women but not in men. <i>Diabetologia</i> , 2012, 55, 73-79.	6.3	16
27	Many patients with Type 1 diabetes estimate their prandial insulin need inappropriately. <i>Journal of Diabetes</i> , 2010, 2, 194-202.	1.8	55
28	Depression is associated with the metabolic syndrome among patients with type 1 diabetes. <i>Annals of Medicine</i> , 2010, 42, 495-501.	3.8	29
29	Health-related quality of life in patients with type 1 diabetes--association with diabetic complications (the FinnDiane Study). <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 1903-1908.	0.7	56
30	The cross-sectional associations between sense of coherence and diabetic microvascular complications, glycaemic control, and patients' conceptions of type 1 diabetes. <i>Health and Quality of Life Outcomes</i> , 2010, 8, 142.	2.4	22