

# Arumugam Nagalingam

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9598893/publications.pdf>

Version: 2024-02-01

9  
papers

545  
citations

1170033  
9  
h-index

1637695  
9  
g-index

9  
all docs

9  
docs citations

9  
times ranked

1019  
citing authors

#	ARTICLE	IF	CITATIONS
1	A Procarcinogenic Colon Microbe Promotes Breast Tumorigenesis and Metastatic Progression and Concomitantly Activates Notch and $\beta$ -Catenin Axes. <i>Cancer Discovery</i> , 2021, 11, 1138-1157.	7.7	88
2	Hyperleptinemia in obese state renders luminal breast cancers refractory to tamoxifen by coordinating a crosstalk between Med1, miR205 and ErbB. <i>Npj Breast Cancer</i> , 2021, 7, 105.	2.3	12
3	Honokiol abrogates leptin-induced tumor progression by inhibiting Wnt1-MTA1- $\beta$ -catenin signaling axis in a microRNA-34a dependent manner. <i>Oncotarget</i> , 2015, 6, 16396-16410.	0.8	50
4	Honokiol activates LKB1-miR-34a axis and antagonizes the oncogenic actions of leptin in breast cancer. <i>Oncotarget</i> , 2015, 6, 29947-29962.	0.8	49
5	Honokiol inhibits epithelial-mesenchymal transition in breast cancer cells by targeting signal transducer and activator of transcription 3/Zeb1/E-cadherin axis. <i>Molecular Oncology</i> , 2014, 8, 565-580.	2.1	85
6	Mechanistic Elucidation of the Antitumor Properties of Withaferin A in Breast Cancer. <i>Cancer Research</i> , 2014, 74, 2617-2629.	0.4	77
7	Integral Role of PTP1B in Adiponectin-Mediated Inhibition of Oncogenic Actions of Leptin in Breast Carcinogenesis. <i>Neoplasia</i> , 2013, 15, 23-IN11.	2.3	55
8	Med1 plays a critical role in the development of tamoxifen resistance. <i>Carcinogenesis</i> , 2012, 33, 918-930.	1.3	54
9	Survivin upregulation, dependent on leptin-EGFR-Notch1 axis, is essential for leptin-induced migration of breast carcinoma cells. <i>Endocrine-Related Cancer</i> , 2011, 18, 413-428.	1.6	75