

# Toyohiro Naito

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9602054/publications.pdf>

Version: 2024-02-01

37  
papers

454  
citations

687363

13  
h-index

752698

20  
g-index

37  
all docs

37  
docs citations

37  
times ranked

592  
citing authors

#	ARTICLE	IF	CITATIONS
1	Development of a microfluidic dispensing device for multivariate data acquisition and application in molecularly imprinting hydrogel preparation. <i>Journal of Materials Chemistry B</i> , 2022, 10, 6664-6672.	5.8	1
2	Poly(ethylene glycol) Hydrogels with a Boronic Acid Monomer via Molecular Imprinting for Selective Removal of Quinic Acid Gamma-Lactone in Coffee. <i>ACS Applied Polymer Materials</i> , 2021, 3, 226-232.	4.4	6
3	Fluorescent detection of target proteins via a molecularly imprinted hydrogel. <i>Analytical Methods</i> , 2021, 13, 3086-3091.	2.7	4
4	Simple chemical detection based on a surface-modified electroosmotic pump <i>via</i> interval immobilization. <i>Analytical Methods</i> , 2021, 13, 1559-1564.	2.7	0
5	Development of Microfluidic Techniques for Onsite Analysis. <i>Bunseki Kagaku</i> , 2021, 70, 341-349.	0.2	1
6	Label-Free Cancer Stem-like Cell Assay Conducted at a Single Cell Level Using Microfluidic Mechanotyping Devices. <i>Analytical Chemistry</i> , 2021, 93, 14409-14416.	6.5	10
7	Separation of halogenated benzenes enabled by investigation of halogen-π interactions with carbon materials. <i>Chemical Science</i> , 2020, 11, 409-418.	7.4	17
8	Separation of saccharides using fullerene-bonded silica monolithic columns via π interactions in liquid chromatography. <i>Scientific Reports</i> , 2020, 10, 13850.	3.3	8
9	Development of Microfluidic Components for Micro Total Analysis Systems. <i>Chromatography</i> , 2020, 41, 31-37.	1.7	6
10	Tunable Liquid Chromatographic Separation of H/D Isotopologues Enabled by Aromatic π Interactions. <i>Analytical Chemistry</i> , 2020, 92, 4065-4072.	6.5	10
11	Controllable Molecular Sieving by <i>copoly</i> (Poly(ethylene glycol) Acrylate/Poly(ethylene glycol)) Tj ETQq1 1 0.784314 rgBT /Over Materials, 2020, 2, 3886-3893.	4.4	6
12	Analysis and Survey of PM<sub>2.5</sub> from a Biological Viewpoint at Kyushu University Ito Campus. <i>Bunseki Kagaku</i> , 2020, 69, 741-746.	0.2	1
13	Engineering Nanowire-Mediated Cell Lysis for Microbial Cell Identification. <i>ACS Nano</i> , 2019, 13, 2262-2273.	14.6	17
14	Differentiating π Interactions by Constructing Concave/Convex Surfaces Using a Bucky Bowl Molecule, Corannulene in Liquid Chromatography. <i>Analytical Chemistry</i> , 2019, 91, 2439-2446.	6.5	17
15	Efficient extraction of estrogen receptor-activated compounds from environmental surface water via a receptor-mimic adsorbent, a hydrophilic PEG-based molecularly imprinted polymer. <i>Chemosphere</i> , 2019, 217, 204-212.	8.2	19
16	Magnetic Field Stimuli-Sensitive Drug Release Using a Magnetic Thermal Seed Coated with Thermal-Responsive Molecularly Imprinted Polymer. <i>ACS Biomaterials Science and Engineering</i> , 2019, 5, 759-767.	5.2	33
17	Detection of Molecular Adsorbate in Aqueous Solution Based on Electroosmosis. <i>Sensors and Materials</i> , 2019, 31, 45.	0.5	3
18	Suppression of Hydrophobicity and Optimizations of a Ligand-Immobilization for Effective Affinity Chromatography Using a Spongy Monolith. <i>Chromatography</i> , 2018, 39, 113-118.	1.7	4

#	ARTICLE	IF	CITATIONS
19	Quantitative Evaluation of Dielectric Breakdown of Silicon Micro- and Nanofluidic Devices for Electrophoretic Transport of a Single DNA Molecule. <i>Micromachines</i> , 2018, 9, 180.	2.9	3
20	Isotope Effects on Hydrogen Bonding and CH/CD <sup>13</sup> C Interaction. <i>Journal of Physical Chemistry C</i> , 2018, 122, 15026-15032.	3.1	18
21	Selective adsorption of carbohydrates and glycoproteins via molecularly imprinted hydrogels: application to visible detection by a boronic acid monomer. <i>Chemical Communications</i> , 2017, 53, 7290-7293.	4.1	16
22	New platform for simple and rapid protein-based affinity reactions. <i>Scientific Reports</i> , 2017, 7, 178.	3.3	18
23	Identification and characterization of a thermally cleaved fragment of monoclonal antibody-A detected by sodium dodecyl sulfate-capillary gel electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 140, 98-104.	2.8	20
24	Tunable separations based on a molecular size effect for biomolecules by poly(ethylene glycol) gel-based capillary electrophoresis. <i>Journal of Chromatography A</i> , 2017, 1523, 107-113.	3.7	13
25	Competitive ELISA-like Label-free Detection of Lysozyme by Using a Fluorescent Monomer-doped Molecularly Imprinted Hydrogel. <i>Analytical Sciences</i> , 2017, 33, 1311-1315.	1.6	7
26	Development of a C <sub>70</sub> -Fullerene Bonded Silica-Monolithic Capillary and Its Retention Characteristics in Liquid Chromatography. <i>Chromatography</i> , 2017, 38, 45-51.	1.7	12
27	Validation of Capillary Zone Electrophoretic Method for Evaluating Monoclonal Antibodies and Antibody-Drug Conjugates. <i>Chromatography</i> , 2016, 37, 117-124.	1.7	11
28	Three-Dimensional Fabrication for Microfluidics by Conventional Techniques and Equipment Used in Mass Production. <i>Micromachines</i> , 2016, 7, 82.	2.9	11
29	Specific Intermolecular Interactions by the Localized $\pi$ -Electrons in C <sub>70</sub> -fullerene. <i>ChemistrySelect</i> , 2016, 1, 5900-5904.	1.5	11
30	Simple Preparation and Characterization of Viscoelastic Gels Induced by Multiple Intermolecular Interactions Using Low-Molecular-Weight Species. <i>Bulletin of the Chemical Society of Japan</i> , 2015, 88, 1575-1580.	3.2	0
31	Unique Separation Behavior of a C <sub>60</sub> -Fullerene-Bonded Silica Monolith Prepared by an Effective Thermal Coupling Agent. <i>Chemistry - A European Journal</i> , 2015, 21, 18095-18098.	3.3	18
32	C <sub>60</sub> -Fullerene Bonded Silica Monolithic Capillary for Specific Separations of Aromatic Compounds. <i>Chromatography</i> , 2015, 36, 105-113.	1.7	12
33	Molecularly Imprinted Polymers for Selective Adsorption of Lysozyme and Cytochrome <i>c</i> Using a PEG-Based Hydrogel: Selective Recognition for Different Conformations Due to pH Conditions. <i>Macromolecules</i> , 2015, 48, 4081-4087.	4.8	49
34	Simple and Effective Label-Free Capillary Electrophoretic Analysis of Sugars by Complexation Using Quinoline Boronic Acids. <i>Analytical Chemistry</i> , 2015, 87, 5068-5073.	6.5	7
35	Tunable Molecular Sieving in Gel Electrophoresis Using a Poly(ethylene glycol)-Based Hydrogel. <i>Chromatography</i> , 2014, 35, 81-86.	1.7	5
36	Effective determination of a pharmaceutical, sulphiride, in river water by online SPE-LC-MS using a molecularly imprinted polymer as a pre-concentration medium. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 89, 111-117.	2.8	33

#	ARTICLE	IF	CITATIONS
37	Development of a C60-fullerene bonded open-tubular capillary using a photo/thermal active agent for liquid chromatographic separations by $\pi$ - $\pi$ interactions. Journal of Chromatography A, 2014, 1323, 174-178.	3.7	27