

Zhaojin Yu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9615806/publications.pdf>

Version: 2024-02-01

33
papers

780
citations

623734

14
h-index

526287

27
g-index

37
all docs

37
docs citations

37
times ranked

1279
citing authors

#	ARTICLE	IF	CITATIONS
1	Advances of research of Fc-fusion protein that activate NK cells for tumor immunotherapy. <i>International Immunopharmacology</i> , 2022, 109, 108783.	3.8	8
2	Exosome loaded genipin crosslinked hydrogel facilitates full thickness cutaneous wound healing in rat animal model. <i>Drug Delivery</i> , 2021, 28, 884-893.	5.7	37
3	Peptide-based therapeutic cancer vaccine: Current trends in clinical application. <i>Cell Proliferation</i> , 2021, 54, e13025.	5.3	68
4	lncRNA-Xist/miR-101-3p/KLF6/C/EBP β axis promotes TAM polarization to regulate cancer cell proliferation and migration. <i>Molecular Therapy - Nucleic Acids</i> , 2021, 23, 536-551.	5.1	80
5	HLA-A2.1-restricted ECM1-derived epitope LA through DC cross-activation priming CD8+ T and NK cells: a novel therapeutic tumour vaccine. <i>Journal of Hematology and Oncology</i> , 2021, 14, 71.	17.0	11
6	A Study of the Identification, Fragmentation Mode and Metabolic Pathways of Imatinib in Rats Using UHPLC-Q-TOF-MS/MS. <i>Journal of Analytical Methods in Chemistry</i> , 2021, 2021, 1-15.	1.6	0
7	Integrated microenvironment-associated genomic profiles identify LRRC15 mediating recurrent glioblastoma-associated macrophages infiltration. <i>Journal of Cellular and Molecular Medicine</i> , 2021, 25, 5534-5546.	3.6	7
8	Down-Regulation of an Autophagy-Related Gene SERPINA1 as a Superior Prognosis Biomarker Associates with Relapse and Distant Metastasis in Colon Adenocarcinoma. <i>OncoTargets and Therapy</i> , 2021, Volume 14, 3861-3872.	2.0	5
9	Mannose-modified liposome designed for epitope peptide drug delivery in cancer immunotherapy. <i>International Immunopharmacology</i> , 2021, 101, 108148.	3.8	10
10	Recent progress on MHC-I epitope prediction in tumor immunotherapy. <i>American Journal of Cancer Research</i> , 2021, 11, 2401-2416.	1.4	0
11	Hypoxia-related prognostic model in bladder urothelial reflects immune cell infiltration. <i>American Journal of Cancer Research</i> , 2021, 11, 5076-5093.	1.4	0
12	Bioinformatics Analysis Suggests the Combined Expression of AURKB and KIF18B Being an Important Event in the Development of Clear Cell Renal Cell Carcinoma. <i>Pathology and Oncology Research</i> , 2020, 26, 1583-1594.	1.9	8
13	TEX19 promotes ovarian carcinoma progression and is a potential target for epitope vaccine immunotherapy. <i>Life Sciences</i> , 2020, 241, 117171.	4.3	7
14	Silencing KIF18B enhances radiosensitivity: identification of a promising therapeutic target in sarcoma. <i>EBioMedicine</i> , 2020, 61, 103056.	6.1	7
15	Anterior gradient 2-derived peptide upregulates major histocompatibility complex class II-related chains A/B in hepatocellular carcinoma cells. <i>Life Sciences</i> , 2020, 246, 117396.	4.3	5
16	Designing high affinity target-binding peptides to HLA-E: a key membrane antigen of multiple myeloma. <i>Aging</i> , 2020, 12, 20457-20470.	3.1	5
17	Prognostic value of CXCL17 and CXCR8 expression in patients with colon cancer. <i>Oncology Letters</i> , 2020, 20, 2711-2720.	1.8	8
18	SNORD89 promotes stemness phenotype of ovarian cancer cells by regulating Notch1-c-Myc pathway. <i>Journal of Translational Medicine</i> , 2019, 17, 259.	4.4	43

#	ARTICLE	IF	CITATIONS
19	The Analysis of Key Factors Related to ADCs Structural Design. <i>Frontiers in Pharmacology</i> , 2019, 10, 373.	3.5	45
20	Prognostic relevance of miR-137 and its liver microenvironment regulatory target gene AFM in hepatocellular carcinoma. <i>Journal of Cellular Physiology</i> , 2019, 234, 11888-11899.	4.1	11
21	Identification of cancer/testis antigen 1/2 gene as a potential hepatocellular carcinoma therapeutic target by hub gene screening with topological analysis. <i>Oncology Letters</i> , 2019, 18, 4778-4788.	1.8	6
22	Moesin is an independent prognostic marker for ER ⁺ positive breast cancer. <i>Oncology Letters</i> , 2018, 17, 1921-1933.	1.8	12
23	Poly(ADP-ribose) polymerase-3 overexpression is associated with poor prognosis in patients with breast cancer following chemotherapy. <i>Oncology Letters</i> , 2018, 16, 5621-5630.	1.8	7
24	The Hedgehog signaling pathway is associated with poor prognosis in breast cancer patients with the CD44 ⁺ /CD24 ^{low} phenotype. <i>Molecular Medicine Reports</i> , 2016, 14, 5261-5270.	2.4	13
25	MiR-487a Promotes TGF- β 1-induced EMT, the Migration and Invasion of Breast Cancer Cells by Directly Targeting MAGI2. <i>International Journal of Biological Sciences</i> , 2016, 12, 397-408.	6.4	51
26	HDAC2 overexpression is a poor prognostic factor of breast cancer patients with increased multidrug resistance-associated protein expression who received anthracyclines therapy. <i>Japanese Journal of Clinical Oncology</i> , 2016, 46, 893-902.	1.3	35
27	MicroRNA-100 suppresses the migration and invasion of breast cancer cells by targeting FZD-8 and inhibiting Wnt/ β -catenin signaling pathway. <i>Tumor Biology</i> , 2016, 37, 5001-5011.	1.8	74
28	The Hedgehog signalling pathway mediates drug response of MCF-7 mammosphere cells in breast cancer patients. <i>Clinical Science</i> , 2015, 129, 809-822.	4.3	46
29	Ano1/TMEM16A Overexpression Is Associated with Good Prognosis in PR-Positive or HER2-Negative Breast Cancer Patients following Tamoxifen Treatment. <i>PLoS ONE</i> , 2015, 10, e0126128.	2.5	39
30	Combined expression of ezrin and E-cadherin is associated with lymph node metastasis and poor prognosis in breast cancer. <i>Oncology Reports</i> , 2015, 34, 165-174.	2.6	20
31	Combined expression of aldehyde dehydrogenase 1A1 and β -catenin is associated with lymph node metastasis and poor survival in breast cancer patients following cyclophosphamide treatment. <i>Oncology Reports</i> , 2015, 34, 3163-3173.	2.6	5
32	BRCA1 promoter hypermethylation in sporadic epithelial ovarian carcinoma: Association with low expression of BRCA1, improved survival and co-expression of DNA methyltransferases. <i>Oncology Letters</i> , 2014, 7, 1088-1096.	1.8	21
33	Expression of BAMBI and its combination with Smad7 correlates with tumor invasion and poor prognosis in gastric cancer. <i>Tumor Biology</i> , 2014, 35, 7047-7056.	1.8	23