

Thomas Andlid

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9653378/publications.pdf>

Version: 2024-02-01

44
papers

1,619
citations

236925

25
h-index

302126

39
g-index

44
all docs

44
docs citations

44
times ranked

1886
citing authors

#	ARTICLE	IF	CITATIONS
1	Extracts of Digested Berries Increase the Survival of <i>Saccharomyces cerevisiae</i> during H ₂ O ₂ Induced Oxidative Stress. <i>Molecules</i> , 2021, 26, 1057.	3.8	4
2	Isolation, identification, and selection of strains as candidate probiotics and starters for fermentation of Swedish legumes. <i>Food and Nutrition Research</i> , 2020, 64, .	2.6	8
3	Effects of Pulsed Electric Field-Assisted Osmotic Dehydration and Edible Coating on the Recovery of Anthocyanins from In Vitro Digested Berries. <i>Foods</i> , 2019, 8, 505.	4.3	11
4	Tailoring bilberry powder functionality through processing: Effects of drying and fractionation on the stability of total polyphenols and anthocyanins. <i>Food Science and Nutrition</i> , 2019, 7, 1017-1026.	3.4	8
5	Appetite and Subsequent Food Intake Were Unaffected by the Amount of Sourdough and Rye in Soft Breadâ€”A Randomized Cross-Over Breakfast Study. <i>Nutrients</i> , 2018, 10, 1594.	4.1	5
6	Supercritical Fluid Extraction of Berry Seeds: Chemical Composition and Antioxidant Activity. <i>Journal of Food Quality</i> , 2018, 2018, 1-10.	2.6	25
7	<i>Bifidobacterium callitrichidarum</i> sp. nov. from the faeces of the emperor tamarin (<i>Saguinus</i>) Tj ETQq1 1 0.784314 r _g BT /Overlock 10 T _g BT	2.7	25
8	Strain improvement of <i>Pichia kudriavzevii</i> TY13 for raised phytase production and reduced phosphate repression. <i>Microbial Biotechnology</i> , 2017, 10, 341-353.	4.2	17
9	Isolation, Identification and Characterization of Yeasts from Fermented Goat Milk of the Yaghnob Valley in Tajikistan. <i>Frontiers in Microbiology</i> , 2016, 7, 1690.	3.5	38
10	Secretion of non-cell-bound phytase by the yeast <i>Pichia kudriavzevii</i> TY13. <i>Journal of Applied Microbiology</i> , 2015, 118, 1126-1136.	3.1	18
11	Assessing phytase activity. <i>Journal of Biological Methods</i> , 2015, 2, e16.	0.6	33
12	Biosynthesis and cellular content of folate in bifidobacteria across host species with different diets. <i>Anaerobe</i> , 2014, 30, 169-177.	2.1	25
13	Degradation of phytate by <i>Pichia kudriavzevii</i> TY13 and <i>Hanseniaspora guilliermondii</i> TY14 in Tanzanian togwa. <i>International Journal of Food Microbiology</i> , 2012, 153, 73-77.	4.7	29
14	The potential of bifidobacteria as a source of natural folate. <i>Journal of Applied Microbiology</i> , 2012, 112, 975-984.	3.1	85
15	Biodiversity and phytase capacity of yeasts isolated from Tanzanian togwa. <i>International Journal of Food Microbiology</i> , 2010, 136, 352-358.	4.7	50
16	Determination of Fe ²⁺ and Fe ³⁺ in Aqueous Solutions Containing Food Chelators by Differential Pulse Anodic Stripping Voltammetry. <i>Electroanalysis</i> , 2010, 22, 1090-1096.	2.9	40
17	Phytate degradation by human gut isolated <i>Bifidobacterium pseudocatenulatum</i> ATCC27919 and its probiotic potential. <i>International Journal of Food Microbiology</i> , 2009, 135, 7-14.	4.7	48
18	Blocking Peptides Decrease Tissue Transglutaminase Processing of Gliadin in Vitro. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 10150-10155.	5.2	7

#	ARTICLE	IF	CITATIONS
19	Impaired uptake of β -carotene by Caco-2 human intestinal cells in the presence of iron. <i>International Journal of Food Sciences and Nutrition</i> , 2009, 60, 125-135.	2.8	9
20	Production of folates by yeasts in Tanzanian fermented togwa. <i>FEMS Yeast Research</i> , 2008, 8, 781-787.	2.3	42
21	Growth rate and medium composition strongly affect folate content in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Food Microbiology</i> , 2008, 123, 93-100.	4.7	37
22	Biofortification of folates in white wheat bread by selection of yeast strain and process. <i>International Journal of Food Microbiology</i> , 2008, 127, 32-36.	4.7	41
23	Effect of Nutrient Starvation on the Cellular Composition and Metabolic Capacity of <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2007, 73, 4839-4848.	3.1	40
24	Improved extracellular phytase activity in <i>Saccharomyces cerevisiae</i> by modifications in the PHO system. <i>International Journal of Food Microbiology</i> , 2006, 108, 60-67.	4.7	20
25	Lactic acid fermentation stimulated iron absorption by Caco-2 cells is associated with increased soluble iron content in carrot juice. <i>British Journal of Nutrition</i> , 2006, 96, 705-11.	2.3	31
26	Improved iron solubility in carrot juice fermented by homo- and hetero-fermentative lactic acid bacteria. <i>Food Microbiology</i> , 2005, 22, 53-61.	4.2	50
27	Degradation of Phytate by High-Phytase <i>Saccharomyces cerevisiae</i> Strains during Simulated Gastrointestinal Digestion. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 5438-5444.	5.2	42
28	Lactic Acid Decreases Fe(II) and Fe(III) Retention but Increases Fe(III) Transepithelial Transfer by Caco-2 Cells. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 6919-6923.	5.2	14
29	Inherent biodiversity of folate content and composition in yeasts. <i>Trends in Food Science and Technology</i> , 2005, 16, 311-316.	15.1	52
30	Development of a Simplified Method for the Determination of Folates in Baker's Yeast by HPLC with Ultraviolet and Fluorescence Detection. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 2406-2411.	5.2	82
31	Metabolism of extracellular inositol hexaphosphate (phytate) by <i>Saccharomyces cerevisiae</i> . <i>International Journal of Food Microbiology</i> , 2004, 97, 157-169.	4.7	46
32	Phytate content is reduced and β -glucanase activity suppressed in malted barley steeped with lactic acid at high temperature. <i>Journal of the Science of Food and Agriculture</i> , 2004, 84, 653-662.	3.5	34
33	Microbial leaching of uranium and other trace elements from shale mine tailings at Ranstad. <i>Geoderma</i> , 2004, 122, 177-194.	5.1	78
34	Prolonged Transit Time through the Stomach and Small Intestine Improves Iron Dialyzability and Uptake in Vitro. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 5131-5136.	5.2	23
35	Combined Impact of pH and Organic Acids on Iron Uptake by Caco-2 Cells. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 7820-7824.	5.2	45
36	Mobilisation of Radionuclides by Ligands Produced by Bacteria from the Deep Subsurface. <i>Materials Research Society Symposia Proceedings</i> , 2003, 807, 433.	0.1	3

#	ARTICLE	IF	CITATIONS
37	Organic Acids Influence Iron Uptake in the Human Epithelial Cell Line Caco-2. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 6233-6238.	5.2	90
38	Phytogenic and microbial phytases in human nutrition. <i>International Journal of Food Science and Technology</i> , 2002, 37, 823-833.	2.7	98
39	Phytate degradation by micro-organisms in synthetic media and pea flour. <i>Journal of Applied Microbiology</i> , 2002, 93, 197-204.	3.1	38
40	Inositol Hexaphosphate Hydrolysis by Baker's Yeast. Capacity, Kinetics, and Degradation Products. <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 100-104.	5.2	68
41	Characterization of <i>Saccharomyces cerevisiae</i> CBS 7764 Isolated From Rainbow Trout Intestine. <i>Systematic and Applied Microbiology</i> , 1999, 22, 145-155.	2.8	20
42	Yeast colonizing the intestine of rainbow trout (<i>Salmo gairdneri</i>) and turbot (<i>Scophthalmus maximus</i>). <i>Microbial Ecology</i> , 1995, 30, 321-34.	2.8	96
43	Cell surface hydrophobicity and its relation to adhesion of yeasts isolated from fish gut. <i>Colloids and Surfaces B: Biointerfaces</i> , 1994, 2, 199-208.	5.0	22
44	The expression of potential colonization factors of yeasts isolated from fish during different growth conditions. <i>Canadian Journal of Microbiology</i> , 1993, 39, 1135-1141.	1.7	22