

# Jiro Ikeda

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9692797/publications.pdf>

Version: 2024-02-01

15  
papers

417  
citations

840119

11  
h-index

996533

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

805  
citing authors

#	ARTICLE	IF	CITATIONS
1	Macrophage Jak2 deficiency accelerates atherosclerosis through defects in cholesterol efflux. <i>Communications Biology</i> , 2022, 5, 132.	2.0	4
2	Endothelial HMGB1 Is a Critical Regulator of LDL Transcytosis via an SREBP2-SR-BI Axis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021, 41, 200-216.	1.1	26
3	Radiation Impacts Early Atherosclerosis by Suppressing Intimal LDL Accumulation. <i>Circulation Research</i> , 2021, 128, 530-543.	2.0	12
4	Role of myeloid-derived chemokine CCL5/RANTES at an early stage of atherosclerosis. <i>Journal of Molecular and Cellular Cardiology</i> , 2021, 156, 69-78.	0.9	17
5	Suppression of abdominal aortic aneurysm formation by AR-R17779, an agonist for the $\alpha 7$ nicotinic acetylcholine receptor. <i>Atherosclerosis</i> , 2016, 244, 113-120.	0.4	8
6	PPAR $\alpha$ Agonists Attenuate Palmitate-Induced ER Stress through Up-Regulation of SCD-1 in Macrophages. <i>PLoS ONE</i> , 2015, 10, e0128546.	1.1	41
7	Suppression of abdominal aortic aneurysm formation by inhibition of prolyl hydroxylase domain protein through attenuation of inflammation and extracellular matrix disruption. <i>Clinical Science</i> , 2014, 126, 671-678.	1.8	27
8	Stimulation of $\alpha 7$ nicotinic acetylcholine receptor by AR-R17779 suppresses atherosclerosis and aortic aneurysm formation in apolipoprotein E-deficient mice. <i>Vascular Pharmacology</i> , 2014, 61, 49-55.	1.0	32
9	Deletion of <i>Phd2</i> in Myeloid Lineage Attenuates Hypertensive Cardiovascular Remodeling. <i>Journal of the American Heart Association</i> , 2013, 2, e000178.	1.6	30
10	Prolyl Hydroxylase Domain Protein 2 Plays a Critical Role in Diet-Induced Obesity and Glucose Intolerance. <i>Circulation</i> , 2013, 127, 2078-2087.	1.6	65
11	Acetylcholinesterase inhibitors attenuate angiogenesis. <i>Clinical Science</i> , 2012, 123, 241-249.	1.8	11
12	Beraprost sodium, a stable prostacyclin analogue, improves insulin resistance in high-fat diet-induced obese mice. <i>Journal of Endocrinology</i> , 2012, 213, 285-291.	1.2	19
13	Inhibition of Prolyl Hydroxylase Domain-Containing Protein Downregulates Vascular Angiotensin II Type 1 Receptor. <i>Hypertension</i> , 2011, 58, 386-393.	1.3	23
14	Inhibition of MDM2 attenuates neointimal hyperplasia via suppression of vascular proliferation and inflammation. <i>Cardiovascular Research</i> , 2011, 91, 711-719.	1.8	33
15	Inhibition of Prolyl Hydroxylase Domain-Containing Protein Suppressed Lipopolysaccharide-Induced TNF- $\alpha$ Expression. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009, 29, 2132-2137.	1.1	69