

Roman Fiedler

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9696001/publications.pdf>

Version: 2024-02-01

19
papers

278
citations

1163117

8
h-index

940533

16
g-index

19
all docs

19
docs citations

19
times ranked

518
citing authors

#	ARTICLE	IF	CITATIONS
1	Glycation Alters the Fatty Acid Binding Capacity of Human Serum Albumin. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 3033-3046.	5.2	10
2	FC020: The Uraemic Toxin Indoxyl Sulfate is an Activator Of Caspase-4 but not of Caspase-1 in Monocyte-Like THP-1 Cells. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, .	0.7	0
3	Serum Testosterone Levels Are Not Modified by Vitamin D Supplementation in Dialysis Patients and Healthy Subjects. <i>Nephron</i> , 2021, 145, 481-485.	1.8	3
4	Pyroptosis: A Common Feature of Immune Cells of Haemodialysis Patients. <i>Toxins</i> , 2021, 13, 839.	3.4	4
5	No significant relation of proinflammatory slanDCs with uremic pruritus. <i>European Journal of Inflammation</i> , 2020, 18, 205873922092685.	0.5	0
6	NLRP3 Inflammasome Activation in Hemodialysis and Hypertensive Patients with Intact Kidney Function. <i>Toxins</i> , 2020, 12, 675.	3.4	8
7	P0924IMMUNE DISTURBANCES INDUCED BY HYPERVOLEMIA DO NOT ACTIVATE THE NLRP3 INFLAMMASOME IN HAEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
8	Hypervolemia-Induced Immune Disturbances Do Not Involve IL-1 β but IL-6 and IL-10 Activation in Haemodialysis Patients. <i>Toxins</i> , 2020, 12, 159.	3.4	6
9	Circulating miR-421 Targeting Leucocytic Angiotensin Converting Enzyme 2 Is Elevated in Patients with Chronic Kidney Disease. <i>Nephron</i> , 2019, 141, 61-74.	1.8	26
10	Influence of Transketolase-Catalyzed Reactions on the Formation of Glycolaldehyde and Glyoxal Specific Posttranslational Modifications under Physiological Conditions. <i>Journal of Agricultural and Food Chemistry</i> , 2018, 66, 1498-1508.	5.2	5
11	Modulation of leucocytic angiotensin-converting enzymes expression in patients maintained on high-permeable haemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 34-43.	0.7	9
12	Monocytic angiotensin-converting enzyme 2 relates to atherosclerosis in patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfw206.	0.7	42
13	Detection of Free Advanced Glycation End Products in Vivo during Hemodialysis. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 930-937.	5.2	25
14	Association between autonomic nervous dysfunction and cellular inflammation in end-stage renal disease. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 210.	1.7	8
15	Uremic Conditions Drive Human Monocytes to Pro-Atherogenic Differentiation via an Angiotensin-Dependent Mechanism. <i>PLoS ONE</i> , 2014, 9, e102137.	2.5	20
16	Randomized Controlled Pilot Study of 2 Weeks' Treatment With High Cutoff Membrane for Hemodialysis Patients With Elevated C-reactive Protein. <i>Artificial Organs</i> , 2012, 36, 886-893.	1.9	26
17	Vitamin D Deficiency, Mortality, and Hospitalization in Hemodialysis Patients with or without Protein-Energy Wasting. <i>Nephron Clinical Practice</i> , 2011, 119, c220-c226.	2.3	20
18	Hemoglobin Cycling in Hemodialysis Patients. <i>Nephrology Research & Reviews</i> , 2010, 2, 1-5.	0.2	4

#	ARTICLE	IF	CITATIONS
19	Clinical nutrition scores are superior for the prognosis of haemodialysis patients compared to lab markers and bioelectrical impedance. Nephrology Dialysis Transplantation, 2009, 24, 3812-3817.	0.7	62