## Xing Wu

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9749916/publications.pdf

Version: 2024-02-01

		566801	500791
31	857	15	28
papers	citations	h-index	g-index
38	38	38	1347
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Epidemiology of Traumatic Brain Injury in Eastern China, 2004: A Prospective Large Case Study. Journal of Trauma, 2008, 64, 1313-1319.	2.3	121
2	How are different neural networks related to consciousness?. Annals of Neurology, 2015, 78, 594-605.	2.8	102
3	In vivo tracking of superparamagnetic iron oxide nanoparticle–labeled mesenchymal stem cell tropism to malignant gliomas using magnetic resonance imaging. Journal of Neurosurgery, 2008, 108, 320-329.	0.9	101
4	Coagulopathy in Traumatic Brain Injury and Its Correlation with Progressive Hemorrhagic Injury: A Systematic Review and Meta-Analysis. Journal of Neurotrauma, 2016, 33, 1279-1291.	1.7	72
5	Prognostic value of coagulation tests for in-hospital mortality in patients with traumatic brain injury. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2018, 26, 3.	1.1	44
6	N-Terminal Pro-B-Type Natriuretic Peptide in Patients With Isolated Traumatic Brain Injury: A Prospective Cohort Study. Journal of Trauma, 2011, 71, 820-825.	2.3	28
7	Circulating neutrophil-to-lymphocyte ratio at admission predicts the long-term outcome in acute traumatic cervical spinal cord injury patients. BMC Musculoskeletal Disorders, 2020, 21, 548.	0.8	26
8	Geniposide ameliorates chronic unpredictable mild stress induced depression-like behavior through inhibition of ceramide-PP2A signaling via the PI3K/Akt/GSK3 $\hat{I}^2$ axis. Psychopharmacology, 2021, 238, 2789-2800.	1.5	26
9	Prognostic Value of Neutrophil-to-Lymphocyte Ratio in Predicting the 6-Month Outcome of Patients with Traumatic Brain Injury: A Retrospective Study. World Neurosurgery, 2019, 124, e411-e416.	0.7	24
10	Prevalence of severe hypokalaemia in patients with traumatic brain injury. Injury, 2015, 46, 35-41.	0.7	23
11	Low-dose recombinant factor VIIa for reversing coagulopathy in patients with isolated traumatic brain injury. Journal of Critical Care, 2015, 30, 116-120.	1.0	23
12	Intracranial Pressure Monitoring-Aided Management Associated with Favorable Outcomes in Patients with Hypertension-Related Spontaneous Intracerebral Hemorrhage. Translational Stroke Research, 2020, 11, 1253-1263.	2.3	22
13	Effects of Deferoxamine Mesylate on Hematoma and Perihematoma Edema after Traumatic Intracerebral Hemorrhage. Journal of Neurotrauma, 2017, 34, 2753-2759.	1.7	19
14	Empirical Combination Antibiotic Therapy Improves the Outcome of Nosocomial Meningitis or Ventriculitis in Neuro-Critical Care Unit Patients. Surgical Infections, 2016, 17, 465-472.	0.7	18
15	Diagnosis and Management of Combined Central Diabetes Insipidus and Cerebral Salt Wasting Syndrome After Traumatic Brain Injury. World Neurosurgery, 2016, 88, 483-487.	0.7	18
16	ISO-alpha-acids improve the hematoma resolution and prevent peri-hematoma inflammations by transforming microglia via PPARgamma-CD36 axis in ICH rats. International Immunopharmacology, 2020, 83, 106396.	1.7	18
17	<scp>GABA<sub>A</sub></scp> receptor deficits predict recovery in patients with disorders of consciousness: A preliminary multimodal [ <sup>11</sup> C]Flumazenil <scp>PET</scp> and f <scp>MRI</scp> study. Human Brain Mapping, 2015, 36, 3867-3877.	1.9	17
18	Impact of fibrinogen level on the prognosis of patients with traumatic brain injury: a single-center analysis of 2570 patients. World Journal of Emergency Surgery, 2020, 15, 54.	2.1	17

#	Article	IF	Citations
19	Activity of factor VII in patients with isolated blunt traumatic brain injury. Journal of Trauma and Acute Care Surgery, 2014, 76, 114-120.	1.1	16
20	Early post-traumatic seizures are associated with valproic acid plasma concentrations and UGT1A6/CYP2C9 genetic polymorphisms in patients with severe traumatic brain injury. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2017, 25, 85.	1.1	16
21	Intensive blood pressure control reduces the risk of progressive hemorrhage in patients with acute hypertensive intracerebral hemorrhage: A retrospective observational study. Clinical Neurology and Neurosurgery, 2019, 180, 1-6.	0.6	14
22	Prognostic significance of stem cell marker CD133 determined by promoter methylation but not by immunohistochemical expression in malignant gliomas. Journal of Neuro-Oncology, 2016, 127, 221-232.	1.4	13
23	p53 participates in the protective effects of ischemic post-conditioning against OGD-reperfusion injury in primary cultured spinal cord neurons. Neuroscience Letters, 2017, 638, 129-134.	1.0	13
24	Posterior cingulate cross-hemispheric functional connectivity predicts the level of consciousness in traumatic brain injury. Scientific Reports, 2017, 7, 387.	1.6	12
25	FVIIa prevents the progressive hemorrhaging of a brain contusion by protecting microvessels via formation of the TF–FVIIa–FXa complex. Neuroscience, 2017, 348, 114-125.	1.1	11
26	Prevention options for ventriculoperitoneal shunt infections: a retrospective analysis during a five-year period. International Journal of Clinical and Experimental Medicine, 2015, 8, 19775-80.	1.3	10
27	Admission circulating monocytes level is an independent predictor of outcome in traumatic brain injury. Brain Injury, 2018, 32, 515-522.	0.6	9
28	Diagnosis and Management for Secondary Low- or Negative-Pressure Hydrocephalus and a New Hydrocephalus Classification Based on Ventricular Pressure. World Neurosurgery, 2019, 124, e510-e516.	0.7	7
29	Trends of Intra-Cranial Bacterial Infection in Patients Requiring Emergency Neurosurgery. Surgical Infections, 2020, 21, 677-683.	0.7	5
30	Contribution of factor VII polymorphisms to coagulopathy in patients with isolated traumatic brain injury. Clinical Neurology and Neurosurgery, 2021, 208, 106836.	0.6	5
31	The First 24 h Hemodynamic Management in NICU after Revascularization Surgery in Moyamoya Disease. Behavioural Neurology, 2021, 2021, 1-7.	1.1	3