## Ralf Dieckmann

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9750220/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Genetic but No Phenotypic Associations between Biocide Tolerance and Antibiotic Resistance in Escherichia coli from German Broiler Fattening Farms. Microorganisms, 2021, 9, 651.	3.6	21
2	Backtracking and forward checking of human listeriosis clusters identified a multiclonal outbreak linked to <i>Listeria monocytogenes</i> in meat products of a single producer. Emerging Microbes and Infections, 2020, 9, 1600-1608.	6.5	27
3	Microbially competent 3D skin: a test system that reveals insight into host–microbe interactions and their potential toxicological impact. Archives of Toxicology, 2020, 94, 3487-3502.	4.2	12
4	A Proof of Principle for the Detection of Viable Brucella spp. in Raw Milk by qPCR Targeting Bacteriophages. Microorganisms, 2020, 8, 1326.	3.6	3
5	MALDI-TOF MS and genomic analysis can make the difference in the clarification of canine brucellosis outbreaks. Scientific Reports, 2020, 10, 19246.	3.3	9
6	Evaluation of a Newly Developed Vacuum Dried Microtiter Plate for Rapid Biocide Susceptibility Testing of Clinical Enterococcus faecium Isolates. Microorganisms, 2020, 8, 551.	3.6	4
7	Benzalkonium Chloride Induces a VBNC State in Listeria monocytogenes. Microorganisms, 2020, 8, 184.	3.6	28
8	Microbiological Safety of Non-Food Products: What Can We Learn from the RAPEX Database?. International Journal of Environmental Research and Public Health, 2019, 16, 1599.	2.6	8
9	Overview of validated alternative methods for the detection of foodborne bacterial pathogens. Trends in Food Science and Technology, 2017, 62, 113-118.	15.1	87
10	Differential detection of pathogenic Yersinia spp. by fluorescence in situ hybridization. Food Microbiology, 2017, 62, 39-45.	4.2	17
11	Molecular Tracing to Find Source of Protracted Invasive Listeriosis Outbreak, Southern Germany, 2012–2016. Emerging Infectious Diseases, 2017, 23, 1680-1683.	4.3	47
12	The Risk of Bacterial Infection After Tattooing. Deutsches Ärzteblatt International, 2016, 113, 665-671.	0.9	41
13	Rapid characterisation of Klebsiella oxytoca isolates from contaminated liquid hand soap using mass spectrometry, FTIR and Raman spectroscopy. Faraday Discussions, 2016, 187, 353-375.	3.2	29
14	Peptaibol, Secondaryâ€Metabolite, and Hydrophobin Pattern of Commercial Biocontrol Agents Formulated with Species of the <i>Trichoderma harzianum</i> Complex. Chemistry and Biodiversity, 2015, 12, 662-684.	2.1	57
15	Virulence Profiles of Vibrio vulnificus in German Coastal Waters, a Comparison of North Sea and Baltic Sea Isolates. International Journal of Environmental Research and Public Health, 2015, 12, 15943-15959.	2.6	14
16	Sampling and Homogenization Strategies Significantly Influence the Detection of Foodborne Pathogens in Meat. BioMed Research International, 2015, 2015, 1-8.	1.9	17
17	FISHing for bacteria in food – A promising tool for the reliable detection of pathogenic bacteria?. Food Microbiology, 2015, 46, 395-407	4.2	84
18	Characterization of trh2 Harbouring Vibrio parahaemolyticus Strains Isolated in Germany. PLoS ONE, 2015, 10, e0118559.	2.5	15

RALF DIECKMANN

#	Article	IF	CITATIONS
19	Evaluation of molecular methods to discriminate the closely related species Vibrio fluvialis and Vibrio furnissii. International Journal of Medical Microbiology, 2014, 304, 851-857.	3.6	22
20	Pathogenic vibrios in environmental, seafood and clinical sources in Germany. International Journal of Medical Microbiology, 2014, 304, 843-850.	3.6	124
21	Cell-free synthesis of functional thermostable direct hemolysins of Vibrio parahaemolyticus. Toxicon, 2013, 76, 132-142.	1.6	35
22	Genotypic Diversity and Virulence Characteristics of Clinical and Environmental Vibrio vulnificus Isolates from the Baltic Sea Region. Applied and Environmental Microbiology, 2013, 79, 3570-3581.	3.1	34
23	Spread of a Distinct Stx2-Encoding Phage Prototype among Escherichia coli O104:H4 Strains from Outbreaks in Germany, Norway, and Georgia. Journal of Virology, 2012, 86, 10444-10455.	3.4	39
24	The Production of Multiple Small Peptaibol Families by Single 14â€Module Peptide Synthetases in <i>Trichoderma</i> / <i>Hypocrea</i> . Chemistry and Biodiversity, 2012, 9, 499-535.	2.1	66
25	Rapid Screening of Epidemiologically Important Salmonella enterica subsp. enterica Serovars by Whole-Cell Matrix-Assisted Laser Desorption Ionization–Time of Flight Mass Spectrometry. Applied and Environmental Microbiology, 2011, 77, 4136-4146.	3.1	145
26	Characterization of Yersinia Using MALDI-TOF Mass Spectrometry and Chemometrics. Analytical Chemistry, 2010, 82, 8464-8475.	6.5	60
27	Differential Regulation and Posttranslational Processing of the Class II Hydrophobin Genes from the Biocontrol Fungus <i>Hypocrea atroviridis</i> . Applied and Environmental Microbiology, 2009, 75, 3222-3229.	3.1	23
28	Polymerase Chain Reaction for the Rapid Detection and Serovar Identification of Salmonella in Food and Feeding Stuff. Food Analytical Methods, 2009, 2, 81-95.	2.6	47
29	The Trichoderma brevicompactum clade: a separate lineage with new species, new peptaibiotics, and mycotoxins. Mycological Progress, 2008, 7, 177-219.	1.4	136
30	Spongiibacter marinus gen. nov., sp. nov., a halophilic marine bacterium isolated from the boreal sponge Haliclona sp. 1. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, 585-590.	1.7	32
31	Spongiispira norvegica gen. nov., sp. nov., a marine bacterium isolated from the boreal sponge Isops phlegraei. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, 1815-1820.	1.7	18
32	Rapid Classification and Identification of Salmonellae at the Species and Subspecies Levels by Whole-Cell Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry. Applied and Environmental Microbiology, 2008, 74, 7767-7778.	3.1	236
33	Formation of Atroviridin by <i>Hypocrea atroviridis</i> Is Conidiation Associated and Positively Regulated by Blue Light and the G Protein GNA3. Eukaryotic Cell, 2007, 6, 2332-2342.	3.4	48
34	Intact-cell MALDI-TOF mass spectrometry analysis of peptaibol formation by the genus Trichoderma/Hypocrea: can molecular phylogeny of species predict peptaibol structures?. Microbiology (United Kingdom), 2007, 153, 3417-3437.	1.8	77
35	Peptaibol Production bySepedonium Strains Parasitizing Boletales. Chemistry and Biodiversity, 2007, 4, 1103-1115.	2.1	32
36	Direct identification of hydrophobins and their processing in Trichoderma using intact-cell MALDI-TOF MS. FEBS Journal, 2007, 274, 841-852.	4.7	49

RALF DIECKMANN

#	Article	IF	CITATIONS
37	Nonribosomal Peptide Synthesis in Schizosaccharomyces pombe and the Architectures of Ferrichrome-Type Siderophore Synthetases in Fungi. ChemBioChem, 2006, 7, 612-622.	2.6	83
38	Qualitative variation of alkaloids in color morphs of Cystodytes (Ascidiacea). Biochemical Systematics and Ecology, 2005, 33, 1107-1119.	1.3	38
39	Hassallidin A, a Glycosylated Lipopeptide with Antifungal Activity from the CyanobacteriumHassalliasp Journal of Natural Products, 2005, 68, 695-700.	3.0	97
40	Algicide production by the filamentous cyanobacterium Fischerella sp. CENA 19. Journal of Applied Phycology, 2004, 16, 237-243.	2.8	43
41	ATPase activity of non-ribosomal peptide synthetases. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2004, 1696, 83-91.	2.3	5
42	Nonribosomal peptide synthetases-evidence for a second ATP-binding site. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2002, 1601, 93-99.	2.3	8
43	Barettin, revisited?. Tetrahedron Letters, 2002, 43, 3385-3386.	1.4	51
44	Synthesis of (di)adenosine polyphosphates by non-ribosomal peptide synthetases (NRPS). BBA - Proteins and Proteomics, 2001, 1546, 234-241.	2.1	21
45	Probing the domain structure and ligand-induced conformational changes by limited proteolysis of tyrocidine synthetase 1. Journal of Molecular Biology, 1999, 288, 129-140.	4.2	49
46	Editing of non-cognate aminoacyl adenylates by peptide synthetases. Biochemical Journal, 1999, 342, 715-719.	3.7	15
47	Editing of non-cognate aminoacyl adenylates by peptide synthetases. Biochemical Journal, 1999, 342, 715.	3.7	6
48	ACV Synthetase: Expression of Amino Acid Activating Domains of thePenicillium chrysogenumEnzyme inAspergillus nidulans. Biochemical and Biophysical Research Communications, 1997, 237, 166-169.	2.1	11
49	The Adenylation Domain of Tyrocidine Synthetase 1. Structural and Functional Role of the Interdomain Linker Region and the (S/T)GT(T/S)GXPKG Core Sequence. FEBS Journal, 1997, 247, 1074-1082.	0.2	23