

# Yasuhiro Kuroda

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9783143/publications.pdf>

Version: 2024-02-01

12  
papers

166  
citations

1163117

8  
h-index

1199594

12  
g-index

12  
all docs

12  
docs citations

12  
times ranked

245  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Vaccination with profilin encapsulated in oligomannose-coated liposomes induces significant protective immunity against <i>Toxoplasma gondii</i> . <i>Vaccine</i> , 2014, 32, 1781-1785.  | 3.8 | 26        |
| 2  | Immunization with <i>Toxoplasma gondii</i> peroxiredoxin 1 induces protective immunity against toxoplasmosis in mice. <i>PLoS ONE</i> , 2017, 12, e0176324.   | 2.5 | 24        |
| 3  | A Critical Role for Sialylation in Cryoglobulin Activity of Murine IgG3 Monoclonal Antibodies. <i>Journal of Immunology</i> , 2005, 175, 1056-1061.   | 0.8 | 22        |
| 4  | Neospora GRA6 possesses immune-stimulating activity and confers efficient protection against <i>Neospora caninum</i> infection in mice. <i>Veterinary Parasitology</i> , 2019, 267, 61-68.  | 1.8 | 19        |
| 5  | Structural studies on IgG oligosaccharides of patients with primary Sjögren's syndrome. <i>Glycoconjugate Journal</i> , 2002, 19, 23-31.  | 2.7 | 16        |
| 6  | Abnormal IgG galactosylation and arthritis in MRL-Fas <sup>lpr</sup> or MRL-Fas <sup>Lgld</sup> mice are under the control of the MRL genetic background. <i>FEBS Letters</i> , 2001, 507, 210-214.   | 2.8 | 14        |
| 7  | Critical role of TLR2 in triggering protective immunity with cyclophilin entrapped in oligomannose-coated liposomes against <i>Neospora caninum</i> infection in mice. <i>Vaccine</i> , 2019, 37, 937-944.                                    | 3.8 | 14        |
| 8  | Requirement of TLR4 signaling for the induction of a Th1 immune response elicited by oligomannose-coated liposomes. <i>Immunology Letters</i> , 2016, 178, 61-67.   | 2.5 | 9         |
| 9  | Comparison of the carbohydrate preference of SIGNR1 as a phagocytic receptor with the preference as an adhesion molecule. <i>International Immunopharmacology</i> , 2014, 19, 27-36.  | 3.8 | 7         |
| 10 | In vitro activation and maturation of human mononuclear phagocytes by stimulation with liposomes coated with a neoglycolipid containing $\alpha$ 3, $\alpha$ 6-mannotriose. <i>Glycoconjugate Journal</i> , 2019, 36, 185-197. <sup>2,7</sup> |     | 6         |
| 11 | Importance of particle size of oligomannose-coated liposomes for induction of Th1 immunity. <i>International Immunopharmacology</i> , 2021, 99, 108068.   | 3.8 | 5         |
| 12 | In vitro uptake of oligomannose-coated liposomes leads to differentiation of inflammatory monocytes into mature antigen-presenting cells that can activate T cells. <i>International Immunopharmacology</i> , 2018, 57, 102-111.              | 3.8 | 4         |