

Hiroko Nagafuchi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9801273/publications.pdf>

Version: 2024-02-01

34
papers

1,082
citations

361413

20
h-index

454955

30
g-index

34
all docs

34
docs citations

34
times ranked

1218
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinical features of Behçet's disease patients with joint symptoms in Japan: A national multicenter study. <i>Modern Rheumatology</i> , 2022, 32, 1146-1152.	1.8	1
2	Pregnancy outcomes in patients with rheumatoid arthritis who discontinue methotrexate treatment to conceive. <i>Clinical Rheumatology</i> , 2022, 41, 669-675.	2.2	1
3	Rheumatoid arthritis relapse in patients with other iatrogenic immunodeficiency-associated lymphoproliferative disorders and its treatment. <i>Modern Rheumatology</i> , 2021, 31, 1087-1093.	1.8	1
4	Establishing clinical remission criteria and the framework of a treat-to-target algorithm for Takayasu arteritis: Results of a Delphi exercise carried out by an expert panel of the Japan Research Committee of the Ministry of Health, Labour and Welfare for intractable vasculitis. <i>Modern Rheumatology</i> , 2021, 31, 1094-1103.	1.8	5
5	Lack of partial renal response by 12 weeks after induction therapy predicts poor renal response and systemic damage accrual in lupus nephritis class III or IV. <i>Arthritis Research and Therapy</i> , 2017, 19, 4.	3.5	18
6	Psoriatic Arthritis with Annular Pustular Psoriasis. <i>Internal Medicine</i> , 2016, 55, 519-521.	0.7	0
7	Recurrent Bilateral Focal Myositis. <i>Internal Medicine</i> , 2016, 55, 3369-3374.	0.7	3
8	Clinical subsets associated with different anti-aminoacyl transfer RNA synthetase antibodies and their association with coexisting anti-Ro52. <i>Modern Rheumatology</i> , 2016, 26, 403-409.	1.8	38
9	Retrospective analysis of long-term outcome of chronic progressive neurological manifestations in Behçet's disease. <i>Journal of the Neurological Sciences</i> , 2015, 349, 143-148.	0.6	32
10	Long-term safety and efficacy of rituximab in 7 Japanese patients with ANCA-associated vasculitis. <i>Modern Rheumatology</i> , 2015, 25, 603-608.	1.8	21
11	Serum level of soluble triggering receptor expressed on myeloid cells-1 as a biomarker of disease activity in relapsing polychondritis. <i>Modern Rheumatology</i> , 2014, 24, 129-136.	1.8	32
12	Assessment of the Birmingham vasculitis activity score in patients with MPO-ANCA-associated vasculitis: sub-analysis from a study by the Japanese Study Group for MPO-ANCA-associated vasculitis. <i>Modern Rheumatology</i> , 2014, 24, 304-309.	1.8	12
13	Analysis of various factors on the relapse of acute neurological attacks in Behçet's disease. <i>Modern Rheumatology</i> , 2014, 24, 961-965.	1.8	8
14	Comparison of Phenotype and Outcome in Microscopic Polyangiitis Between Europe and Japan. <i>Journal of Rheumatology</i> , 2014, 41, 325-333.	2.0	83
15	Assessment of the Birmingham vasculitis activity score in patients with MPO-ANCA-associated vasculitis: sub-analysis from a study by the Japanese Study Group for MPO-ANCA-associated vasculitis. <i>Modern Rheumatology</i> , 2013, 23, 1.	1.8	1
16	Severity-based treatment for Japanese patients with MPO-ANCA-associated vasculitis: the JMAAV study. <i>Modern Rheumatology</i> , 2012, 22, 394-404.	1.8	77
17	Clinical characteristics of neuro-Behçet's disease in Japan: a multicenter retrospective analysis. <i>Modern Rheumatology</i> , 2012, 22, 405-413.	1.8	73
18	Severity-based treatment for Japanese patients with MPO-ANCA-associated vasculitis: the JMAAV study. <i>Modern Rheumatology</i> , 2012, 22, 394-404.	1.8	45

#	ARTICLE	IF	CITATIONS
19	Clinical characteristics of neuro-Behçet's disease in Japan: a multicenter retrospective analysis. <i>Modern Rheumatology</i> , 2012, 22, 405-413.	1.8	48
20	Cutaneous polyarteritis nodosa associated with HLA-B39-positive undifferentiated spondyloarthritis in a Japanese patient. <i>Modern Rheumatology</i> , 2012, 22, 783-786.	1.8	2
21	AC13, a C-terminal fragment of apolipoprotein A-II, is a candidate biomarker for microscopic polyangiitis. <i>Arthritis and Rheumatism</i> , 2011, 63, 3613-3624.	6.7	10
22	Lupus antibodies to the HMGB1 chromosomal protein: epitope mapping and association with disease activity. <i>Modern Rheumatology</i> , 2009, 19, 283-292.	1.8	37
23	Lupus antibodies to the HMGB1 chromosomal protein: epitope mapping and association with disease activity. <i>Modern Rheumatology</i> , 2009, 19, 283-292.	1.8	32
24	Clinical and histopathologic features of 8 patients with microscopic polyangiitis including two with a slowly progressive clinical course. <i>Journal of the American Academy of Dermatology</i> , 2007, 57, 840-848.	1.2	22
25	The role of HMGB-1 on the development of necrosis during hepatic ischemia and hepatic ischemia/reperfusion injury in mice. <i>Journal of Surgical Research</i> , 2005, 124, 59-66.	1.6	87
26	Abnormal killer inhibitory receptor expression on natural killer cells in patients with Behçet's disease. <i>Rheumatology International</i> , 2004, 24, 212-216.	3.0	41
27	Anatomical and functional recovery by embryonic stem cell-derived neural tissue of a mouse model of brain damage. <i>Journal of the Neurological Sciences</i> , 2004, 219, 107-117.	0.6	42
28	Txk, a Member of Nonreceptor Tyrosine Kinase of Tec Family, Acts as a Th1 Cell-Specific Transcription Factor and Regulates IFN- γ Gene Transcription. <i>Journal of Immunology</i> , 2002, 168, 2365-2370.	0.8	64
29	Flk1+ cells derived from mouse embryonic stem cells reconstitute hematopoiesis in vivo in SCID mice. <i>Experimental Hematology</i> , 2002, 30, 1444-1453.	0.4	34
30	Characterization of Tissue Outgrowth Developed in vitro in Patients with Rheumatoid Arthritis: Involvement of T Cells in the Development of Tissue Outgrowth. <i>International Archives of Allergy and Immunology</i> , 2000, 121, 68-79.	2.1	11
31	Txk, a Nonreceptor Tyrosine Kinase of the Tec Family, Is Expressed in T Helper Type 1 Cells and Regulates Interferon γ Production in Human T Lymphocytes. <i>Journal of Experimental Medicine</i> , 1999, 190, 1147-1154.	8.5	82
32	Etiopathology of Behçet's disease: immunological aspects. <i>Yonsei Medical Journal</i> , 1997, 38, 350.	2.2	67
33	Methyl-vitamin B12 blocks the CD28 co-stimulatory pathway in human T cells and its possible therapeutic application for T cell-mediated diseases, including rheumatoid arthritis. <i>Japanese Journal of Rheumatology</i> , 1997, 7, 35-45.	0.0	0
34	Fine antigen specificity of human $\gamma\delta$ T cell lines ($V\delta 9+$) established by repetitive stimulation with a serotype (KTH-1) of a gram-positive bacterium, <i>Streptococcus sanguis</i> . <i>European Journal of Immunology</i> , 1994, 24, 1536-1543.	2.9	52