

Peter W Reddien

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9811057/peter-w-reddien-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

67
papers

7,704
citations

41
h-index

74
g-index

74
ext. papers

9,241
ext. citations

15.4
avg, IF

6.61
L-index

#	Paper	IF	Citations
67	The planarian wound epidermis gene equinox is required for blastema formation in regeneration.. <i>Nature Communications</i> , 2022 , 13, 2726	17.4	3
66	activin-2 is required for regeneration of polarity on the planarian anterior-posterior axis. <i>PLoS Genetics</i> , 2021 , 17, e1009466	6	4
65	Planarian stem cells specify fate yet retain potency during the cell cycle. <i>Cell Stem Cell</i> , 2021 , 28, 1307-1322.e511	33.3	11
64	Principles of regeneration revealed by the planarian eye. <i>Current Opinion in Cell Biology</i> , 2021 , 73, 19-25	9	2
63	Muscle and neuronal guidepost-like cells facilitate planarian visual system regeneration. <i>Science</i> , 2020 , 368,	33.3	9
62	Acoel genome reveals the regulatory landscape of whole-body regeneration. <i>Science</i> , 2019 , 363,	33.3	61
61	Muscle functions as a connective tissue and source of extracellular matrix in planarians. <i>Nature Communications</i> , 2019 , 10, 1592	17.4	26
60	The cells of regeneration. <i>Science</i> , 2019 , 365, 314-316	33.3	1
59	A small set of conserved genes, including sp5 and Hox, are activated by Wnt signaling in the posterior of planarians and acoels. <i>PLoS Genetics</i> , 2019 , 15, e1008401	6	11
58	Nuclear receptor NR4A is required for patterning at the ends of the planarian anterior-posterior axis. <i>ELife</i> , 2019 , 8,	8.9	9
57	A small set of conserved genes, including sp5 and Hox, are activated by Wnt signaling in the posterior of planarians and acoels 2019 , 15, e1008401		
56	A small set of conserved genes, including sp5 and Hox, are activated by Wnt signaling in the posterior of planarians and acoels 2019 , 15, e1008401		
55	A small set of conserved genes, including sp5 and Hox, are activated by Wnt signaling in the posterior of planarians and acoels 2019 , 15, e1008401		
54	A small set of conserved genes, including sp5 and Hox, are activated by Wnt signaling in the posterior of planarians and acoels 2019 , 15, e1008401		
53	Cell type transcriptome atlas for the planarian. <i>Science</i> , 2018 , 360,	33.3	202
52	Self-organization and progenitor targeting generate stable patterns in planarian regeneration. <i>Science</i> , 2018 , 360, 404-409	33.3	23
51	Clonal Analysis of Planarian Stem Cells by Subtotal Irradiation and Single-Cell Transplantation. <i>Methods in Molecular Biology</i> , 2018 , 1774, 479-495	1.4	6

50	FoxF-1 Controls Specification of Non-body Wall Muscle and Phagocytic Cells in Planarians. <i>Current Biology</i> , 2018 , 28, 3787-3801.e6	6.3	21
49	Cellular and Molecular Responses Unique to Major Injury Are Dispensable for Planarian Regeneration. <i>Cell Reports</i> , 2018 , 25, 2577-2590.e3	10.6	21
48	The Cellular and Molecular Basis for Planarian Regeneration. <i>Cell</i> , 2018 , 175, 327-345	56.2	130
47	Landmarks in Existing Tissue at Wounds Are Utilized to Generate Pattern in Regenerating Tissue. <i>Current Biology</i> , 2017 , 27, 733-742	6.3	28
46	Eye Absence Does Not Regulate Planarian Stem Cells during Eye Regeneration. <i>Developmental Cell</i> , 2017 , 40, 381-391.e3	10.2	28
45	Planarian Epidermal Stem Cells Respond to Positional Cues to Promote Cell-Type Diversity. <i>Developmental Cell</i> , 2017 , 40, 491-504.e5	10.2	47
44	Acoel regeneration mechanisms indicate an ancient role for muscle in regenerative patterning. <i>Nature Communications</i> , 2017 , 8, 1260	17.4	22
43	Orthogonal muscle fibres have different instructive roles in planarian regeneration. <i>Nature</i> , 2017 , 551, 623-628	50.4	61
42	Hedgehog signaling regulates gene expression in planarian glia. <i>ELife</i> , 2016 , 5,	8.9	42
41	A widely employed germ cell marker is an ancient disordered protein with reproductive functions in diverse eukaryotes. <i>ELife</i> , 2016 , 5,	8.9	35
40	Two FGFR-LWnt circuits organize the planarian anteroposterior axis. <i>ELife</i> , 2016 , 5,	8.9	69
39	teashirt is required for head-versus-tail regeneration polarity in planarians. <i>Development (Cambridge)</i> , 2015 , 142, 1062-72	6.6	21
38	A Generic and Cell-Type-Specific Wound Response Precedes Regeneration in Planarians. <i>Developmental Cell</i> , 2015 , 35, 632-645	10.2	139
37	Single-cell analysis reveals functionally distinct classes within the planarian stem cell compartment. <i>Cell Stem Cell</i> , 2014 , 15, 326-339	18	191
36	Neoblast specialization in regeneration of the planarian <i>Schmidtea mediterranea</i> . <i>Stem Cell Reports</i> , 2014 , 3, 339-52	8	140
35	Whole-body acoel regeneration is controlled by Wnt and Bmp-Admp signaling. <i>Current Biology</i> , 2014 , 24, 1107-13	6.3	113
34	A forkhead transcription factor is wound-induced at the planarian midline and required for anterior pole regeneration. <i>PLoS Genetics</i> , 2014 , 10, e1003999	6	59
33	Lin28: time for tissue repair. <i>Cell</i> , 2013 , 155, 738-9	56.2	7

32	pbx is required for pole and eye regeneration in planarians. <i>Development (Cambridge)</i> , 2013 , 140, 719-296.6	4.2
31	Muscle cells provide instructions for planarian regeneration. <i>Cell Reports</i> , 2013 , 4, 633-41	10.6 159
30	Specialized progenitors and regeneration. <i>Development (Cambridge)</i> , 2013 , 140, 951-7	6.6 88
29	Tissue absence initiates regeneration through follistatin-mediated inhibition of activin signaling. <i>ELife</i> , 2013 , 2, e00247	8.9 106
28	Transcriptome analysis of the planarian eye identifies ovo as a specific regulator of eye regeneration. <i>Cell Reports</i> , 2012 , 2, 294-307	10.6 134
27	Genetic regulators of a pluripotent adult stem cell system in planarians identified by RNAi and clonal analysis. <i>Cell Stem Cell</i> , 2012 , 10, 299-311	18 140
26	A molecular wound response program associated with regeneration initiation in planarians. <i>Genes and Development</i> , 2012 , 26, 988-1002	12.6 163
25	The cellular basis for animal regeneration. <i>Developmental Cell</i> , 2011 , 21, 172-85	10.2 374
24	Constitutive gene expression and the specification of tissue identity in adult planarian biology. <i>Trends in Genetics</i> , 2011 , 27, 277-85	8.5 76
23	A Bmp/Admp regulatory circuit controls maintenance and regeneration of dorsal-ventral polarity in planarians. <i>Current Biology</i> , 2011 , 21, 294-9	6.3 81
22	Polarized notum activation at wounds inhibits Wnt function to promote planarian head regeneration. <i>Science</i> , 2011 , 332, 852-5	33.3 168
21	Clonogenic neoblasts are pluripotent adult stem cells that underlie planarian regeneration. <i>Science</i> , 2011 , 332, 811-6	33.3 433
20	A regulatory program for excretory system regeneration in planarians. <i>Development (Cambridge)</i> , 2011 , 138, 4387-98	6.6 114
19	dlx and sp6-9 Control optic cup regeneration in a prototypic eye. <i>PLoS Genetics</i> , 2011 , 7, e1002226	6 99
18	The Zn finger protein Iguana impacts Hedgehog signaling by promoting ciliogenesis. <i>Developmental Biology</i> , 2010 , 337, 148-56	3.1 81
17	Planarian regeneration involves distinct stem cell responses to wounds and tissue absence. <i>Developmental Biology</i> , 2010 , 344, 979-91	3.1 227
16	The Mi-2-like Smed-CHD4 gene is required for stem cell differentiation in the planarian <i>Schmidtea mediterranea</i> . <i>Development (Cambridge)</i> , 2010 , 137, 1231-41	6.6 77
15	A wound-induced Wnt expression program controls planarian regeneration polarity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 17061-6	11.5 181

14	Wnt signaling and the polarity of the primary body axis. <i>Cell</i> , 2009 , 139, 1056-68	56.2	354
13	Smed-betacatenin-1 is required for anteroposterior blastema polarity in planarian regeneration. <i>Science</i> , 2008 , 319, 327-30	33.3	284
12	Gene nomenclature guidelines for the planarian <i>Schmidtea mediterranea</i> . <i>Developmental Dynamics</i> , 2008 , 237, 3099-101	2.9	20
11	BMP signaling regulates the dorsal planarian midline and is needed for asymmetric regeneration. <i>Development (Cambridge)</i> , 2007 , 134, 4043-51	6.6	135
10	DPL-1 DP, LIN-35 Rb and EFL-1 E2F act with the MCD-1 zinc-finger protein to promote programmed cell death in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2007 , 175, 1719-33	4	29
9	Identification of genes needed for regeneration, stem cell function, and tissue homeostasis by systematic gene perturbation in planaria. <i>Developmental Cell</i> , 2005 , 8, 635-49	10.2	337
8	SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. <i>Science</i> , 2005 , 310, 1327-30	33.3	463
7	Fundamentals of planarian regeneration. <i>Annual Review of Cell and Developmental Biology</i> , 2004 , 20, 725-57	12.6	659
6	The engulfment process of programmed cell death in <i>caenorhabditis elegans</i> . <i>Annual Review of Cell and Developmental Biology</i> , 2004 , 20, 193-221	12.6	197
5	Ingestion of bacterially expressed double-stranded RNA inhibits gene expression in planarians. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100 Suppl 1, 11861-5	11.5	226
4	Phagocytosis promotes programmed cell death in <i>C. elegans</i> . <i>Nature</i> , 2001 , 412, 198-202	50.4	284
3	CED-2/Crkl and CED-10/Rac control phagocytosis and cell migration in <i>Caenorhabditis elegans</i> . <i>Nature Cell Biology</i> , 2000 , 2, 131-6	23.4	353
2	Mutational analysis of the <i>Caenorhabditis elegans</i> cell-death gene <i>ced-3</i> . <i>Genetics</i> , 1999 , 153, 1655-71	4	71
1	m6A is required for resolving progenitor identity during planarian stem cell differentiation		1