

# Viktoria Zeisler-Diehl

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/981496/publications.pdf>

Version: 2024-02-01

20  
papers

741  
citations

687363

13  
h-index

752698

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

1026  
citing authors

#	ARTICLE	IF	CITATIONS
1	Interaction of surfactants with barley leaf surfaces: time-dependent recovery of contact angles is due to foliar uptake of surfactants. <i>Planta</i> , 2022, 255, 1.	3.2	23
2	Increased cuticular wax deposition does not change residual foliar transpiration. <i>Plant, Cell and Environment</i> , 2022, 45, 1157-1171.	5.7	13
3	Alcohol Ethoxylates Enhancing the Cuticular Uptake of Lipophilic Epoxiconazole Do Not Increase the Rates of Cuticular Transpiration of Leaf and Fruit Cuticles. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 777-784.	5.2	2
4	Non-Coding RNA Analyses of Seasonal Cambium Activity in <i>Populus tomentosa</i> . <i>Cells</i> , 2022, 11, 640.	4.1	10
5	CESTâ€2 overexpression alters lipid metabolism and extends longevity of mitochondrial mutants. <i>EMBO Reports</i> , 2022, 23, e52606.	4.5	5
6	Analysis of Extracellular Cell Wall Lipids: Wax, Cutin, and Suberin in Leaves, Roots, Fruits, and Seeds. <i>Methods in Molecular Biology</i> , 2021, 2295, 275-293.	0.9	9
7	Asymmetric water transport in dense leaf cuticles and cuticle-inspired compositionally graded membranes. <i>Nature Communications</i> , 2021, 12, 1267.	12.8	19
8	ABCG transporters export cutin precursors for the formation of the plant cuticle. <i>Current Biology</i> , 2021, 31, 2111-2123.e9.	3.9	28
9	Abscisic acid applied to sweet cherry at fruit set increases amounts of cell wall and cuticular wax components at the ripe stage. <i>Scientia Horticulturae</i> , 2021, 283, 110097.	3.6	15
10	Russetting in Apple is Initiated after Exposure to Moisture Ends: Molecular and Biochemical Evidence. <i>Plants</i> , 2021, 10, 65.	3.5	16
11	Seminal roots of wild and cultivated barley differentially respond to osmotic stress in gene expression, suberization, and hydraulic conductivity. <i>Plant, Cell and Environment</i> , 2020, 43, 344-357.	5.7	39
12	Osmotic stress enhances suberization of apoplastic barriers in barley seminal roots: analysis of chemical, transcriptomic and physiological responses. <i>New Phytologist</i> , 2019, 221, 180-194.	7.3	89
13	Epicuticular wax on leaf cuticles does not establish the transpiration barrier, which is essentially formed by intracuticular wax. <i>Journal of Plant Physiology</i> , 2018, 227, 66-74.	3.5	72
14	Root cortical senescence decreases root respiration, nutrient content and radial water and nutrient transport in barley. <i>Plant, Cell and Environment</i> , 2017, 40, 1392-1408.	5.7	79
15	The composite water and solute transport of barley ( <i>Hordeum vulgare</i> ) roots: effect of suberized barriers. <i>Annals of Botany</i> , 2017, 119, mcw252.	2.9	32
16	Overexpression of the Novel Arabidopsis Gene At5g02890 Alters Inflorescence Stem Wax Composition and Affects Phytohormone Homeostasis. <i>Frontiers in Plant Science</i> , 2017, 8, 68.	3.6	13
17	Epicuticular wax on cherry laurel ( <i>Prunus laurocerasus</i> ) leaves does not constitute the cuticular transpiration barrier. <i>Planta</i> , 2016, 243, 65-81.	3.2	59
18	Wax and cutin mutants of Arabidopsis: Quantitative characterization of the cuticular transport barrier in relation to chemical composition. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 1336-1344.	2.4	22

#	ARTICLE	IF	CITATIONS
19	FAX1, a Novel Membrane Protein Mediating Plastid Fatty Acid Export. PLoS Biology, 2015, 13, e1002053.	5.6	162
20	Association between the concentration of n -alkanes and tolerance to cracking in commercial varieties of sweet cherry fruits. Scientia Horticulturae, 2015, 197, 57-65.	3.6	34