Glyn H Lewis

List of Publications by Citations

Source: https://exaly.com/author-pdf/98499/glyn-h-lewis-publications-by-citations.pdf
Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

703 papers	40,546 citations	97 h-index	174 g-index
758 ext. papers	48,497 ext. citations	7 avg, IF	7.41 L-index

#	Paper	IF	Citations
703	Cannabis use and risk of psychotic or affective mental health outcomes: a systematic review. *Lancet, The*, **2007**, 370, 319-28	40	1554
702	Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychological Medicine*, **1992**, 22, 465-86	6.9	1362
701	Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nature Genetics*, **2018**, 50, 668-681	36.3	1301
700	Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry,the*, **2020**, 7, 611-627	23.3	1004
699	A mega-analysis of genome-wide association studies for major depressive disorder. *Molecular Psychiatry*, **2013**, 18, 497-511	15.1	853
698	Identification of common genetic risk variants for autism spectrum disorder. *Nature Genetics*, **2019**, 51, 431-444	36.3	746
697	Self reported cannabis use as a risk factor for schizophrenia in Swedish conscripts of 1969: historical cohort study. *BMJ, The*, **2002**, 325, 1199	5.9	543
696	Association of serum interleukin 6 and C-reactive protein in childhood with depression and psychosis in young adult life: a population-based longitudinal study. *JAMA Psychiatry*, **2014**, 71, 1121-8	14.5	476
695	Gene X environment interactions at the serotonin transporter locus. *Biological Psychiatry*, **2009**, 65, 211-9	7.9	475
694	Psychosis, victimisation and childhood disadvantage: evidence from the second British National Survey of Psychiatric Morbidity. *British Journal of Psychiatry*, **2004**, 185, 220-6	5.4	414
693	Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. *Cell*, **2019**, 179, 1469-1482.e11	56.2	402
692	Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psychopharmacology guidelines. *Journal of Psychopharmacology*, **2015**, 29, 459-525	4.6	381
691	Maternal depression during pregnancy and the postnatal period: risks and possible mechanisms for offspring depression at age 18 years. *JAMA Psychiatry*, **2013**, 70, 1312-9	14.5	371
690	A longitudinal study of premorbid IQ Score and risk of developing schizophrenia, bipolar disorder, severe depression, and other nonaffective psychoses. *Archives of General Psychiatry*, **2004**, 61, 354-60		360
689	Evidence-based guidelines for treating depressive disorders with antidepressants: a revision of the 2000 British Association for Psychopharmacology guidelines. *Journal of Psychopharmacology*, **2008**, 22, 343-96	4.6	341
688	IQ and risk for schizophrenia: a population-based cohort study. *Psychological Medicine*, **1997**, 27, 1311-23	6.9	316
687	Personality disorder: the patients psychiatrists dislike. *British Journal of Psychiatry*, **1988**, 153, 44-9	5.4	311

686	Antidepressant activity of anti-cytokine treatment: a systematic review and meta-analysis of clinical trials of chronic inflammatory conditions. *Molecular Psychiatry*, **2018**, 23, 335-343	15.1	310
685	The epidemiology of fatigue: more questions than answers. *Journal of Epidemiology and Community Health*, **1992**, 46, 92-7	5.1	310
684	Schizophrenia and city life. *Lancet, The*, **1992**, 340, 137-40	40	300
683	Poverty, unemployment, and common mental disorders: population based cohort study. *BMJ: British Medical Journal*, **1998**, 317, 115-9		295
682	Prenatal maternal infection, neurodevelopment and adult schizophrenia: a systematic review of population-based studies. *Psychological Medicine*, **2013**, 43, 239-57	6.9	284
681	Effectiveness and cost-effectiveness of mindfulness-based cognitive therapy compared with maintenance antidepressant treatment in the prevention of depressive relapse or recurrence (PREVENT): a randomised controlled trial. *Lancet, The*, **2015**, 386, 63-73	40	278
680	Therapist-delivered Internet psychotherapy for depression in primary care: a randomised controlled trial. *Lancet, The*, **2009**, 374, 628-34	40	271
679	The National Psychiatric Morbidity surveys of Great Britain--initial findings from the household survey. *Psychological Medicine*, **1997**, 27, 775-89	6.9	263
678	Prospective study of peer victimization in childhood and psychotic symptoms in a nonclinical population at age 12 years. *Archives of General Psychiatry*, **2009**, 66, 527-36		260
677	The influence of age and sex on the prevalence of depressive conditions: report from the National Survey of Psychiatric Morbidity. *Psychological Medicine*, **1998**, 28, 9-19	6.9	252
676	Cross sectional study of symptom attribution and recognition of depression and anxiety in primary care. *BMJ: British Medical Journal*, **1999**, 318, 436-9		249
675	Parental depression, maternal antidepressant use during pregnancy, and risk of autism spectrum disorders: population based case-control study. *BMJ, The*, **2013**, 346, f2059	5.9	226
674	Suicide, deprivation, and unemployment: record linkage study. *BMJ: British Medical Journal*, **1998**, 317, 1283-6		226
673	Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: results of the CoBalT randomised controlled trial. *Lancet, The*, **2013**, 381, 375-84	40	223
672	Paternal age and schizophrenia: a population based cohort study. *BMJ, The*, **2004**, 329, 1070	5.9	218
671	Developmental precursors of affective illness in a general population birth cohort. *Archives of General Psychiatry*, **1997**, 54, 625-31		213
670	Premorbid adjustment and personality in people with schizophrenia. *British Journal of Psychiatry*, **1998**, 172, 308-13; discussion 314-5	5.4	213
669	Nicotine, alcohol and drug dependence and psychiatric comorbidity. Results of a national household survey. *British Journal of Psychiatry*, **2001**, 179, 432-7	5.4	212

668	Violence against women by their intimate partner during pregnancy and postnatal depression: a prospective cohort study. *Lancet, The*, **2010**, 376, 903-10	40	211
667	Psychotic experiences and psychotic disorders at age 18 in relation to psychotic experiences at age 12 in a longitudinal population-based cohort study. *American Journal of Psychiatry*, **2013**, 170, 742-50	11.9	206
666	Obsessive-compulsive disorder: prevalence, comorbidity, impact, and help-seeking in the British National Psychiatric Morbidity Survey of 2000. *American Journal of Psychiatry*, **2006**, 163, 1978-85	11.9	202
665	Social adversity in childhood and the risk of developing psychosis: a national cohort study. *American Journal of Psychiatry*, **2005**, 162, 1652-7	11.9	200
664	Effects of cannabis use on outcomes of psychotic disorders: systematic review. *British Journal of Psychiatry*, **2008**, 193, 357-63	5.4	194
663	Self-reported psychotic symptoms in the general population: results from the longitudinal study of the British National Psychiatric Morbidity Survey. *British Journal of Psychiatry*, **2006**, 188, 519-26	5.4	191
662	Epilepsy and psychiatric comorbidity: a nationally representative population-based study. *Epilepsia*, **2012**, 53, 1095-103	6.4	189
661	IQ and non-clinical psychotic symptoms in 12-year-olds: results from the ALSPAC birth cohort. *British Journal of Psychiatry*, **2008**, 193, 185-91	5.4	187
660	The patient-doctor relationship: a synthesis of the qualitative literature on patients' perspectives. *British Journal of General Practice*, **2009**, 59, e116-33	1.6	180
659	Pharmacological and psychotherapeutic interventions for management of obsessive-compulsive disorder in adults: a systematic review and network meta-analysis. *Lancet Psychiatry,the*, **2016**, 3, 730-739	33.3	176
658	Unequal access and unmet need: neurotic disorders and the use of primary care services. *Psychological Medicine*, **2000**, 30, 1359-67	6.9	169
657	Infections in the CNS during childhood and the risk of subsequent psychotic illness: a cohort study of more than one million Swedish subjects. *American Journal of Psychiatry*, **2008**, 165, 59-65	11.9	168
656	Assessment of Bidirectional Relationships Between Physical Activity and Depression Among Adults: A 2-Sample Mendelian Randomization Study. *JAMA Psychiatry*, **2019**, 76, 399-408	14.5	165
655	Prevalence of low-grade inflammation in depression: a systematic review and meta-analysis of CRP levels. *Psychological Medicine*, **2019**, 49, 1958-1970	6.9	162
654	Common mental disorders in Santiago, Chile: prevalence and socio-demographic correlates. *British Journal of Psychiatry*, **2001**, 178, 228-33	5.4	161
653	Childhood infection and adult schizophrenia: a meta-analysis of population-based studies. *Schizophrenia Research*, **2012**, 139, 161-8	3.6	159
652	Minimal clinically important difference on the Beck Depression Inventory--II according to the patient's perspective. *Psychological Medicine*, **2015**, 45, 3269-79	6.9	151
651	Socioeconomic status, standard of living, and neurotic disorder. *Lancet, The*, **1998**, 352, 605-9	40	150

650	Contribution of common genetic variants to antidepressant response. *Biological Psychiatry*, **2013**, 73, 679-82	7.9	146
649	Phenotypic Manifestation of Genetic Risk for Schizophrenia During Adolescence in the General Population. *JAMA Psychiatry*, **2016**, 73, 221-8	14.5	145
648	The Shona Symptom Questionnaire: the development of an indigenous measure of common mental disorders in Harare. *Acta Psychiatrica Scandinavica*, **1997**, 95, 469-75	6.5	144
647	Aftercare and clinical characteristics of people with mental illness who commit suicide: a case-control study. *Lancet, The*, **1999**, 353, 1397-400	40	143
646	Perceptions of social capital and the built environment and mental health. *Social Science and Medicine*, **2006**, 62, 3072-83	5.1	138
645	Factors influencing the development and amelioration of suicidal thoughts in the general population. Cohort study. *British Journal of Psychiatry*, **2004**, 185, 385-93	5.4	138
644	Clinical and social outcomes of adolescent self harm: population based birth cohort study. *BMJ, The* , **2014**, 349, g5954	5.9	137
643	Parental socioeconomic status and risk of offspring autism spectrum disorders in a Swedish population-based study. *Journal of the American Academy of Child and Adolescent Psychiatry*, **2012**, 51, 467-476.e6	7.2	136
642	Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. *BMJ, The*, **2013**, 347, f4913	5.9	136
641	Psychiatric and social aspects of suicidal behaviour in prisons. *Psychological Medicine*, **2005**, 35, 257-69	6.9	133
640	Signs of asphyxia at birth and risk of schizophrenia. Population-based case-control study. *British Journal of Psychiatry*, **2001**, 179, 403-8	5.4	132
639	Randomised controlled trials investigating pharmacological and psychological interventions for treatment-refractory depression. Systematic review. *British Journal of Psychiatry*, **2002**, 181, 284-94	5.4	132
638	Pathways between childhood victimization and psychosis-like symptoms in the ALSPAC birth cohort. *Schizophrenia Bulletin*, **2013**, 39, 1045-55	1.3	130
637	Strategies for preventing suicide. *British Journal of Psychiatry*, **1997**, 171, 351-354	5.4	129
636	Paternal age and risk for schizophrenia. *British Journal of Psychiatry*, **2003**, 183, 405-8	5.4	128
635	Adolescent alcohol and tobacco use and early socioeconomic position: the ALSPAC birth cohort. *Pediatrics*, **2011**, 127, e948-55	7.4	125
634	Substance misuse and psychiatric comorbidity: an overview of the OPCS National Psychiatric Morbidity Survey. *Addictive Behaviors*, **1998**, 23, 909-18	4.2	125
633	Income inequality and the prevalence of common mental disorders in Britain. *British Journal of Psychiatry*, **2001**, 178, 222-7	5.4	125

632	Are British psychiatrists racist?. *British Journal of Psychiatry*, **1990**, 157, 410-5	5.4	125
631	Treatment of postnatal depression in low-income mothers in primary-care clinics in Santiago, Chile: a randomised controlled trial. *Lancet, The*, **2007**, 370, 1629-37	40	124
630	Detection of depression and anxiety in primary care: follow up study. *BMJ, The*, **2002**, 325, 1016-7	5.9	124
629	Refugee migration and risk of schizophrenia and other non-affective psychoses: cohort study of 1.3 million people in Sweden. *BMJ, The*, **2016**, 352, i1030	5.9	123
628	Socio-economic position and common mental disorders. Longitudinal study in the general population in the UK. *British Journal of Psychiatry*, **2006**, 189, 109-17	5.4	123
627	Pragmatic randomised controlled trials in psychiatry. *British Journal of Psychiatry*, **1999**, 175, 217-23	5.4	123
626	Factors affecting survival in Alzheimer's disease. *Psychological Medicine*, **1991**, 21, 363-70	6.9	123
625	Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. *BMJ, The*, **2012**, 344, e2758	5.9	122
624	Education and income: which is more important for mental health?. *Journal of Epidemiology and Community Health*, **2003**, 57, 501-5	5.1	121
623	Material standard of living, social class, and the prevalence of the common mental disorders in Great Britain. *Journal of Epidemiology and Community Health*, **1998**, 52, 8-14	5.1	121
622	Neurotic disorders and the receipt of psychiatric treatment. *Psychological Medicine*, **2000**, 30, 1369-76	6.9	120
621	The Aberfan disaster: 33-year follow-up of survivors. *British Journal of Psychiatry*, **2003**, 182, 532-6	5.4	119
620	Discontinuation rates of SSRIs and tricyclic antidepressants: a meta-analysis and investigation of heterogeneity. *British Journal of Psychiatry*, **1997**, 170, 120-7	5.4	112
619	Comparison of the General Health Questionnaire and the Hospital Anxiety and Depression Scale. *British Journal of Psychiatry*, **1990**, 157, 860-4	5.4	110
618	Genetic Association of Major Depression With Atypical Features and Obesity-Related Immunometabolic Dysregulations. *JAMA Psychiatry*, **2017**, 74, 1214-1225	14.5	109
617	The influence of age and sex on the prevalence of depressive conditions: report from the National Survey of Psychiatric Morbidity. *International Review of Psychiatry*, **2003**, 15, 74-83	3.6	107
616	Psychosis in the community and in prisons: a report from the British National Survey of psychiatric morbidity. *American Journal of Psychiatry*, **2005**, 162, 774-80	11.9	107
615	Patterns of alcohol use and multiple risk behaviour by gender during early and late adolescence: the ALSPAC cohort. *Journal of Public Health*, **2012**, 34 Suppl 1, i20-30	3.5	106

614	Resuscitation at birth and cognition at 8 years of age: a cohort study. *Lancet, The*, **2009**, 373, 1615-22	40	106
613	The development of a computerized assessment for minor psychiatric disorder. *Psychological Medicine*, **1988**, 18, 737-45	6.9	105
612	Tolerance of suicide, religion and suicide rates: an ecological and individual study in 19 Western countries. *Psychological Medicine*, **1997**, 27, 1165-71	6.9	103
611	Individuals, schools, and neighborhood: a multilevel longitudinal study of variation in incidence of psychotic disorders. *Archives of General Psychiatry*, **2010**, 67, 914-22		102
610	Birth weight and psychological distress at age 45-51 years: results from the Aberdeen Children of the 1950s cohort study. *British Journal of Psychiatry*, **2005**, 187, 21-8	5.4	102
609	Assessing psychiatric disorder with a human interviewer or a computer. *Journal of Epidemiology and Community Health*, **1994**, 48, 207-10	5.1	102
608	Cannabis, COMT and psychotic experiences. *British Journal of Psychiatry*, **2011**, 199, 380-5	5.4	100
607	Investigating the association between cigarette smoking and schizophrenia in a cohort study. *American Journal of Psychiatry*, **2003**, 160, 2216-21	11.9	100
606	Sibling bullying and risk of depression, anxiety, and self-harm: a prospective cohort study. *Pediatrics* , **2014**, 134, e1032-9	7.4	97
605	Comparison of two self administered psychiatric questionnaires (GHQ-12 and SRQ-20) in primary care in Chile. *Social Psychiatry and Psychiatric Epidemiology*, **1992**, 27, 168-73	4.5	97
604	Mood, anxiety and psychotic phenomena measure a common psychopathological factor. *Psychological Medicine*, **2015**, 45, 1483-93	6.9	96
603	Peer victimization during adolescence and risk for anxiety disorders in adulthood: a prospective cohort study. *Depression and Anxiety*, **2014**, 31, 574-82	8.4	96
602	Psychopathy among prisoners in England and Wales. *International Journal of Law and Psychiatry*, **2009**, 32, 134-41	2.6	96
601	Bullying in elementary school and psychotic experiences at 18 years: a longitudinal, population-based cohort study. *Psychological Medicine*, **2014**, 44, 2199-211	6.9	95
600	Violence and psychiatric morbidity in a national household population--a report from the British Household Survey. *American Journal of Epidemiology*, **2006**, 164, 1199-208	3.8	95
599	Violence and psychiatric morbidity in the national household population of Britain: public health implications. *British Journal of Psychiatry*, **2006**, 189, 12-9	5.4	95
598	Genetic predictors of response to serotonergic and noradrenergic antidepressants in major depressive disorder: a genome-wide analysis of individual-level data and a meta-analysis. *PLoS Medicine*, **2012**, 9, e1001326	11.6	94
597	Trends in service use and treatment for mental disorders in adults throughout Great Britain. *British Journal of Psychiatry*, **2004**, 185, 378-84	5.4	93

596	Psychosis and drug dependence: results from a national survey of prisoners. *British Journal of Psychiatry*, **2002**, 181, 393-8	5.4	93
595	Being bullied during childhood and the prospective pathways to self-harm in late adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, **2013**, 52, 608-18.e2	7.2	92
594	Contextual risk factors for the common mental disorders in Britain: a multilevel investigation of the effects of place. *Journal of Epidemiology and Community Health*, **2003**, 57, 616-21	5.1	90
593	Drug use and initiation in prison: results from a national prison survey in England and Wales. *Addiction*, **2002**, 97, 1551-60	4.6	90
592	The National Psychiatric Morbidity surveys of Great Britain--strategy and methods. *Psychological Medicine*, **1997**, 27, 765-74	6.9	89
591	Heritability of social cognitive skills in children and adolescents. *British Journal of Psychiatry*, **1999**, 175, 559-64	5.4	88
590	Are SSRIs a cost-effective alternative to tricyclics?. *British Journal of Psychiatry*, **1996**, 168, 404-9	5.4	88
589	Marriage and risk of dementia: systematic review and meta-analysis of observational studies. *Journal of Neurology, Neurosurgery and Psychiatry*, **2018**, 89, 231-238	5.5	88
588	Investigating whether adverse prenatal and perinatal events are associated with non-clinical psychotic symptoms at age 12 years in the ALSPAC birth cohort. *Psychological Medicine*, **2009**, 39, 1457-67	6.9	87
587	Is socioeconomic status in early life associated with drug use? A systematic review of the evidence. *Drug and Alcohol Review*, **2009**, 28, 142-53	3.2	86
586	Rural/non-rural differences in rates of common mental disorders in Britain: prospective multilevel cohort study. *British Journal of Psychiatry*, **2006**, 188, 51-7	5.4	86
585	Overcoming the effects of false positives and threshold bias in graph theoretical analyses of neuroimaging data. *NeuroImage*, **2015**, 118, 313-33	7.9	85
584	Validity of the Short Mood and Feelings Questionnaire in late adolescence. *Psychological Assessment*, **2014**, 26, 752-62	5.3	85
583	Lone mothers, social exclusion and depression. *Psychological Medicine*, **2003**, 33, 715-22	6.9	83
582	Prevalence, psychosocial correlates and service utilization of depressive and anxiety disorders in Hong Kong: the Hong Kong Mental Morbidity Survey (HKMMS). *Social Psychiatry and Psychiatric Epidemiology*, **2015**, 50, 1379-88	4.5	82
581	'Junk food' diet and childhood behavioural problems: results from the ALSPAC cohort. *European Journal of Clinical Nutrition*, **2009**, 63, 491-8	5.2	82
580	Fetal growth and childhood behavioral problems: results from the ALSPAC cohort. *American Journal of Epidemiology*, **2006**, 163, 829-37	3.8	82
579	Differences in risk factors for self-harm with and without suicidal intent: findings from the ALSPAC cohort. *Journal of Affective Disorders*, **2014**, 168, 407-14	6.6	81

578	Characterisation of zuclopenthixol metabolism by in vitro and therapeutic drug monitoring studies. *Acta Psychiatrica Scandinavica*, **2010**, 122, 444-53	6.5	81
577	Negative self-schemas and the onset of depression in women: longitudinal study. *British Journal of Psychiatry*, **2005**, 186, 302-7	5.4	81
576	Minimal phenotyping yields genome-wide association signals of low specificity for major depression. *Nature Genetics*, **2020**, 52, 437-447	36.3	80
575	Adolescent self-harm and suicidal thoughts in the ALSPAC cohort: a self-report survey in England. *BMC Psychiatry*, **2012**, 12, 69	4.2	80
574	Prevalence of treatment-resistant depression in primary care: cross-sectional data. *British Journal of General Practice*, **2013**, 63, e852-8	1.6	80
573	Course of Cognitive Development From Infancy to Early Adulthood in the Psychosis Spectrum. *JAMA Psychiatry*, **2018**, 75, 270-279	14.5	79
572	Genetic Differences in the Immediate Transcriptome Response to Stress Predict Risk-Related Brain Function and Psychiatric Disorders. *Neuron*, **2015**, 86, 1189-202	13.9	79
571	Cost-effectiveness of therapist-delivered online cognitive-behavioural therapy for depression: randomised controlled trial. *British Journal of Psychiatry*, **2010**, 197, 297-304	5.4	78
570	Classroom based cognitive behavioural therapy in reducing symptoms of depression in high risk adolescents: pragmatic cluster randomised controlled trial. *BMJ, The*, **2012**, 345, e6058	5.9	78
569	Reboxetine versus other antidepressive agents for depression. *The Cochrane Library*,	5.2	78
568	Cognitive behavioural therapies versus other psychological therapies for depression. *The Cochrane Library*, **2018**,	5.2	78
567	Cognitive behavioural therapies versus treatment as usual for depression. *The Cochrane Library*, **2018**,	5.2	78
566	Peer victimisation during adolescence and its impact on depression in early adulthood: prospective cohort study in the United Kingdom. *BMJ, The*, **2015**, 350, h2469	5.9	76
565	Geographic variation in the prevalence of common mental disorders in Britain: a multilevel investigation. *American Journal of Epidemiology*, **2003**, 157, 730-7	3.8	76
564	Timing of menarche and depressive symptoms in adolescent girls from a UK cohort. *British Journal of Psychiatry*, **2011**, 198, 17-23, sup 1-2	5.4	75
563	A cohort study of low Apgar scores and cognitive outcomes. *Archives of Disease in Childhood: Fetal and Neonatal Edition*, **2008**, 93, F115-20	4.7	75
562	Temporal relations between unexplained fatigue and depression: longitudinal data from an international study in primary care. *Psychosomatic Medicine*, **2004**, 66, 330-5	3.7	75
561	Suicide acceptability in African- and white Americans: the role of religion. *Journal of Nervous and Mental Disease*, **1998**, 186, 12-6	1.8	75

560	Putting trials on trial--the costs and consequences of small trials in depression: a systematic review of methodology. *Journal of Epidemiology and Community Health*, **1997**, 51, 354-8	5.1	73
559	Stigma revisited, disclosure of emotional problems in primary care consultations in Wales. *Social Science and Medicine*, **2003**, 56, 2191-200	5.1	72
558	Treatment seeking by individuals with obsessive-compulsive disorder from the british psychiatric morbidity survey of 2000. *Psychiatric Services*, **2007**, 58, 977-82	3.3	71
557	Psychiatric disorder in veterans of the Persian Gulf War of 1991: Systematic review. *British Journal of Psychiatry*, **2003**, 182, 391-403	5.4	71
556	Are there neurological and sensory risk factors for schizophrenia?. *Schizophrenia Research*, **1995**, 14, 247-51	3.1	71
555	Associations of cannabis and cigarette use with psychotic experiences at age 18: findings from the Avon Longitudinal Study of Parents and Children. *Psychological Medicine*, **2014**, 44, 3435-44	6.9	70
554	Efficacy and acceptability of selective serotonin reuptake inhibitors for the treatment of depression in Parkinson's disease: a systematic review and meta-analysis of randomized controlled trials. *BMC Neurology*, **2010**, 10, 49	3.1	70
553	Antidepressants for the treatment of depression in people with schizophrenia: a systematic review. *Psychological Medicine*, **2003**, 33, 589-99	6.9	70
552	Alcohol consumption as a risk factor for anxiety and depression: results from the longitudinal follow-up of the National Psychiatric Morbidity Survey. *British Journal of Psychiatry*, **2005**, 187, 544-51	5.4	70
551	An observation tool to assist with the assessment of urban residential environments. *Journal of Environmental Psychology*, **2005**, 25, 293-305	6.7	70
550	Long-term effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: follow-up of the CoBalT randomised controlled trial. *Lancet Psychiatry,the*, **2016**, 3, 137-44	23.3	69
549	GPs' experiences of primary care mental health research: a qualitative study of the barriers to recruitment. *Family Practice*, **2007**, 24, 518-25	1.9	69
548	An Analysis of Two Genome-wide Association Meta-analyses Identifies a New Locus for Broad Depression Phenotype. *Biological Psychiatry*, **2017**, 82, 322-329	7.9	68
547	Maternal tobacco, cannabis and alcohol use during pregnancy and risk of adolescent psychotic symptoms in offspring. *British Journal of Psychiatry*, **2009**, 195, 294-300	5.4	68
546	Psychological distress and risk of long-term disability: population-based longitudinal study. *Journal of Epidemiology and Community Health*, **2012**, 66, 586-92	5.1	66
545	Mental health inequalities in Wales, UK: multi-level investigation of the effect of area deprivation. *British Journal of Psychiatry*, **2005**, 186, 417-22	5.4	66
544	Socio-economic status in childhood and later alcohol use: a systematic review. *Addiction*, **2007**, 102, 1546-63	4.3	65
543	Clarifying the relationship between unexplained chronic fatigue and psychiatric morbidity: results from a community survey in Great Britain. *American Journal of Psychiatry*, **2000**, 157, 1492-8	11.9	65

542	Association between a functional interleukin 6 receptor genetic variant and risk of depression and psychosis in a population-based birth cohort. *Brain, Behavior, and Immunity*, **2018**, 69, 264-272	16.6	65
541	The reluctance to seek treatment for neurotic disorders. *International Review of Psychiatry*, **2003**, 15, 123-8	3.6	64
540	'Third wave' cognitive and behavioural therapies versus other psychological therapies for depression. *The Cochrane Library*, **2013**, CD008704	5.2	63
539	Cannabis and suicide: longitudinal study. *British Journal of Psychiatry*, **2009**, 195, 492-7	5.4	62
538	Suicide, religion, and socioeconomic conditions. An ecological study in 26 countries, 1990. *Journal of Epidemiology and Community Health*, **1999**, 53, 204-10	5.1	62
537	A systematic review of the clinical effectiveness and cost-effectiveness of pharmacological and psychological interventions for the management of obsessive-compulsive disorder in children/adolescents and adults. *Health Technology Assessment*, **2016**, 20, 1-392	4.4	62
536	The association between early autistic traits and psychotic experiences in adolescence. *Schizophrenia Research*, **2012**, 135, 164-9	3.6	61
535	Patients' views of physical activity as treatment for depression: a qualitative study. *British Journal of General Practice*, **2011**, 61, 149-56	1.6	61
534	Computerized assessment of common mental disorders in primary care: effect on clinical outcome. *Family Practice*, **1996**, 13, 120-6	1.9	61
533	A qualitative study exploring how GPs decide to prescribe antidepressants. *British Journal of General Practice*, **2005**, 55, 755-62	1.6	61
532	Assessing mental health in primary care research using standardized scales: can it be carried out over the telephone?. *Psychological Medicine*, **2004**, 34, 157-62	6.9	60
531	A population-based study of genetic variation and psychotic experiences in adolescents. *Schizophrenia Bulletin*, **2014**, 40, 1254-62	1.3	59
530	The prevalence and costs of psychiatric disorders and learning disabilities. *British Journal of Psychiatry*, **1995**, 166, 9-18	5.4	59
529	A pragmatic randomised controlled trial to compare antidepressants with a community-based psychosocial intervention for the treatment of women with postnatal depression: the RESPOND trial. *Health Technology Assessment*, **2010**, 14, iii-iv, ix-xi, 1-153	4.4	59
528	Polygenic scores predict alcohol problems in an independent sample and show moderation by the environment. *Genes*, **2014**, 5, 330-46	4.2	58
527	Are cities bad for your mental health?. *Psychological Medicine*, **1994**, 24, 913-5	6.9	58
526	Use of self-administered instruments to assess psychiatric disorders in older people: validity of the General Health Questionnaire, the Center for Epidemiologic Studies Depression Scale and the self-completion version of the revised Clinical Interview Schedule. *Psychological Medicine*, **2013**, 43, 2649-56	6.9	57
525	Are female children more vulnerable to the long-term effects of maternal depression during pregnancy?. *Journal of Affective Disorders*, **2016**, 189, 329-35	6.6	56

524	Paternal and maternal ages at conception and risk of bipolar affective disorder in their offspring. *Psychological Medicine*, **2010**, 40, 477-85	6.9	56
523	Geographical variation in rates of common mental disorders in Britain: prospective cohort study. *British Journal of Psychiatry*, **2005**, 187, 29-34	5.4	56
522	Association of social contact with dementia and cognition: 28-year follow-up of the Whitehall II cohort study. *PLoS Medicine*, **2019**, 16, e1002862	11.6	55
521	A longitudinal study of factors predicting change in cognitive test scores over time, in an older hypertensive population. *Psychological Medicine*, **1996**, 26, 555-68	6.9	55
520	Pharmacogenetics of antidepressant response: A polygenic approach. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, **2017**, 75, 128-134	5.5	54
519	Integrated analysis of environmental and genetic influences on cord blood DNA methylation in new-borns. *Nature Communications*, **2019**, 10, 2548	17.4	54
518	Depression in paediatric chronic fatigue syndrome. *Archives of Disease in Childhood*, **2013**, 98, 425-8	2.2	54
517	Non-psychotic psychiatric disorder and subsequent risk of schizophrenia. Cohort study. *British Journal of Psychiatry*, **2000**, 177, 416-20	5.4	54
516	Longitudinal association between inflammatory markers and specific symptoms of depression in a prospective birth cohort. *Brain, Behavior, and Immunity*, **2019**, 76, 74-81	16.6	53
515	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. *Lancet Psychiatry,the*, **2019**, 6, 903-914	23.3	52
514	Accuracy of general hospital dementia diagnoses in England: Sensitivity, specificity, and predictors of diagnostic accuracy 2008-2016. *Alzheimerks and Dementia*, **2018**, 14, 933-943	1.2	52
513	Associations of cannabis and cigarette use with depression and anxiety at age 18: findings from the Avon Longitudinal Study of Parents and Children. *PLoS ONE*, **2015**, 10, e0122896	3.7	52
512	A population-based study of atopic disorders and inflammatory markers in childhood before psychotic experiences in adolescence. *Schizophrenia Research*, **2014**, 152, 139-45	3.6	52
511	Physical activity and common mental disorder: results from the Caerphilly study. *American Journal of Epidemiology*, **2007**, 165, 946-54	3.8	52
510	The National Psychiatric Morbidity Surveys of Great Britain--initial findings from the household survey. *International Review of Psychiatry*, **2003**, 15, 29-42	3.6	52
509	Regional differences in mental health in Great Britain. *Journal of Epidemiology and Community Health*, **1992**, 46, 608-11	5.1	52
508	Childhood Epstein-Barr Virus infection and subsequent risk of psychotic experiences in adolescence: a population-based prospective serological study. *Schizophrenia Research*, **2014**, 158, 19-24	3.6	51
507	Polymorphism of the 5-HT transporter and response to antidepressants: randomised controlled trial. *British Journal of Psychiatry*, **2011**, 198, 464-71	5.4	51

506	40,000 memories in young teenagers: psychometric properties of the Autobiographical Memory Test in a UK cohort study. *Memory*, **2012**, 20, 300-20	1.8	51
505	Unexplained fatigue syndromes in a multinational primary care sample: specificity of definition and prevalence and distinctiveness from depression and generalized anxiety. *American Journal of Psychiatry*, **2003**, 160, 785-7	11.9	51
504	Income inequality and self rated health in Britain. *Journal of Epidemiology and Community Health*, **2002**, 56, 436-41	5.1	51
503	Religious identity and comfort beliefs in three groups of psychiatric patients and a group of medical controls. *International Journal of Social Psychiatry*, **1994**, 40, 124-34	8.5	51
502	Clinical effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: the CoBalT randomised controlled trial. *Health Technology Assessment*, **2014**, 18, 1-167, vii-viii	4.4	51
501	The association between loneliness and depressive symptoms among adults aged 50 years and older: a 12-year population-based cohort study. *Lancet Psychiatry,the*, **2021**, 8, 48-57	23.3	51
500	Schizophrenia-like topological changes in the structural connectome of individuals with subclinical psychotic experiences. *Human Brain Mapping*, **2015**, 36, 2629-43	5.9	50
499	Antidepressants during pregnancy and autism in offspring: population based cohort study. *BMJ, The*, **2017**, 358, j2811	5.9	50
498	Outcomes of childhood conduct problem trajectories in early adulthood: findings from the ALSPAC study. *European Child and Adolescent Psychiatry*, **2014**, 23, 539-49	5.5	50
497	Effects of stressful life events, maternal depression and 5-HTTLPR genotype on emotional symptoms in pre-adolescent children. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **2009**, 150B, 670-82	3.5	50
496	Common mental disorders and the built environment in Santiago, Chile. *British Journal of Psychiatry* , **2007**, 190, 394-401	5.4	50
495	Social roles and gender difference in the prevalence of common mental disorders. *British Journal of Psychiatry*, **1998**, 173, 489-93	5.4	50
494	Association between pubertal development and depressive symptoms in girls from a UK cohort. *Psychological Medicine*, **2012**, 42, 2579-89	6.9	49
493	Patterns of alcohol use in early adolescence predict problem use at age 16. *Alcohol and Alcoholism*, **2012**, 47, 169-77	3.5	49
492	Effects of emotion perception training on mood in undergraduate students: randomised controlled trial. *British Journal of Psychiatry*, **2012**, 201, 71-2	5.4	49
491	Examining interactions between risk factors for psychosis. *British Journal of Psychiatry*, **2010**, 197, 207-11	5.4	49
490	Investigating if psychosis-like symptoms (PLIKS) are associated with family history of schizophrenia or paternal age in the ALSPAC birth cohort. *Schizophrenia Research*, **2008**, 104, 279-86	3.6	49
489	Physical activity and emotional problems amongst adolescents : a longitudinal study. *Social Psychiatry and Psychiatric Epidemiology*, **2008**, 43, 765-72	4.5	49

488	Does Childhood Trauma Moderate Polygenic Risk for Depression? A Meta-analysis of 5765 Subjects From the Psychiatric Genomics Consortium. *Biological Psychiatry*, **2018**, 84, 138-147	7.9	48
487	Developmental pathways from childhood conduct problems to early adult depression: findings from the ALSPAC cohort. *British Journal of Psychiatry*, **2014**, 205, 17-23	5.4	48
486	Childhood parasomnias and psychotic experiences at age 12 years in a United Kingdom birth cohort. *Sleep*, **2014**, 37, 475-82	1.1	48
485	Physical activity and depression in adolescents: cross-sectional findings from the ALSPAC cohort. *Social Psychiatry and Psychiatric Epidemiology*, **2012**, 47, 1023-33	4.5	48
484	The association of serum 25-hydroxyvitamin D3 and D2 with depressive symptoms in childhood--a prospective cohort study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, **2012**, 53, 757-66	7.9	48
483	Effectiveness of an integrated telehealth service for patients with depression: a pragmatic randomised controlled trial of a complex intervention. *Lancet Psychiatry,the*, **2016**, 3, 515-25	23.3	48
482	Genome-wide gene-environment analyses of major depressive disorder and reported lifetime traumatic experiences in UK Biobank. *Molecular Psychiatry*, **2020**, 25, 1430-1446	15.1	47
481	The association between depressive symptoms from early to late adolescence and later use and harmful use of alcohol. *European Child and Adolescent Psychiatry*, **2014**, 23, 1219-30	5.5	46
480	Impairments in cognition across the spectrum of psychiatric disorders: evidence from a Swedish conscript cohort. *Schizophrenia Bulletin*, **2008**, 34, 1035-41	1.3	46
479	Obsessive-compulsive disorder and personality disorder: evidence from the British National Survey of Psychiatric Morbidity 2000. *Social Psychiatry and Psychiatric Epidemiology*, **2006**, 41, 862-7	4.5	46
478	Somatic presentation of psychiatric morbidity in general practice. *British Journal of General Practice*, **1995**, 45, 143-7	1.6	46
477	A Population-Based Cohort Study Examining the Incidence and Impact of Psychotic Experiences From Childhood to Adulthood, and Prediction of Psychotic Disorder. *American Journal of Psychiatry*, **2020**, 177, 308-317	11.9	46
476	Using Data Linkage to Investigate Inconsistent Reporting of Self-Harm and Questionnaire Non-Response. *Archives of Suicide Research*, **2016**, 20, 113-41	2.3	45
475	Adverse effects from antidepressant treatment: randomised controlled trial of 601 depressed individuals. *Psychopharmacology*, **2014**, 231, 2921-31	4.7	45
474	TElehealth in CHronic disease: mixed-methods study to develop the TECH conceptual model for intervention design and evaluation. *BMJ Open*, **2015**, 5, e006448	3	45
473	Primary group size, social support, gender and future mental health status in a prospective study of people living in private households throughout Great Britain. *Psychological Medicine*, **2005**, 35, 705-14	6.9	45
472	The comparison of latent variable models of non-psychotic psychiatric morbidity in four culturally diverse populations. *Psychological Medicine*, **1998**, 28, 145-52	6.9	45
471	Factors associated with differential response to online cognitive behavioural therapy. *Social Psychiatry and Psychiatric Epidemiology*, **2012**, 47, 827-33	4.5	44

470	Common mental disorders, subthreshold symptoms and disability: longitudinal study. *British Journal of Psychiatry*, **2010**, 197, 411-2	5.4	44
469	Informal work and common mental disorders. *Social Psychiatry and Psychiatric Epidemiology*, **2003**, 38, 485-9	4.5	44
468	A pragmatic randomised controlled trial to evaluate the cost-effectiveness of a physical activity intervention as a treatment for depression: the treating depression with physical activity (TREAD) trial. *Health Technology Assessment*, **2012**, 16, 1-164, iii-iv	4.4	44
467	Longitudinal association between different levels of alcohol consumption and a new onset of depression and generalized anxiety disorder: Results from an international study in primary care. *Psychiatry Research*, **2016**, 243, 30-4	9.9	43
466	Childhood conduct disorder trajectories, prior risk factors and cannabis use at age 16: birth cohort study. *Addiction*, **2013**, 108, 2129-38	4.6	43
465	Effect of cytochrome CYP2C19 metabolizing activity on antidepressant response and side effects: Meta-analysis of data from genome-wide association studies. *European Neuropsychopharmacology*, **2018**, 28, 945-954	1.2	43
464	Association of Late Adolescent Personality With Risk for Subsequent Serious Mental Illness Among Men in a Swedish Nationwide Cohort Study. *JAMA Psychiatry*, **2017**, 74, 703-711	14.5	42
463	The association of fetal and childhood growth with risk of schizophrenia. Cohort study of 720,000 Swedish men and women. *Schizophrenia Research*, **2005**, 79, 315-22	3.6	42
462	Social support networks and type of neurotic symptom among adults in British households. *Psychological Medicine*, **2003**, 33, 307-18	6.9	42
461	A cross-sectional study of the prevalence of psychopathology in adults with congenital heart disease. *Journal of Psychosomatic Research*, **2002**, 52, 65-8	4.1	42
460	Selective serotonin reuptake inhibitors versus tricyclic and heterocyclic antidepressants: comparison of drug adherence. *Cochrane Database of Systematic Reviews*, **2000**, CD002791		42
459	A systematic review and meta-analysis of clinical predictors of lithium response in bipolar disorder. *Acta Psychiatrica Scandinavica*, **2019**, 140, 94-115	6.5	41
458	Prenatal and early life exposure to stressful life events and risk of autism spectrum disorders: population-based studies in Sweden and England. *PLoS ONE*, **2012**, 7, e38893	3.7	41
457	Do socio-economic risk factors predict the incidence and maintenance of psychiatric disorder in primary care?. *Psychological Medicine*, **1997**, 27, 73-80	6.9	41
456	Psychiatric disorder and dysfunction in the UK National Survey of Psychiatric Morbidity. *Social Psychiatry and Psychiatric Epidemiology*, **2000**, 35, 191-7	4.5	41
455	Socioeconomic status and alcohol-related behaviors in mid- to late adolescence in the Avon Longitudinal Study of Parents and Children. *Journal of Studies on Alcohol and Drugs*, **2014**, 75, 541-5	1.9	41
454	If cannabis caused schizophrenia--how many cannabis users may need to be prevented in order to prevent one case of schizophrenia? England and Wales calculations. *Addiction*, **2009**, 104, 1856-61	4.6	40
453	Thinking life is not worth living. A population survey of Great Britain. *Social Psychiatry and Psychiatric Epidemiology*, **2002**, 37, 351-6	4.5	40

452	Nicotine, alcohol and drug dependence, and psychiatric comorbidity--results of a national household survey. *International Review of Psychiatry*, **2003**, 15, 50-6	3.6	40
451	Mental health of British farmers. *Occupational and Environmental Medicine*, **2003**, 60, 181-5; discussion 185-6	2.1	40
450	Cortisol awakening response and subsequent depression: prospective longitudinal study. *British Journal of Psychiatry*, **2014**, 204, 137-43	5.4	39
449	The association between lower educational attainment and depression owing to shared genetic effects? Results in ~25,000 subjects. *Molecular Psychiatry*, **2015**, 20, 735-43	15.1	39
448	Lack of association between COMT and working memory in a population-based cohort of healthy young adults. *Neuropsychopharmacology*, **2013**, 38, 1253-63	8.7	39
447	Risk factors and the prevalence of neurosis and psychosis in ethnic groups in Great Britain. *Social Psychiatry and Psychiatric Epidemiology*, **2004**, 39, 939-46	4.5	39
446	Childhood inflammatory markers and intelligence as predictors of subsequent persistent depressive symptoms: a longitudinal cohort study. *Psychological Medicine*, **2018**, 48, 1514-1522	6.9	39
445	Lifetime Prevalence and Correlates of Schizophrenia-Spectrum, Affective, and Other Non-affective Psychotic Disorders in the Chinese Adult Population. *Schizophrenia Bulletin*, **2017**, 43, 1280-1290	1.3	38
444	Adolescent alcohol use is positively associated with later depression in a population-based U.K. cohort. *Journal of Studies on Alcohol and Drugs*, **2014**, 75, 758-65	1.9	38
443	The association between bullying and early stages of suicidal ideation in late adolescents in Greece. *BMC Psychiatry*, **2011**, 11, 22	4.2	38
442	Height and body mass index in young adulthood and risk of schizophrenia: a longitudinal study of 1 347 520 Swedish men. *Acta Psychiatrica Scandinavica*, **2007**, 116, 378-85	6.5	38
441	Gender differences in reoffending after discharge from medium-secure units. National cohort study in England and Wales. *British Journal of Psychiatry*, **2006**, 189, 168-72	5.4	38
440	Social roles and the gender difference in rates of the common mental disorders in Britain: a 7-year, population-based cohort study. *Psychological Medicine*, **2001**, 31, 1055-64	6.9	38
439	Apolipoprotein E-epsilon 4 allele and Alzheimer's disease. *Lancet, The*, **1993**, 342, 1308-9	40	38
438	The Avon Longitudinal Study of Parents and Children (ALSPAC) birth cohort as a resource for studying psychopathology in childhood and adolescence: a summary of findings for depression and psychosis. *Social Psychiatry and Psychiatric Epidemiology*, **2015**, 50, 1017-27	4.5	37
437	Cross-cultural patterns of the association between varying levels of alcohol consumption and the common mental disorders of depression and anxiety: secondary analysis of the WHO Collaborative Study on Psychological Problems in General Health Care. *Drug and Alcohol Dependence*, **2013**, 133, 825-31	4.9	37
436	Patient-doctor depth-of-relationship scale: development and validation. *Annals of Family Medicine*, **2011**, 9, 538-45	2.9	37
435	Cross-cultural differences in the epidemiology of unexplained fatigue syndromes in primary care. *British Journal of Psychiatry*, **2003**, 182, 205-9	5.4	37

434	Protocol for the insight study: a randomised controlled trial of single-dose tocilizumab in patients with depression and low-grade inflammation. *BMJ Open*, **2018**, 8, e025333	3	37
433	Defective processing speed and nonclinical psychotic experiences in children: longitudinal analyses in a large birth cohort. *American Journal of Psychiatry*, **2013**, 170, 550-7	11.9	36
432	Ethnic differences in prisoners. 1: criminality and psychiatric morbidity. *British Journal of Psychiatry*, **2002**, 181, 473-80	5.4	36
431	Age-period-cohort analysis of the incidence of schizophrenia in Scotland. *Psychological Medicine*, **1996**, 26, 963-73	6.9	36
430	Proactive and Reactive Response Inhibition across the Lifespan. *PLoS ONE*, **2015**, 10, e0140383	3.7	36
429	A cluster randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of classroom-based cognitive-behavioural therapy (CBT) in reducing symptoms of depression in high-risk adolescents. *Health Technology Assessment*, **2013**, 17, vii-xvii, 1-109	4.4	36
428	The effectiveness and cost-effectiveness of mindfulness-based cognitive therapy compared with maintenance antidepressant treatment in the prevention of depressive relapse/recurrence: results of a randomised controlled trial (the PREVENT study). *Health Technology Assessment*, **2015**, 19, 1-124	4.4	36
427	Association of Maternal and Paternal Depression in the Postnatal Period With Offspring Depression at Age 18 Years. *JAMA Psychiatry*, **2019**, 76, 290-296	14.5	36
426	'Third wave' cognitive and behavioural therapies versus treatment as usual for depression. *The Cochrane Library*, **2013**, CD008705	5.2	35
425	The association between paternal and adolescent depressive symptoms: evidence from two population-based cohorts. *Lancet Psychiatry,the*, **2017**, 4, 920-926	23.3	35
424	Association between locus of control in childhood and psychotic symptoms in early adolescence: results from a large birth cohort. *Cognitive Neuropsychiatry*, **2011**, 16, 385-402	2	35
423	Risk of schizophrenia and other non-affective psychosis among individuals exposed to head injury: case control study. *Schizophrenia Research*, **2006**, 88, 119-26	3.6	35
422	Systematic review of multi-symptom conditions in Gulf War veterans. *Psychological Medicine*, **2006**, 36, 735-47	6.9	35
421	Dynamical features in fetal and postnatal zinc-copper metabolic cycles predict the emergence of autism spectrum disorder. *Science Advances*, **2018**, 4, eaat1293	14.3	35
420	The longitudinal association between psychotic experiences, depression and suicidal behaviour in a population sample of adolescents. *Social Psychiatry and Psychiatric Epidemiology*, **2015**, 50, 1809-17	4.5	34
419	Childhood sleep disturbance and risk of psychotic experiences at 18: UK birth cohort. *British Journal of Psychiatry*, **2015**, 207, 23-9	5.4	34
418	Dimensions of parental alcohol use/problems and offspring temperament, externalizing behaviors, and alcohol use/problems. *Alcoholism: Clinical and Experimental Research*, **2013**, 37, 2118-27	3.7	34
417	Mental health and quality of residential environment. *British Journal of Psychiatry*, **2007**, 191, 500-5	5.4	34

416	Antidepressants for people with both schizophrenia and depression. *The Cochrane Library*, **2002**, CD002305	3.5	34
415	Self-reported school experience as a predictor of self-harm during adolescence: a prospective cohort study in the South West of England (ALSPAC). *Journal of Affective Disorders*, **2015**, 173, 163-9	6.6	33
414	Father absence and timing of menarche in adolescent girls from a UK cohort: the mediating role of maternal depression and major financial problems. *Journal of Adolescence*, **2014**, 37, 291-301	3.4	33
413	Hair Cortisol in Twins: Heritability and Genetic Overlap with Psychological Variables and Stress-System Genes. *Scientific Reports*, **2017**, 7, 15351	4.9	33
412	Prevalence and risk factors for HIV, hepatitis B, and hepatitis C in people with severe mental illness: a total population study of Sweden. *Lancet Psychiatry,the*, **2017**, 4, 685-693	23.3	33
411	Early life socio-economic position and later alcohol use: birth cohort study. *Addiction*, **2013**, 108, 516-25	4.6	33
410	Cost-effectiveness of collaborative care for depression in UK primary care: economic evaluation of a randomised controlled trial (CADET). *PLoS ONE*, **2014**, 9, e104225	3.7	33
409	Association of Combined Patterns of Tobacco and Cannabis Use in Adolescence With Psychotic Experiences. *JAMA Psychiatry*, **2018**, 75, 240-246	14.5	32
408	Family socioeconomic position in early life and onset of depressive symptoms and depression: a prospective cohort study. *Social Psychiatry and Psychiatric Epidemiology*, **2017**, 52, 95-103	4.5	32
407	A population-based longitudinal study of childhood neurodevelopmental disorders, IQ and subsequent risk of psychotic experiences in adolescence. *Psychological Medicine*, **2014**, 44, 3229-38	6.9	32
406	Misconceptions about gene–environment interactions in psychiatry. *Evidence-Based Mental Health*, **2011**, 13, 65-68	11.1	32
405	Association between serum C-reactive protein and DSM-IV generalized anxiety disorder in adolescence: Findings from the ALSPAC cohort. *Neurobiology of Stress*, **2016**, 4, 55-61	7.6	32
404	Parental mental illness and eating disorders in offspring. *International Journal of Eating Disorders*, **2015**, 48, 383-91	6.3	31
403	Investigating the genetic architecture of general and specific psychopathology in adolescence. *Translational Psychiatry*, **2018**, 8, 145	8.6	31
402	Blood-Based Protein Changes in Childhood Are Associated With Increased Risk for Later Psychotic Disorder: Evidence From a Nested Case-Control Study of the ALSPAC Longitudinal Birth Cohort. *Schizophrenia Bulletin*, **2018**, 44, 297-306	1.3	31
401	Factors associated with being a false positive on the General Health Questionnaire. *Social Psychiatry and Psychiatric Epidemiology*, **2005**, 40, 402-7	4.5	31
400	Exploring neural dysfunction in 'clinical high risk' for psychosis: a quantitative review of fMRI studies. *Journal of Psychiatric Research*, **2015**, 61, 122-34	5.2	30
399	Perinatal maternal life events and psychotic experiences in children at twelve years in a birth cohort study. *Schizophrenia Research*, **2014**, 152, 158-63	3.6	30

398	Adolescent alcohol use is predicted by childhood temperament factors before age 5, with mediation through personality and peers. *Alcoholism: Clinical and Experimental Research*, **2013**, 37, 2108-17	3.7	30
397	No pain, no gain: depressed clients' experiences of cognitive behavioural therapy. *British Journal of Clinical Psychology*, **2013**, 52, 347-64	3.6	30
396	Social anxiety is associated with general but not specific biases in emotion recognition. *Psychiatry Research*, **2013**, 210, 199-207	9.9	30
395	Trajectories of change in self-reported psychotic-like experiences in childhood and adolescence. *Schizophrenia Research*, **2012**, 140, 104-9	3.6	30
394	Early menarche and depressive symptoms from adolescence to young adulthood in a UK cohort. *Journal of the American Academy of Child and Adolescent Psychiatry*, **2013**, 52, 591-8.e2	7.2	30
393	Association between maternal depressogenic cognitive style during pregnancy and offspring cognitive style 18 years later. *American Journal of Psychiatry*, **2013**, 170, 434-41	11.9	30
392	Panic disorder and subthreshold panic in the UK general population: epidemiology, comorbidity and functional limitation. *European Psychiatry*, **2011**, 26, 354-62	6	30
391	Involving patients with depression in research: survey of patients' attitudes to participation. *British Journal of General Practice*, **2011**, 61, 134-41	1.6	30
390	Psychiatric morbidity in primary health care in Santiago, Chile. Preliminary findings. *British Journal of Psychiatry*, **1994**, 165, 530-3	5.4	30
389	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. *Health Technology Assessment*, **2016**, 20, 1-192	4.4	30
388	Psychological therapies for treatment-resistant depression in adults. *The Cochrane Library*, **2018**, 5, CD010558	5.2	30
387	Ethnicity and coercion among involuntarily detained psychiatric in-patients. *British Journal of Psychiatry*, **2010**, 196, 75-6	5.4	29
386	Study protocol for a randomized controlled trial comparing mindfulness-based cognitive therapy with maintenance anti-depressant treatment in the prevention of depressive relapse/recurrence: the PREVENT trial. *Trials*, **2010**, 11, 99	2.8	29
385	Alcohol consumption as a risk factor for non-recovery from common mental disorder: results from the longitudinal follow-up of the National Psychiatric Morbidity Survey. *Psychological Medicine*, **2008**, 38, 451-5	6.9	29
384	Ethnic differences in prisoners. 2: risk factors and psychiatric service use. *British Journal of Psychiatry*, **2002**, 181, 481-7	5.4	29
383	Links between social class and common mental disorders in Northeast Brazil. *Social Psychiatry and Psychiatric Epidemiology*, **2001**, 36, 101-7	4.5	29
382	Mirtazapine added to SSRIs or SNRIs for treatment resistant depression in primary care: phase III randomised placebo controlled trial (MIR). *BMJ, The*, **2018**, 363, k4218	5.9	29
381	Assessing negative cognitive style: Development and validation of a Short-Form version of the Cognitive Style Questionnaire. *Personality and Individual Differences*, **2012**, 52, 581-585	3.3	28

380	Rationale and development of the physical activity counselling intervention for a pragmatic TRial of Exercise and Depression in the UK (TREAD-UK). *Mental Health and Physical Activity*, **2010**, 3, 85-91	5	28
379	Depression and anxiety in children with CFS/ME: cause or effect?. *Archives of Disease in Childhood*, **2011**, 96, 211-4	2.2	28
378	Schizophrenia sans frontieres: concepts of schizophrenia among French and British psychiatrists. *BMJ: British Medical Journal*, **1993**, 307, 489-92		28
377	Reward-Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A Systematic Review and Meta-analysis. *JAMA Psychiatry*, **2020**, 77, 1286-1295	14.5	28
376	Association of Hydroxylmethyl Glutaryl Coenzyme A Reductase Inhibitors, L-Type Calcium Channel Antagonists, and Biguanides With Rates of Psychiatric Hospitalization and Self-Harm in Individuals With Serious Mental Illness. *JAMA Psychiatry*, **2019**, 76, 382-390	14.5	28
375	The longitudinal association between external locus of control, social cognition and adolescent psychopathology. *Social Psychiatry and Psychiatric Epidemiology*, **2017**, 52, 643-655	4.5	27
374	Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. *British Journal of General Practice*, **2016**, 66, e78-84	1.6	27
373	Identification of a plasma signature of psychotic disorder in children and adolescents from the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort. *Translational Psychiatry*, **2017**, 7, e1240	8.6	27
372	Common versus psychopathology-specific risk factors for psychotic experiences and depression during adolescence. *Psychological Medicine*, **2014**, 44, 2557-66	6.9	27
371	Cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment resistant depression in primary care: the CoBalT randomised controlled trial protocol. *Contemporary Clinical Trials*, **2012**, 33, 312-9	2.3	27
370	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depression--study protocol. *BMC Health Services Research*, **2009**, 9, 188	2.9	27
369	Inequities in mental health care after health care system reform in Chile. *American Journal of Public Health*, **2006**, 96, 109-13	5.1	27
368	Psychiatric morbidity in prisoners and solitary cellular confinement, I: disciplinary segregation. *Journal of Forensic Psychiatry and Psychology*, **2003**, 14, 298-319	0.9	27
367	The case-control study in psychiatry. *British Journal of Psychiatry*, **1990**, 157, 197-207	5.4	27
366	Scientific rigor and the art of motorcycle maintenance. *Nature Biotechnology*, **2014**, 32, 871-3	44.5	26
365	Parental socio-economic position during childhood as a determinant of self-harm in adolescence. *Social Psychiatry and Psychiatric Epidemiology*, **2014**, 49, 193-203	4.5	26
364	Socio-economic status, standard of living, and neurotic disorder. *International Review of Psychiatry*, **2003**, 15, 91-6	3.6	26
363	Clarifying the relationship between unexplained chronic fatigue and psychiatric morbidity: results from a community survey in Great Britain. *International Review of Psychiatry*, **2003**, 15, 57-64	3.6	26

362	Is there a gender difference on the association between informal work and common mental disorders?. *Social Psychiatry and Psychiatric Epidemiology*, **2005**, 40, 622-7	4.5	26
361	Longitudinal associations between adolescent psychotic experiences and depressive symptoms. *PLoS ONE*, **2014**, 9, e105758	3.7	26
360	New insights into the pharmacogenomics of antidepressant response from the GENDEP and STAR*D studies: rare variant analysis and high-density imputation. *Pharmacogenomics Journal*, **2018** , 18, 413-421	3.5	26
359	Mediation of Developmental Risk Factors for Psychosis by White Matter Microstructure in Young Adults With Psychotic Experiences. *JAMA Psychiatry*, **2016**, 73, 396-406	14.5	25
358	Childhood internalizing symptoms are negatively associated with early adolescent alcohol use. *Alcoholism: Clinical and Experimental Research*, **2014**, 38, 1680-8	3.7	25
357	Exploring patients' reasons for declining contact in a cognitive behavioural therapy randomised controlled trial in primary care. *British Journal of General Practice*, **2012**, 62, e371-7	1.6	25
356	Conduct problem trajectories and alcohol use and misuse in mid to late adolescence. *Drug and Alcohol Dependence*, **2013**, 133, 100-7	4.9	25
355	Is the General Health Questionnaire (12 item) a culturally biased measure of psychiatric disorder?. *Social Psychiatry and Psychiatric Epidemiology*, **1995**, 30, 20-5	4.5	25
354	Observer bias in the assessment of anxiety and depression. *Social Psychiatry and Psychiatric Epidemiology*, **1991**, 26, 265-72	4.5	25
353	Serum 25-hydroxyvitamin D3 and D2 and non-clinical psychotic experiences in childhood. *PLoS ONE*, **2012**, 7, e41575	3.7	25
352	Variation in the recall of socially rewarding information and depressive symptom severity: a prospective cohort study. *Acta Psychiatrica Scandinavica*, **2017**, 135, 489-498	6.5	24
351	Association of measures of fetal and childhood growth with non-clinical psychotic symptoms in 12-year-olds: the ALSPAC cohort. *British Journal of Psychiatry*, **2009**, 194, 521-6	5.4	24
350	Risk of low Apgar score and socioeconomic position: a study of Swedish male births. *Acta Paediatrica, International Journal of Paediatrics*, **2008**, 97, 1275-80	3.1	24
349	Clinical judgement and the standardized interview in psychiatry. *Psychological Medicine*, **1989**, 19, 971-9	6.9	24
348	The association of 25-hydroxyvitamin D3 and D2 with behavioural problems in childhood. *PLoS ONE* , **2012**, 7, e40097	3.7	24
347	Psychiatric disorder in veterans of the Persian Gulf War of 1991. Systematic review. *British Journal of Psychiatry*, **2003**, 182, 391-403	5.4	24
346	The process and delivery of CBT for depression in adults: a systematic review and network meta-analysis. *Psychological Medicine*, **2019**, 49, 1937-1947	6.9	23
345	The influence of school on whether girls develop eating disorders. *International Journal of Epidemiology*, **2016**, 45, 480-8	7.8	23

344	Testing a level of response to alcohol-based model of heavy drinking and alcohol problems in 1,905 17-year-olds. *Alcoholism: Clinical and Experimental Research*, **2011**, 35, 1897-904	3.7	23
343	One-year outcome of unexplained fatigue syndromes in primary care: results from an international study. *Psychological Medicine*, **2003**, 33, 857-66	6.9	23
342	Substance misuse and psychiatric comorbidity: an overview of the OPCS National Psychiatric Morbidity Survey. *International Review of Psychiatry*, **2003**, 15, 43-9	3.6	23
341	Unequal access and unmet need: neurotic disorders and the use of primary care services. *International Review of Psychiatry*, **2003**, 15, 115-22	3.6	23
340	Comparing sex inequalities in common affective disorders across countries: Great Britain and Chile. *Social Science and Medicine*, **2005**, 60, 1693-703	5.1	23
339	Obstetric complications and risk of schizophrenia. Effect of gender, age at diagnosis and maternal history of psychosis. *British Journal of Psychiatry*, **2001**, 179, 409-14	5.4	23
338	Another British disease? A recent increase in the prevalence of psychiatric morbidity. *Journal of Epidemiology and Community Health*, **1993**, 47, 358-61	5.1	23
337	Discrimination on the grounds of diagnosis. *Addiction*, **1990**, 85, 883-890	4.6	23
336	Screening for depression in primary care. *British Journal of General Practice*, **2005**, 55, 659-60	1.6	23
335	Psychological processes mediating the association between developmental trauma and specific psychotic symptoms in adults: a systematic review and meta-analysis. *World Psychiatry*, **2021**, 20, 107-123	14.4	23
334	Trajectories of Neighborhood Cohesion in Childhood, and Psychotic and Depressive Symptoms at Age 13 and 18 Years. *Journal of the American Academy of Child and Adolescent Psychiatry*, **2017**, 56, 570-577	7.2	22
333	Hospitalisation of people with dementia: evidence from English electronic health records from 2008 to 2016. *European Journal of Epidemiology*, **2019**, 34, 567-577	12.1	22
332	Pain in veterans of the Gulf War of 1991: a systematic review. *BMC Musculoskeletal Disorders*, **2006**, 7, 74	2.8	22
331	Dysfunctional attitudes and the common mental disorders in primary care. *Journal of Affective Disorders*, **2003**, 75, 269-78	6.6	22
330	Misconceptions about gene-environment interactions in psychiatry. *Evidence-Based Mental Health*, **2010**, 13, 65-8	11.1	22
329	Depression and self-harm from adolescence to young adulthood in sexual minorities compared with heterosexuals in the UK: a population-based cohort study. *The Lancet Child and Adolescent Health*, **2019**, 3, 91-98	14.5	22
328	How data science can advance mental health research. *Nature Human Behaviour*, **2019**, 3, 24-32	12.8	22
327	Childhood interleukin-6, C-reactive protein and atopic disorders as risk factors for hypomanic symptoms in young adulthood: a longitudinal birth cohort study. *Psychological Medicine*, **2017**, 47, 23-33	6.9	21

326	Risk of depression and self-harm in teenagers identifying with goth subculture: a longitudinal cohort study. *Lancet Psychiatry,the*, **2015**, 2, 793-800	23.3	21
325	Age- and puberty-dependent association between IQ score in early childhood and depressive symptoms in adolescence. *Psychological Medicine*, **2011**, 41, 333-43	6.9	21
324	Neurotic disorders and the receipt of psychiatric treatment. *International Review of Psychiatry*, **2003**, 15, 108-14	3.6	21
323	Dimensions of neurosis. *Psychological Medicine*, **1992**, 22, 1011-8	6.9	21
322	Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depressive disorder		21
321	Suicide risk among refugees compared with non-refugee migrants and the Swedish-born majority population. *British Journal of Psychiatry*, **2020**, 217, 686-692	5.4	21
320	Do eating disorders in parents predict eating disorders in children? Evidence from a Swedish cohort. *Acta Psychiatrica Scandinavica*, **2015**, 132, 51-9	6.5	20
319	Sleep disturbances and depressive symptoms: an investigation of their longitudinal association in a representative sample of the UK general population. *Psychological Medicine*, **2013**, 43, 329-39	6.9	20
318	A Randomized Controlled Trial of Cognitive Behavioural Therapy as an Adjunct to Pharmacotherapy in Primary Care Based Patients with Treatment Resistant Depression: A Pilot Study. *Behavioural and Cognitive Psychotherapy*, **2008**, 36, 21-33	2.1	20
317	GENetic and clinical predictors of treatment response in depression: the GenPod randomised trial protocol. *Trials*, **2008**, 9, 29	2.8	20
316	Prenatal and childhood adverse life events, inflammation and depressive symptoms across adolescence. *Journal of Affective Disorders*, **2020**, 260, 577-582	6.6	20
315	The association between pubertal status and depressive symptoms and diagnoses in adolescent females: A population-based cohort study. *PLoS ONE*, **2018**, 13, e0198804	3.7	20
314	Psychotic Experiences, Working Memory, and the Developing Brain: A Multimodal Neuroimaging Study. *Cerebral Cortex*, **2015**, 25, 4828-38	5.1	19
313	Systematic review and meta-analysis of serotonin transporter genotype and discontinuation from antidepressant treatment. *European Neuropsychopharmacology*, **2013**, 23, 1143-50	1.2	19
312	Cost-effectiveness of cognitive-behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: economic evaluation of the CoBalT Trial. *British Journal of Psychiatry*, **2014**, 204, 69-76	5.4	19
311	Cost-effectiveness of classroom-based cognitive behaviour therapy in reducing symptoms of depression in adolescents: a trial-based analysis. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, **2014**, 55, 1390-7	7.9	19
310	The study of gene-environment interactions in psychiatry: limited gains at a substantial cost?. *Psychological Medicine*, **2010**, 40, 711-716	6.9	19
309	Emotional symptoms in children: The effect of maternal depression, life events, and COMT genotype. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **2009**, 150B, 209-18	3.5	19

308	Computerised patient-specific guidelines for management of common mental disorders in primary care: a randomised controlled trial. *British Journal of General Practice*, **2004**, 54, 832-7	1.6	19
307	Risk of schizophrenia, schizoaffective, and bipolar disorders by migrant status, region of origin, and age-at-migration: a national cohort study of 1.8 million people. *Psychological Medicine*, **2019**, 49, 2354-2363	6.9	19
306	Copy number variants and therapeutic response to antidepressant medication in major depressive disorder. *Pharmacogenomics Journal*, **2014**, 14, 395-9	3.5	18
305	Individual- and area-level influence on suicide risk: a multilevel longitudinal study of Swedish schoolchildren. *Psychological Medicine*, **2014**, 44, 267-77	6.9	18
304	General Practitioners beliefs about physical activity for managing depression in primary care. *Mental Health and Physical Activity*, **2012**, 5, 13-19	5	18
303	Examining the association between maternal analgesic use during pregnancy and risk of psychotic symptoms during adolescence. *Schizophrenia Research*, **2011**, 126, 220-5	3.6	18
302	Is low IQ associated with an increased risk of developing suicidal thoughts? A cohort study based on an 18-month follow-up of the national psychiatric morbidity survey. *Social Psychiatry and Psychiatric Epidemiology*, **2009**, 44, 34-8	4.5	18
301	The association between birth condition and neuropsychological functioning and educational attainment at school age: a cohort study. *Archives of Disease in Childhood*, **2011**, 96, 30-7	2.2	18
300	Severity of depression and response to antidepressants: GENPOD randomised controlled trial. *British Journal of Psychiatry*, **2012**, 200, 130-6	5.4	18
299	Multi-symptom illnesses, unexplained illness and Gulf War Syndrome. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **2006**, 361, 543-51	5.8	18
298	Palliative day care: what does it cost to run a centre and does attendance affect use of other services?. *Palliative Medicine*, **2003**, 17, 628-37	5.5	18
297	Investigating the effect of demographic and socioeconomic variables on misclassification by the SRQ-20 compared with a psychiatric interview. *Social Psychiatry and Psychiatric Epidemiology*, **2005**, 40, 36-41	4.5	18
296	Patients' views on outcome following head injury: a qualitative study. *BMC Family Practice*, **2005**, 6, 30	2.6	18
295	Psychotic Experiences and Working Memory: A Population-Based Study Using Signal-Detection Analysis. *PLoS ONE*, **2016**, 11, e0153148	3.7	18
294	Complement pathway changes at age 12 are associated with psychotic experiences at age 18 in a longitudinal population-based study: evidence for a role of stress. *Molecular Psychiatry*, **2021**, 26, 524-533	15.1	18
293	Strategies for preventing suicide. *British Journal of Psychiatry*, **1997**, 171, 351-4	5.4	18
292	A population-based prospective birth cohort study of childhood neurocognitive and psychological functioning in healthy survivors of early life meningitis. *Annals of Epidemiology*, **2015**, 25, 236-42	6.4	17
291	Association of polygenic score for major depression with response to lithium in patients with bipolar disorder. *Molecular Psychiatry*, **2021**, 26, 2457-2470	15.1	17

290	Prevention of anxiety and depression in Chinese: a randomized clinical trial testing the effectiveness of a stepped care program in primary care. *Journal of Affective Disorders*, **2014**, 169, 212-20	6.6	17
289	Theory of mind and social functioning in first episode psychosis. *Cognitive Neuropsychiatry*, **2013**, 18, 219-42	2	17
288	Long-term impact of poor birth condition on social and economic outcomes in early adulthood. *Pediatrics*, **2011**, 127, e1498-504	7.4	17
287	Smoking and common mental disorders: a population-based survey in Santiago, Chile. *Social Psychiatry and Psychiatric Epidemiology*, **2007**, 42, 874-80	4.5	17
286	DSM-IV. Diagnostic and Statistical Manual of Mental Disorders, 4th edn. By the American Psychiatric Association. (Pp. 886; £34.95.) APA: Washington, DC.1994.. *Psychological Medicine*, **1996**, 26, 651-652	6.9	17
285	Symptoms of generalized anxiety disorder but not panic disorder at age 15 years increase the risk of depression at 18 years in the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort study. *Psychological Medicine*, **2016**, 46, 73-85	6.9	17
284	Trajectories of autistic social traits in childhood and adolescence and disordered eating behaviours at age 14 years: A UK general population cohort study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, **2021**, 62, 75-85	7.9	17
283	Polygenic risk for schizophrenia, disordered eating behaviours and body mass index in adolescents. *British Journal of Psychiatry*, **2019**, 215, 428-433	5.4	16
282	The relationship between psychotic symptoms and social functioning in a non-clinical population of 12year olds. *Schizophrenia Research*, **2013**, 150, 404-9	3.6	16
281	Investigation of thyroid dysfunction is more likely in patients with high psychological morbidity. *Family Practice*, **2012**, 29, 163-7	1.9	16
280	Mindfulness-based 'third wave' cognitive and behavioural therapies versus other psychological therapies for depression. *Cochrane Database of Systematic Reviews*, **2010**,		16
279	Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. *Health Technology Assessment*, **2018**, 22, 1-136	4.4	16
278	Maintenance or Discontinuation of Antidepressants in Primary Care. *New England Journal of Medicine*, **2021**, 385, 1257-1267	59.2	16
277	The Hong Kong mental morbidity survey: background and study design. *East Asian Archives of Psychiatry*, **2014**, 24, 30-6	1.1	16
276	Cognitive styles and future depressed mood in early adulthood: the importance of global attributions. *Journal of Affective Disorders*, **2015**, 171, 60-7	6.6	15
275	Early adolescent outcomes of joint developmental trajectories of problem behavior and IQ in childhood. *European Child and Adolescent Psychiatry*, **2018**, 27, 1595-1605	5.5	15
274	Adolescent over-general memory, life events and mental health outcomes: Findings from a UK cohort study. *Memory*, **2016**, 24, 348-63	1.8	15
273	Directional relationships between alcohol use and antisocial behavior across adolescence. *Alcoholism: Clinical and Experimental Research*, **2014**, 38, 2024-33	3.7	15

272	The longitudinal association between social functioning and theory of mind in first-episode psychosis. *Cognitive Neuropsychiatry*, **2014**, 19, 58-80	2	15
271	Social inference and social anxiety: evidence of a fear-congruent self-referential learning bias. *Journal of Behavior Therapy and Experimental Psychiatry*, **2012**, 43, 1082-7	2.6	15
270	Toxoplasma and coxiella infection and psychiatric morbidity: a retrospective cohort analysis. *BMC Psychiatry*, **2004**, 4, 32	4.2	15
269	Randomised controlled trial of gradual antipsychotic reduction and discontinuation in people with schizophrenia and related disorders: the RADAR trial (Research into Antipsychotic Discontinuation and Reduction). *BMJ Open*, **2019**, 9, e030912	3	15
268	Antidepressant Controlled Trial For Negative Symptoms In Schizophrenia (ACTIONS): a double-blind, placebo-controlled, randomised clinical trial. *Health Technology Assessment*, **2016**, 20, 1-46	4.4	15
267	Examining reward-seeking, negative self-beliefs and over-general autobiographical memory as mechanisms of change in classroom prevention programs for adolescent depression. *Journal of Affective Disorders*, **2015**, 186, 320-7	6.6	14
266	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. *Psychological Medicine*, **2020**, 1-8	6.9	14
265	Maternal perinatal depressive symptoms and offspring psychotic experiences at 18 years of age: a longitudinal study. *Lancet Psychiatry,the*, **2020**, 7, 431-440	23.3	14
264	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. *Trials*, **2017**, 18, 496	2.8	14
263	IQ and adolescent self-harm behaviours in the ALSPAC birth cohort. *Journal of Affective Disorders*, **2014**, 152-154, 175-82	6.6	14
262	Childhood traumatic events and adolescent overgeneral autobiographical memory: findings in a U.K. cohort. *Journal of Behavior Therapy and Experimental Psychiatry*, **2014**, 45, 330-8	2.6	14
261	An empirical investigation of the structure of anxiety and depressive symptoms in late adolescence: cross-sectional study using the Greek version of the revised Clinical Interview Schedule. *Psychiatry Research*, **2011**, 186, 419-26	9.9	14
260	Childhood diet and behavioural problems: results from the ALSPAC cohort. *European Journal of Clinical Nutrition*, **2011**, 65, 720-6	5.2	14
259	Exploring the relationship between cannabis use and psychosis. *Addiction*, **2004**, 99, 1353-5	4.6	14
258	"Patient knows best"-detection of common mental disorders in Santiago, Chile: cross sectional study. *BMJ: British Medical Journal*, **2001**, 322, 79-81		14
257	Effectiveness of a Mobile App Intervention for Anxiety and Depression Symptoms in University Students: Randomized Controlled Trial. *JMIR MHealth and UHealth*, **2020**, 8, e15418	5.5	14
256	Association between childhood psychiatric disorders and psychotic experiences in adolescence: A population-based longitudinal study. *Comprehensive Psychiatry*, **2016**, 69, 45-52	7.3	14
255	Pharmacological interventions for treatment-resistant depression in adults. *The Cochrane Library*, **2019**, 12, CD010557	5.2	14

254	Developmental cascades of internalising symptoms, externalising problems and cognitive ability from early childhood to middle adolescence. *European Psychiatry*, **2019**, 57, 61-69	6	13
253	Integrated Lipidomics and Proteomics Point to Early Blood-Based Changes in Childhood Preceding Later Development of Psychotic Experiences: Evidence From the Avon Longitudinal Study of Parents and Children. *Biological Psychiatry*, **2019**, 86, 25-34	7.9	13
252	Moderators of response to cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care. *Journal of Affective Disorders*, **2015**, 174, 272-80	6.6	13
251	Is smoking heaviness causally associated with alcohol use? A Mendelian randomization study in four European cohorts. *International Journal of Epidemiology*, **2018**, 47, 1098-1105	7.8	13
250	The relative responsiveness of test instruments can be estimated using a meta-analytic approach: an illustration with treatments for depression. *Journal of Clinical Epidemiology*, **2016**, 77, 68-77	5.7	13
249	Visual Acuity in Late Adolescence and Future Psychosis Risk in a Cohort of 1 Million Men. *Schizophrenia Bulletin*, **2019**, 45, 571-578	1.3	13
248	Genetic predictors of antidepressant side effects: a grouped candidate gene approach in the Genome-Based Therapeutic Drugs for Depression (GENDEP) study. *Journal of Psychopharmacology*, **2014**, 28, 142-50	4.6	13
247	Participants' experiences of facilitated physical activity for the management of depression in primary care. *Journal of Health Psychology*, **2014**, 19, 1430-42	3.1	13
246	Genetic susceptibility for bipolar disorder and response to antidepressants in major depressive disorder. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **2014**, 165B, 77-83	3.5	13
245	Perinatal folate-related exposures and risk of psychotic symptoms in the ALSPAC birth cohort. *Schizophrenia Research*, **2010**, 120, 177-83	3.6	13
244	Physical activity as a treatment for depression: the TREAD randomised trial protocol. *Trials*, **2010**, 11, 105	2.8	13
243	Protocol for a randomised controlled trial of a school based cognitive behaviour therapy (CBT) intervention to prevent depression in high risk adolescents (PROMISE). *Trials*, **2010**, 11, 114	2.8	13
242	Prevalence of reported pain, widespread pain, and pain symmetry in veterans of the Persian Gulf War (1990-1991): the use of pain manikins in Persian Gulf War health research. *Military Medicine*, **2006**, 171, 1181-6	1.3	13
241	The National Survey of Psychiatric Morbidity among prisoners and the future of prison healthcare. *Medicine, Science and the Law*, **2002**, 42, 245-50	1.1	13
240	Leisure activity participation and risk of dementia: An 18-year follow-up of the Whitehall II Study. *Neurology*, **2020**, 95, e2803-e2815	6.5	13
239	The Community Navigator Study: Results from a feasibility randomised controlled trial of a programme to reduce loneliness for people with complex anxiety or depression. *PLoS ONE*, **2020**, 15, e0233535	3.7	12
238	The incidence and associated risk factors for sudden unexplained death in psychiatric in-patients in England and Wales. *Journal of Psychopharmacology*, **2011**, 25, 1533-42	4.6	12
237	Childhood facial emotion recognition and psychosis-like symptoms in a nonclinical population at 12 years of age: results from the ALSPAC birth cohort. *Cognitive Neuropsychiatry*, **2011**, 16, 136-57	2	12

236	What's in a name? A comparison of four psychiatric assessments. *Social Psychiatry and Psychiatric Epidemiology*, **1998**, 33, 27-31	4.5	12
235	Patients' opinions of the use of psychiatric case-finding questionnaires in general practice. *Health Expectations*, **2002**, 5, 282-8	3.7	12
234	British Psychiatric Morbidity Survey. *International Review of Psychiatry*, **2003**, 15, 14-8	3.6	12
233	Psychiatric morbidity in prisoners and solitary cellular confinement, II: special ('strip') cells. *Journal of Forensic Psychiatry and Psychology*, **2003**, 14, 320-340	0.9	12
232	The National Psychiatric Morbidity Surveys of Great Britain--strategy and methods. *International Review of Psychiatry*, **2003**, 15, 5-13	3.6	12
231	Effect of early life experiences and personality on the reporting of psychosocial distress in general practice. A preliminary investigation. *British Journal of Psychiatry*, **1996**, 168, 116-20	5.4	12
230	Minimal phenotyping yields GWAS hits of reduced specificity for major depression		12
229	A longitudinal investigation of childhood communication ability and adolescent psychotic experiences in a community sample. *Schizophrenia Research*, **2016**, 173, 54-61	3.6	12
228	Association of Whole-Genome and NETRIN1 Signaling Pathway-Derived Polygenic Risk Scores for Major Depressive Disorder and White Matter Microstructure in the UK Biobank. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, **2019**, 4, 91-100	3.4	12
227	Changes in the Prevalence and Correlates of Weight-Control Behaviors and Weight Perception in Adolescents in the UK, 1986-2015. *JAMA Pediatrics*, **2021**, 175, 267-275	8.3	12
226	Prevalence of anxiety disorders in community dwelling older adults in Hong Kong. *International Psychogeriatrics*, **2017**, 29, 259-267	3.4	11
225	Association of neighbourhood migrant density and risk of non-affective psychosis: a national, longitudinal cohort study. *Lancet Psychiatry,the*, **2020**, 7, 327-336	23.3	11
224	Why are there discrepancies between depressed patients' Global Rating of Change and scores on the Patient Health Questionnaire depression module? A qualitative study of primary care in England. *BMJ Open*, **2017**, 7, e014519	3	11
223	Predictors of mental health service utilization in the 12 months before imprisonment: analysis of results from a national prisons survey. *Australian and New Zealand Journal of Psychiatry*, **2006**, 40, 548-53	2.6	11
222	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. *Wellcome Open Research*, **2019** , 4, 69	4.8	11
221	Fear of negative evaluation biases social evaluation inference: evidence from a probabilistic learning task. *PLoS ONE*, **2015**, 10, e0119456	3.7	11
220	Neighborhood Characteristics at Birth and Positive and Negative Psychotic Symptoms in Adolescence: Findings From the ALSPAC Birth Cohort. *Schizophrenia Bulletin*, **2020**, 46, 581-591	1.3	11
219	Applying polygenic risk scoring for psychiatric disorders to a large family with bipolar disorder and major depressive disorder. *Communications Biology*, **2018**, 1, 163	6.7	11

218	Role of age, gender and marital status in prognosis for adults with depression: An individual patient data meta-analysis. *Epidemiology and Psychiatric Sciences*, **2021**, 30, e42	5.1	11
217	Curiosity killed the cat: no evidence of an association between cat ownership and psychotic symptoms at ages 13 and 18 years in a UK general population cohort. *Psychological Medicine*, **2017**, 47, 1659-1667	6.9	10
216	Longitudinal associations between psychotic experiences and disordered eating behaviours in adolescence: a UK population-based study. *The Lancet Child and Adolescent Health*, **2018**, 2, 591-599	14.5	10
215	Update to the study protocol for a randomized controlled trial comparing mindfulness-based cognitive therapy with maintenance anti-depressant treatment depressive relapse/recurrence: the PREVENT trial. *Trials*, **2014**, 15, 217	2.8	10
214	Allowing for non-adherence to treatment in a randomized controlled trial of two antidepressants (citalopram versus reboxetine): an example from the GENPOD trial. *Psychological Medicine*, **2014**, 44, 2855-66	6.9	10
213	Patients' experiences of participating in a large-scale trial of cognitive behavioural therapy for depression: a mixed methods study. *Family Practice*, **2013**, 30, 705-11	1.9	10
212	Borna disease virus and mental health: a cross-sectional study. *QJM - Monthly Journal of the Association of Physicians*, **2005**, 98, 247-54	2.7	10
211	Psychiatric and alcohol disorders as risk factors for drug abuse. A case-control study among adults in Rio de Janeiro, Brazil. *Social Psychiatry and Psychiatric Epidemiology*, **1996**, 31, 355-63	4.5	10
210	Suicide and psychiatric services. *Lancet, The*, **1994**, 344, 822	40	10
209	Association of Primary Care Consultation Patterns With Early Signs and Symptoms of Psychosis. *JAMA Network Open*, **2018**, 1, e185174	10.4	10
208	Stressful life events, inflammation and emotional and behavioural problems in children: A population-based study. *Brain, Behavior, and Immunity*, **2019**, 80, 66-72	16.6	9
207	Social factors associated with readiness for sexual activity in adolescents: a population-based cohort study. *Archives of Sexual Behavior*, **2015**, 44, 669-78	3.5	9
206	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. *Trials*, **2016**, 17, 66	2.8	9
205	Peer victimisation during adolescence and its impact on depression in early adulthood: prospective cohort study in the United Kingdom. *British Journal of Sports Medicine*, **2016**, 50, 176-83	10.3	9
204	A U-shaped relationship between systolic blood pressure and panic symptoms: the HUNT study. *Psychological Medicine*, **2012**, 42, 1969-76	6.9	9
203	Behavioural therapies versus other psychological therapies for depression. *Cochrane Database of Systematic Reviews*, **2010**,		9
202	Cognitive behavioural therapies versus treatment as usual for depression. *The Cochrane Library*, **2010**,	5.2	9
201	Cannabis and psychosis Authors' reply. *Lancet, The*, **2007**, 370, 1539-1540	40	9

200	Globalization and Mental Health57-78		9
199	Classification, disability and the public health agenda. *British Medical Bulletin*, **2001**, 57, 3-15	5.4	9
198	Possible Association of APOE Genotype with Working Memory in Young Adults. *PLoS ONE*, **2015**, 10, e0135894	3.7	9
197	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. *Wellcome Open Research*, **2019** , 4, 69	4.8	9
196	Volumetric, relaxometric and diffusometric correlates of psychotic experiences in a non-clinical sample of young adults. *NeuroImage: Clinical*, **2016**, 12, 550-558	5.3	9
195	Combined influence of depressive symptoms and systemic inflammation on all-cause and cardiovascular mortality: evidence for differential effects by gender in the English Longitudinal Study of Ageing. *Psychological Medicine*, **2019**, 49, 1521-1531	6.9	9
194	Classical Human Leukocyte Antigen Alleles and C4 Haplotypes Are Not Significantly Associated With Depression. *Biological Psychiatry*, **2020**, 87, 419-430	7.9	9
193	Suicide, self-harm and thoughts of suicide or self-harm in infectious disease epidemics: a systematic review and meta-analysis. *Epidemiology and Psychiatric Sciences*, **2021**, 30, e32	5.1	9
192	Effects of exposure to bodies of different sizes on perception of and satisfaction with own body size: two randomized studies. *Royal Society Open Science*, **2018**, 5, 171387	3.3	8
191	Variation in recognition of happy and sad facial expressions and self-reported depressive symptom severity: A prospective cohort study. *Journal of Affective Disorders*, **2019**, 257, 461-469	6.6	8
190	World region of origin and common mental disorders among migrant women in Spain. *Journal of Immigrant and Minority Health*, **2014**, 16, 1111-20	2.2	8
189	Disability pension for psychiatric disorders: regional differences in Norway 1988-2000. *Nordic Journal of Psychiatry*, **2006**, 60, 255-62	2.3	8
188	Family networks during migration and risk of non-affective psychosis: A population-based cohort study. *Schizophrenia Research*, **2019**, 208, 268-275	3.6	8
187	Cognitive Impairment and Depression-Cause, Consequence, or Coincidence?. *JAMA Psychiatry*, **2019** , 76, 239-240	14.5	8
186	Childhood and Parental Asthma, Future Risk of Bipolar Disorder and Schizophrenia Spectrum Disorders: A Population-Based Cohort Study. *Schizophrenia Bulletin*, **2019**, 45, 360-368	1.3	8
185	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpRession): study protocol for a	2.8	7
184	The Community Navigator Study: a feasibility randomised controlled trial of an intervention to increase community connections and reduce loneliness for people with complex anxiety or depression. *Trials*, **2017**, 18, 493	2.8	7
183	Association between childhood infection, serum inflammatory markers and intelligence: findings from a population-based prospective birth cohort study. *Epidemiology and Infection*, **2018**, 146, 256-264	4.3	7

182	Patient-reported outcome measures for monitoring primary care patients with depression: PROMDEP feasibility randomised trial. *BMJ Open*, **2017**, 7, e015266	3	7
181	Mindfulness-based 'third wave' cognitive and behavioural therapies versus treatment as usual for depression. *Cochrane Database of Systematic Reviews*, **2010**,		7
180	Frequency of psychosis-like symptoms in a non-clinical population of 12 year olds: Results from the Alspac birth cohort. *European Psychiatry*, **2008**, 23, S282	6	7
179	The contribution of depressive 'disorder characteristics' to determinations of prognosis for adults with depression: an individual patient data meta-analysis. *Psychological Medicine*, **2021**, 51, 1068-1081	6.9	7
178	Predicting prognosis for adults with depression using individual symptom data: a comparison of modelling approaches. *Psychological Medicine*, **2021**, 1-11	6.9	7
177	Management of treatment-resistant depression in primary care: a mixed-methods study. *British Journal of General Practice*, **2018**, 68, e673-e681	1.6	7
176	The Genetic Architecture of Depression in Individuals of East Asian Ancestry: A Genome-Wide Association Study. *JAMA Psychiatry*, **2021**, 78, 1258-1269	14.5	7
175	Individuals' Long Term Use of Cognitive Behavioural Skills to Manage their Depression: A Qualitative Study. *Behavioural and Cognitive Psychotherapy*, **2017**, 45, 46-57	2.1	6
174	Hemispheric language asymmetry in first episode psychosis and schizotypy: the role of cannabis consumption and cognitive disorganization. *Schizophrenia Bulletin*, **2015**, 41 Suppl 2, S455-64	1.3	6
173	REDUCE (Reviewing long-term antidepressant use by careful monitoring in everyday practice) internet and telephone support to people coming off long-term antidepressants: protocol for a randomised controlled trial. *Trials*, **2020**, 21, 419	2.8	6
172	Trends in levels of self-reported psychological distress among individuals who seek psychiatric services over eight years: a comparison between age groups in three population surveys in Stockholm County. *BMC Psychiatry*, **2017**, 17, 345	4.2	6
171	Do disordered eating behaviours in girls vary by school characteristics? A UK cohort study. *European Child and Adolescent Psychiatry*, **2018**, 27, 1473-1481	5.5	6
170	Psychotic experiences and social functioning: a longitudinal study. *Social Psychiatry and Psychiatric Epidemiology*, **2013**, 48, 1053-65	4.5	6
169	Postnatal depression in women after unsuccessful attempted abortion. *British Journal of Psychiatry*, **2011**, 198, 237-8	5.4	6
168	Cognitive behavioural therapies versus other psychological therapies for depression. *The Cochrane Library*, **2010**,	5.2	6
167	Defining Replication: A Response to Kaufman and Colleagues. *Biological Psychiatry*, **2010**, 67, e21-e23	7.9	6
166	Interpersonal, cognitive analytic and other integrative therapies versus treatment as usual for depression. *The Cochrane Library*, **2010**,	5.2	6
165	Continuum of self-destructiveness. *Lancet, The*, **1998**, 351, 81-2	40	6

164	Is the association of hypertension and panic disorder explained by clustering of autonomic panic symptoms in hypertensive patients?. *Journal of Affective Disorders*, **2008**, 111, 344-50	6.6	6
163	Psychiatric and neuropsychiatric syndromes and COVID-19 - Authors' reply. *Lancet Psychiatry,the*, **2020**, 7, 664-665	23.3	6
162	MRI Indices of Cortical Development in Young People With Psychotic Experiences: Influence of Genetic Risk and Persistence of Symptoms. *Schizophrenia Bulletin*, **2019**, 45, 169-179	1.3	6
161	Comparison between self-administered depression questionnaires and patients' own views of changes in their mood: a prospective cohort study in primary care. *Psychological Medicine*, **2021**, 51, 853-860	6.9	6
160	Patient-reported outcome measures for monitoring primary care patients with depression (PROMDEP): study protocol for a randomised controlled trial. *Trials*, **2020**, 21, 441	2.8	5
159	Transcranial magnetic stimulation for depression. *Lancet, The*, **2018**, 391, 1639-1640	40	5
158	The association between maternal-reported responses to infant crying at 4 weeks and 6 months and offspring depression at 18: a longitudinal study. *Archives of Womenks Mental Health*, **2016**, 19, 401-8	5	5
157	Getting the measure of outcomes in clinical practice. *Advances in Psychiatric Treatment*, **2014**, 20, 165-171		5
156	Mothers' own recollections of being parented and risk of offspring depression 18 years later: a prospective cohort study. *Depression and Anxiety*, **2014**, 31, 38-43	8.4	5
155	Common mental disorders in late pregnancy in women who wanted or attempted an abortion. *Psychological Medicine*, **2010**, 40, 1467-73	6.9	5
154	Does baseline fatigue influence treatment response to reboxetine or citalopram in depression? An open label randomized controlled trial. *Journal of Psychopharmacology*, **2012**, 26, 663-9	4.6	5
153	Detection of patient psychological distress and longitudinal patient-doctor relationships: a cross-sectional study. *British Journal of General Practice*, **2012**, 62, e167-73	1.6	5
152	Depression and public health. *International Review of Psychiatry*, **1996**, 8, 289-294	3.6	5
151	Phenotypic variation between parent-offspring trios and non-trios in genetic studies of schizophrenia. *Journal of Psychiatric Research*, **2006**, 40, 622-6	5.2	5
150	Epidemiology in community psychiatric research: common uses and methodological issues. *Epidemiologia E Psichiatria Sociale*, **2001**, 10, 18-26		5
149	Influence of theta-burst transcranial magnetic stimulation over the dorsolateral prefrontal cortex on emotion processing in healthy volunteers. *Cognitive, Affective and Behavioral Neuroscience*, **2020** , 20, 1278-1293	3.5	5
148	Is social support pre-treatment associated with prognosis for adults with depression in primary care?. *Acta Psychiatrica Scandinavica*, **2021**, 143, 392-405	6.5	5
147	Effective dose 50 method as the minimal clinically important difference: Evidence from depression trials. *Journal of Clinical Epidemiology*, **2021**, 137, 200-208	5.7	5

146	Risk of low Apgar scores and socioeconomic status over a 30-year period. *Journal of Maternal-Fetal and Neonatal Medicine*, **2014**, 27, 603-7	2	4
145	Evaluating the role of a galanin enhancer genotype on a range of metabolic, depressive and addictive phenotypes. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **2014**, 165B, 654-64	3.5	4
144	Reboxetine versus other antidepressive agents for depression. *The Cochrane Library*, **2009**,	5.2	4
143	New Evidence is Required. *British Journal of Psychiatry*, **1997**, 171, 227-227	5.4	4
142	Premorbid neuropsychological performance as a risk factor for schizophrenia: 13-Year follow-up of 50 000 conscripts. *Schizophrenia Research*, **1995**, 15, 114	3.6	4
141	The computer will see you now. *BMJ: British Medical Journal*, **1989**, 299, 138-9		4
140	Longitudinal population subgroups of CRP and risk of depression in the ALSPAC birth cohort. *Comprehensive Psychiatry*, **2020**, 96, 152143	7.3	4
139	Longitudinal associations between circulating interleukin-6 and C-reactive protein in childhood, and eating disorders and disordered eating in adolescence. *Brain, Behavior, and Immunity*, **2020**, 89, 491-500	16.6	4
138	Prediction of individuals at high risk of chronic kidney disease during treatment with lithium for bipolar disorder. *BMC Medicine*, **2021**, 19, 99	11.4	4
137	Network meta-analyses and treatment recommendations for obsessive-compulsive disorder - Authors' reply. *Lancet Psychiatry,the*, **2016**, 3, 921-922	23.3	4
136	The shrouded visibility of eating disorders research. *Lancet Psychiatry,the*, **2021**, 8, 91-92	23.3	4
135	The mental health of lesbian, gay, and bisexual adults compared with heterosexual adults: results of two nationally representative English household probability samples. *Psychological Medicine*, **2021**, 1-10	6.9	4
134	Identifying the Common Genetic Basis of Antidepressant Response. *Biological Psychiatry Global Open Science*, **2021**,		4
133	Validation of the Chinese Version of the Revised Clinical Interview Schedule: Findings from Hong Kong Mental Morbidity Survey. *East Asian Archives of Psychiatry*, **2017**, 27, 3-10	1.1	4
132	Newer-Generation Antidepressants and Suicide Risk. *Psychotherapy and Psychosomatics*, **2019**, 88, 371-372	9.4	3
131	Common mechanisms for type 2 diabetes and psychosis: Findings from a prospective birth cohort. *Schizophrenia Research*, **2020**, 223, 227-235	3.6	3
130	The role of gender inequalities in adolescent depression. *Lancet Psychiatry,the*, **2020**, 7, 471-472	23.3	3
129	Cognitive behavioural therapy for treatment-resistant depression - Authors' reply. *Lancet, The*, **2013**, 381, 1814-5	40	3

128	Pharmacological interventions for treatment-resistant depression in adults. *The Cochrane Library*, **2013**,	5.2	3
127	Behavioural therapies versus treatment as usual for depression. *The Cochrane Library*, **2010**,	5.2	3
126	Humanistic therapies versus treatment as usual for depression. *The Cochrane Library*, **2010**,	5.2	3
125	Psychiatric morbidity, social development and cognition, and later schizophrenia: Premorbid adjustment or prodrome?. *Schizophrenia Research*, **1997**, 24, 249	3.6	3
124	Can access to psychiatric health care explain regional differences in disability pension with psychiatric disorders?. *Social Psychiatry and Psychiatric Epidemiology*, **2007**, 42, 366-71	4.5	3
123	Mental health after head injury. *Journal of Neurology, Neurosurgery and Psychiatry*, **2001**, 71, 431	5.5	3
122	"You don't give me flowers anymore": an analysis of gift-giving to medical and psychiatric inpatients. *Social Psychiatry and Psychiatric Epidemiology*, **1999**, 34, 136-40	4.5	3
121	Catatonia: demographic, clinical and laboratory associations.. *Psychological Medicine*, **2021**, 1-11	6.9	3
120	Computerized assessments of psychiatric disorder using PROQSY: discussion paper. *Journal of the Royal Society of Medicine*, **1992**, 85, 403-6	2.3	3
119	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. *Programme Grants for Applied Research*, **2019**, 7, 1-108	1.5	3
118	Alcohol consumption and internalising disorders in young adults of ALSPAC: a population-based study. *Journal of Epidemiology and Community Health*, **2020**, 74, 1023-1027	5.1	3
117	Inhibitory control of positive and negative information and adolescent depressive symptoms: a population-based cohort study. *Psychological Medicine*, **2020**, 1-11	6.9	3
116	Prevalence of symptoms, ever having received a diagnosis and treatment of depression and anxiety, and associations with health service use amongst the general population in two Russian cities. *BMC Psychiatry*, **2020**, 20, 537	4.2	3
115	Association between circulating levels of C-reactive protein and positive and negative symptoms of psychosis in adolescents in a general population birth cohort. *Journal of Psychiatric Research*, **2021**, 143, 534-542	5.2	3
114	Device-measured sedentary behaviour and anxiety symptoms during adolescence: a 6-year prospective cohort study. *Psychological Medicine*, **2020**, 1-10	6.9	3
113	No evidence that CBT is less effective than antidepressants in moderate to severe depression. *Evidence-Based Mental Health*, **2016**, 19, 125	11.1	3
112	Assessment of the Role of IQ in Associations Between Population Density and Deprivation and Nonaffective Psychosis. *JAMA Psychiatry*, **2020**, 77, 729-736	14.5	3
111	Risk-taking to obtain reward: sex differences and associations with emotional and depressive symptoms in a nationally representative cohort of UK adolescents. *Psychological Medicine*, **2021**, 1-9	6.9	3

110	Longitudinal association between binge eating and metabolic syndrome in adults: Findings from the ELSA-Brasil cohort. *Acta Psychiatrica Scandinavica*, **2021**, 144, 464-474	6.5	3
109	Older adults respond better to psychological therapy than working-age adults: evidence from a large sample of mental health service attendees. *Journal of Affective Disorders*, **2021**, 294, 85-93	6.6	3
108	Materials used to support cognitive behavioural therapy for depression: a survey of therapists' clinical practice and views. *Cognitive Behaviour Therapy*, **2019**, 48, 463-481	4.4	2
107	Association between homicide rates and suicide rates: a countrywide longitudinal analysis of 5507 Brazilian municipalities. *BMJ Open*, **2020**, 10, e040069	3	2
106	Prenatal and childhood adversity and inflammation in children: A population-based longitudinal study. *Brain, Behavior, and Immunity*, **2020**, 87, 524-530	16.6	2
105	Paternal and maternal depression and offspring risk: additive effects or worse? - Authors' reply. *Lancet Psychiatry,the*, **2018**, 5, 108	23.3	2
104	Antidepressants During Pregnancy and Autism in Offspring: Population Based Cohort Study. *Obstetrical and Gynecological Survey*, **2017**, 72, 689-692	2.4	2
103	Authors reply to Davies and colleagues, Donnelly, and Pilling and Anderson. *BMJ, The*, **2012**, 345, e4500-e4500	5.9	2
102	Psychological therapies for treatment-resistant depression in adults. *The Cochrane Library*, **2013**,	5.2	2
101	Psychodynamic therapies versus other psychological therapies for depression. *The Cochrane Library* , **2010**,	5.2	2
100	Understanding emotion: lessons from anxiety. *Behavioral and Brain Sciences*, **2012**, 35, 145	0.9	2
99	Humanistic therapies versus other psychological therapies for depression. *The Cochrane Library*, **2010**, 2010,	5.2	2
98	Psychodynamic therapies versus treatment as usual for depression. *The Cochrane Library*, **2010**,	5.2	2
97	Interpersonal, cognitive analytic and other integrative therapies versus other psychological therapies for depression. *The Cochrane Library*, **2010**,	5.2	2
96	Strategies for the prevention of psychiatric disorder in primary care in south London. *Journal of Epidemiology and Community Health*, **1997**, 51, 304-9	5.1	2
95	Asphyxia at birth and schizophrenia. *British Journal of Psychiatry*, **2002**, 180, 465	5.4	2
94	Computers in primary care. *International Review of Psychiatry*, **1992**, 4, 307-310	3.6	2
93	A Patient Stratification Approach to Identifying the Likelihood of Continued Chronic Depression and Relapse Following Treatment for Depression.. *Journal of Personalized Medicine*, **2021**, 11,	3.6	2

92	Development and psychometric properties of the Carer - Head Injury Neurobehavioral Assessment Scale (C-HINAS) and the Carer - Head Injury Participation Scale (C-HIPS): patient and family determined outcome scales. *Neuropsychiatric Disease and Treatment*, **2007**, 3, 389-408	3.1	2
91	Development and psychometric properties of the Patient-Head Injury Participation Scale (P-HIPS) and the Patient-Head Injury Neurobehavioral Assessment Scale (P-HINAS): patient and family determined outcomes scales. *Neuropsychiatric Disease and Treatment*, **2007**, 3, 373-88	3.1	2
90	An evidence-based approach to the use of telehealth in long-term health conditions: development of an intervention and evaluation through pragmatic randomised controlled trials in patients with depression or raised cardiovascular risk. *Programme Grants for Applied Research*, **2017**, 5, 1-468	1.5	2
89	Testing the Independent and Joint Contribution of Exposure to Neurodevelopmental Adversity and Childhood Trauma to Risk of Psychotic Experiences in Adulthood. *Schizophrenia Bulletin*, **2021**, 47, 776-784	4.2	2
88	Cost-Effectiveness of Different Formats for Delivery of Cognitive Behavioral Therapy for Depression: A Systematic Review Based Economic Model. *Value in Health*, **2020**, 23, 1662-1670	3.3	2
87	Trends in method-specific suicide in Brazil from 2000 to 2017. *Social Psychiatry and Psychiatric Epidemiology*, **2021**, 56, 1779-1790	4.5	2
86	The importance of transdiagnostic symptom level assessment to understanding prognosis for depressed adults: analysis of data from six randomised control trials. *BMC Medicine*, **2021**, 19, 109	11.4	2
85	Associations between psychosis and visual acuity impairment: A systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*, **2021**, 144, 6-27	6.5	2
84	Further research needed for severe and chronic depression in specialist mental health services. *Lancet Psychiatry,the*, **2016**, 3, 792-3	23.3	2
83	Eighteen-year trajectories of depressive symptoms in mothers with a lifetime eating disorder: findings from the ALSPAC cohort. *British Journal of Psychiatry*, **2020**, 216, 90-96	5.4	2
82	The Role of Primary School Composition in the Trajectories of Internalising and Externalising Problems across Childhood and Adolescence. *Journal of Abnormal Child Psychology*, **2020**, 48, 197-211	4	2
81	Internalising symptoms mediate the longitudinal association between childhood inflammation and psychotic-like experiences in adulthood. *Schizophrenia Research*, **2020**, 215, 424-429	3.6	2
80	Suicide, self-harm and thoughts of suicide or self-harm in infectious disease epidemics: a systematic review and meta-analysis CORRIGENDUM. *Epidemiology and Psychiatric Sciences*, **2021**, 30,	5.1	2
79	Risk factors for complex posttraumatic stress disorder in UK police. *Occupational Medicine*, **2021**, 71, 351-357	2.1	2
78	Ability-grouping and problem behavior trajectories in childhood and adolescence: Results from a U.K. population-based sample. *Child Development*, **2021**,	4.9	2
77	Socioeconomic Indicators of Treatment Prognosis for Adults With Depression: A Systematic Review and Individual Patient Data Meta-analysis.. *JAMA Psychiatry*, **2022**,	14.5	2
76	An analysis of views about supported reduction or discontinuation of antipsychotic treatment among people with schizophrenia and other psychotic disorders.. *BMC Psychiatry*, **2022**, 22, 185	4.2	2
75	Identifying and treating hearing loss in mild cognitive impairment: a feasibility study for an interventional trial with hearing aids. *International Psychogeriatrics*, **2020**, 32, 537-538	3.4	1

74	Are refugees at increased risk of suicide compared with non-refugee migrants and the host populationAnna-Clara Hollander. *European Journal of Public Health*, **2017**, 27,	2.1	1
73	Network Meta-Analysis of the Relative Efficacy of Pharmacological and Psychological Interventions in Adults with Obsessive Compulsive Disorder. *Value in Health*, **2014**, 17, A454-5	3.3	1
72	Are multiple physical symptoms a poor prognostic factor or just a marker of depression severity? Secondary analysis of the GenPod trial. *Journal of Affective Disorders*, **2014**, 163, 40-6	6.6	1
71	. *Australian and New Zealand Journal of Psychiatry*, **2006**, 40, 548-553	2.6	1
70	Parental age difference and schizophrenia. *British Journal of Psychiatry*, **2004**, 184, 540; author reply 540-1	5.4	1
69	Obstetric complications and schizophrenia. *BMJ: British Medical Journal*, **1993**, 306, 268; author reply 268-9		1
68	Demise of the rotational training scheme. *Psychiatric Bulletin*, **1990**, 14, 681-681		1
67	Racism and psychiatry. *British Journal of Psychiatry*, **1991**, 158, 432-3	5.4	1
66	Effect of detection of depression in general practice. Study had methodological limitations. *BMJ: British Medical Journal*, **1996**, 312, 512-3		1
65	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. *Applied Health Economics and Health Policy*, **2021**, 1	3.4	1
64	Precursors and correlates of transient and persistent longitudinal profiles of psychotic experiences from late childhood through early adulthood.. *British Journal of Psychiatry*, **2021**, 1-9	5.4	1
63	Genetic variation in the Major Histocompatibility Complex and association with depression		1
62	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. *Wellcome Open Research*,4, 69	4.8	1
61	Trajectories of internalising and externalising symptoms and inflammation in the general child population. *Psychoneuroendocrinology*, **2020**, 118, 104723	5	1
60	Population-based needs assessment. *Current Opinion in Psychiatry*, **1999**, 12, 191-194	4.9	1
59	Risk taking to obtain reward: gender differences and associations with emotional and depressive symptoms in a nationally representative cohort of UK adolescents		1
58	Cost-Effectiveness of Sertraline in Primary Care According to Initial Severity and Duration of Depressive Symptoms: Findings from the PANDA RCT. *PharmacoEconomics - Open*, **2020**, 4, 427-438	2.1	1
57	Does anxiety moderate the effectiveness of mirtazapine in patients with treatment-resistant depression? A secondary analysis of the MIR trial. *Journal of Psychopharmacology*, **2020**, 34, 1342-1349	4.6	1

56	The effect of sertraline on emotional processing: secondary analyses of the PANDA randomised controlled trial. *Psychological Medicine*, **2021**, 1-8	6.9	1
55	The association between paternal depressogenic cognitive styles during pregnancy and offspring depressogenic cognitive styles: an 18-year prospective cohort study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, **2018**, 59, 604-614	7.9	1
54	Multimedia-delivered cognitive behavioural therapy versus face-to-face cognitive behavioural therapy for depression in adults. *The Cochrane Library*, **2018**,	5.2	1
53	Transcriptome-based polygenic score links depression-related corticolimbic gene expression changes to sex-specific brain morphology and depression risk. *Neuropsychopharmacology*, **2021**, 46, 2304-2311	8.7	1
52	Cognitive style and drinking to cope: A prospective cohort study. *Addiction*, **2021**,	4.6	1
51	P.3.19 The effects of cannabidiol on cerebral blood flow and its relationship to memory: an arterial spin labelling study. *European Neuropsychopharmacology*, **2019**, 29, S693	1.2	0
50	Re: Accuracy of general hospital dementia diagnoses in England: Sensitivity, specificity, and predictors of diagnostic accuracy 2008-2016. *Alzheimerks and Dementia*, **2019**, 15, 313-314	1.2	0
49	Depressive symptoms at age 9-13 and chronic disabling fatigue at age 16: A longitudinal study. *Journal of Adolescence*, **2019**, 75, 123-129	3.4	0
48	Republished research: Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. *British Journal of Sports Medicine*, **2013**, 47, 629-629	10.3	0
47	Mental health of UK Afghan and Iraq veterans. *Lancet, The*, **2010**, 375, 1758-60	40	0
46	Lithium augmentation in treatment-refractory unipolar depression. *British Journal of Psychiatry*, **2003**, 182, 456; author reply 456-7	5.4	0
45	Stigmatising views towards individuals with eating disorders: trends and associations from 1998 to 2008 using a repeated cross-sectional design.. *British Journal of Psychiatry*, **2021**, 1-7	5.4	0
44	Antidepressant medication to prevent depression relapse in primary care: the ANTLER RCT. *Health Technology Assessment*, **2021**, 25, 1-62	4.4	0
43	Life events and treatment prognosis for depression: A systematic review and individual patient data meta-analysis.. *Journal of Affective Disorders*, **2021**, 299, 298-308	6.6	0
42	Inflammatory and blood gas markers of COVID-19 delirium compared to non-COVID-19 delirium: a cross-sectional study. *Aging and Mental Health*, **2021**, 1-8	3.5	0
41	Sertraline in primary care: comments on the PANDA trial - Authors' reply. *Lancet Psychiatry,the*, **2020**, 7, 19	23.3	0
40	Recall bias during adolescence: Gender differences and associations with depressive symptoms. *Journal of Affective Disorders*, **2021**, 282, 299-307	6.6	0
39	Association of Emotion Regulation Trajectories in Childhood With Anorexia Nervosa and Atypical Anorexia Nervosa in Early Adolescence. *JAMA Psychiatry*, **2021**, 78, 1249-1257	14.5	0

38	Association between prenatal maternal infection and disordered eating behaviours in adolescence: a UK population-based prospective birth cohort study. *Psychological Medicine*, **2020**, 50, 927-935	6.9	0
37	Sex differences in depressive symptoms and tolerability after treatment with selective serotonin reuptake inhibitor antidepressants: Secondary analyses of the GENPOD trial. *Journal of Psychopharmacology*, **2021**, 35, 919-927	4.6	0
36	Clinical Neuroimaging Findings in Catatonia: Neuroradiological Reports of MRI Scans of Psychiatric Inpatients With and Without Catatonia.. *Journal of Neuropsychiatry and Clinical Neurosciences*, **2022**, appneuropsych2107	2.7	0
35	The Development and Internal Evaluation of a Predictive Model to Identify for Whom Mindfulness-Based Cognitive Therapy Offers Superior Relapse Prevention for Recurrent Depression Versus Maintenance Antidepressant Medication. *Clinical Psychological Science*,216770262210768	6	0
34	Review of assessment tools for psychotic-like experiences misses the psychosis-like symptoms semi-structured interview (PLIKSi) in ALSPAC. *Microbial Biotechnology*, **2017**, 11, 269-270	3.3	
33	Childhood interleukin-6, C-reactive protein and atopic disorders as risk factors for hypomanic symptoms in young adulthood: a longitudinal birth cohort study. *Psychological Medicine*, **2017**, 47, 984-986	6.9	
32	The influence of school in the development of eating disorders: a record-linkage study. *Lancet, The*, **2015**, 385 Suppl 1, S24	40	
31	Prescribing Anti-Depressants by Baseline Severity: Evidence Synthesis, Economic Model and Value of Information Analysis. *Value in Health*, **2014**, 17, A458	3.3	
30	P.2.a.021 Genome-wide association study of antidepressant-induced sexual dysfunction in depressed males. *European Neuropsychopharmacology*, **2014**, 24, S373-S374	1.2	
29	Meta-analysis of CYP2C19 association with efficacy and side effects of citalopram and escitalopram. *European Neuropsychopharmacology*, **2017**, 27, S582-S583	1.2	
28	OP24 Associations of cannabis and cigarette use with depression and anxiety at age 18: findings from the avon longitudinal study of parents and children. *Journal of Epidemiology and Community Health*, **2015**, 69, A19.2-A19	5.1	
27	PP37 Use of mendelian randomisation to assess whether smoking is causally associated with alcohol consumption. *Journal of Epidemiology and Community Health*, **2015**, 69, A68.2-A68	5.1	
26	Authors' reply to Males. *BMJ, The*, **2015**, 350, h3648	5.9	
25	Associations of Cannabis and Tobacco Use with Depression at Age 18: Findings from the Avon Longitudinal Study of Parents and Children.. *International Journal of Epidemiology*, **2015**, 44, i282-i283	7.8	
24	Commentary on Griffith-Lendering et al. (2013): cross-lagging cannabis and psychosis vulnerability. *Addiction*, **2013**, 108, 741-2	4.6	
23	P2-327 Physical activity and depression in adolescents: cross-sectional results from the ALSPAC cohort. *Journal of Epidemiology and Community Health*, **2011**, 65, A312-A313	5.1	
22	P.2.c.015 Efficacy of SSRIs in treating depression in patients with ischaemic heart disease, a meta-analysis. *European Neuropsychopharmacology*, **2009**, 19, S412	1.2	
21	Fetal Growth and Childhood Behavioural Problems. *American Journal of Epidemiology*, **2006**, 163, S18-S18	3.8	

20	Physical Activity and Common Mental Disorder in Men: The Caerphilly Cohort. *American Journal of Epidemiology*, **2006**, 163, S218-S218	3.8
19	Investigating the Association between Cigarette Smoking and Risk of Developing Schizophrenia in a Cohort Study. *Clinical Science*, **2003**, 104, 41P-41P	
18	Non-psychotic, non-major affective psychiatric disorders in adolescent boys were associated with greater schizophrenia risk. *Evidence-Based Mental Health*, **2002**, 5, 60	11.1
17	SES01.01 Developmental precursors of affective illness in a general population birth cohort. *European Psychiatry*, **2000**, 15, 213s-213s	6
16	FC13.06 The cross-national epidemiology of unexplained chronic fatigue. *European Psychiatry*, **2000** , 15, 313s-313s	6
15	Expert opinion. *Psychiatric Bulletin*, **1990**, 14, 124-124	
14	Fluoxetine Supplement; Journal September 1988, 153. *Psychiatric Bulletin*, **1989**, 13, 565-565	
13	You don't read my papers anymore an investigation into the use of Barbra Streisand song titles in the psychiatric literature. *Psychiatric Bulletin*, **2005**, 29, 35-35	
12	Unity is strength. Commentary on Research in the real world. *Psychiatric Bulletin*, **2006**, 30, 46-47	
11	From the Mental Patient to the Person. By P. Barham and R. Hayward. London: Tavistock/Routledge. 1991. 162 pp. £35.00.. *British Journal of Psychiatry*, **1993**, 162, 867-867	5.4
10	Introduction to Epidemiologic Research Methods1-8	
9	Reprint of: Internalising symptoms mediate the longitudinal association between childhood inflammation and psychotic-like experiences in adulthood. *Schizophrenia Research*, **2020**, 226, 24-29	3.6
8	Conditional Beliefs of Primary-Care Patients with Treatment-Resistant Depression. *Behavioural and Cognitive Psychotherapy*, **2016**, 44, 513-26	2.1
7	P.3.03 The acute effects of cannabidiol on emotional processing and anxiety: A neurocognitive imaging study. *European Neuropsychopharmacology*, **2019**, 29, S680	1.2
6	The Epidemiology of Mental Illness12-23	
5	70SEVERITY OF DEMENTIA AT DEATH: COHORT STUDY USING ROUTINELY COLLECTED SECONDARY CARE DATA. *Age and Ageing*, **2018**, 47, iii26-iii27	3
4	Reply to: To the editor: how statistics killed the cat E. F. Torrey, R. H Yolken. *Psychological Medicine*, **2018**, 48, 692-692	6.9
3	Closing the gap between research and practice. New evidence is required. *British Journal of Psychiatry*, **1997**, 171, 227	5.4

2	Local understandings of PTSD and complex PTSD among health professionals working with adolescents in violent neighbourhoods of São Paulo city, Brazil.. *BMC Psychiatry*, **2022**, 22, 196	4.2
1	An open letter to the President. *Psychiatric Bulletin*, **1988**, 12, 554-554	