Lie Zheng

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9853484/publications.pdf

Version: 2024-02-01

		933447	940533
19	524	10	16
papers	citations	h-index	g-index
19	19	19	639
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Efficacy of adjuvant EGFR inhibitors and impact of clinical factors in resected <i>EGFR </i> -mutated non-small-cell lung cancer: a meta-analysis. Future Oncology, 2022, 18, 1159-1169.	2.4	6
2	Clinical efficacy and safety of individualized pembrolizumab administration based on pharmacokinetic in advanced non-small cell lung cancer: An open-label, single-arm, exploratory clinical trial Journal of Clinical Oncology, 2022, 40, e21161-e21161.	1.6	O
3	Postoperative adjuvant hepatic arterial infusion chemotherapy (HAIC) with FOLFOX to improve outcomes of patients with hepatocellular carcinoma with microvascular invasion: A prospective multicenter, phase 3, randomized, controlled clinical trial Journal of Clinical Oncology, 2022, 40, 4013-4013.	1.6	1
4	Segmental resection is associated with decreased survival in patients with stage IA non-small cell lung cancer with a tumor size of 21–30 mm. Translational Lung Cancer Research, 2021, 10, 900-913.	2.8	7
5	Anti-PD-1 Immunotherapy Improves the Efficacy of Hepatic Artery Infusion Chemotherapy in Advanced Hepatocellular Carcinoma. Journal of Hepatocellular Carcinoma, 2021, Volume 8, 167-176.	3.7	30
6	Neoadjuvant transarterial infusion chemotherapy with FOLFOX could improve outcomes of resectable BCLC stage A/B hepatocellular carcinoma patients beyond Milan criteria: An interim analysis of a multi-center, phase 3, randomized, controlled clinical trial Journal of Clinical Oncology, 2021, 39, 4008-4008.	1.6	13
7	HN1 promotes tumor associated lymphangiogenesis and lymph node metastasis via NF-κB signaling activation in cervical carcinoma. Biochemical and Biophysical Research Communications, 2020, 530, 87-94.	2.1	10
8	Postoperative Adjuvant Transarterial Infusion Chemotherapy with FOLFOX Could Improve Outcomes of Hepatocellular Carcinoma Patients with Microvascular Invasion: A Preliminary Report of a Phase III, Randomized Controlled Clinical Trial. Annals of Surgical Oncology, 2020, 27, 5183-5190.	1.5	25
9	PET/CT alone versus PET/CT and MRI-guided therapy in patients with upper aerodigestive tract natural killer/T-cell lymphoma, nasal type: A prospective study Journal of Clinical Oncology, 2019, 37, 7537-7537.	1.6	0
10	Hepatic arterial infusion of oxaliplatin plus fluorouracil/leucovorin vs. sorafenib for advanced hepatocellular carcinoma. Journal of Hepatology, 2018, 69, 60-69.	3.7	153
11	FOXAI: a phase II trial evaluating the efficacy and safety of hepatic arterial infusion of oxaliplatin plus fluorouracil/leucovorin for advanced hepatocellular carcinoma. Gut, 2018, 67, 395.1-396.	12.1	68
12	Adjuvant transcatheter arterial chemoembolization after curative resection for hepatocellular carcinoma patients with solitary tumor and microvascular invasion: a randomized clinical trial of efficacy and safety. Cancer Communications, 2018, 38, 1-12.	9.2	92
13	Neoadjuvant chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in nasopharyngeal carcinoma patients with cervical nodal necrosis. Scientific Reports, 2017, 7, 42624.	3.3	14
14	Advantage of PET/CT in Target Delineation of MRI-negative Cervical Lymph Nodes In Intensity-Modulated Radiation Therapy Planning for Nasopharyngeal Carcinoma. Journal of Cancer, 2017, 8, 4117-4123.	2.5	10
15	Prognostic Value of Cavernous Sinus Invasion in Patients with Nasopharyngeal Carcinoma Treated with Intensity-Modulated Radiotherapy. PLoS ONE, 2016, 11, e0146787.	2.5	6
16	Small cell carcinoma of the rectum: A report of imaging results from four cases. Oncology Letters, 2016, 11, 1671-1676.	1.8	3
17	Elevated plasma fibrinogen level shows superior prognostic value than Epstein-Barr virus DNA load for stage IVA/B nasopharyngeal carcinoma patients in the intensity-modulated radiotherapy era. Oncotarget, 2016, 7, 46242-46252.	1.8	10
18	Prognostic value of plasma fibrinogen level and cervical nodal necrosis in stage IVA/B nasopharyngeal carcinoma patients who had positive cervical nodal metastasis Journal of Clinical Oncology, 2016, 34, 6039-6039.	1.6	0

#	Article	lF	CITATIONS
19	Prognostic Value of Cervical Nodal Necrosis in Nasopharyngeal Carcinoma: Analysis of 1800 Patients with Positive Cervical Nodal Metastasis at MR Imaging. Radiology, 2015, 276, 536-544.	7.3	76