

# Sharon Micallef

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/9854819/sharon-micallef-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

12  
papers

273  
citations

7  
h-index

14  
g-index

14  
ext. papers

511  
ext. citations

14.5  
avg, IF

1.9  
L-index

#	Paper	IF	Citations
12	Procalcitonin algorithm in critically ill adults with undifferentiated infection or suspected sepsis. A randomized controlled trial. <i>American Journal of Respiratory and Critical Care Medicine</i> , <b>2014</b> , 190, 1102-1110	10.2	130
11	Effect of 12 mg vs 6 mg of Dexamethasone on the Number of Days Alive Without Life Support in Adults With COVID-19 and Severe Hypoxemia: The COVID STEROID 2 Randomized Trial. <i>JAMA - Journal of the American Medical Association</i> , <b>2021</b> , 326, 1807-1817	27.4	46
10	Level of agreement on resuscitation decisions among hospital specialists and barriers to documenting do not attempt resuscitation (DNAR) orders in ward patients. <i>Resuscitation</i> , <b>2011</b> , 82, 815-818	4	31
9	Balanced Multielectrolyte Solution versus Saline in Critically Ill Adults.. <i>New England Journal of Medicine</i> , <b>2022</b> ,	59.2	18
8	Higher vs lower doses of dexamethasone in patients with COVID-19 and severe hypoxia (COVID STEROID 2) trial: Protocol and statistical analysis plan. <i>Acta Anaesthesiologica Scandinavica</i> , <b>2021</b> , 65, 834-845	1.9	12
7	Dexamethasone 12mg versus 6mg for patients with COVID-19 and severe hypoxaemia: a pre-planned, secondary Bayesian analysis of the COVID STEROID 2 trial. <i>Intensive Care Medicine</i> , <b>2021</b> , 48, 45	14.5	11
6	Higher vs Lower Doses of Dexamethasone in Patients with COVID-19 and Severe Hypoxia (COVID STEROID 2) trial: Protocol for a secondary Bayesian analysis. <i>Acta Anaesthesiologica Scandinavica</i> , <b>2021</b> , 65, 702-710	1.9	8
5	Health-related quality of life in survivors of septic shock: 6-month follow-up from the ADRENAL trial. <i>Intensive Care Medicine</i> , <b>2020</b> , 46, 1696-1706	14.5	7
4	Balanced Crystalloids versus Saline in Critically Ill Adults [A Systematic Review with Meta-Analysis] <b>2022</b> , 1,		6
3	Dexamethasone 12 mg versus 6 mg for patients with COVID-19 and severe hypoxia: an international, randomized, blinded trial		2
2	Balanced crystalloids compared to normal saline for fluid therapy in critically ill adult patients: Systematic review and meta-analysis protocol		1
1	Long-term outcomes of dexamethasone 12mg versus 6mg in patients with COVID-19 and severe hypoxaemia.. <i>Intensive Care Medicine</i> , <b>2022</b> , 1	14.5	1