

# Pål Berg-Hansen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9891270/publications.pdf>

Version: 2024-02-01

23  
papers

389  
citations

840776

11  
h-index

839539

18  
g-index

24  
all docs

24  
docs citations

24  
times ranked

717  
citing authors

#	ARTICLE	IF	CITATIONS
1	Stereotyped Bâ€cell responses are linked to IgG constant region polymorphisms in multiple sclerosis. <i>European Journal of Immunology</i> , 2022, 52, 550-565.	2.9	10
2	Sensor-based gait analyses of the six-minute walk test identify qualitative improvement in gait parameters of people with multiple sclerosis after rehabilitation. <i>Journal of Neurology</i> , 2022, 269, 3723-3734.	3.6	6
3	The influence of socioeconomic factors on access to disease modifying treatment in a Norwegian multiple sclerosis cohort. <i>Multiple Sclerosis and Related Disorders</i> , 2022, 61, 103759.	2.0	3
4	Exploring Retinal Blood Vessel Diameters as Biomarkers in Multiple Sclerosis. <i>Journal of Clinical Medicine</i> , 2022, 11, 3109.	2.4	3
5	Serum neurofilament light chain concentration predicts disease worsening in multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2022, 28, 1859-1870.	3.0	14
6	Fatigue in multiple sclerosis is associated with socioeconomic factors. <i>Multiple Sclerosis and Related Disorders</i> , 2022, 64, 103955.	2.0	5
7	The course of multiple sclerosis rewritten: a Norwegian population-based study on disease demographics and progression. <i>Journal of Neurology</i> , 2021, 268, 1330-1341.	3.6	17
8	High prevalence of fatigue in contemporary patients with multiple sclerosis. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2021, 7, 205521732199982.	1.0	18
9	Early High Efficacy Treatment in Multiple Sclerosis Is the Best Predictor of Future Disease Activity Over 1 and 2 Years in a Norwegian Population-Based Registry. <i>Frontiers in Neurology</i> , 2021, 12, 693017.	2.4	45
10	Maternal education has significant influence on progression in multiple sclerosis. <i>Multiple Sclerosis and Related Disorders</i> , 2021, 53, 103052.	2.0	6
11	No significant differences in absenteeism or academic achievements in a Norwegian multiple sclerosis case control study. <i>Multiple Sclerosis and Related Disorders</i> , 2021, 54, 103141.	2.0	2
12	Hospitalization following influenza infection and pandemic vaccination in multiple sclerosis patients: a nationwide population-based registry study from Norway. <i>European Journal of Epidemiology</i> , 2020, 35, 355-362.	5.7	22
13	Prevalence of multiple sclerosis in rural and urban districts in Telemark county, Norway. <i>Multiple Sclerosis and Related Disorders</i> , 2020, 45, 102352.	2.0	8
14	LesionQuant for Assessment of MRI in Multiple Sclerosisâ€”A Promising Supplement to the Visual Scan Inspection. <i>Frontiers in Neurology</i> , 2020, 11, 546744.	2.4	9
15	The diagnostic value of IgG index versus oligoclonal bands in cerebrospinal fluid of patients with multiple sclerosis. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2020, 6, 205521731990129.	1.0	18
16	No differential gene expression for CD4+ T cells of MS patients and healthy controls. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2019, 5, 205521731985690.	1.0	9
17	Quantitative proteomic analyses of CD4+ and CD8+ T cells reveal differentially expressed proteins in multiple sclerosis patients and healthy controls. <i>Clinical Proteomics</i> , 2019, 16, 19.	2.1	24
18	Restriction spectrum imaging of white matter and its relation to neurological disability in multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2019, 25, 687-698.	3.0	8

#	ARTICLE	IF	CITATIONS
19	Increased DNA methylation of SLFN12 in CD4+ and CD8+ T cells from multiple sclerosis patients. PLoS ONE, 2018, 13, e0206511.	2.5	37
20	Genetic variants are major determinants of CSF antibody levels in multiple sclerosis. Brain, 2015, 138, 632-643.	7.6	54
21	Prevalence of multiple sclerosis among immigrants in Norway. Multiple Sclerosis Journal, 2015, 21, 695-702.	3.0	43
22	Oligoclonal band phenotypes in MS differ in their HLA class II association, while specific KIR ligands at HLA class I show association to MS in general. Journal of Neuroimmunology, 2014, 274, 174-179.	2.3	7
23	Calprotectin levels in the cerebrospinal fluid reflect disease activity in multiple sclerosis. Journal of Neuroimmunology, 2009, 216, 98-102.	2.3	21