

Yifen Shi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9921038/publications.pdf>

Version: 2024-02-01

19
papers

611
citations

840776

11
h-index

752698

20
g-index

22
all docs

22
docs citations

22
times ranked

933
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Multiomics data identified <scp>TP53</scp> and <scp>LRP1B</scp> as key regulatory gene related to immune phenotypes via <scp>EPCAM</scp> in <scp>HCC</scp>. <i>Cancer Medicine</i> , 2022, 11, 2145-2158. | 2.8 | 2 |
| 2 | Iron overload adversely effects bone marrow haematogenesis via SIRT-SOD2-mROS in a process ameliorated by curcumin. <i>Cellular and Molecular Biology Letters</i> , 2021, 26, 2. | 7.0 | 12 |
| 3 | Long non-coding RNA-based signature for predicting prognosis of hepatocellular carcinoma. <i>Bioengineered</i> , 2021, 12, 673-681. | 3.2 | 10 |
| 4 | Activation-induced cytidine deaminase overexpression in double-hit lymphoma: potential target for novel anticancer therapy. <i>Scientific Reports</i> , 2020, 10, 14164. | 3.3 | 6 |
| 5 | CD9 knockdown suppresses cell proliferation, adhesion, migration and invasion, while promoting apoptosis and the efficacy of chemotherapeutic drugs and imatinib in Ph+ ALL SUPâ€B15 cells. <i>Molecular Medicine Reports</i> , 2020, 22, 2791-2800. | 2.4 | 1 |
| 6 | Peptide-immobilized starch/PEG sponge with rapid shape recovery and dual-function for both uncontrolled and noncompressible hemorrhage. <i>Acta Biomaterialia</i> , 2019, 99, 220-235. | 8.3 | 64 |
| 7 | Fabricating antimicrobial peptide-immobilized starch sponges for hemorrhage control and antibacterial treatment. <i>Carbohydrate Polymers</i> , 2019, 222, 115012. | 10.2 | 69 |
| 8 | Exosome secreted from adipose-derived stem cells attenuates diabetic nephropathy by promoting autophagy flux and inhibiting apoptosis in podocyte. <i>Stem Cell Research and Therapy</i> , 2019, 10, 95. | 5.5 | 211 |
| 9 | Unrelated Donor Peripheral Blood Stem Cell Transplantation for Patients with Î²-Thalassemia Major Based on a Novel Conditioning Regimen. <i>Biology of Blood and Marrow Transplantation</i> , 2019, 25, 1592-1596. | 2.0 | 22 |
| 10 | Benzene induces haematotoxicity by promoting deacetylation and autophagy. <i>Journal of Cellular and Molecular Medicine</i> , 2019, 23, 1022-1033. | 3.6 | 22 |
| 11 | Effect of initial body mass index on survival outcome of patients with myelodysplastic syndrome: a single-center retrospective study. <i>Leukemia and Lymphoma</i> , 2018, 59, 129-137. | 1.3 | 5 |
| 12 | Mean platelet volume predicts prognosis in patients with diffuse large Bâ€cell lymphoma. <i>Hematological Oncology</i> , 2018, 36, 104-109. | 1.7 | 13 |
| 13 | Benzene metabolite hydroquinone induces apoptosis of bone marrow mononuclear cells through inhibition of Î²-catenin signaling. <i>Toxicology in Vitro</i> , 2018, 46, 361-369. | 2.4 | 29 |
| 14 | Prognostic Significance of Pretreatment Neutrophil/Lymphocyte Ratio and Platelet/Lymphocyte Ratio in Patients with Diffuse Large B-Cell Lymphoma. <i>BioMed Research International</i> , 2018, 2018, 1-8. | 1.9 | 28 |
| 15 | lncRNA KRAL reverses 5-fluorouracil resistance in hepatocellular carcinoma cells by acting as a ceRNA against miR-141. <i>Cell Communication and Signaling</i> , 2018, 16, 47. | 6.5 | 62 |
| 16 | Maintenance therapy with all-trans retinoic acid and arsenic trioxide improves relapse-free survival in adults with low- to intermediate-risk acute promyelocytic leukemia who have achieved complete remission after consolidation therapy. <i>OncoTargets and Therapy</i> , 2017, Volume 10, 2305-2313. | 2.0 | 13 |
| 17 | Peripheral blood lymphocyte to monocyte ratio recovery from low levels at diagnosis after completion of first line therapy predicts good clinical outcomes in patients with diffuse large B-cell lymphoma. <i>Oncotarget</i> , 2017, 8, 19556-19565. | 1.8 | 12 |
| 18 | Prognostic significance of the red blood cell distribution width in diffuse large B-cell lymphoma patients. <i>Oncotarget</i> , 2017, 8, 40724-40731. | 1.8 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Histone Deacetylase Inhibitors Trichostatin A and MCP30 Relieve Benzene-Induced Hematotoxicity via Restoring Topoisomerase II α . PLoS ONE, 2016, 11, e0153330. | 2.5 | 9 |