

Davide Lombardi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9927761/publications.pdf>

Version: 2024-02-01

16
papers

296
citations

1307594

7
h-index

888059

17
g-index

17
all docs

17
docs citations

17
times ranked

623
citing authors

#	ARTICLE	IF	CITATIONS
1	Ultrasound of benign thyroid nodules: A 120 months follow-up study. <i>Clinical Endocrinology</i> , 2021, 94, 866-871.	2.4	4
2	KIR-HLA Functional Repertoire Influences Trastuzumab Efficiency in Patients With HER2-Positive Breast Cancer. <i>Frontiers in Immunology</i> , 2021, 12, 791958.	4.8	2
3	Ultrasound Microvascular Blood Flow Evaluation: A New Tool for the Management of Thyroid Nodule?. <i>International Journal of Endocrinology</i> , 2019, 2019, 1-6.	1.5	18
4	Adjuvant Tamoxifen Plus Ovarian Function Suppression Versus Tamoxifen Alone in Premenopausal Women With Early Breast Cancer: Patient-Reported Outcomes in the Suppression of Ovarian Function Trial. <i>Journal of Clinical Oncology</i> , 2016, 34, 1601-1610.	1.6	100
5	Anthracycline-free neoadjuvant therapy induces pathological complete responses by exploiting immune proficiency in HER2+ breast cancer patients. <i>BMC Cancer</i> , 2014, 14, 954.	2.6	9
6	Epirubicin and docetaxel as neoadjuvant treatment of hormone receptor positive, HER-2 negative breast cancer: findings from two successive phase II studies. <i>Radiology and Oncology</i> , 2013, 47, 57-62.	1.7	2
7	Lapatinib-Based Therapy in Heavily Pretreated HER2-Positive Metastatic Breast Cancer: A Single Institution Experience. <i>Tumori</i> , 2012, 98, 33-38.	1.1	4
8	A different immunologic profile characterizes patients with HER-2-overexpressing and HER-2-negative locally advanced breast cancer: implications for immune-based therapies. <i>Breast Cancer Research</i> , 2011, 13, R117.	5.0	30
9	Epirubicin and Docetaxel as Neoadjuvant Treatment of Locally Advanced Breast Cancer: A Phase II Study. <i>Tumori</i> , 2010, 96, 229-233.	1.1	3
10	Epirubicin and docetaxel as neoadjuvant treatment of locally advanced breast cancer: a phase II study. <i>Tumori</i> , 2010, 96, 229-33.	1.1	2
11	Incidence of Palmar-Plantar Erythrodysesthesia in Pretreated and Unpretreated Patients Receiving Pegylated Liposomal Doxorubicin. <i>Tumori</i> , 2009, 95, 687-690.	1.1	3
12	Selecting for BRCA1 testing using a combination of homogeneous selection criteria and immunohistochemical characteristics of breast cancers. <i>BMC Cancer</i> , 2009, 9, 360.	2.6	3
13	Familial breast cancer: characteristics and outcome of BRCA 1&2 positive and negative cases. <i>BMC Cancer</i> , 2005, 5, 70.	2.6	73
14	New oral drugs in older patients: a review of idarubicin in elderly patients. <i>Critical Reviews in Oncology/Hematology</i> , 2004, 49, 153-163.	4.4	29
15	Idarubicin. <i>Tumori</i> , 2002, 88, S73-S74.	1.1	2
16	Continuous Infusion Fluorouracil in the Management of Advanced Breast Cancer: A Phase II Study. <i>Tumori</i> , 2000, 86, 42-45.	1.1	10