

Yvonne Ingje Verkuil

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/995635/publications.pdf>

Version: 2024-02-01

33
papers

887
citations

687363

13
h-index

526287

27
g-index

35
all docs

35
docs citations

35
times ranked

1396
citing authors

#	ARTICLE	IF	CITATIONS
1	A supergene determines highly divergent male reproductive morphs in the ruff. <i>Nature Genetics</i> , 2016, 48, 79-83.	21.4	411
2	A global threats overview for Numeniini populations: synthesising expert knowledge for a group of declining migratory birds. <i>Bird Conservation International</i> , 2017, 27, 6-34.	1.3	87
3	Losing a staging area: Eastward redistribution of Afro-Eurasian ruffs is associated with deteriorating fuelling conditions along the western flyway. <i>Biological Conservation</i> , 2012, 149, 51-59.	4.1	46
4	A novel mitochondrial gene order in shorebirds (Scolopacidae, Charadriiformes). <i>Molecular Phylogenetics and Evolution</i> , 2010, 57, 411-416.	2.7	32
5	A global population redistribution in a migrant shorebird detected with continent-wide qualitative breeding survey data. <i>Diversity and Distributions</i> , 2011, 17, 144-151.	4.1	31
6	The interplay between habitat availability and population differentiation: a case study on genetic and morphological structure in an inland wader (Charadriiformes). <i>Biological Journal of the Linnean Society</i> , 2012, 106, 641-656.	1.6	22
7	New primers for molecular sex identification of waders. <i>Wader Study</i> , 2017, 124, .	0.4	21
8	Characterization of MHC class I in a long-distance migrant shorebird suggests multiple transcribed genes and intergenic recombination. <i>Immunogenetics</i> , 2013, 65, 211-225.	2.4	19
9	Prenatal Transfer of Gut Bacteria in Rock Pigeon. <i>Microorganisms</i> , 2020, 8, 61.	3.6	19
10	DNA metabarcoding quantifies the relative biomass of arthropod taxa in songbird diets: Validation with camera-recorded diets. <i>Ecology and Evolution</i> , 2022, 12, e8881.	1.9	19
11	Wind effects on prey availability: How northward migrating waders use brackish and hypersaline lagoons in the sivash, Ukraine. <i>Journal of Sea Research</i> , 1993, 31, 359-374.	1.0	18
12	Spring Migration of Ruffs (<i>Philomachus pugnax</i>) in Fryslân: Estimates of Staging Duration Using Resighting Data. <i>Ardea</i> , 2010, 98, 21-33.	0.6	16
13	Low genetic differentiation between Greenlandic and Siberian Sanderling populations implies a different phylogeographic history than found in Red Knots. <i>Journal of Ornithology</i> , 2016, 157, 325-332.	1.1	16
14	Biometrics of Ruffs (<i>Philomachus pugnax</i>) migrating in spring through southern Belarus with special emphasis on the occurrence of "faeders". <i>Ringling and Migration</i> , 2007, 23, 134-140.	0.4	13
15	Detection of earthworm prey by Ruff (<i>Philomachus pugnax</i>). <i>Ibis</i> , 2017, 159, 647-656.	1.9	11
16	When a typical jumper skips: itineraries and staging habitats used by Red Knots (<i>Calidris canutus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 1235-1251.	1.9	10
17	Fast fuelling but light flight in Broad-billed Sandpipers <i>Limicola falcinellus</i> : stopover ecology at a final take-off site in spring (Sivash, Ukraine). <i>Ibis</i> , 2006, 148, 211-220.	1.9	9
18	No evident spatial genetic structuring in the rapidly declining Black-tailed Godwit <i>Limosa limosa</i> in The Netherlands. <i>Conservation Genetics</i> , 2011, 12, 629-636.	1.5	9

#	ARTICLE	IF	CITATIONS
19	Non-breeding fÅ der RuffsPhilomachus pugnaxassociate according to sex, not morphology. Bird Study, 2008, 55, 241-246.	1.0	8
20	Genetic variation in nuclear and mitochondrial markers supports a large sex difference in lifetime reproductive skew in a lekking species. Ecology and Evolution, 2014, 4, 3626-3632.	1.9	8
21	Discovery of a morphologically and genetically distinct population of Black-tailed Godwits in the East Asian-Australasian Flyway. Ibis, 2021, 163, 448-462.	1.9	8
22	Apparent annual survival of staging ruffs during a period of population decline: insights from sex and site-use related differences. Population Ecology, 2015, 57, 613-624.	1.2	7
23	Size, shape and sex differences in three subspecies of Black-tailed Godwits <i>Limosa limosa</i>. Bird Study, 2020, 67, 45-52.	1.0	7
24	Global flyway evolution in red knots <i>Calidris canutus</i> and genetic evidence for a Nearctic refugium. Molecular Ecology, 2022, 31, 2124-2139.	3.9	7
25	Growth, maturity, and diet of the pearl whiplay (<i>Fontitrygon margaritella</i>) from the BijagÃ³s Archipelago, GuÃinea-Bissau. PeerJ, 2022, 10, e12894.	2.0	6
26	Use of Agricultural Fields by Ruffs Staging in Southwest Friesland in 2003-2013. Ardea, 2016, 104, 23-32.	0.6	5
27	Characterization of polymorphic microsatellite DNA markers in the black-tailed godwit (<i>Limosa) Tj ETQq1 1 0.784314 rgBT /Over 4.8 4	1.0	4
28	Molecular identification of temperate Cricetidae and Muridae rodent species using fecal samples collected in a natural habitat. Mammal Research, 2018, 63, 379-385.	1.3	3
29	Central-West Siberian-breeding Bar-tailed Godwits (<i>Limosa lapponica</i>) segregate in two morphologically distinct flyway populations. Ibis, 2022, 164, 468-485.	1.9	3
30	Within- and between-Year Variation in the Presence of Individually Marked Ruff Calidris pugnax at a Stopover Site during Northward Migration. Ardea, 2022, 110, .	0.6	3
31	Intertidal Flats of East and Southeast Asia. , 2018, , 1865-1874.		1
32	Intertidal Flats of East and Southeast Asia. , 2016, , 1-10.		1
33	British Ornithologistsâ€™ Union. Ibis, 2021, 163, 309-311.	1.9	0