

Megumu Yano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9997094/publications.pdf>

Version: 2024-02-01

15
papers

73
citations

1478505
6
h-index

1588992
8
g-index

15
all docs

15
docs citations

15
times ranked

55
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of serum orosomuroid with 30-min plasma glucose and glucose excursion during oral glucose tolerance tests in non-obese young Japanese women. <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000508.	2.8	14
2	Determinants and correlates of adipose tissue insulin resistance index in Japanese women without diabetes and obesity. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001686.	2.8	9
3	Relationship Between Dietary Variety and Frailty in Older Japanese Women During the Period of Restriction on Outings Due to COVID-19. <i>Journals of Gerontology - Series B Psychological Sciences and Social Sciences</i> , 2021, 76, e256-e262.	3.9	9
4	Reduced Birth Weight, Decreased Early-Phase Insulin Secretion, and Increased Glucose Concentrations after Oral Glucose Tolerance Test in Japanese Women Aged 20 Years with Family History of Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2020, 2020, 1-6.	2.3	7
5	Effects of 4G-beta-D-Galactosylsucrose in patients with depression: A randomized, double-blinded, placebo-controlled, parallel-group comparative study. <i>Journal of Psychiatric Research</i> , 2022, 148, 110-120.	3.1	7
6	Urinary sodium-to-potassium ratio and serum asymmetric dimethylarginine levels in patients with type 2 diabetes. <i>Hypertension Research</i> , 2018, 41, 913-922.	2.7	6
7	Higher circulating adiponectin and lower orosomuroid were associated with postload glucose ≥ 70 mg/dL, a possible inverse marker for dysglycemia, in young Japanese women. <i>BMJ Open Diabetes Research and Care</i> , 2019, 7, e000596.	2.8	4
8	Relationship Between Social Activity and Frailty in Japanese Older Women During Restriction on Outings due to COVID-19. <i>Canadian Geriatrics Journal</i> , 2021, 24, 320-324.	1.2	4
9	Elevated serum adiponectin and tumor necrosis factor- α and decreased transthyretin in Japanese elderly women with low grip strength and preserved muscle mass and insulin sensitivity. <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000537.	2.8	3
10	Higher circulating orosomuroid and lower early-phase insulin secretion in midlife Japanese with slower glucose disposal during oral glucose tolerance tests. <i>Diabetology International</i> , 2020, 11, 27-32.	1.4	3
11	Birth weight was associated positively with gluteofemoral fat mass and inversely with 2-h postglucose insulin concentrations, a marker of insulin resistance, in young normal-weight Japanese women. <i>Diabetology International</i> , 2022, 13, 375-380.	1.4	3
12	Association of family history of type 2 diabetes with blood pressure and resting heart rate in young normal weight Japanese women. <i>Diabetology International</i> , 2022, 13, 220-225.	1.4	2
13	Association of Age and Anemia With Adiponectin Serum Levels in Normal-Weight Japanese Women. <i>Journal of Clinical Medicine Research</i> , 2019, 11, 367-374.	1.2	2
14	Associations of serum transthyretin with triglyceride in non-obese elderly Japanese women independently of insulin resistance, HDL cholesterol, and adiponectin. <i>Diabetology International</i> , 2021, 12, 405-411.	1.4	0
15	Lower dietary variety is associated with worse sleep quality in community-dwelling elderly Japanese women. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2021, 30, 275-282.	0.4	0