

Participation in physical activity decreased more in people with rheumatoid arthritis than in the general population during the COVID-19 lockdown:

Rheumatology International

42, 241-250

DOI: [10.1007/s00296-021-05054-4](https://doi.org/10.1007/s00296-021-05054-4)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Identifying the Psychometric Properties of the Malay Version of the WHOQOL-BREF among Employees with Obesity Problem. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 7542.	1.2	2
2	Experiences of patients with rheumatoid arthritis during and after COVID-19-induced quarantine in terms of physical activity and health status: A qualitative study. <i>Journal of Nursing Management</i> , 0, , .	1.4	0
3	Social participation and mental health of immunocompromised individuals before and after COVID-19 vaccination—Results of a longitudinal observational study over three time points. <i>Frontiers in Psychiatry</i> , 0, 13, .	1.3	6
4	Physical activity in rheumatoid arthritis—“is it time to push the pace of change?”. <i>Rheumatology Advances in Practice</i> , 2022, 7, .	0.3	1
5	Online Delivery of the Individualized Reduction of Falls Intervention for Persons With Multiple Sclerosis Who Use a Wheelchair or Scooter Full-time: A Pilot Study. <i>International Journal of MS Care</i> , 2023, 25, 82-90.	0.4	2