

CITATION REPORT

List of articles citing

Okadaic acid mimics multiple changes in early protein phosphorylation and gene expression induced by tumor necrosis factor or interleukin-1

DOI: PM/1370482

Journal of Biological Chemistry, 1992, 267, 1846-52.

Source: <https://exaly.com/paper-pdf/126578217/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
97	Myosin II phosphorylation and the dynamics of stress fibers in serum-deprived and stimulated fibroblasts. <i>Molecular Biology of the Cell</i> , 1993 , 3, 1037-48	3.5	48
96	Okadaic acid as an inducer of the 78-kDa glucose-regulated protein in 9L rat brain tumor cells. <i>Journal of Cellular Biochemistry</i> , 1993 , 51, 91-101	4.7	15
95	Okadaic acid, a phosphatase inhibitor, enhances the phorbol ester-induced interleukin-1 beta expression via an AP-1-mediated mechanism. <i>Scandinavian Journal of Immunology</i> , 1993 , 38, 570-4	3.4	15
94	Transmembrane signaling in periodontal mesenchymal cells: the linkage between stimulus and response. <i>Periodontology 2000</i> , 1993 , 3, 76-98	12.9	4
93	Tumor necrosis factor activities and cancer therapy--a perspective. 1993 , 57, 79-128		87
92	Effect of okadaic acid on apo B and apo A-I secretion by CaCo-2 cells. <i>Lipids and Lipid Metabolism</i> , 1993 , 1168, 130-43		16
91	Inhibitors of protein phosphatase-1 and -2A; two of the major serine/threonine protein phosphatases involved in cellular regulation. <i>Current Opinion in Structural Biology</i> , 1993 , 3, 934-943	8.1	56
90	Cytoplasmic phospholipase A2 activity and gene expression are stimulated by tumor necrosis factor: dexamethasone blocks the induced synthesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993 , 90, 4475-9	11.5	187
89	Tumor promotion by inhibitors of protein phosphatases 1 and 2A: the okadaic acid class of compounds. <i>Advances in Cancer Research</i> , 1993 , 61, 143-94	5.9	235
88	Ethylene Signal Is Transduced via Protein Phosphorylation Events in Plants. <i>Plant Cell</i> , 1993 , 5, 523-530	11.6	148
87	Okadaic acid regulation of the retinoblastoma gene product is correlated with the inhibition of growth factor-induced cell proliferation in mouse fibroblasts. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993 , 90, 5460-3	11.5	18
86	Signal transduction by tumor necrosis factor mediated by JNK protein kinases. <i>Molecular and Cellular Biology</i> , 1994 , 14, 8376-84	4.8	433
85	Effects of cytokines on synthesis and function of the hepatic asialoglycoprotein receptor. <i>Journal of Cellular Physiology</i> , 1994 , 158, 527-34	7	22
84	Transforming growth factor beta 1 inhibits interleukin-1-induced but enhances ionomycin-induced interferon-gamma production in a T cell lymphoma: comparison with the effects of rapamycin. <i>Journal of Cellular Physiology</i> , 1994 , 160, 141-53	7	17
83	Proinflammatory cytokines (interleukin-1 beta and tumor necrosis factor-alpha) down regulate synthesis and secretion of thrombospondin by human endothelial cells. <i>Journal of Cellular Physiology</i> , 1994 , 160, 367-77	7	30
82	Analysis of cellular phosphoproteins by two-dimensional gel electrophoresis: applications for cell signaling in normal and cancer cells. <i>Electrophoresis</i> , 1994 , 15, 417-40	3.6	78
81	Involvement of a putative protein-tyrosine phosphatase and I kappa B-alpha serine phosphorylation in nuclear factor kappa B activation by tumor necrosis factor. <i>Journal of Biological Chemistry</i> , 1995 , 270, 18881-7	5.4	44

80	The protein phosphatase inhibitor calyculin A stimulates chemokine production by human synovial cells. <i>Biochemical Journal</i> , 1995 , 311 (Pt 1), 89-95	3.8	8
79	Interleukin-1 signal transduction. <i>International Journal of Clinical and Laboratory Research</i> , 1995 , 25, 169-77		35
78	IL-1 beta regulates rat mesangial cyclooxygenase II gene expression by tyrosine phosphorylation. <i>Kidney International</i> , 1995 , 47, 1354-63	9.9	30
77	Interleukin-1-induced IL-8 and IL-6 gene expression and production in human mesangial cells is differentially regulated by cAMP. <i>Kidney International</i> , 1995 , 48, 1767-77	9.9	26
76	Detection of interleukin-1 signal transduction inhibitors: action of protein kinase inhibitors. <i>International Journal of Immunopharmacology</i> , 1995 , 17, 887-94		1
75	Requirement of tyrosine- and serine/threonine kinases in the transcriptional activation of the mammalian grp78/BiP promoter by thapsigargin. <i>Journal of Biological Chemistry</i> , 1995 , 270, 494-502	5.4	44
74	Okadaic acid increases nerve growth factor secretion, mRNA stability, and gene transcription in primary cultures of cortical astrocytes. <i>Journal of Biological Chemistry</i> , 1995 , 270, 5994-9	5.4	44
73	Tumor necrosis factor alpha-induced phosphorylation of insulin receptor substrate-1 (IRS-1). Possible mechanism for suppression of insulin-stimulated tyrosine phosphorylation of IRS-1. <i>Journal of Biological Chemistry</i> , 1995 , 270, 23780-4	5.4	325
72	Towards an understanding of the signal transduction pathways for interleukin 1. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1995 , 1266, 31-44	4.9	95
71	Okadaic acid increased annexin I and induced differentiation of human promyelocytic leukemia cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1995 , 1266, 23-30	4.9	5
70	Protein phosphorylation in apoptosis. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1995 , 1269, 187-99	4.9	81
69	The mRNA overexpression of inflammatory enzymes, phospholipase A2 and cyclooxygenase, in the large bowel mucosa and neoplasms of F344 rats treated with naturally occurring carcinogen, 1-hydroxyanthraquinone. <i>Cancer Letters</i> , 1995 , 97, 75-82	9.9	16
68	Naturally-Derived Tumor Promoters and Inhibitors of Carcinogenesis. <i>Toxin Reviews</i> , 1996 , 15, 129-156		4
67	Interleukin-1 signal transduction. <i>Life Sciences</i> , 1996 , 59, 61-83	6.8	121
66	Okadaic acid-like toxin in systemic lupus erythematosus patients: hypothesis for toxin-induced pathology, immune dysregulation, and transactivation of herpesviruses. <i>Medical Hypotheses</i> , 1996 , 47, 217-25	3.8	1
65	Regulation of rat interstitial collagenase gene expression in growth cartilage and chondrocytes by vitamin D3, interleukin-1 and okadaic acid. <i>Journal of Cellular Biochemistry</i> , 1996 , 63, 395-409	4.7	21
64	T1/ST2 signaling establishes it as a member of an expanding interleukin-1 receptor family. <i>Journal of Biological Chemistry</i> , 1996 , 271, 5777-83	5.4	165
63	Inhibition of tumor necrosis factor signal transduction in endothelial cells by dimethylaminopurine. <i>Journal of Biological Chemistry</i> , 1996 , 271, 28624-9	5.4	28

62	Phosphorylation of protein-tyrosine phosphatase PTP-1B on identical sites suggests activation of a common signaling pathway during mitosis and stress response in mammalian cells. <i>Journal of Biological Chemistry</i> , 1997 , 272, 2957-62	5.4	38
61	Phosphorylation of alphaB-crystallin in response to various types of stress. <i>Journal of Biological Chemistry</i> , 1997 , 272, 29934-41	5.4	165
60	Interleukin-1-induced growth inhibition of human melanoma cells. Interleukin-1-induced antizyme expression is responsible for ornithine decarboxylase activity down-regulation. <i>Journal of Biological Chemistry</i> , 1997 , 272, 3376-83	5.4	28
59	Stimulation of interleukin-8 production by okadaic acid and vanadate in a human promyelocyte cell line, an HL-60 subline. Possible role of mitogen-activated protein kinase on the okadaic acid-induced NF-kappaB activation. <i>Journal of Biological Chemistry</i> , 1997 , 272, 15366-72	5.4	82
58	Tumor necrosis factor. <i>Growth Factors and Cytokines in Health and Disease</i> , 1997 , 2, 669-726		4
57	Tumor promotion and TNF- α gene expression by the okadaic acid class tumor promoters. <i>Journal of Environmental Science and Health, Part C: Environmental Carcinogenesis and Ecotoxicology Reviews</i> , 1997 , 15, 1-40	4.5	13
56	Effect of a protein phosphatase inhibitor, okadaic acid, on thrombomodulin expression in cultured human umbilical vein endothelial cells. <i>Thrombosis Research</i> , 1997 , 85, 169-76	8.2	1
55	Inhibition of protein tyrosine phosphatases causes phosphorylation of tyrosine-331 in the p60 TNF receptor and inactivates the receptor-associated kinase. <i>FEBS Letters</i> , 1997 , 410, 361-7	3.8	12
54	Tumor necrosis factor-alpha and interleukin-1 induce activation of MAP kinase and SAP kinase in human neuroma fibroblasts. <i>Neurochemistry International</i> , 1997 , 30, 401-10	4.4	48
53	Tumor-necrosis factor-alpha modulates mitogen-activated protein kinase activity of epidermal-growth-factor-stimulated MCF-7 breast cancer cells. <i>FEBS Journal</i> , 1997 , 249, 421-6		13
52	Multisite phosphorylation and the integration of stress signals at p53. <i>Cellular Signalling</i> , 1998 , 10, 159-66	6.9	171
51	Differential dose-dependent effects of epidermal growth factor on gene expression in A431 cells: evidence for a signal transduction pathway that can bypass Raf-1 activation. <i>Endocrinology</i> , 1998 , 139, 2382-91	4.8	19
50	H441 pulmonary epithelial cell mitogenic effects and signaling pathways in response to HGF and TGF-alpha. <i>Experimental Lung Research</i> , 1998 , 24, 27-39	2.3	18
49	Molecular dissection of nucleolin's role in growth and cell proliferation: new insights. <i>FASEB Journal</i> , 1999 , 13, 1911-1922	0.9	410
48	Ras mutation, irrespective of cell type and p53 status, determines a cell's destiny to undergo apoptosis by okadaic acid, an inhibitor of protein phosphatase 1 and 2A. <i>Molecular Pharmacology</i> , 1999 , 56, 515-25	4.3	44
47	Striking multiplicity of eIF4E-BP1 phosphorylated isoforms identified by 2D gel electrophoresis regulation by heat shock. <i>FEBS Journal</i> , 1999 , 265, 728-43		21
46	Tumour necrosis factor-alpha activates a calcium sensitization pathway in guinea-pig bronchial smooth muscle. <i>Journal of Physiology</i> , 1999 , 518 (Pt 2), 561-9	3.9	61
45	Regulation of protein kinase cascades by protein phosphatase 2A. <i>Trends in Biochemical Sciences</i> , 1999 , 24, 186-91	10.3	689

44	Transformation is associated with an increase in sensitivity to TNF-mediated lysis as a result of an increase in TNF-induced protein tyrosine phosphatase activity. <i>International Journal of Cancer</i> , 1999 , 81, 141-7	7.5	3
43	Okadaic acid stimulates the expression of vascular endothelial growth factor gene. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 265, 584-8	3.4	8
42	Induction of protein phosphatase type 2A in response to disruption of cell-matrix interactions. <i>Journal of Cellular Physiology</i> , 2000 , 182, 88-96	7	7
41	Effects of serine/threonine protein phosphatases on ion channels in excitable membranes. <i>Physiological Reviews</i> , 2000 , 80, 173-210	47.9	229
40	IL-1 beta protects human chondrocytes from CD95-induced apoptosis. <i>Journal of Immunology</i> , 2000 , 164, 2233-9	5.3	60
39	Phosphorylation of 130- and 95-kDa substrates associated with tumor necrosis factor-alpha receptor CD120a (p55). <i>Journal of Biological Chemistry</i> , 2000 , 275, 793-800	5.4	2
38	Tumor necrosis factor-induced microtubule stabilization mediated by hyperphosphorylated oncoprotein 18 promotes cell death. <i>Journal of Biological Chemistry</i> , 2000 , 275, 33876-82	5.4	28
37	Role of serine/threonine protein phosphatase 2A in cancer. <i>Cancer Letters</i> , 2001 , 170, 1-13	9.9	155
36	EGF receptor crosstalks with cytokine receptors leading to the activation of c-Jun kinase in response to UV irradiation in human keratinocytes. <i>Cellular Signalling</i> , 2001 , 13, 139-44	4.9	55
35	Phosphorylation-independent association of CXCR2 with the protein phosphatase 2A core enzyme. <i>Journal of Biological Chemistry</i> , 2001 , 276, 16960-8	5.4	51
34	Tumor necrosis factor-induced modulation of glyoxalase I activities through phosphorylation by PKA results in cell death and is accompanied by the formation of a specific methylglyoxal-derived AGE. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 949-54	11.5	86
33	A role for protein phosphatase-2A in p38 mitogen-activated protein kinase-mediated regulation of the c-Jun NH(2)-terminal kinase pathway in human neutrophils. <i>Journal of Biological Chemistry</i> , 2002 , 277, 40687-96	5.4	74
32	Isolation and synthesis of TNF-alpha release inhibitors from Fijian kava (Piper methysticum). <i>Phytomedicine</i> , 2003 , 10, 309-17	6.5	46
31	Identification of nucleolin as an AU-rich element binding protein involved in bcl-2 mRNA stabilization. <i>Journal of Biological Chemistry</i> , 2004 , 279, 10855-63	5.4	132
30	Increased AP-1 and NF-kappaB activation and recruitment with the combination of the proinflammatory cytokines IL-1beta, tumor necrosis factor alpha and IL-17 in rheumatoid synoviocytes. <i>Arthritis Research</i> , 2004 , 6, R190-8		58
29	Reversal of stathmin-mediated resistance to paclitaxel and vinblastine in human breast carcinoma cells. <i>Molecular Pharmacology</i> , 2007 , 71, 1233-40	4.3	85
28	Hepatic gene expression changes in mice associated with prolonged sublethal microcystin exposure. <i>Toxicologic Pathology</i> , 2007 , 35, 594-605	2.1	44
27	Intoxication alimentaire par les phycotoxines marines. <i>EMC - Médecine D Urgence</i> , 2007 , 2, 1-9		

26	Activation of nuclear factor kappa B in human neuroblastoma cell lines. <i>Journal of Neurochemistry</i> , 1994 , 62, 1716-26	6	21
25	Chronic microcystin exposure induces hepatocyte proliferation with increased expression of mitotic and cyclin-associated genes in P53-deficient mice. <i>Toxicologic Pathology</i> , 2008 , 36, 190-203	2.1	28
24	p53 Plays an important role in cell fate determination after exposure to microcystin-LR. <i>Environmental Health Perspectives</i> , 2010 , 118, 1292-8	8.4	44
23	The Use of Proteomics in the Study of Molecular Responses and Toxicity Pathways in Biological Systems. <i>Advances in Molecular Toxicology</i> , 2011 , 5, 45-109	0.4	2
22	Genotoxicity and potential carcinogenicity of cyanobacterial toxins - a review. <i>Mutation Research - Reviews in Mutation Research</i> , 2011 , 727, 16-41	7	225
21	Time-course analysis of injured skeletal muscle suggests a critical involvement of ERK1/2 signaling in the acute inflammatory response. <i>Muscle and Nerve</i> , 2012 , 45, 552-61	3.4	10
20	Oxidative stress induces inactivation of protein phosphatase 2A, promoting proinflammatory NF- κ B in aged rat kidney. <i>Free Radical Biology and Medicine</i> , 2013 , 61, 206-17	7.8	43
19	Tumor promoters: from chemicals to inflammatory proteins. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 1603-14	4.9	41
18	Microcystin-LR induces anoikis resistance to the hepatocyte uptake transporter OATP1B3-expressing cell lines. <i>Toxicology</i> , 2014 , 326, 53-61	4.4	3
17	Stathmin involvement in the maternal embryonic leucine zipper kinase pathway in glioblastoma. <i>Proteome Science</i> , 2016 , 14, 6	2.6	11
16	Phorbol esters in seed oil of <i>Jatropha curcas</i> L. (sabooodam in Thai) and their association with cancer prevention: from the initial investigation to the present topics. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017 , 143, 1359-1369	4.9	8
15	The concept of the okadaic acid class of tumor promoters is revived in endogenous protein inhibitors of protein phosphatase 2A, SET and CIP2A, in human cancers. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018 , 144, 2339-2349	4.9	20
14	Expression and phosphorylation of mammalian small heat shock proteins. <i>Progress in Molecular and Subcellular Biology</i> , 2002 , 28, 129-50	3	19
13	Dephosphorylation of the small heat shock protein Hsp27 in vivo by protein phosphatase 2A.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 9176-9183	5.4	81
12	Tumor necrosis factor-induced activation and increased tyrosine phosphorylation of mitogen-activated protein (MAP) kinase in human fibroblasts.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 18994-18999	5.4	105
11	Protein phosphatase inhibitors induce the sustained expression of the Egr-1 gene and the hyperphosphorylation of its gene product.. <i>Journal of Biological Chemistry</i> , 1992 , 267, 12991-12997	5.4	51
10	Inhibitors of serine/threonine phosphatases enhance phosphorylation of the interferon-gamma receptor while selectively attenuating interferon-gamma-induced gene expression in human peripheral-blood monocytes. <i>Biochemical Journal</i> , 1994 , 299 (Pt 3), 799-803	3.8	6
9	Activation of Protein Kinases and the Inactivation of Protein Phosphatase 2A in Tumour Necrosis Factor and Interleukin-1 Signal-Transduction Pathways. <i>FEBS Journal</i> , 1995 , 229, 503-511		34

8	Interleukin-1beta expression after inhibition of protein phosphatases in endotoxin-tolerant cells. <i>Vaccine Journal</i> , 1998 , 5, 281-7		15
7	Signal transduction by tumor necrosis factor mediated by JNK protein kinases. <i>Molecular and Cellular Biology</i> , 1994 , 14, 8376-8384	4.8	138
6	Cell-specific regulation of type II phospholipase A2 expression in rat mesangial cells. <i>Journal of Clinical Investigation</i> , 1993 , 92, 2524-32	15.9	28
5	Interleukin 1 alpha causes rapid activation of cytosolic phospholipase A2 by phosphorylation in rat mesangial cells. <i>Journal of Clinical Investigation</i> , 1994 , 93, 1224-33	15.9	93
4	Tumor necrosis factor-alpha inhibits stem cell factor-induced proliferation of human bone marrow progenitor cells in vitro. Role of p55 and p75 tumor necrosis factor receptors. <i>Journal of Clinical Investigation</i> , 1994 , 94, 165-72	15.9	49
3	Stathmin 1 in normal and malignant hematopoiesis. <i>BMB Reports</i> , 2014 , 47, 660-5	5.5	30
2	Encyclopedia of Cancer. 2014 , 1-3		
1	Encyclopedia of Cancer. 2017 , 3186-3188		