

CITATION REPORT

List of articles citing

A simple and efficient procedure for transformation of yeasts

DOI: PM/1503765
BioTechniques, 1992, 13, 18-20.

Source: <https://exaly.com/paper-pdf/126627443/citation-report.pdf>

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
525	CDC46/MCM5, a yeast protein whose subcellular localization is cell cycle-regulated, is involved in DNA replication at autonomously replicating sequences. 1992 , 89, 10459-63		87
524	Cloning and manipulation of the Schizosaccharomyces pombe his7+ gene as a new selectable marker for molecular genetic studies. 1993 , 24, 491-5		67
523	Parameters affecting lithium acetate-mediated transformation of Saccharomyces cerevisiae and development of a rapid and simplified procedure. 1993 , 24, 455-9		124
522	Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. 1993 , 73, 1197-206		969
521	Alteration of the Saccharomyces cerevisiae COX2 mRNA 5' untranslated leader by mitochondrial gene replacement and functional interaction with the translational activator protein PET111. 1993 , 4, 1327-35		73
520	Metabolic instability and constitutive endocytosis of STE6, the α -factor transporter of Saccharomyces cerevisiae. 1994 , 5, 1185-98		96
519	T7 RNA polymerase-dependent expression of COXII in yeast mitochondria. <i>Molecular and Cellular Biology</i> , 1994 , 14, 4643-52	4.8	37
518	Reduced but accurate translation from a mutant AUA initiation codon in the mitochondrial COX2 mRNA of Saccharomyces cerevisiae. 1994 , 242, 383-90		40
517	Isolation of an Arabidopsis thaliana casein kinase II beta subunit by complementation in Saccharomyces cerevisiae. 1994 , 25, 649-58		41
516	Phosphoribosylpyrophosphate synthetase (PRS): a new gene family in Saccharomyces cerevisiae. 1994 , 10, 1031-44		37
515	Effect of upstream sequences of the ADH1 promoter on the expression of Hormoconis resinae glucoamylase P by Saccharomyces cerevisiae. 1994 , 121, 229-35		3
514	Structure and heterologous expression of the Ustilago maydis viral toxin KP4. 1994 , 11, 155-64		54
513	TEL+CEN antagonism on plasmids involves telomere repeat sequences tracts and gene products that interact with chromosomal telomeres. 1994 , 103, 237-50		18
512	Expression of functional Raphanus sativus antifungal protein in yeast. 1994 , 348, 228-32		24
511	Glucose repression of yeast mitochondrial transcription: kinetics of derepression and role of nuclear genes. <i>Molecular and Cellular Biology</i> , 1994 , 14, 1160-70	4.8	75
510	The peroxisome proliferator-activated receptor interacts with the retinoid X receptor in vivo. 1994 , 148, 327-30		66
509	Cell cycle-dependent transcription of CLN2 is conferred by multiple distinct cis-acting regulatory elements. <i>Molecular and Cellular Biology</i> , 1994 , 14, 4788-801	4.8	70

508	Interactions between DNA-bound trimers of the yeast heat shock factor. <i>Molecular and Cellular Biology</i> , 1994 , 14, 501-8	4.8	74
507	Mutations in the <i>Saccharomyces cerevisiae</i> CDC1 gene affect double-strand-break-induced intrachromosomal recombination. <i>Molecular and Cellular Biology</i> , 1994 , 14, 8037-50	4.8	19
506	Interactions among three proteins that specifically activate translation of the mitochondrial COX3 mRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994 , 14, 1045-53	4.8	66
505	A novel role for DNA photolyase: binding to DNA damaged by drugs is associated with enhanced cytotoxicity in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994 , 14, 8071-7	4.8	46
504	The mitochondrial receptor complex: the small subunit Mom8b/Isp6 supports association of receptors with the general insertion pore and transfer of preproteins. <i>Molecular and Cellular Biology</i> , 1995 , 15, 6196-205	4.8	107
503	KIN28 encodes a C-terminal domain kinase that controls mRNA transcription in <i>Saccharomyces cerevisiae</i> but lacks cyclin-dependent kinase-activating kinase (CAK) activity. <i>Molecular and Cellular Biology</i> , 1995 , 15, 2983-92	4.8	197
502	TFIIIB placement on a yeast U6 RNA gene in vivo is directed primarily by TFIIIC rather than by sequence-specific DNA contacts. <i>Molecular and Cellular Biology</i> , 1995 , 15, 1455-66	4.8	65
501	Genetic and biochemical interactions between an essential kinetochore protein, Cbf2p/Ndc10p, and the CDC34 ubiquitin-conjugating enzyme. <i>Molecular and Cellular Biology</i> , 1995 , 15, 4835-42	4.8	29
500	Endoproteolytic processing of recombinant proalbumin variants by the yeast Kex2 protease. <i>Biochemical Journal</i> , 1995 , 308 (Pt 1), 321-5	3.8	18
499	Regulation of nuclear genes encoding mitochondrial proteins in <i>Saccharomyces cerevisiae</i> . 1995 , 177, 6836-43		18
498	Molecular cloning of the plasma membrane H(+)-ATPase from <i>Kluyveromyces lactis</i> : a single nucleotide substitution in the gene confers ethidium bromide resistance and deficiency in K ⁺ uptake. 1995 , 177, 2360-7		20
497	Enhanced plasmin inhibition by a reactive center lysine mutant of the Kunitz-type protease inhibitor domain of the amyloid beta-protein precursor. <i>Journal of Biological Chemistry</i> , 1995 , 270, 22827-30	5.4	20
496	Isolation of <i>Schizosaccharomyces pombe</i> isopentenyl diphosphate isomerase cDNA clones by complementation and synthesis of the enzyme in <i>Escherichia coli</i> . <i>Journal of Biological Chemistry</i> , 1995 , 270, 11298-303	5.4	31
495	High level expression, partial purification, and functional reconstitution of the human AE1 anion exchanger in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1995 , 270, 21028-34	5.4	32
494	A Ca ²⁺ -binding chimera of human lysozyme and bovine alpha-lactalbumin that can form a molten globule. <i>Journal of Biological Chemistry</i> , 1995 , 270, 10514-24	5.4	29
493	Mutated yeast heat shock transcription factor exhibits elevated basal transcriptional activation and confers metal resistance. <i>Journal of Biological Chemistry</i> , 1995 , 270, 25079-86	5.4	49
492	The cytoplasmic tail domain of the vacuolar protein sorting receptor Vps10p and a subset of VPS gene products regulate receptor stability, function, and localization. 1995 , 6, 1089-102		140
491	Identification of the yeast nuclear gene for the mitochondrial homologue of bacterial ribosomal protein L16. 1995 , 23, 3673-7		13

490	Glucose-dependent turnover of the mRNAs encoding succinate dehydrogenase peptides in <i>Saccharomyces cerevisiae</i> : sequence elements in the 5' untranslated region of the Ip mRNA play a dominant role. 1995 , 6, 1125-43		66
489	Inducible expression of a heterologous protein in <i>Hansenula polymorpha</i> using the alcohol oxidase 1 promoter of <i>Pichia pastoris</i> . 1996 , 177, 163-7		30
488	Crystal structure and mutational analysis of the human CDK2 kinase complex with cell cycle-regulatory protein CksHs1. 1996 , 84, 863-74		216
487	A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. 1996 , 87, 391-404		805
486	Initiation of glycogen synthesis in yeast. Requirement of multiple tyrosine residues for function of the self-glucosylating Glg proteins in vivo. <i>Journal of Biological Chemistry</i> , 1996 , 271, 26554-60	5-4	23
485	Activation and repression by nuclear hormone receptors: hormone modulates an equilibrium between active and repressive states. <i>Molecular and Cellular Biology</i> , 1996 , 16, 3807-13	4-8	120
484	Identification and analysis of a functional human homolog of the SPT4 gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996 , 16, 2848-56	4-8	32
483	Human protein Sam68 relocalization and interaction with poliovirus RNA polymerase in infected cells. 1996 , 93, 2296-301		140
482	cDNA cloning and characterisation of a maize homologue of the MCM proteins required for the initiation of DNA replication. 1996 , 252, 125-36		31
481	Isolation and characterization of SUG2. A novel ATPase family component of the yeast 26 S proteasome. <i>Journal of Biological Chemistry</i> , 1996 , 271, 32810-7	5-4	56
480	Localization of a yeast early Golgi mannosyltransferase, Och1p, involves retrograde transport. 1996 , 132, 985-98		137
479	Assembly of the ER to Golgi SNARE complex requires Uso1p. 1996 , 132, 755-67		177
478	BED1, a gene encoding a galactosyltransferase homologue, is required for polarized growth and efficient bud emergence in <i>Saccharomyces cerevisiae</i> . 1996 , 132, 137-51		36
477	Isolation and characterization of cDNAs corresponding to two human calcium, calmodulin-regulated, 3',5'-cyclic nucleotide phosphodiesterases. <i>Journal of Biological Chemistry</i> , 1996 , 271, 796-806	5-4	102
476	Heterologous complementation of a Rieske iron-sulfur protein-deficient <i>Saccharomyces cerevisiae</i> by the Rip1 gene of <i>Schizosaccharomyces pombe</i> . <i>Journal of Biological Chemistry</i> , 1996 , 271, 15341-5	5-4	21
475	Two classes of plant cDNA clones differentially complement yeast calcineurin mutants and increase salt tolerance of wild-type yeast. <i>Journal of Biological Chemistry</i> , 1996 , 271, 12859-66	5-4	146
474	<i>Saccharomyces cerevisiae</i> homologs of mammalian B and B α subunits of protein phosphatase 2A direct the enzyme to distinct cellular functions. <i>Journal of Biological Chemistry</i> , 1997 , 272, 8256-62	5-4	87
473	Endosome to Golgi retrieval of the vacuolar protein sorting receptor, Vps10p, requires the function of the VPS29, VPS30, and VPS35 gene products. 1997 , 137, 79-92		325

472	A novel membrane-associated metalloprotease, Ste24p, is required for the first step of NH2-terminal processing of the yeast a-factor precursor. 1997 , 136, 271-85		133
471	Genomic footprinting of Mig1p in the MAL62 promoter. Binding is dependent upon carbon source and competitive with the Mal63p activator. <i>Journal of Biological Chemistry</i> , 1997 , 272, 4613-22	5-4	17
470	Sec31 encodes an essential component of the COPII coat required for transport vesicle budding from the endoplasmic reticulum. 1997 , 8, 205-17		127
469	Characterization of a novel yeast SNARE protein implicated in Golgi retrograde traffic. 1997 , 8, 2659-76		96
468	Functional analysis of ribosomal protein L2 in yeast mitochondria. <i>Journal of Biological Chemistry</i> , 1997 , 272, 8165-71	5-4	14
467	Mutations in the CYS4 gene provide evidence for regulation of the yeast vacuolar H ⁺ -ATPase by oxidation and reduction in vivo. <i>Journal of Biological Chemistry</i> , 1997 , 272, 28149-57	5-4	46
466	The unfolded protein response coordinates the production of endoplasmic reticulum protein and endoplasmic reticulum membrane. 1997 , 8, 1805-14		318
465	Biogenesis of the <i>Saccharomyces cerevisiae</i> mating pheromone a-factor. 1997 , 136, 251-69		112
464	A genetic system for Archaea of the genus <i>Methanosarcina</i> : liposome-mediated transformation and construction of shuttle vectors. 1997 , 94, 2626-31		195
463	A lesion in the DNA replication initiation factor Mcm10 induces pausing of elongation forks through chromosomal replication origins in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997 , 17, 3261-71	4.8	129
462	Evidence that Spt3 functionally interacts with Mot1, TFIIA, and TATA-binding protein to confer promoter-specific transcriptional control in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997 , 17, 287-95	4.8	101
461	Stress proteins on the yeast cell surface determine resistance to osmotin, a plant antifungal protein. 1997 , 94, 7082-7		120
460	Modification of seed oil content and acyl composition in the brassicaceae by expression of a yeast sn-2 acyltransferase gene. 1997 , 9, 909-23		246
459	The GCN4 leucine zipper can functionally substitute for the heat shock transcription factor α trimerization domain. 1997 , 273, 61-74		28
458	A new family of K ⁺ transporters from <i>Arabidopsis</i> that are conserved across phyla. 1997 , 415, 206-11		135
457	Na ⁺ -K ⁺ -ATPase alpha-subunit containing Q905-V930 of gastric H ⁺ -K ⁺ -ATPase alpha preferentially assembles with H ⁺ -K ⁺ -ATPase beta. 1997 , 272, C923-30		36
456	Identification of the FKS1 gene of <i>Candida albicans</i> as the essential target of 1,3-beta-D-glucan synthase inhibitors. 1997 , 41, 2471-9		261
455	Molecular genetic and biochemical analysis of <i>Brassica napus</i> proliferating cell nuclear antigen function. 1997 , 34, 693-700		2

454	Two syntaxin homologues in the TGN/endosomal system of yeast. <i>EMBO Journal</i> , 1998 , 17, 113-26	13	217
453	Accumulation of mitochondrially synthesized <i>Saccharomyces cerevisiae</i> Cox2p and Cox3p depends on targeting information in untranslated portions of their mRNAs. <i>EMBO Journal</i> , 1998 , 17, 5796-804	13	96
452	Hus1p, a conserved fission yeast checkpoint protein, interacts with Rad1p and is phosphorylated in response to DNA damage. <i>EMBO Journal</i> , 1998 , 17, 2055-66	13	104
451	<i>Kluyveromyces lactis</i> SEF1 and its <i>Saccharomyces cerevisiae</i> homologue bypass the unknown essential function, but not the mitochondrial RNase P function, of the <i>S. cerevisiae</i> RPM2 gene. 1998 , 14, 77-87		19
450	Identification and analysis of homologues of <i>Saccharomyces cerevisiae</i> Spt3 suggest conserved functional domains. 1998 , 14, 409-17		6
449	An improved protocol for the preparation of yeast cells for transformation by electroporation. 1998 , 14, 565-71		156
448	Cloning of the <i>Candida albicans</i> nucleoside transporter by complementation of nucleoside transport-deficient <i>Saccharomyces</i> . 1998 , 14, 1257-1265		8
447	Two dehydration-inducible transcripts from the resurrection plant <i>Craterostigma plantagineum</i> encode interacting homeodomain-leucine zipper proteins. <i>Plant Journal</i> , 1998 , 15, 413-21	6.9	79
446	Control of mRNA turnover as a mechanism of glucose repression in <i>Saccharomyces cerevisiae</i> . 1998 , 30, 1175-93		42
445	Osmotin, a plant antifungal protein, subverts signal transduction to enhance fungal cell susceptibility. 1998 , 1, 807-17		101
444	Functional identification of a fatty acid delta5 desaturase gene from <i>Caenorhabditis elegans</i> . 1998 , 439, 215-8		69
443	Functional characterization of missense mutations in ATP7B: Wilson disease mutation or normal variant?. 1998 , 63, 1663-74		150
442	Characterization of a human homologue of the <i>Saccharomyces cerevisiae</i> transcription factor spt3 (SUPT3H). 1998 , 53, 90-6		16
441	Expression in <i>Saccharomyces cerevisiae</i> of antigenically and enzymatically active recombinant glutamic acid decarboxylase. 1998 , 61, 57-68		16
440	New alleles of the yeast MPS1 gene reveal multiple requirements in spindle pole body duplication. 1998 , 9, 759-74		53
439	The syntaxin Tlg1p mediates trafficking of chitin synthase III to polarized growth sites in yeast. 1998 , 9, 3383-97		101
438	cut11(+): A gene required for cell cycle-dependent spindle pole body anchoring in the nuclear envelope and bipolar spindle formation in <i>Schizosaccharomyces pombe</i> . 1998 , 9, 2839-55		137
437	Ste6p mutants defective in exit from the endoplasmic reticulum (ER) reveal aspects of an ER quality control pathway in <i>Saccharomyces cerevisiae</i> . 1998 , 9, 2767-84		99

436	Valine 904, tyrosine 898, and cysteine 908 in Na,K-ATPase alpha subunits are important for assembly with beta subunits. <i>Journal of Biological Chemistry</i> , 1998 , 273, 29400-5	5-4	17
435	Yeast Clk-1 homologue (Coq7/Cat5) is a mitochondrial protein in coenzyme Q synthesis. <i>Journal of Biological Chemistry</i> , 1998 , 273, 3351-7	5-4	109
434	A novel calmodulin-regulated Ca ²⁺ -ATPase (ACA2) from Arabidopsis with an N-terminal autoinhibitory domain. <i>Journal of Biological Chemistry</i> , 1998 , 273, 1099-106	5-4	128
433	AtKUP1: an Arabidopsis gene encoding high-affinity potassium transport activity. 1998 , 10, 51-62		285
432	Identification and characterization of an unusual double serine/threonine protein phosphatase 2C in the malaria parasite Plasmodium falciparum. <i>Journal of Biological Chemistry</i> , 1998 , 273, 11241-7	5-4	40
431	Sec35p, a novel peripheral membrane protein, is required for ER to Golgi vesicle docking. 1998 , 141, 1107-19		127
430	Characterization of a temperature-sensitive yeast vacuolar ATPase mutant with defects in actin distribution and bud morphology. <i>Journal of Biological Chemistry</i> , 1998 , 273, 18470-80	5-4	34
429	Isolation of a Delta5-fatty acid desaturase gene from <i>Mortierella alpina</i> . <i>Journal of Biological Chemistry</i> , 1998 , 273, 19055-9	5-4	134
428	Dual roles for Ste24p in yeast a-factor maturation: NH ₂ -terminal proteolysis and COOH-terminal CAAX processing. 1998 , 142, 635-49		113
427	Iqg1p, a yeast homologue of the mammalian IQGAPs, mediates cdc42p effects on the actin cytoskeleton. 1998 , 142, 443-55		89
426	The <i>Saccharomyces cerevisiae</i> prenylcysteine carboxyl methyltransferase Ste14p is in the endoplasmic reticulum membrane. 1998 , 9, 2231-47		95
425	imp2, a new component of the actin ring in the fission yeast <i>Schizosaccharomyces pombe</i> . 1998 , 143, 415-27		60
424	An essential domain within Cdc34p is required for binding to a complex containing Cdc4p and Cdc53p in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998 , 273, 4040-5	5-4	48
423	Mammalian prenylcysteine carboxyl methyltransferase is in the endoplasmic reticulum. <i>Journal of Biological Chemistry</i> , 1998 , 273, 15030-4	5-4	234
422	Role for the ubiquitin-proteasome system in the vacuolar degradation of Ste6p, the a-factor transporter in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998 , 18, 779-89	4-8	61
421	Identification and analysis of Mot3, a zinc finger protein that binds to the retrotransposon Ty long terminal repeat (delta) in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998 , 18, 1879-90	4-8	22
420	Mutations in the yeast KEX2 gene cause a Vma(-)-like phenotype: a possible role for the Kex2 endoprotease in vacuolar acidification. <i>Molecular and Cellular Biology</i> , 1998 , 18, 1534-43	4-8	30
419	A membrane coat complex essential for endosome-to-Golgi retrograde transport in yeast. 1998 , 142, 665-81		553

418	Identification of a <i>Caenorhabditis elegans</i> Delta6-fatty-acid-desaturase by heterologous expression in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 1998 , 330 (Pt 2), 611-4	3.8	133
417	RNA polymerase-specific nucleosome disruption by transcription in vivo. <i>Journal of Biological Chemistry</i> , 1999 , 274, 16431-6	5.4	20
416	The <i>aroC</i> gene of <i>Aspergillus nidulans</i> codes for a monofunctional, allosterically regulated chorismate mutase. <i>Journal of Biological Chemistry</i> , 1999 , 274, 22275-82	5.4	23
415	A novel family of yeast chaperons involved in the distribution of V-ATPase and other membrane proteins. <i>Journal of Biological Chemistry</i> , 1999 , 274, 26885-93	5.4	70
414	A <i>Selaginella lepidophylla</i> trehalose-6-phosphate synthase complements growth and stress-tolerance defects in a yeast <i>tps1</i> mutant. 1999 , 119, 1473-82		138
413	Identification of a calmodulin-regulated Ca ²⁺ -ATPase in the endoplasmic reticulum. 1999 , 119, 1165-76		120
412	S-Adenosylmethionine-dependent methylation in <i>Saccharomyces cerevisiae</i> . Identification of a novel protein arginine methyltransferase. <i>Journal of Biological Chemistry</i> , 1999 , 274, 814-24	5.4	181
411	Sec34p, a protein required for vesicle tethering to the yeast Golgi apparatus, is in a complex with Sec35p. 1999 , 147, 729-42		118
410	A protein phosphatase 2C gene, <i>LjNPP2C1</i> , from <i>Lotus japonicus</i> induced during root nodule development. 1999 , 96, 1738-43		26
409	Ser-534 in the hinge 1 region of <i>Arabidopsis</i> nitrate reductase is conditionally required for binding of 14-3-3 proteins and in vitro inhibition. <i>Journal of Biological Chemistry</i> , 1999 , 274, 4160-5	5.4	37
408	Apyrase functions in plant phosphate nutrition and mobilizes phosphate from extracellular ATP. 1999 , 119, 543-52		93
407	Role of the copper-binding domain in the copper transport function of ATP7B, the P-type ATPase defective in Wilson disease. <i>Journal of Biological Chemistry</i> , 1999 , 274, 12408-13	5.4	127
406	Genetic analysis and enzyme activity suggest the existence of more than one minimal functional unit capable of synthesizing phosphoribosyl pyrophosphate in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999 , 274, 12480-7	5.4	21
405	Yeast SMF1 mediates H(+)-coupled iron uptake with concomitant uncoupled cation currents. <i>Journal of Biological Chemistry</i> , 1999 , 274, 35089-94	5.4	125
404	The <i>Arabidopsis thaliana</i> TAG1 mutant has a mutation in a diacylglycerol acyltransferase gene. <i>Plant Journal</i> , 1999 , 19, 645-53	6.9	285
403	Development of a yeast-based assay system for monitoring microsatellite instability. 1999 , 176, 205-12		1
402	Induction of autophagy and inhibition of tumorigenesis by beclin 1. 1999 , 402, 672-6		2674
401	Cohabitation of insulators and silencing elements in yeast subtelomeric regions. <i>EMBO Journal</i> , 1999 , 18, 2522-37	13	190

400	Overexpression of the gene for Rpb7 subunit of yeast RNA polymerase II rescues the phenotypes associated with absence of the largest, nonessential subunit Rpb4. 1999 , 78, 149-156		13
399	Isolation of a <i>Histoplasma capsulatum</i> cDNA that complements a mitochondrial NAD(+)-isocitrate dehydrogenase subunit I-deficient mutant of <i>Saccharomyces cerevisiae</i> . 1999 , 15, 799-804		2
398	N-terminal chimeric constructs improve the expression of sarcoplasmic reticulum Ca(2+)-ATPase in yeast. 1999 , 1461, 83-95		10
397	Functional characterization of <i>rpn3</i> uncovers a distinct 19S proteasomal subunit requirement for ubiquitin-dependent proteolysis of cell cycle regulatory proteins in budding yeast. <i>Molecular and Cellular Biology</i> , 1999 , 19, 6872-90	4.8	38
396	The abundance of cell cycle regulatory protein Cdc4p is controlled by interactions between its F box and Skp1p. <i>Molecular and Cellular Biology</i> , 1999 , 19, 1759-67	4.8	45
395	Three yeast proteins related to the human candidate tumor suppressor p33(ING1) are associated with histone acetyltransferase activities. <i>Molecular and Cellular Biology</i> , 2000 , 20, 3807-16	4.8	133
394	In vivo targeting of a sunflower oil body protein in yeast secretory (sec) mutants. <i>Plant Journal</i> , 2000 , 23, 159-70	6.9	44
393	Crystal structure and mutational analysis of the <i>Saccharomyces cerevisiae</i> cell cycle regulatory protein Cks1: implications for domain swapping, anion binding and protein interactions. 2000 , 8, 841-50		33
392	Modulation at a distance of proton conductance through the <i>Saccharomyces cerevisiae</i> mitochondrial F1F0-ATP synthase by variants of the oligomycin sensitivity-conferring protein containing substitutions near the C-terminus. 2000 , 32, 595-607		7
391	Hexose transporters of tomato: molecular cloning, expression analysis and functional characterization. 2000 , 44, 687-97		43
390	Overexpression of <i>afpc10+</i> in fission yeast can suppress the temperature sensitivity of <i>fnuc2-663</i> mutant but not its sterility. 2000 , 34, 684-690		1
389	Characterization of a foldase, protein disulfide isomerase A, in the protein secretory pathway of <i>Aspergillus niger</i> . 2000 , 66, 775-82		85
388	Isolation and functional expression of human COQ3, a gene encoding a methyltransferase required for ubiquinone biosynthesis. <i>Journal of Biological Chemistry</i> , 2000 , 275, 12381-7	5.4	67
387	The family of SMF metal ion transporters in yeast cells. <i>Journal of Biological Chemistry</i> , 2000 , 275, 33388-94	5.4	102
386	Tea2p is a kinesin-like protein required to generate polarized growth in fission yeast. 2000 , 151, 15-28		116
385	PNG1, a yeast gene encoding a highly conserved peptide:N-glycanase. 2000 , 149, 1039-52		186
384	Mutational analysis of subunit G (Vma10p) of the yeast vacuolar H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2000 , 275, 37232-9	5.4	44
383	Coevolution of transcriptional and allosteric regulation at the chorismate metabolic branch point of <i>Saccharomyces cerevisiae</i> . 2000 , 97, 13585-90		18

382	Altered distribution of the yeast plasma membrane H ⁺ -ATPase as a feature of vacuolar H ⁺ -ATPase null mutants. <i>Journal of Biological Chemistry</i> , 2000 , 275, 40088-95	5-4	25
381	A role for ectophosphatase in xenobiotic resistance. 2000 , 12, 519-33		132
380	HARO7 encodes chorismate mutase of the methylotrophic yeast <i>Hansenula polymorpha</i> and is derepressed upon methanol utilization. 2000 , 182, 4188-97		23
379	The interaction of nitric oxide (NO) with the yeast transcription factor Ace1: A model system for NO-protein thiol interactions with implications to metal metabolism. 2000 , 97, 2491-6		54
378	Reconstitution of the Ste24p-dependent N-terminal proteolytic step in yeast a-factor biogenesis. <i>Journal of Biological Chemistry</i> , 2000 , 275, 6227-33	5-4	47
377	Geranylgeranyltransferase I of <i>Candida albicans</i> : null mutants or enzyme inhibitors produce unexpected phenotypes. 2000 , 182, 704-13		35
376	The abundance of Met30p limits SCF(Met30p) complex activity and is regulated by methionine availability. <i>Molecular and Cellular Biology</i> , 2000 , 20, 7845-52	4.8	26
375	Identification of a calmodulin-regulated soybean Ca(2 ⁺)-ATPase (SCA1) that is located in the plasma membrane. 2000 , 12, 1393-407		88
374	Effects of nitric oxide on the copper-responsive transcription factor Ace1 in <i>Saccharomyces cerevisiae</i> : cytotoxic and cytoprotective actions of nitric oxide. 2000 , 377, 296-303		28
373	Chimeras of Delta6-fatty acid and Delta8-sphingolipid desaturases. 2000 , 279, 779-85		38
372	Cloning and expression of cDNAs encoding plant V-ATPase subunits in the corresponding yeast null mutants. 2000 , 1459, 489-98		11
371	Functional identification of a delta8-sphingolipid desaturase from <i>Borago officinalis</i> . 2001 , 388, 293-8		48
370	Regulation of hisHF transcription of <i>Aspergillus nidulans</i> by adenine and amino acid limitation. 2001 , 32, 21-31		11
369	The <i>Candida albicans</i> gene encoding the cytoplasmic leucyl-tRNA synthetase: implications for the evolution of CUG codon reassignment. 2001 , 275, 133-40		5
368	AtBS14a and AtBS14b, two Bet1/Sft1-like SNAREs from <i>Arabidopsis thaliana</i> that complement mutations in the yeast SFT1 gene. 2001 , 500, 177-82		17
367	Features of V-ATPases that distinguish them from F-ATPases. 2001 , 504, 223-8		22
366	Efficient large scale purification of his-tagged proton translocating NADH:ubiquinone oxidoreductase (complex I) from the strictly aerobic yeast <i>Yarrowia lipolytica</i> . 2001 , 1504, 363-70		72
365	Identification of functional regions of Cbp3p, an enzyme-specific chaperone required for the assembly of ubiquinol-cytochrome c reductase in yeast mitochondria. 2001 , 1506, 103-16		10

364	Vesicular monoamine transporters heterologously expressed in the yeast <i>Saccharomyces cerevisiae</i> display high-affinity tetrabenazine binding. 2001 , 1510, 426-41		17
363	Ouabain and substrate affinities of human Na(+)-K(+)-ATPase alpha(1)beta(1), alpha(2)beta(1), and alpha(3)beta(1) when expressed separately in yeast cells. 2001 , 281, C1355-64		62
362	Nodule-specific regulation of phosphatidylinositol transfer protein expression in <i>Lotus japonicus</i> . 2001 , 13, 1369-82		51
361	Screening of cadmium-responsive genes in <i>Arabidopsis thaliana</i> . 2001 , 24, 1177-1188		121
360	Resistance to the plant PR-5 protein osmotin in the model fungus <i>Saccharomyces cerevisiae</i> is mediated by the regulatory effects of SSD1 on cell wall composition. <i>Plant Journal</i> , 2001 , 25, 271-80	6.9	48
359	<i>Arabidopsis</i> polyamine biosynthesis: absence of ornithine decarboxylase and the mechanism of arginine decarboxylase activity. <i>Plant Journal</i> , 2001 , 27, 551-60	6.9	204
358	Dsl1p, an essential protein required for membrane traffic at the endoplasmic reticulum/Golgi interface in yeast. 2001 , 2, 212-31		36
357	Seryl-tRNA synthetase is not responsible for the evolution of CUG codon reassignment in <i>Candida albicans</i> . 2001 , 18, 313-22		6
356	The role of haustoria in sugar supply during infection of broad bean by the rust fungus <i>Uromyces fabae</i> . 2001 , 98, 8133-8		277
355	Selective formation of Sed5p-containing SNARE complexes is mediated by combinatorial binding interactions. 2001 , 12, 521-38		51
354	MGA2 is involved in the low-oxygen response element-dependent hypoxic induction of genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001 , 21, 6161-9	4.8	77
353	Identification and characterization of a low oxygen response element involved in the hypoxic induction of a family of <i>Saccharomyces cerevisiae</i> genes. Implications for the conservation of oxygen sensing in eukaryotes. <i>Journal of Biological Chemistry</i> , 2001 , 276, 14374-84	5.4	85
352	A defect in coenzyme Q biosynthesis is responsible for the respiratory deficiency in <i>Saccharomyces cerevisiae</i> abc1 mutants. <i>Journal of Biological Chemistry</i> , 2001 , 276, 18161-8	5.4	99
351	Herpes simplex virus virion host shutoff protein requires a mammalian factor for efficient in vitro endoribonuclease activity. 2001 , 75, 1172-85		26
350	New roles for the Snp1 and Exo84 proteins in yeast pre-mRNA splicing. <i>Journal of Biological Chemistry</i> , 2001 , 276, 31004-15	5.4	23
349	Pho23 is associated with the Rpd3 histone deacetylase and is required for its normal function in regulation of gene expression and silencing in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001 , 276, 24068-74	5.4	53
348	Vps26p, a component of retromer, directs the interactions of Vps35p in endosome-to-Golgi retrieval. 2001 , 12, 3242-56		85
347	Rad23 provides a link between the Png1 deglycosylating enzyme and the 26 S proteasome in yeast. <i>Journal of Biological Chemistry</i> , 2001 , 276, 21601-7	5.4	98

346	Structure-function analysis of fission yeast Hus1-Rad1-Rad9 checkpoint complex. 2001 , 12, 3744-58		43
345	Two related kinesins, klp5+ and klp6+, foster microtubule disassembly and are required for meiosis in fission yeast. 2001 , 12, 3919-32		117
344	The coatamer-interacting protein Dsl1p is required for Golgi-to-endoplasmic reticulum retrieval in yeast. <i>Journal of Biological Chemistry</i> , 2001 , 276, 39150-60	5-4	79
343	pkl1(+)and klp2(+): Two kinesins of the Kar3 subfamily in fission yeast perform different functions in both mitosis and meiosis. 2001 , 12, 3476-88		96
342	The yeast a-factor transporter Ste6p, a member of the ABC superfamily, couples ATP hydrolysis to pheromone export. <i>Journal of Biological Chemistry</i> , 2001 , 276, 29007-11	5-4	30
341	Characterization of the promoter of PRS1 in <i>Saccharomyces cerevisiae</i> identifies three regions potentially involved in control of expression. 2001 , 183, 795-9		0
340	Requirement of the N-terminal extension for vacuolar trafficking and transport activity of yeast Ycf1p, an ATP-binding cassette transporter. 2002 , 13, 4443-55		53
339	Analysis of RGS proteins in <i>Saccharomyces cerevisiae</i> . 2002 , 344, 617-31		67
338	Mutational analysis of the subunit C (Vma5p) of the yeast vacuolar H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2002 , 277, 8979-88	5-4	49
337	A <i>Saccharomyces cerevisiae</i> gene required for heterologous fatty acid elongase activity encodes a microsomal beta-keto-reductase. <i>Journal of Biological Chemistry</i> , 2002 , 277, 11481-8	5-4	69
336	Rice phosphate transporters include an evolutionarily divergent gene specifically activated in arbuscular mycorrhizal symbiosis. 2002 , 99, 13324-9		490
335	Mga2p processing by hypoxia and unsaturated fatty acids in <i>Saccharomyces cerevisiae</i> : impact on LORE-dependent gene expression. 2002 , 1, 481-90		42
334	Functional dissection of a Rice Dr1/DrAp1 transcriptional repression complex. 2002 , 14, 181-95		17
333	The Sec34/Sec35p complex, a Ypt1p effector required for retrograde intra-Golgi trafficking, interacts with Golgi SNAREs and COPI vesicle coat proteins. 2002 , 157, 631-43		183
332	Identification of the functional domains of yeast sorting nexins Vps5p and Vps17p. 2002 , 13, 2826-40		82
331	Novel vacuolar H ⁺ -ATPase complexes resulting from overproduction of Vma5p and Vma13p. <i>Journal of Biological Chemistry</i> , 2002 , 277, 2716-24	5-4	41
330	Host cell factor-1 interacts with and antagonizes transactivation by the cell cycle regulatory factor Miz-1. <i>Journal of Biological Chemistry</i> , 2002 , 277, 46799-808	5-4	25
329	Glycopeptide export from the endoplasmic reticulum into cytosol is mediated by a mechanism distinct from that for export of misfolded glycoprotein. 2002 , 12, 803-11		12

328	The Ran GTPase system in fission yeast affects microtubules and cytokinesis in cells that are competent for nucleocytoplasmic protein transport. <i>Molecular and Cellular Biology</i> , 2002 , 22, 8491-505	4.8	28
327	Identification of modulators of mammalian G-protein signaling by functional screens in the yeast <i>Saccharomyces cerevisiae</i> . 2002 , 344, 153-68		11
326	Truncation of <i>Arabidopsis thaliana</i> and <i>Selaginella lepidophylla</i> trehalose-6-phosphate synthase unlocks high catalytic activity and supports high trehalose levels on expression in yeast. <i>Biochemical Journal</i> , 2002 , 366, 63-71	3.8	66
325	Soybean ascorbate peroxidase suppresses Bax-induced apoptosis in yeast by inhibiting oxygen radical generation. 2002 , 290, 457-62		58
324	Identification of a cDNA encoding a novel C18-Delta(9) polyunsaturated fatty acid-specific elongating activity from the docosahexaenoic acid (DHA)-producing microalga, <i>Isochrysis galbana</i> . 2002 , 510, 159-65		103
323	Replication of a linear mini-chromosome with terminal inverted repeats from the <i>Kluyveromyces lactis</i> linear DNA plasmid k2 in the cytoplasm of <i>Saccharomyces cerevisiae</i> . 2002 , 48, 13-23		4
322	Differential expression and function of <i>Arabidopsis thaliana</i> NHX Na ⁺ /H ⁺ antiporters in the salt stress response. <i>Plant Journal</i> , 2002 , 30, 529-39	6.9	404
321	Characterization of five novel dehydration-responsive homeodomain leucine zipper genes from the resurrection plant <i>Craterostigma plantagineum</i> . 2002 , 49, 601-10		74
320	Functional characterisation of two cytochrome b5-fusion desaturases from <i>Anemone leveillei</i> : the unexpected identification of a fatty acid Delta6-desaturase. 2003 , 217, 983-92		19
319	The upstream regulator, Rsr1p, and downstream effectors, Gic1p and Gic2p, of the Cdc42p small GTPase coordinately regulate initiation of budding in <i>Saccharomyces cerevisiae</i> . 2003 , 8, 235-50		24
318	The dihydroceramide desaturase is not essential for cell viability in <i>Schizosaccharomyces pombe</i> . 2003 , 538, 192-6		14
317	Identification of primula fatty acid delta 6-desaturases with n-3 substrate preferences. 2003 , 542, 100-4		63
316	The diverse bacterial origins of the <i>Arabidopsis</i> polyamine biosynthetic pathway. 2003 , 549, 26-30		60
315	Overproduction of polypeptides corresponding to the amino terminus of the F-box proteins Cdc4p and Met30p inhibits ubiquitin ligase activities of their SCF complexes. 2003 , 2, 123-33		19
314	She4p/Dim1p interacts with the motor domain of unconventional myosins in the budding yeast, <i>Saccharomyces cerevisiae</i> . 2003 , 14, 2237-49		59
313	A single amino acid substitution in herpes simplex virus type 1 VP16 inhibits binding to the virion host shutoff protein and is incompatible with virus growth. 2003 , 77, 2892-902		24
312	Genetic, biochemical, and morphological evidence for the involvement of N-glycosylation in biosynthesis of the cell wall beta1,6-glucan of <i>Saccharomyces cerevisiae</i> . 2003 , 100, 15381-6		24
311	Insight into functional aspects of Stt3p, a subunit of the oligosaccharyl transferase. Evidence for interaction of the N-terminal domain of Stt3p with the protein kinase C cascade. <i>Journal of Biological Chemistry</i> , 2003 , 278, 51441-7	5.4	21

310	Cdc50p, a conserved endosomal membrane protein, controls polarized growth in <i>Saccharomyces cerevisiae</i> . 2003 , 14, 730-47		42
309	Dsl1p, an essential component of the Golgi-endoplasmic reticulum retrieval system in yeast, uses the same sequence motif to interact with different subunits of the COPI vesicle coat. <i>Journal of Biological Chemistry</i> , 2003 , 278, 51722-34	5.4	76
308	AtFACE-2, a functional prenylated protein protease from <i>Arabidopsis thaliana</i> related to mammalian Ras-converting enzymes. <i>Journal of Biological Chemistry</i> , 2003 , 278, 42091-7	5.4	42
307	Yeast V1-ATPase: affinity purification and structural features by electron microscopy. <i>Journal of Biological Chemistry</i> , 2003 , 278, 47299-306	5.4	53
306	The <i>Saccharomyces cerevisiae</i> COQ6 gene encodes a mitochondrial flavin-dependent monooxygenase required for coenzyme Q biosynthesis. <i>Journal of Biological Chemistry</i> , 2003 , 278, 25308-16	5.4	58
305	A region within a luminal loop of <i>Saccharomyces cerevisiae</i> Ycf1p directs proteolytic processing and substrate specificity. 2003 , 2, 588-98		32
304	Identification, functional expression and enzymic analysis of two distinct CaaX proteases from <i>Caenorhabditis elegans</i> . <i>Biochemical Journal</i> , 2003 , 370, 1047-54	3.8	25
303	A striking quality control subcompartment in <i>Saccharomyces cerevisiae</i> : the endoplasmic reticulum-associated compartment. 2004 , 15, 908-21		73
302	<i>Arabidopsis</i> formin AtFH6 is a plasma membrane-associated protein upregulated in giant cells induced by parasitic nematodes. 2004 , 16, 2529-40		120
301	Simultaneous yet independent regulation of actin cytoskeletal organization and translation initiation by glucose in <i>Saccharomyces cerevisiae</i> . 2004 , 15, 1544-56		40
300	Intracellular trafficking of the human Wilson protein: the role of the six N-terminal metal-binding sites. <i>Biochemical Journal</i> , 2004 , 380, 805-13	3.8	94
299	Septin ring assembly requires concerted action of polarisome components, a PAK kinase Cla4p, and the actin cytoskeleton in <i>Saccharomyces cerevisiae</i> . 2004 , 15, 5329-45		87
298	The amino-terminal portion of the F-box protein Met30p mediates its nuclear import and assimilation into an SCF complex. <i>Journal of Biological Chemistry</i> , 2004 , 279, 6674-82	5.4	7
297	HOR7, a multicopy suppressor of the Ca ²⁺ -induced growth defect in sphingolipid mannosyltransferase-deficient yeast. <i>Journal of Biological Chemistry</i> , 2004 , 279, 36390-6	5.4	7
296	Distinct machinery is required in <i>Saccharomyces cerevisiae</i> for the endoplasmic reticulum-associated degradation of a multispinning membrane protein and a soluble luminal protein. <i>Journal of Biological Chemistry</i> , 2004 , 279, 38369-78	5.4	207
295	Aep3p stabilizes the mitochondrial bicistronic mRNA encoding subunits 6 and 8 of the H ⁺ -translocating ATP synthase of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004 , 279, 15728-33	5.4	48
294	Cdc50p, a protein required for polarized growth, associates with the Drs2p P-type ATPase implicated in phospholipid translocation in <i>Saccharomyces cerevisiae</i> . 2004 , 15, 3418-32		210
293	Yellow stripe1. Expanded roles for the maize iron-phytosiderophore transporter. 2004 , 135, 112-20		159

292	Protein sorting in the late Golgi of <i>Saccharomyces cerevisiae</i> does not require mannosylated sphingolipids. <i>Journal of Biological Chemistry</i> , 2004 , 279, 1020-9	5.4	22
291	Two genes encoding <i>Arabidopsis halleri</i> MTP1 metal transport proteins co-segregate with zinc tolerance and account for high MTP1 transcript levels. <i>Plant Journal</i> , 2004 , 39, 425-39	6.9	249
290	The <i>Aspergillus fumigatus</i> transcriptional activator CpcA contributes significantly to the virulence of this fungal pathogen. 2004 , 52, 785-99		103
289	The internalization of yeast Ste6p follows an ordered series of events involving phosphorylation, ubiquitination, recognition and endocytosis. 2004 , 5, 165-80		26
288	Kinetic properties of a micronutrient transporter from <i>Pisum sativum</i> indicate a primary function in Fe uptake from the soil. 2004 , 218, 784-92		102
287	A novel beta-glucosidase in <i>Uromyces fabae</i> : feast or fight?. 2004 , 45, 96-103		15
286	Complementation of <i>Saccharomyces cerevisiae</i> ccc2 mutant by a putative P1B-ATPase from <i>Brassica napus</i> supports a copper-transporting function. 2004 , 566, 218-22		11
285	A tRNA(TRP) gene mediates the suppression of cbs2-223 previously attributed to ABC1/COQ8. 2004 , 317, 648-53		33
284	Functional assessment of the carboxy-terminus of the Wilson disease copper-transporting ATPase, ATP7B. 2004 , 83, 473-81		18
283	Identification and characterization of a novel Delta6-fatty acid desaturase gene from <i>Rhizopus arrhizus</i> . 2004 , 556, 81-5		20
282	Characterization of a novel NADP(+)-dependent D-arabitol dehydrogenase from the plant pathogen <i>Uromyces fabae</i> . <i>Biochemical Journal</i> , 2005 , 389, 289-95	3.8	46
281	Detecting protein-protein interaction in live yeast by flow cytometry. 2005 , 63, 77-86		18
280	Identification of residues in the WD-40 repeat motif of the F-box protein Met30p required for interaction with its substrate Met4p. 2005 , 273, 361-70		5
279	F1F0-ATP synthase complex interactions in vivo can occur in the absence of the dimer specific subunit e. 2005 , 37, 55-66		39
278	A rapid and simple procedure for high-efficiency lithium acetate transformation of cryopreserved <i>Schizosaccharomyces pombe</i> cells. 2005 , 22, 799-804		35
277	Cell-based assays to detect inhibitors of fungal mRNA capping enzymes and characterization of sinefungin as a cap methyltransferase inhibitor. 2005 , 10, 355-64		20
276	The isoprenoid substrate specificity of isoprenylcysteine carboxymethyltransferase: development of novel inhibitors. <i>Journal of Biological Chemistry</i> , 2005 , 280, 29454-61	5.4	33
275	Alteration of a novel dispensable mitochondrial ribosomal small-subunit protein, Rsm28p, allows translation of defective COX2 mRNAs. 2005 , 4, 337-45		13

274	Defects in structural integrity of ergosterol and the Cdc50p-Drs2p putative phospholipid translocase cause accumulation of endocytic membranes, onto which actin patches are assembled in yeast. 2005 , 16, 5592-609		38
273	Specific substitutions in the echinocandin target Fks1p account for reduced susceptibility of rare laboratory and clinical <i>Candida</i> sp. isolates. 2005 , 49, 3264-73		328
272	Structural and functional separation of the N- and C-terminal domains of the yeast V-ATPase subunit H. <i>Journal of Biological Chemistry</i> , 2005 , 280, 36978-85	5-4	44
271	Possible roles for mannitol and mannitol dehydrogenase in the biotrophic plant pathogen <i>Uromyces fabae</i> . 2005 , 137, 190-8		129
270	Subunits of the translocon interact with components of the oligosaccharyl transferase complex. <i>Journal of Biological Chemistry</i> , 2005 , 280, 22917-24	5-4	53
269	Yeast as a tractable genetic system for functional studies of the insulin-degrading enzyme. <i>Journal of Biological Chemistry</i> , 2005 , 280, 27481-90	5-4	22
268	In vivo self-interaction of nodavirus RNA replicase protein a revealed by fluorescence resonance energy transfer. 2005 , 79, 8909-19		43
267	CBT1 interacts genetically with CBP1 and the mitochondrially encoded cytochrome b gene and is required to stabilize the mature cytochrome b mRNA of <i>Saccharomyces cerevisiae</i> . 2005 , 171, 949-57		8
266	Purification, functional reconstitution, and characterization of the <i>Saccharomyces cerevisiae</i> isoprenylcysteine carboxymethyltransferase Ste14p. <i>Journal of Biological Chemistry</i> , 2005 , 280, 7336-45	5-4	36
265	Mutational analysis of the stator subunit E of the yeast V-ATPase. <i>Journal of Biological Chemistry</i> , 2005 , 280, 18393-402	5-4	22
264	A dual upstream open reading frame-based autoregulatory circuit controlling polyamine-responsive translation. <i>Journal of Biological Chemistry</i> , 2005 , 280, 39229-37	5-4	113
263	Meiotic telomere clustering requires actin for its formation and cohesin for its resolution. 2005 , 170, 213-23		124
262	Alpha-synuclein targets the plasma membrane via the secretory pathway and induces toxicity in yeast. 2005 , 170, 47-59		94
261	Screening for mutants defective in secretory protein maturation and ER quality control. 2005 , 35, 366-72		4
260	Osmotin is a homolog of mammalian adiponectin and controls apoptosis in yeast through a homolog of mammalian adiponectin receptor. 2005 , 17, 171-80		155
259	Molecular and functional characterization of a fructose specific transporter from the gray mold fungus <i>Botrytis cinerea</i> . 2005 , 42, 601-10		63
258	ATP7B mediates vesicular sequestration of copper: insight into biliary copper excretion. 2006 , 130, 493-506		104
257	<i>Saccharomyces cerevisiae</i> -based molecular tool kit for manipulation of genes from gram-negative bacteria. 2006 , 72, 5027-36		282

256	Mutational analysis of the Lem3p-Dnf1p putative phospholipid-translocating P-type ATPase reveals novel regulatory roles for Lem3p and a carboxyl-terminal region of Dnf1p independent of the phospholipid-translocating activity of Dnf1p in yeast. 2006 , 344, 323-31		26
255	A common genetic system for functional studies of pitrilysin and related M16A proteases. <i>Biochemical Journal</i> , 2006 , 398, 145-52	3.8	10
254	Regulation of chronological aging in <i>Schizosaccharomyces pombe</i> by the protein kinases Pka1 and Sck2. 2006 , 5, 345-57		91
253	The <i>Aspergillus nidulans</i> F-box protein GrrA links SCF activity to meiosis. 2006 , 61, 76-88		64
252	Substrate specificity of acyl-Delta(6)-desaturases from Continental versus Macaronesian <i>Echium</i> species. 2006 , 67, 540-4		19
251	Xylitol dehydrogenase from <i>Candida tropicalis</i> : molecular cloning of the gene and structural analysis of the protein. 2006 , 73, 631-9		7
250	Identification of interacting proteins using the yeast two-hybrid screen. 2006 , 332, 211-31		1
249	Identification of mouse palmitoyl-coenzyme A Delta9-desaturase. 2006 , 47, 700-4		81
248	A cytoplasmic peptide: N-glycanase. 2006 , 415, 46-55		25
247	Independent and simultaneous translocation of two substrates by a nucleotide sugar transporter. 2006 , 103, 16176-81		30
246	Mutational analysis of the ras converting enzyme reveals a requirement for glutamate and histidine residues. <i>Journal of Biological Chemistry</i> , 2006 , 281, 4596-605	5.4	40
245	Subclassification of the RBCC/TRIM superfamily reveals a novel motif necessary for microtubule binding. <i>Journal of Biological Chemistry</i> , 2006 , 281, 8970-80	5.4	208
244	Identification of the yeast R-SNARE Nyv1p as a novel longin domain-containing protein. 2006 , 17, 4282-99		36
243	A functional role for the GCC185 golgin in mannose 6-phosphate receptor recycling. 2006 , 17, 4353-63		104
242	Identification of novel mutations in ACT1 and SLA2 that suppress the actin-cable-overproducing phenotype caused by overexpression of a dominant active form of Bni1p in <i>Saccharomyces cerevisiae</i> . 2006 , 173, 527-39		7
241	Dimeric organization of the yeast oligosaccharyl transferase complex. 2006 , 103, 8947-52		34
240	Identification of a domain in the V0 subunit d that is critical for coupling of the yeast vacuolar proton-translocating ATPase. <i>Journal of Biological Chemistry</i> , 2006 , 281, 30001-14	5.4	33
239	Cysteine-disulfide cross-linking to monitor SNARE complex assembly during endoplasmic reticulum-Golgi transport. <i>Journal of Biological Chemistry</i> , 2006 , 281, 2281-8	5.4	19

238	Small-molecule inhibitors of the Rce1p CaaX protease. 2007 , 12, 983-93	54
237	Diploids heterozygous for a vma13Delta mutation in <i>Saccharomyces cerevisiae</i> highlight the importance of V-ATPase subunit balance in supporting vacuolar acidification and silencing cytosolic V1-ATPase activity. <i>Journal of Biological Chemistry</i> , 2007 , 282, 8521-32	54 16
236	Involvement of a Golgi-resident GPI-anchored protein in maintenance of the Golgi structure. 2007 , 18, 1261-71	15
235	The <i>Medicago truncatula</i> DMI1 protein modulates cytosolic calcium signaling. 2007 , 145, 192-203	85
234	Copper binding to the N-terminal metal-binding sites or the CPC motif is not essential for copper-induced trafficking of the human Wilson protein (ATP7B). <i>Biochemical Journal</i> , 2007 , 401, 143-53 ^{3,8}	45
233	Conservation of the salt overly sensitive pathway in rice. 2007 , 143, 1001-12	400
232	Endocytic recycling in yeast is regulated by putative phospholipid translocases and the Ypt31p/32p-Rcy1p pathway. 2007 , 18, 295-312	128
231	Phototrophic Fe(II) oxidation promotes organic carbon acquisition by <i>Rhodobacter capsulatus</i> SB1003. 2007 , 73, 6150-8	17
230	Cloning, functional identification and structural modelling of <i>Vitis vinifera</i> S-adenosylmethionine decarboxylase. 2007 , 164, 1208-19	7
229	Convenient transformation of anamorphic basidiomycetous yeasts belonging to genus <i>pseudozyma</i> induced by electroporation. 2007 , 104, 517-20	17
228	Probing the SERCA1a sarcoplasmic reticulum Ca ²⁺ -ATPase phosphorylation-site mutant D351E with inorganic phosphate. 2007 , 40, 1323-32	
227	An unusual S-adenosylmethionine synthetase gene from dinoflagellate is methylated. 2007 , 8, 87	12
226	Differential regulation of cadmium-inducible expression of iron-deficiency-responsive genes in tobacco and barley. 2007 , 129, 622-634	36
225	Inhibition of the CaaX proteases Rce1p and Ste24p by peptidyl (acyloxy)methyl ketones. 2007 , 1773, 853-62	33
224	TOR-induced resistance to toxic adenosine analogs in <i>Leishmania</i> brought about by the internalization and degradation of the adenosine permease. 2007 , 313, 1963-78	14
223	Overexpression of AeNHX1, a root-specific vacuolar Na ⁺ /H ⁺ antiporter from <i>Agropyron elongatum</i> , confers salt tolerance to <i>Arabidopsis</i> and <i>Festuca</i> plants. 2007 , 26, 1663-72	51
222	Cloning and molecular characterisation of a Delta8-sphingolipid-desaturase from <i>Nicotiana tabacum</i> closely related to Delta6-acyl-desaturases. 2007 , 64, 241-50	17
221	A bifunctional TPS-TPP enzyme from yeast confers tolerance to multiple and extreme abiotic-stress conditions in transgenic <i>Arabidopsis</i> . 2007 , 226, 1411-21	158

220	Cloning and functional characterization of the cis-aconitic acid decarboxylase (CAD) gene from <i>Aspergillus terreus</i> . 2008 , 80, 223-9	94
219	Novel interactions of fission yeast kinesin 8 revealed through in vivo expression of truncation alleles. 2008 , 65, 626-40	6
218	Sequence variation in the ATP-binding domain of the Wilson disease transporter, ATP7B, affects copper transport in a yeast model system. 2008 , 29, 491-501	20
217	In planta biocatalysis screen of P450s identifies 8-methoxypsoralen as a substrate for the CYP82C subfamily, yielding original chemical structures. 2008 , 15, 149-56	30
216	The syntaxins SYP31 and SYP81 control ER-Golgi trafficking in the plant secretory pathway. 2008 , 9, 1629-52	68
215	<i>Fasciola hepatica</i> and <i>Fasciola gigantica</i> : cloning and characterisation of 70 kDa heat-shock proteins reveals variation in HSP70 gene expression between parasite species recovered from sheep. 2008 , 118, 536-42	16
214	Structure of the oligosaccharyl transferase complex at 12 Å resolution. 2008 , 16, 432-40	45
213	Expression of a Vacuolar Na ⁺ /H ⁺ Antiporter Gene of Alfalfa Enhances Salinity Tolerance in Transgenic Arabidopsis. 2008 , 34, 557-564	8
212	Functional characterization of a wheat plasma membrane Na ⁺ /H ⁺ antiporter in yeast. 2008 , 473, 8-15	85
211	Identification and characterization of novel SNPs in CHEK2 in Ashkenazi Jewish men with prostate cancer. 2008 , 270, 173-80	15
210	High sensibility to reactivation by acidic lipids of the recombinant human plasma membrane Ca ²⁺ -ATPase isoform 4xb purified from <i>Saccharomyces cerevisiae</i> . 2008 , 1778, 2757-64	9
209	Identification of cytochrome c oxidase subunit 6A1 as a suppressor of Bax-induced cell death by yeast-based functional screening. 2008 , 373, 58-63	18
208	The nuclear migration protein NUDF/LIS1 forms a complex with NUDC and BNFA at spindle pole bodies. 2008 , 7, 1041-52	16
207	Iron-induced turnover of the Arabidopsis IRON-REGULATED TRANSPORTER1 metal transporter requires lysine residues. 2008 , 146, 1964-73	106
206	Protein kinases Fpk1p and Fpk2p are novel regulators of phospholipid asymmetry. 2008 , 19, 1783-97	65
205	A plant Ca ²⁺ pump, ACA2, relieves salt hypersensitivity in yeast. Modulation of cytosolic calcium signature and activation of adaptive Na ⁺ homeostasis. <i>Journal of Biological Chemistry</i> , 2008 , 283, 3497-3506	43
204	Calnexin is involved in apoptosis induced by endoplasmic reticulum stress in the fission yeast. 2008 , 19, 4404-20	54
203	Mrd1p binds to pre-rRNA early during transcription independent of U3 snoRNA and is required for compaction of the pre-rRNA into small subunit processomes. 2008 , 36, 4364-80	18

202	The catalytic activity of protein-disulfide isomerase requires a conformationally flexible molecule. <i>Journal of Biological Chemistry</i> , 2008 , 283, 33630-40	5-4	69
201	The <i>Schizosaccharomyces pombe</i> Hsp104 disaggregase is unable to propagate the [PSI] prion. <i>PLoS ONE</i> , 2009 , 4, e6939	3-7	16
200	Subunit interactions and requirements for inhibition of the yeast V1-ATPase. <i>Journal of Biological Chemistry</i> , 2009 , 284, 13316-13325	5-4	41
199	HOS3, an ELO-like gene, inhibits effects of ABA and implicates a S-1-P/ceramide control system for abiotic stress responses in <i>Arabidopsis thaliana</i> . 2009 , 2, 138-51		37
198	The tether connecting cytosolic (N terminus) and membrane (C terminus) domains of yeast V-ATPase subunit a (Vph1) is required for assembly of V0 subunit d. <i>Journal of Biological Chemistry</i> , 2009 , 284, 19522-32	5-4	15
197	Controlled enlargement of the glycoprotein vesicle surrounding a volvox embryo requires the InvB nucleotide-sugar transporter and is required for normal morphogenesis. 2009 , 21, 1166-81		32
196	Multiple antibiotic resistance in <i>Arabidopsis</i> is conferred by mutations in a chloroplast-localized transport protein. 2009 , 151, 559-73		57
195	Heterologous expression studies of <i>Saccharomyces cerevisiae</i> reveal two distinct trypanosomatid CaaX protease activities and identify their potential targets. 2009 , 8, 1891-900		17
194	Multicopy suppressor analysis of thermosensitive YIP1 alleles implicates GOT1 in transport from the ER. <i>Journal of Cell Science</i> , 2009 , 122, 1540-50	5-3	21
193	Identification of a novel delta9-fatty acid desaturase gene and its promoter from oil-producing fungus <i>Rhizopus arrhizus</i> . 2009 , 36, 177-86		7
192	Identification and characterization of a novel delta6-fatty acid desaturase gene from <i>Rhizopus nigricans</i> . 2009 , 36, 2291-7		20
191	A new tip homolog, ShTIP, from <i>Salicornia</i> shows a different involvement in salt stress compared to that of TIP from <i>Arabidopsis</i> . 2009 , 53, 271-277		11
190	Molecular Cloning and Functional Analysis of a Na ⁺ /H ⁺ Antiporter Gene ThNHX1 from a Halophytic Plant <i>Thellungiella halophila</i> . 2009 , 27, 1-12		61
189	Cloning and molecular characterization of the acyl-CoA: diacylglycerol acyltransferase 1 (DGAT1) gene from <i>Echium</i> . 2009 , 44, 555-68		25
188	Proteolytic processing of certain CaaX motifs can occur in the absence of the Rce1p and Ste24p CaaX proteases. 2009 , 26, 451-63		11
187	Yeast Ste23p shares functional similarities with mammalian insulin-degrading enzymes. 2009 , 26, 595-610		12
186	Identification of ADP-ribosylation factor 4 as a suppressor of N-(4-hydroxyphenyl)retinamide-induced cell death. 2009 , 276, 53-60		16
185	Functional studies of membrane-bound and purified human Hedgehog receptor Patched expressed in yeast. 2009 , 1788, 1813-21		14

184	Cloning and functional characterization of a fatty acid β -desaturase from <i>Oenothera biennis</i> : Production of β -linolenic acid by heterologous expression in <i>Saccharomyces cerevisiae</i> . 2010 , 57, 568-573		2
183	Modulation of the inhibitor properties of dipeptidyl (acyloxy)methyl ketones toward the CaaX proteases. 2010 , 18, 6230-7		17
182	Cloning and molecular characterization of a glycerol-3-phosphate O-acyltransferase (GPAT) gene from <i>Echium</i> (Boraginaceae) involved in the biosynthesis of cutin polyesters. 2010 , 232, 987-97		15
181	Cloning and Characterization of Functional Trehalose-6-Phosphate Synthase Gene in Maize. 2010 , 53, 134-141		25
180	Regulation of the plant plasma membrane H ⁺ -ATPase by its C-terminal domain: what do we know for sure?. 2010 , 89, 145-51		26
179	The sunflower plastidial omega3-fatty acid desaturase (HaFAD7) contains the signalling determinants required for targeting to, and retention in, the endoplasmic reticulum membrane in yeast but requires co-expressed ferredoxin for activity. 2010 , 71, 1050-8		8
178	Photoaffinity labeling of Ras converting enzyme 1 (Rce1p) using a benzophenone-containing peptide substrate. 2010 , 18, 5675-84		7
177	Chemical inhibition of CaaX protease activity disrupts yeast Ras localization. 2010 , 27, 327-43		21
176	Analysis of a genome-wide set of gene deletions in the fission yeast <i>Schizosaccharomyces pombe</i> . 2010 , 28, 617-623		510
175	A septin from the filamentous fungus <i>A. nidulans</i> induces atypical pseudohyphae in the budding yeast <i>S. cerevisiae</i> . <i>PLoS ONE</i> , 2010 , 5, e9858	3-7	16
174	<i>Candida albicans</i> -produced farnesol stimulates <i>Pseudomonas</i> quinolone signal production in LasR-defective <i>Pseudomonas aeruginosa</i> strains. 2010 , 156, 3096-3107		88
173	Fission yeast <i>lec1-ino80</i> -mediated nucleosome eviction regulates nucleotide and phosphate metabolism. <i>Molecular and Cellular Biology</i> , 2010 , 30, 657-74	4-8	37
172	Use of the plant defense protein osmotin to identify <i>Fusarium oxysporum</i> genes that control cell wall properties. 2010 , 9, 558-68		16
171	Directed evolution reveals hidden properties of VMAT, a neurotransmitter transporter. <i>Journal of Biological Chemistry</i> , 2010 , 285, 5076-84	5-4	19
170	Functional oligomerization of the <i>Saccharomyces cerevisiae</i> isoprenylcysteine carboxyl methyltransferase, Ste14p. <i>Journal of Biological Chemistry</i> , 2010 , 285, 13380-7	5-4	13
169	Initial polarized bud growth by endocytic recycling in the absence of actin cable-dependent vesicle transport in yeast. 2010 , 21, 1237-52		19
168	Stability study of the human G-protein coupled receptor, Smoothened. 2010 , 1798, 1100-10		19
167	Purification of functional human Cl ⁻ /HCO ₃ ⁻ exchanger, AE1, over-expressed in <i>Saccharomyces cerevisiae</i> . 2010 , 74, 106-15		10

166	Expression of the human atypical kinase ADCK3 rescues coenzyme Q biosynthesis and phosphorylation of Coq polypeptides in yeast coq8 mutants. 2011 , 1811, 348-60		83
165	Newly identified essential amino acid residues affecting β -sphingolipid desaturase activity revealed by site-directed mutagenesis. 2011 , 416, 165-71		6
164	Regulating cytoplasmic calcium homeostasis can reduce aluminum toxicity in yeast. <i>PLoS ONE</i> , 2011 , 6, e21148	3.7	18
163	Oligopeptide transport and regulation of extracellular proteolysis are required for growth of <i>Aspergillus fumigatus</i> on complex substrates but not for virulence. 2011 , 82, 917-35		32
162	Photoaffinity labeling of Ras converting enzyme using peptide substrates that incorporate benzoylphenylalanine (Bpa) residues: improved labeling and structural implications. 2011 , 19, 7559-69		10
161	Analysis of trehalose-6-phosphate synthase (TPS) gene family suggests the formation of TPS complexes in rice. 2011 , 76, 507-22		67
160	Regulation of durum wheat Na ⁺ /H ⁺ exchanger TdSOS1 by phosphorylation. 2011 , 76, 545-56		40
159	Attenuation of yeast UPR is essential for survival and is mediated by IRE1 kinase. 2011 , 193, 41-50		73
158	Requirement for Golgi-localized PI(4)P in fusion of COPII vesicles with Golgi compartments. 2011 , 22, 216-29		22
157	A new autophagy-related checkpoint in the degradation of an ERAD-M target. <i>Journal of Biological Chemistry</i> , 2011 , 286, 11479-91	5.4	15
156	Phenazine-1-carboxylic acid promotes bacterial biofilm development via ferrous iron acquisition. 2011 , 193, 3606-17		153
155	Recruitment of the inhibitor Cand1 to the cullin substrate adaptor site mediates interaction to the neddylation site. 2011 , 22, 153-64		22
154	Vacuolar H ⁺ -ATPase works in parallel with the HOG pathway to adapt <i>Saccharomyces cerevisiae</i> cells to osmotic stress. 2012 , 11, 282-91		37
153	Cell-permeable, small-molecule activators of the insulin-degrading enzyme. 2012 , 17, 1348-61		10
152	Inhibitors of V-ATPase proton transport reveal uncoupling functions of tether linking cytosolic and membrane domains of V0 subunit a (Vph1p). <i>Journal of Biological Chemistry</i> , 2012 , 287, 10236-10250	5.4	37
151	Shadows of an absent partner: ATP hydrolysis and phosphoenzyme turnover of the Spf1 (sensitivity to <i>Pichia farinosa</i> killer toxin) P5-ATPase. <i>Journal of Biological Chemistry</i> , 2012 , 287, 30477-84	5.4	15
150	Overexpression of the Coq8 kinase in <i>Saccharomyces cerevisiae</i> coq null mutants allows for accumulation of diagnostic intermediates of the coenzyme Q6 biosynthetic pathway. <i>Journal of Biological Chemistry</i> , 2012 , 287, 23571-81	5.4	66
149	Essential role of the NH2-terminal region of Cdc24 guanine nucleotide exchange factor in its initial polarized localization in <i>Saccharomyces cerevisiae</i> . 2012 , 11, 2-15		3

148	Evolution of a novel lysine decarboxylase in siderophore biosynthesis. 2012 , 86, 485-99		34
147	Vacuolar membrane transporters OsVIT1 and OsVIT2 modulate iron translocation between flag leaves and seeds in rice. <i>Plant Journal</i> , 2012 , 72, 400-10	6.9	223
146	Isolation and functional characterisation of the genes encoding β -sphingolipid desaturase from <i>Brassica rapa</i> . 2012 , 39, 47-59		10
145	A novel Sec14 phospholipid transfer protein from <i>Nicotiana benthamiana</i> is up-regulated in response to <i>Ralstonia solanacearum</i> infection, pathogen associated molecular patterns and effector molecules and involved in plant immunity. 2012 , 169, 1017-22		31
144	Dcm methylation is detrimental to plasmid transformation in <i>Clostridium thermocellum</i> . 2012 , 5, 30		60
143	Expression of wheat Na(+)/H(+) antiporter TNHXS1 and H(+)- pyrophosphatase TVP1 genes in tobacco from a bicistronic transcriptional unit improves salt tolerance. 2012 , 79, 137-55		83
142	Plant ornithine decarboxylase is not post-transcriptionally feedback regulated by polyamines but can interact with a cytosolic ribosomal protein S15 polypeptide. 2012 , 42, 519-27		7
141	A conserved START domain coenzyme Q-binding polypeptide is required for efficient Q biosynthesis, respiratory electron transport, and antioxidant function in <i>Saccharomyces cerevisiae</i> . 2013 , 1831, 776-791		26
140	The thermostability of two kinds of recombinant β -fatty acid desaturase with different N-terminal sequence lengths in low temperature. 2013 , 171, 165-72		3
139	Topology of the yeast Ras converting enzyme as inferred from cysteine accessibility studies. 2013 , 52, 6601-14		9
138	Molecular characterization of a lysophosphatidylcholine acyltransferase gene belonging to the MBOAT family in <i>Ricinus communis</i> L. 2013 , 48, 663-74		15
137	Enhancement of plasmid DNA transformation efficiencies in early stationary-phase yeast cell cultures. 2013 , 30, 191-200		12
136	Cloning and truncation modification of trehalose-6-phosphate synthase gene from <i>Selaginella pulvinata</i> . 2013 , 512, 414-21		2
135	Phospholipid flippases Lem3p-Dnf1p and Lem3p-Dnf2p are involved in the sorting of the tryptophan permease Tat2p in yeast. <i>Journal of Biological Chemistry</i> , 2013 , 288, 3594-608	5.4	29
134	The contribution of the nonhomologous region of Prs1 to the maintenance of cell wall integrity and cell viability. 2013 , 13, 291-301		5
133	Cloning and molecular characterization of a class A lysophosphatidate acyltransferase gene (EgLPAT2) from <i>Echium</i> (Boraginaceae). 2013 , 115, 1334-1346		6
132	Cloning, characterization and functional analysis of two type 1 diacylglycerol acyltransferases (DGAT1s) from <i>Tetraena mongolica</i> . 2013 , 55, 490-503		17
131	Identification of conformationally sensitive residues essential for inhibition of vesicular monoamine transport by the noncompetitive inhibitor tetrabenazine. <i>Journal of Biological Chemistry</i> , 2013 , 288, 32160-32179	5.4	19

130	Inhibition of nucleotide sugar transport in <i>Trypanosoma brucei</i> alters surface glycosylation. <i>Journal of Biological Chemistry</i> , 2013 , 288, 10599-615	5-4	12
129	Identification of a second amphipathic lipid-packing sensor-like motif that contributes to Gcs1p function in the early endosome-to-TGN pathway. 2013 , 153, 573-87		2
128	In vivo packaging of triacylglycerols enhances <i>Arabidopsis</i> leaf biomass and energy density. 2013 , 162, 626-39		78
127	Protective effects of ETC complex III and cytochrome c against hydrogen peroxide-induced apoptosis in yeast. 2014 , 48, 435-44		4
126	Novel insights in genetic transformation of the probiotic yeast <i>Saccharomyces boulardii</i> . 2014 , 5, 21-9		16
125	Interaction of the phospholipid flippase Drs2p with the F-box protein Rcy1p plays an important role in early endosome to trans-Golgi network vesicle transport in yeast. 2014 , 155, 51-62		19
124	Asymmetric distribution of phosphatidylserine is generated in the absence of phospholipid flippases in <i>Saccharomyces cerevisiae</i> . 2014 , 3, 803-21		13
123	Role of phosphatidylserine in phospholipid flippase-mediated vesicle transport in <i>Saccharomyces cerevisiae</i> . 2014 , 13, 363-75		27
122	Three homologous genes encoding functional β -sphingolipid desaturase in <i>Populus tomentosa</i> . 2014 , 36, 293-301		2
121	Endo- β -N-acetylglucosamidases (ENGases) in the fungus <i>Trichoderma atroviride</i> : possible involvement of the filamentous fungi-specific cytosolic ENGase in the ERAD process. 2014 , 449, 256-61		16
120	Identification and biochemical characterization of five long-chain acyl-coenzyme A synthetases from the diatom <i>Phaeodactylum tricornutum</i> . 2014 , 74, 33-41		20
119	Functional screening of a cDNA library from the desiccation-tolerant plant <i>Selaginella lepidophylla</i> in yeast mutants identifies trehalose biosynthesis genes of plant and microbial origin. 2014 , 127, 803-13		12
118	Structure and function of CrACA1, the major PM-type Ca ²⁺ -ATPase, expressed at the peak of the gravity-directed trans-cell calcium current in spores of the fern <i>Ceratopteris richardii</i> . 2014 , 16 Suppl 1, 151-7		14
117	Yeast phosphofructokinase-1 subunit Pfk2p is necessary for pH homeostasis and glucose-dependent vacuolar ATPase reassembly. <i>Journal of Biological Chemistry</i> , 2014 , 289, 19448-57	5-4	20
116	Hyperactivation of the human plasma membrane Ca ²⁺ pump PMCA h4xb by mutation of Glu99 to Lys. <i>Journal of Biological Chemistry</i> , 2014 , 289, 10761-10768	5-4	6
115	NITROGEN LIMITATION ADAPTATION recruits PHOSPHATE2 to target the phosphate transporter PT2 for degradation during the regulation of <i>Arabidopsis</i> phosphate homeostasis. 2014 , 26, 454-64		172
114	The identification of four histidine kinases that influence sporulation in <i>Clostridium thermocellum</i> . 2014 , 28, 109-19		24
113	Lipid droplets and their component triglycerides and steryl esters regulate autophagosome biogenesis. <i>EMBO Journal</i> , 2015 , 34, 2117-31	13	143

112	Correlation of polyunsaturated fatty acids with the cold adaptation of <i>Rhodotorula glutinis</i> . 2015 , 32, 683-90		22
111	Polyglutamine toxicity in yeast induces metabolic alterations and mitochondrial defects. 2015 , 16, 662		13
110	Inositol depletion restores vesicle transport in yeast phospholipid flippase mutants. <i>PLoS ONE</i> , 2015 , 10, e0120108	3-7	6
109	Isolation and molecular characterization of a novel WIN1/SHN1 ethylene-responsive transcription factor TdSHN1 from durum wheat (<i>Triticum turgidum</i> . L. subsp. durum). 2015 , 252, 1461-73		26
108	Fatty acid synthase is preferentially degraded by autophagy upon nitrogen starvation in yeast. 2015 , 112, 1434-9		43
107	The Erv41-Erv46 complex serves as a retrograde receptor to retrieve escaped ER proteins. 2015 , 208, 197-209		28
106	Tobacco PIC1 Mediates Iron Transport and Regulates Chloroplast Development. 2015 , 33, 401-413		16
105	Examination of Sec22 Homodimer Formation and Role in SNARE-dependent Membrane Fusion. <i>Journal of Biological Chemistry</i> , 2015 , 290, 10657-66	5-4	13
104	Development of a regulatable plasmid-based gene expression system for <i>Clostridium thermocellum</i> . 2015 , 99, 7589-99		18
103	Conserved function of the lysine-based KXD/E motif in Golgi retention for endomembrane proteins among different organisms. 2015 , 26, 4280-93		28
102	The C-terminal region of the human p23 chaperone modulates its structure and function. 2015 , 565, 57-67		12
101	A short motif in <i>Arabidopsis</i> CDK inhibitor ICK1 decreases the protein level, probably through a ubiquitin-independent mechanism. <i>Plant Journal</i> , 2016 , 87, 617-28	6-9	6
100	Molecular and biochemical characterization of the OLE-1 high-oleic castor seed (<i>Ricinus communis</i> L.) mutant. 2016 , 244, 245-58		11
99	Genes encoding (B)-sphingolipid desaturase from various plants: identification, biochemical functions, and evolution. 2016 , 129, 979-987		3
98	Molecular cloning and characterization of the genes encoding a microsomal oleate Δ2 desaturase (CsFAD2) and linoleate Δ5 desaturase (CsFAD3) from <i>Camelina sativa</i> . 2016 , 89, 405-415		21
97	Metabolic gene products have evolved to interact with the cell wall integrity pathway in <i>Saccharomyces cerevisiae</i> . 2016 , 16,		7
96	Identification and characterization of a novel bifunctional (Δ12)/(Δ15)-fatty acid desaturase gene from <i>Rhodospiridium kratochvilovae</i> . 2016 , 38, 1155-64		23
95	Phylogenetic and Expression Analysis of Pear Yellow Stripe-Like Transporters and Functional Verification of PbrYSL4 in Pear Pollen. 2016 , 34, 737-747		2

94	The Vacuolar Manganese Transporter MTP8 Determines Tolerance to Iron Deficiency-Induced Chlorosis in Arabidopsis. 2016 , 170, 1030-45	108
93	Comparison of different microbial laccases as tools for industrial uses. 2016 , 33, 387-98	44
92	8-Hydroxyquinoline-based inhibitors of the Rce1 protease disrupt Ras membrane localization in human cells. 2016 , 24, 160-78	35
91	A rapid and efficient method to study the function of crop plant transporters in Arabidopsis. 2017 , 254, 737-747	4
90	Enhanced metal tolerance correlates with heterotypic variation in SpMTL, a metallothionein-like protein from the hyperaccumulator <i>Sedum plumbizincicola</i> . 2017 , 40, 1368-1378	44
89	Cfs1p, a Novel Membrane Protein in the PQ-Loop Family, Is Involved in Phospholipid Flippase Functions in Yeast. 2017 , 7, 179-192	8
88	Metal Tolerance Protein 8 Mediates Manganese Homeostasis and Iron Reallocation during Seed Development and Germination. 2017 , 174, 1633-1647	63
87	Characterization of the Hsp100 disaggregase from sugarcane (SHsp101) for chaperone like activity in a yeast system. 2017 , 26, 478-487	2
86	Expression of a Na/H antiporter RtNHX1 from a recretohalophyte <i>Reaumuria trigyna</i> improved salt tolerance of transgenic <i>Arabidopsis thaliana</i> . 2017 , 218, 109-120	38
85	Role of <i>Saccharomyces cerevisiae</i> TAN1 (tRNA acetyltransferase) in eukaryotic initiation factor 2B (eIF2B)-mediated translation control and stress response. 2017 , 7, 223	1
84	Identification and characterization of an efficient acyl-CoA: diacylglycerol acyltransferase 1 (DGAT1) gene from the microalga <i>Chlorella ellipsoidea</i> . 2017 , 17, 48	24
83	A modular yeast biosensor for low-cost point-of-care pathogen detection. 2017 , 3, e1603221	59
82	A Palmitic Acid Elongase Affects Eicosapentaenoic Acid and Plastidial Monogalactosyldiacylglycerol Levels in <i>Nannochloropsis</i> . 2017 , 173, 742-759	47
81	Two bifunctional enzymes from the marine protist : biochemical characterization of wax ester synthase/acyl-CoA:diacylglycerol acyltransferase activity catalyzing wax ester and triacylglycerol synthesis. 2017 , 10, 185	9
80	The V-ATPase subunit A is essential for salt tolerance through participating in vacuolar Na compartmentalization in <i>Salicornia europaea</i> . 2017 , 246, 1177-1187	17
79	A defensin-like protein drives cadmium efflux and allocation in rice. 2018 , 9, 645	149
78	Efficient farnesylation of an extended C-terminal C() sequence motif expands the scope of the prenylated proteome. <i>Journal of Biological Chemistry</i> , 2018 , 293, 2770-2785	5-4 20
77	Phospholipid flippases and Sfk1p, a novel regulator of phospholipid asymmetry, contribute to low permeability of the plasma membrane. 2018 , 29, 1203-1218	18

76	Newly Identified Essential Amino Acids Affecting DGAT1 Function Revealed by Site-Directed Mutagenesis. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	1
75	Protein Isoprenylation in Yeast Targets COOH-Terminal Sequences Not Adhering to the CaaX Consensus. 2018 , 210, 1301-1316		6
74	PA5339, a RidA Homolog, Is Required for Full Growth in <i>Pseudomonas aeruginosa</i> . 2018 , 200,		7
73	The LmSAP gene isolated from the halotolerant <i>Lobularia maritima</i> improves salt and ionic tolerance in transgenic tobacco lines. 2018 , 45, 378-391		12
72	Screening Arrayed Libraries with DNA and Protein Baits to Identify Interacting Proteins. 2018 , 1794, 131-149		2
71	Structural and functional characterisation of two novel durum wheat annexin genes in response to abiotic stress. 2018 , 45, 542-552		7
70	The NHR1-1 of Prs1 and the pentameric motif 284KKCPK288 of Prs3 permit multi-functionality of the PRPP synthetase in <i>Saccharomyces cerevisiae</i> . 2019 , 19,		0
69	Optimization of a Methodology for Quantification and Removal of Zinc Gives Insights Into the Effect of This Metal on the Stability and Function of the Zinc-Binding Co-chaperone Ydj1. 2019 , 7, 416		4
68	The <i>Arabidopsis</i> defensin gene AtPDF2.5 mediates cadmium tolerance and accumulation. 2019 , 42, 2681-2695		25
67	Conserved Glu-47 and Lys-50 residues are critical for UDP--acetylglucosamine/UMP antiport activity of the mouse Golgi-associated transporter Slc35a3. <i>Journal of Biological Chemistry</i> , 2019 , 294, 10042-10054	5.4	5
66	Human COQ10A and COQ10B are distinct lipid-binding START domain proteins required for coenzyme Q function. 2019 , 60, 1293-1310		22
65	Functional domain analysis of LmSAP protein reveals the crucial role of the zinc-finger A20 domain in abiotic stress tolerance. 2019 , 256, 1333-1344		10
64	The endophyte <i>Serendipita indica</i> reduces the sodium content of <i>Arabidopsis</i> plants exposed to salt stress: fungal ENA ATPases are expressed and regulated at high pH and during plant co-cultivation in salinity. 2019 , 21, 3364		18
63	OsProT1 and OsProT3 Function to Mediate Proline- and β -aminobutyric acid-specific Transport in Yeast and are Differentially Expressed in Rice (<i>Oryza sativa</i> L.). 2019 , 12, 79		4
62	Functional Analysis of Ion Transport Properties and Salt Tolerance Mechanisms of RthKT1 from the Recretohalophyte <i>Reaumuria trigyna</i> . 2019 , 60, 85-106		11
61	The glycerophosphocholine acyltransferase Gpc1 is part of a phosphatidylcholine (PC)-remodeling pathway that alters PC species in yeast. <i>Journal of Biological Chemistry</i> , 2019 , 294, 1189-1201	5.4	6
60	Molecular dissection of the Erv41-Erv46 retrograde receptor reveals a conserved cysteine-rich region in Erv46 required for retrieval activity. 2020 , 31, 209-220		0
59	Boron-dependent regulation of translation through AUGUAA sequence in yeast. 2020 , 37, 638-646		2

58	An optogenetic system to control membrane phospholipid asymmetry through flippase activation in budding yeast. 2020 , 10, 12474		3
57	A complex genetic interaction implicates that phospholipid asymmetry and phosphate homeostasis regulate Golgi functions. <i>PLoS ONE</i> , 2020 , 15, e0236520	3-7	2
56	Arabidopsis K ⁺ transporter HAK5-mediated high-affinity root K ⁺ uptake is regulated by protein kinases CIPK1 and CIPK9. 2020 , 71, 5053-5060		19
55	Design of high-oleic tobacco (<i>Nicotiana tabacum</i> L.) seed oil by CRISPR-Cas9-mediated knockout of NtFAD2-2. 2020 , 20, 233		12
54	Protein Farnesyltransferase Catalyzes Unanticipated Farnesylation and Geranylgeranylation of Shortened Target Sequences. 2020 , 59, 1149-1162		8
53	Insights into the mechanisms of transport and regulation of the arabidopsis high-affinity K ⁺ transporter HAK51. 2021 , 185, 1860-1874		13
52	Shedding Light on the Dynamic Role of the "Target of Rapamycin" Kinase in the Fast-Growing C Species , a Suitable Model for Biomass Crops. 2021 , 12, 637508		2
51	Plasma Membrane Ca Pump PMCA4z Is More Active Than Splicing Variant PMCA4x. 2021 , 15, 668371		
50	Distinct Roles of N-Terminal Fatty Acid Acylation of the Salinity-Sensor Protein SOS3. 2021 , 12, 691124		1
49	Auxin-Induced Downregulates to Inhibit Lateral Root Formation in Hance. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6-3	1
48	Yeast One- and Two-Hybrid High-Throughput Screenings Using Arrayed Libraries. 2017 , 1629, 47-65		3
47	Interactions between cAMP-dependent and SNF1 protein kinases in the control of glycogen accumulation in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1994 , 269, 27907-27913	5-4	92
46	Phosphatidylserine decarboxylase from <i>Saccharomyces cerevisiae</i> . Isolation of mutants, cloning of the gene, and creation of a null allele.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 21416-21424	5-4	180
45	Cloning, sequencing and functional expression of a cDNA encoding porcine pancreatic preprocarboxypeptidase A1. 1999 , 259, 719-25		5
44	Telomere-mediated plasmid segregation in <i>Saccharomyces cerevisiae</i> involves gene products required for transcriptional repression at silencers and telomeres. 1993 , 133, 171-82		27
43	Glucose repression of <i>fbp1</i> transcription of <i>Schizosaccharomyces pombe</i> is partially regulated by adenylate cyclase activation by a G protein alpha subunit encoded by <i>gpa2</i> (<i>git8</i>). 1994 , 138, 39-45		53
42	The yeast translational allosuppressor, SAL6: a new member of the PP1-like phosphatase family with a long serine-rich N-terminal extension. 1994 , 138, 597-608		31
41	<i>sck1</i> , a high copy number suppressor of defects in the cAMP-dependent protein kinase pathway in fission yeast, encodes a protein homologous to the <i>Saccharomyces cerevisiae</i> SCH9 kinase. 1995 , 140, 457-67		71

40	Alterations in ribosomal protein RPS28 can diversely affect translational accuracy in <i>Saccharomyces cerevisiae</i> . 1995 , 140, 1247-58		23
39	Cell cycle arrest in <i>cdc20</i> mutants of <i>Saccharomyces cerevisiae</i> is independent of Ndc10p and kinetochore function but requires a subset of spindle checkpoint genes. 1998 , 148, 1701-13		45
38	A screen for genes involved in the anaphase proteolytic pathway identifies <i>tsm1(+)</i> , a novel <i>Schizosaccharomyces pombe</i> gene important for microtubule integrity. 1998 , 149, 1251-64		9
37	Specific components of the SAGA complex are required for Gcn4- and Gcr1-mediated activation of the <i>his4-912delta</i> promoter in <i>Saccharomyces cerevisiae</i> . 1999 , 151, 1365-78		34
36	The novel adaptor protein, Mti1p, and Vrp1p, a homolog of Wiskott-Aldrich syndrome protein-interacting protein (WIP), may antagonistically regulate type I myosins in <i>Saccharomyces cerevisiae</i> . 2002 , 160, 923-34		40
35	A third fatty acid delta9-desaturase from <i>Mortierella alpina</i> with a different substrate specificity to <i>ole1p</i> and <i>ole2p</i> . 2002 , 148, 1725-1735		30
34	EmrE, a small <i>Escherichia coli</i> multidrug transporter, protects <i>Saccharomyces cerevisiae</i> from toxins by sequestration in the vacuole. 1999 , 181, 949-56		13
33	Herpes simplex virus VP16 forms a complex with the virion host shutoff protein vhs. 1994 , 68, 2339-46		124
32	Identification and characterization of a small modular domain in the herpes simplex virus host shutoff protein sufficient for interaction with VP16. 1996 , 70, 2124-31		31
31	Genetic dissection of interaction between poliovirus 3D polymerase and viral protein 3AB. 1997 , 71, 9490-8		100
30	Interactions between DNA-bound trimers of the yeast heat shock factor. <i>Molecular and Cellular Biology</i> , 1994 , 14, 501-508	4.8	34
29	Mutations in the <i>Saccharomyces cerevisiae</i> CDC1 gene affect double-strand-break-induced intrachromosomal recombination. <i>Molecular and Cellular Biology</i> , 1994 , 14, 8037-8050	4.8	13
28	Interactions among three proteins that specifically activate translation of the mitochondrial COX3 mRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994 , 14, 1045-1053	4.8	34
27	Glucose repression of yeast mitochondrial transcription: kinetics of derepression and role of nuclear genes. <i>Molecular and Cellular Biology</i> , 1994 , 14, 1160-1170	4.8	39
26	T7 RNA polymerase-dependent expression of COXII in yeast mitochondria. <i>Molecular and Cellular Biology</i> , 1994 , 14, 4643-4652	4.8	23
25	Cell cycle-dependent transcription of CLN2 is conferred by multiple distinct cis-acting regulatory elements. <i>Molecular and Cellular Biology</i> , 1994 , 14, 4788-4801	4.8	43
24	The yeast Mcm1 protein is regulated posttranscriptionally by the flux of glycolysis. <i>Molecular and Cellular Biology</i> , 1995 , 15, 4631-9	4.8	22
23	COQ6 mutations in human patients produce nephrotic syndrome with sensorineural deafness. <i>Journal of Clinical Investigation</i> , 2011 , 121, 2013-24	15.9	292

22	Identification of a ubiquitous family of membrane proteins and their expression in mouse brain. <i>Journal of Experimental Biology</i> , 2000 , 203, 447-457	3	21
21	Inter-species complementation of the translocon beta subunit requires only its transmembrane domain. <i>PLoS ONE</i> , 2008 , 3, e3880	3.7	8
20	Elevation of the Yields of Very Long Chain Polyunsaturated Fatty Acids via Minimal Codon Optimization of Two Key Biosynthetic Enzymes. <i>PLoS ONE</i> , 2016 , 11, e0158103	3.7	3
19	The Transient Receptor Potential (TRP) Channel Family in <i>Colletotrichum graminicola</i> : A Molecular and Physiological Analysis. <i>PLoS ONE</i> , 2016 , 11, e0158561	3.7	6
18	Phosphorylation of the 19S regulatory particle ATPase subunit, Rpt6, modifies susceptibility to proteotoxic stress and protein aggregation. <i>PLoS ONE</i> , 2017 , 12, e0179893	3.7	11
17	A shunt pathway limits the CaaX processing of Hsp40 Ydj1p and regulates Ydj1p-dependent phenotypes. <i>ELife</i> , 2016 , 5,	8.9	15
16	The meiotic driver utilizes controlled protein aggregation to generate selective cell death. <i>ELife</i> , 2020 , 9,	8.9	10
15	A novel role for DNA photolyase: binding to DNA damaged by drugs is associated with enhanced cytotoxicity in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994 , 14, 8071-8077	4.8	15
14	Cloning and Expression of Cell Cycle Genes in <i>Schizosaccharomyces pombe</i> . 1996 , 119-131		
13	<i>Saccharomyces cerevisiae</i> ER membrane protein complex subunit 4 (EMC4) plays a crucial role in eIF2B-mediated translation regulation and survival under stress conditions. <i>Journal of Genetic Engineering and Biotechnology</i> , 2020 , 18, 15	3.1	0
12	Phosphatidylserine prevents the generation of a protein-free giant plasma membrane domain in yeast.		
11	Thewtf4meiotic driver utilizes controlled protein aggregation to generate selective cell death.		0
10	MALDI-MS Analysis of Peptide Libraries Expands the Scope of Substrates for Farnesyltransferase. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
9	Genetic analysis of glucose regulation in <i>saccharomyces cerevisiae</i> : control of transcription versus mRNA turnover. <i>EMBO Journal</i> , 1996 , 15, 363-74	13	27
8	The CXXC motif: imperatives for the formation of native disulfide bonds in the cell. <i>EMBO Journal</i> , 1996 , 15, 2659-67	13	64
7	A conserved motif in the yeast nucleolar protein Nop2p contains an essential cysteine residue. <i>Biochemical Journal</i> , 1999 , 337 (Pt 1), 29-35	3.8	14
6	The <i>Schizosaccharomyces pombe</i> pyp1 protein tyrosine phosphatase negatively regulates nutrient monitoring pathways. <i>Journal of Cell Science</i> , 1996 , 109 (Pt 7), 1919-1925	5.3	21
5	The broad range di- and trinucleotide exchanger SLC35B1 displays asymmetrical affinities for ATP transport across the ER membrane.. <i>Journal of Biological Chemistry</i> , 2022 , 101537	5.4	0

- 4 Screening DHHCs of S-acylated Proteins Using an OsDHHC cDNA Library and Bimolecular Fluorescence Complementation in Rice.. *Plant Journal*, **2022**, 6.9 0
- 3 A novel thaumatin-like protein from durum wheat, TdPR-5, is homologous to known plant allergens, responsive to stress exposure, and confers multiple-abiotic stress tolerances to transgenic yeast. *Plant Gene*, **2022**, 100360 3.1 1
- 2 Structure-Guided Identification of Critical Residues in the Vacuolar Na⁺,K⁺/H⁺ Antiporter NHX1 from *Arabidopsis thaliana*.
- 1 Functional classification and validation of yeast prenylation motifs using machine learning and genetic reporters. *PLoS ONE*, **2022**, 17, e0270128 3.7 0