

General practice aspects of self-poisoning and self-injury

Psychological Medicine

6, 571-575

DOI: [10.1017/s0033291700018195](https://doi.org/10.1017/s0033291700018195)

Citation Report

#	ARTICLE	IF	CITATIONS
2	People who deliberately poison or injure themselves: their problems and their contacts with helping agencies. <i>Psychological Medicine</i> , 1977, 7, 289-303.	4.5	88
3	Patterns of psychotropic drug use. <i>Social Science and Medicine</i> , 1983, 17, 845-851.	3.8	11
4	Benzodiazepines in general practice: time for a decision.. <i>BMJ: British Medical Journal</i> , 1985, 290, 1374-1376.	2.3	67
5	The prediction of suicide. <i>Medical Journal of Australia</i> , 1990, 153, 552-554.	1.7	33
6	Gender issues in the treatment of suicidal individuals. <i>Death Studies</i> , 1994, 18, 513-527.	2.7	20
7	Contacts of suicide attempters with GPs prior to the event: a comparison between Stockholm and Bern. <i>Acta Psychiatrica Scandinavica</i> , 1997, 95, 94-99.	4.5	50
8	Aborted Suicide Attempts: A New Classification of Suicidal Behavior. <i>American Journal of Psychiatry</i> , 1998, 155, 385-389.	7.2	75
9	The role of general practitioners in parasuicide: A Western Australia perspective. <i>Archives of Suicide Research</i> , 1999, 5, 207-214.	2.3	15
10	The role of general practitioners in parasuicide: A Western Australia perspective. <i>Archives of Suicide Research</i> , 1999, 5, 207-214.	2.3	7
11	Self-confrontation interview with suicide attempters. <i>Counselling Psychology Quarterly</i> , 2002, 15, 1-22.	2.3	32
12	Lethality of firearms relative to other suicide methods: a population based study. <i>Journal of Epidemiology and Community Health</i> , 2003, 57, 120-124.	3.7	201
13	Self-poisoning patients in the general hospital: the frequency and nature of their consultations with the general practitioner. <i>Irish Journal of Psychological Medicine</i> , 2003, 20, 15-17.	1.0	1
14	Psychological and Psychosocial Interventions After Attempted Suicide. <i>Crisis</i> , 2004, 25, 108-117.	1.2	63
15	Health care contacts before and after attempted suicide among adolescent and young adult versus older suicide attempters. <i>Psychological Medicine</i> , 2004, 34, 313-321.	4.5	48
17	Clinical management following self-harm in a UK-wide primary care cohort. <i>Journal of Affective Disorders</i> , 2016, 197, 182-188.	4.1	29
18	Suicide Prevention and Primary Care. , 0, , 661-674.		8
19	Presuicide Attempt Communications Between Parasuicides and Consulted Caregivers. <i>Suicide and Life-Threatening Behavior</i> , 1992, 22, 289-302.	1.9	18
20	General practitioner contacts with patients before and after deliberate self harm. <i>British Journal of General Practice</i> , 2003, 53, 365-70.	1.4	43

#	ARTICLE	IF	CITATIONS
21	General practice consultation patterns before and after intentional overdose: a matched control study. <i>British Journal of General Practice</i> , 1990, 40, 102-5.	1.4	15
22	Factors influencing general practitioners' management of psychosocial and physical problems: a study using case vignettes. <i>British Journal of General Practice</i> , 1990, 40, 284-8.	1.4	6
23	Which antidepressant?. <i>British Journal of General Practice</i> , 1991, 41, 123-5.	1.4	6
24	Self-harm: 2. Deliberate nonfatal self-harm. <i>Cmaj</i> , 1983, 129, 121-5.	0.1	2
25	Parasuicide in an urban general practice, 1970-1979. <i>The Journal of the Royal College of General Practitioners</i> , 1982, 32, 273-81.	0.3	4
26	Deliberate self-poisoning: implications for psychotropic drug prescribing in general practice. <i>The Journal of the Royal College of General Practitioners</i> , 1977, 27, 560-3.	0.3	4
27	Parasuicide and the family physician. <i>Canadian Family Physician</i> , 1979, 25, 1315-8.	0.4	1