

Poleward distribution of mesopelagic fishes is constrained

Global Ecology and Biogeography

31, 546-561

DOI: [10.1111/geb.13446](https://doi.org/10.1111/geb.13446)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Evidence of temperature control on mesopelagic fish and zooplankton communities at high latitudes. <i>Frontiers in Marine Science</i> , 0, 9, .	2.5	8
2	Daily activity timing in the Anthropocene. <i>Trends in Ecology and Evolution</i> , 2023, 38, 324-336.	8.7	18
3	Species distribution models describe spatial variability in mesopelagic fish abundance in the Southern Ocean. <i>Frontiers in Marine Science</i> , 0, 9, .	2.5	2
4	Feeding ecology of <i>Benthosema glaciale</i> across the North Atlantic. <i>Frontiers in Marine Science</i> , 0, 10, .	2.5	0
5	Mid-summer fish behavior in a high-latitude twilight zone. <i>Limnology and Oceanography</i> , 0, , .	3.1	1
6	Future shock: Ocean acidification and seasonal water temperatures alter the physiology of competing temperate and coral reef fishes. <i>Science of the Total Environment</i> , 2023, 883, 163684.	8.0	5
7	Combined use of carbon, nitrogen and sulfur stable isotopes reveal trophic structure and connections in deep-sea mesopelagic and demersal fish communities from the Southeastern Pacific Ocean. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2023, 197, 104069.	1.4	0