

A preliminary study of zooplankton sound scattering la energetics, and migrations

Sarsia

63, 255-264

DOI: [10.1080/00364827.1978.10411347](https://doi.org/10.1080/00364827.1978.10411347)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Feeding habits of cod, capelin, and herring in Balsfjorden, northern Norway, July-August 1978: The importance of euphausiids. Sarsia, 1979, 64, 269-277.	0.5	36
4	Ecological investigations on the zooplankton community of Balsfjorden, northern Norway: population dynamics of the euphausiids <i>Thysanoessa inermis</i> (Kr��yer), <i>Thysanoessa raschii</i> (M.Sars) and <i>Meganyctiphanes norvegica</i> (M.Sars) in 1976 and 1977. Journal of Plankton Research, 1981, 3, 177-192.	1.8	60
5	Ecological investigations on the zooplankton community of Balsfjorden, northern Norway: Seasonal changes in body weight and the main biochemical composition of <i>thysanoessa inermis</i> (Kr��yer), <i>T. Raschii</i> (M. Sars), and <i>Meganyctiphanes norvegica</i> (M. Sars) in relation to environmental factors. Journal of Experimental Marine Biology and Ecology, 1981, 49, 103-120.	1.5	76
6	Ecological investigations on the zooplankton community in Balsfjorden, Northern Norway: seasonal changes in the lipid class composition of <i>Meganyctiphanes norvegica</i> (M. Sars), <i>Thysanoessa raschii</i> (M. Sars), and <i>T. inermis</i> (Kr��yer). Journal of Experimental Marine Biology and Ecology, 1981, 54, 209-224.	1.5	73
7	Distribution and standing stock of zooplankton sound-scattering layers along the north Norwegian coast in February-March, 1978. Sarsia, 1981, 66, 147-160.	0.5	12
8	Ecological investigations on the zooplankton community in balsfjorden, northern Norway: Lipids and fatty acids in <i>Meganyctiphanes norvegica</i> , <i>Thysanoessa raschi</i> and <i>T. inermis</i> during mid-winter. Marine Biology, 1981, 62, 131-137.	1.5	158
9	Length and Weight Relationships of Euphausiids and Caloric Values of <i>Meganyctiphanes Norvegica</i> (M.) Tj ETQq0 0,0,rgBT /Overlock 10	0,8	21
10	Zooplankton sound scattering layers in north Norwegian fjords: Specifications and functioning of the acoustical, data- and net-sampling systems. Sarsia, 1982, 67, 187-199.	0.5	15
11	Net-zooplankton biomass of the Adriatic Sea. Marine Biology, 1984, 79, 209-218.	1.5	38
12	The diel migrations and distributions within a Mesopelagic community in the North East Atlantic. 1. Introduction and sampling procedures. Progress in Oceanography, 1984, 13, 245-268.	3.2	73
13	Mass occurrence of <i>Salpa fusiformis</i> in the spring of 1984 off Ireland: implications for sedimentation processes. Marine Biology, 1988, 97, 127-135.	1.5	53
14	Observations on sound scattering layers in the upwelling off N.W.Africa and in the North Equatorial Current. Journal of Plankton Research, 1988, 10, 89-100.	1.8	1
15	Metabolism and elemental composition of zooplankton from the Barents Sea during early Arctic summer. Marine Biology, 1989, 100, 173-183.	1.5	84
16	Phytoplankton dynamics and sedimentation processes during spring and summer in Balsfjord, Northern Norway. Polar Biology, 1989, 10, 113.	1.2	43
17	Vertical distribution and trophic interactions of zooplankton and fish in Masfjorden, Norway. Sarsia, 1990, 75, 65-81.	0.5	131
18	The influence of macrozooplankton on vertical particulate flux. Sarsia, 1991, 76, 1-9.	0.5	106
19	The food of the long-rough dab (<i>Hippoglossoides platessoides limandoides</i> Bloch) in Balsfjorden, North Norway. Sarsia, 1993, 78, 17-24.	0.5	13
20	On the trophic fate of <i>Phaeocystis pouchetti</i> (Harriot). V. Trophic relationships between <i>Phaeocystis</i> and zooplankton: an assessment of methods and size dependence. Journal of Plankton Research, 1994, 16, 487-511.	1.8	63

#	ARTICLE	IF	CITATIONS
21	Behavioural differences in relation to pycnoclines during vertical migration of the euphausiids <i>Meganyctiphanes norvegica</i> (M. Sars) and <i>Thysanoessa raschii</i> (M. Sars). <i>Journal of Plankton Research</i> , 1997, 19, 255-261.	1.8	32
22	Dry weight carbon and nitrogen content of some euphausiids from the north Atlantic Ocean and the Celtic Sea. <i>Journal of Plankton Research</i> , 1999, 21, 2053-2066.	1.8	18
23	Soft-bottom macrofauna in the high-latitude ecosystem of Balsfjord, northern Norway: Species composition, community structure and temporal variability. <i>Sarsia</i> , 2000, 85, 1-13.	0.5	31
24	Seasonal variation in Zooplankton and suspended faecal pellets in the subarctic Norwegian Baisfjorden, in 1996. <i>Sarsia</i> , 2000, 85, 439-452.	0.5	25
25	Effects of a decrease in downwelling irradiance on the daytime vertical distribution patterns of zooplankton and micronekton. <i>Marine Biology</i> , 2002, 140, 1181-1193.	1.5	75
26	Aggregation and vertical migration behavior of <i>Euphausia superba</i> . <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2004, 51, 2119-2137.	1.4	91
27	The functional biology and trophic role of krill (<i>Thysanoessa raschii</i>) in a Greenlandic fjord. <i>Marine Biology</i> , 2011, 158, 1387-1402.	1.5	22
28	Physical and biochemical properties of the euphausiids <i>Thysanoessa inermis</i> , <i>Thysanoessa raschii</i> , and <i>Thysanoessa longipes</i> in the eastern Bering Sea. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2012, 65-70, 173-183.	1.4	27
29	Seasonal Variability in the Zooplankton Community Structure in a Sub-Arctic Fjord as Revealed by Morphological and Molecular Approaches. <i>Frontiers in Marine Science</i> , 2021, 8, .	2.5	13
30	Why Biologists are Interested in Fjords. , 1980, , 53-66.		22
31	Diurnal and Horizontal Variations in a Zooplankton Sound Scattering Layer. , 1979, , 375-382.		3