

Gradient elution in high-performance liquid chromatog

Journal of Chromatography A

165, 3-30

DOI: [10.1016/s0021-9673\(00\)85726-x](https://doi.org/10.1016/s0021-9673(00)85726-x)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Mobile phase effects in reversed-phase chromatography. Journal of Chromatography A, 1979, 185, 99-109.	1.8	163
2	Systematic Approach to optimizing resolution in reversed- phase liquid chromatography, with emphasis on the role of temperature. Journal of Chromatography A, 1979, 185, 153-177.	1.8	119
3	Influence of organic modifiers on the retention behaviour in reversed-phase liquid chromatography and its consequences for gradient elution. Journal of Chromatography A, 1979, 185, 179-195.	1.8	390
4	Gradient elution in high-performance liquid chromatography. Journal of Chromatography A, 1979, 165, 31-58.	1.8	196
5	Gradient elution in liquid chromatography. Journal of Chromatography A, 1980, 192, 1-18.	1.8	24
6	Gradient elution in liquid chromatography. Journal of Chromatography A, 1980, 192, 19-36.	1.8	33
7	Mechanism of solute retention in liquidâ€”solid chromatography and the role of the mobile phase in affecting separation. Journal of Chromatography A, 1980, 184, 363-413.	1.8	276
8	Reversed-phase gradient high-performance liquid chromatography of procyanidins and their oxidation products in ciders and wines, optimised by Snyder's procedures. Journal of Chromatography A, 1980, 194, 62-68.	1.8	51
9	â€œSequential isocratic stepâ€”liquid chromatography: A highthroughput, problem-solving approach with sensitive, near-universal detection for wide polarity mixtures. Journal of Chromatography A, 1980, 199, 219-238.	1.8	20
10	HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS. Journal of Clinical Pharmacy and Therapeutics, 1980, 5, 107-144.	0.7	0
11	REVERSED-PHASE CHROMATOGRAPHY. , 1980, , 113-319.		91
12	Effect of Type and Concentration of Organic Modifier in Aqueous Eluents on Retention in Reversed Phase Systems. Journal of Liquid Chromatography and Related Technologies, 1980, 3, 1625-1636.	0.9	11
13	Reverse Phase Gradient Elution for Chemically Bonded C18 Thin Layer Chromatographic Plates. Journal of Chromatographic Science, 1980, 18, 133-135.	0.7	13
14	Quantitative Retention - Eluent Composition Relationships in Liquid Chromatography. Journal of Liquid Chromatography and Related Technologies, 1980, 3, 1781-1806.	0.9	20
15	Analysis of N-acetyl-N,O,S-permethylated peptides by combined liquid chromatography-mass spectrometry. Journal of Chromatography A, 1981, 218, 519-533.	1.8	18
16	Chromatographic Methods for the Analysis of Polycyclic Aromatic Hydrocarbons in Water Systems. CRC Critical Reviews in Analytical Chemistry, 1981, 12, 69-153.	2.3	0
17	Continuous and gradient TLC development of 4-dimethylaminoazobenzene-4â€²-sulfonyl amino acid derivatives. Journal of High Resolution Chromatography, 1981, 4, 115-120.	2.0	6
18	Comparison of solubility of naphthols in mixtures of water and methanol and their retention in reversed-phase liquid/solid systems. Journal of High Resolution Chromatography, 1981, 4, 465-466.	2.0	0

#	ARTICLE	IF	CITATIONS
19	Surface silanols in silica-bonded hydrocarbonaceous stationary phases. <i>Journal of Chromatography A</i> , 1981, 203, 53-63.	1.8	393
20	Surface silanols in silica-bonded hydrocarbonaceous stationary phases. <i>Journal of Chromatography A</i> , 1981, 203, 65-84.	1.8	418
21	Construction and evaluation of a programmable gradient liquid chromatograph. <i>Journal of Chromatography A</i> , 1981, 210, 419-431.	1.8	6
22	Gradient elution in liquid chromatography. <i>Journal of Chromatography A</i> , 1981, 214, 35-46.	1.8	56
23	Peak width in solvent-programmed chromatography. <i>Journal of Chromatography A</i> , 1981, 204, 77-84.	1.8	127
24	Use of gradient elution for rapid selection of isocratic conditions in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1981, 205, 13-30.	1.8	174
25	Computer-controlled single-pump solvent programmer for high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1981, 218, 83-93.	1.8	5
26	Estimation of aqueous solubilities of organic non-electrolytes using liquid chromatographic retention data. <i>Journal of Chromatography A</i> , 1981, 218, 409-425.	1.8	66
27	Structural effects in enthalpy/entropy compensated and non-compensated behaviour in ion-pair reversed-phase high-performance liquid-solid chromatography. <i>Journal of Chromatography A</i> , 1981, 218, 427-442.	1.8	25
28	High-performance liquid chromatography of amino acids, peptides and proteins. <i>Journal of Chromatography A</i> , 1981, 218, 497-507.	1.8	65
29	Determination of hydrophobic parameters for pyridazinone herbicides by liquid-liquid partition and reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1981, 206, 7-15.	1.8	67
30	Retention parameters and eluent composition in thin-layer and high-performance liquid chromatography of aromatic nitro compounds in reversed-phase systems. <i>Journal of Chromatography A</i> , 1981, 205, 451-456.	1.8	12
31	A simple molecular model of adsorption chromatography XIV. RF or RM? Secondary retention effects in thin-layer chromatography. <i>Chromatographia</i> , 1981, 14, 23-31.	0.7	28
32	TLC as a pilot technique for the optimization of gradient HPLC I. Experimental verification of a graphical method for the optimization of stepwise gradient elution. <i>Chromatographia</i> , 1981, 14, 411-414.	0.7	21
33	Retention Behaviour of Aromatic Sulfonic Acids in Ion-Pair Reversed-Phase Column Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 1981, 19, 308-314.	0.7	32
34	The Effect of Bonded Ligand Structure on Solute Retention in Reversed-Phase High Performance Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 1981, 19, 130-136.	0.7	18
35	Classification of multisolvent mobile phase systems in liquid chromatography. <i>Analytical Chemistry</i> , 1982, 54, 2593-2596.	3.2	48
36	Determination of ethylene oxide oligomer distributions in alcohol ethoxylates by HPLC using a rotating disc-flame ionization detector. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1982, 59, 364-373.	0.8	37

#	ARTICLE	IF	CITATIONS
37	Optimization of separation conditions in reversed phase liquid chromatography using complex solvent mixtures. <i>Chromatographia</i> , 1982, 16, 132-137.	0.7	15
38	Comparison of the relationships between the retention of dansyl derivatives of aliphatic amines and eluent composition for normal-and reversed-phase liquid-solid TLC systems. <i>Chromatographia</i> , 1982, 15, 309-311.	0.7	4
39	Mobile phase effects in reversed-phase chromatography IV. Retention by n-alkylbenzenes as a function of column temperature and the nature and concentration of organic co-solvent. <i>Chromatographia</i> , 1982, 15, 611-615.	0.7	37
42	Retention behavior of alkylbenzenes as a function of temperature and mobile phase composition in reversed-phase chromatography. <i>Journal of Chromatography A</i> , 1982, 248, 325-339.	1.8	109
43	Investigation of N-alkylbenzamides by reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1982, 235, 31-41.	1.8	32
44	Chromatographic behaviour of 4,4'-dimethylaminoazobenzene-4-sulphonyl derivatives of amines in liquid-solid systems. <i>Journal of Chromatography A</i> , 1982, 248, 263-270.	1.8	5
45	Behaviour of dihydroxynaphthalenes in a reversed-phase chromatographic system. <i>Journal of Chromatography A</i> , 1983, 281, 35-47.	1.8	21
46	Quantitative structure-activity relationships for herbicides. <i>Journal of Chromatography A</i> , 1983, 261, 329-343.	1.8	183
47	Use of a stationary phase with a silica-bonded electrically neutral non-macrocyclic ligand for HPLC separations of steroids. <i>Chromatographia</i> , 1983, 17, 669-672.	0.7	7
48	Relationship between log Pow, shake-flask values and capacity factors derived from reversed-phase high-performance liquid chromatography for n-alkylbenzenes and some oecd reference substances. <i>Journal of Chromatography A</i> , 1983, 282, 315-332.	1.8	144
49	Multifactor model for the optimization of selectivity in reversed-phase chromatography. <i>Journal of Chromatography A</i> , 1983, 258, 11-22.	1.8	64
50	Influence of organic modifiers on the isocratic reversed-phase high-performance liquid chromatography of flavonol glycosides from brewer's hops. <i>Journal of Chromatography A</i> , 1983, 258, 103-109.	1.8	7
51	Quantitative determination of the degree of capping and structural investigation of octadecylsilane phases by means of pyrolysis-gas chromatography. <i>Journal of Chromatography A</i> , 1983, 262, 183-192.	1.8	14
52	Separation of macromolecules by reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1983, 255, 163-189.	1.8	123
53	Reversed-phase ion-pair chromatography of anthracyclines. <i>Journal of Chromatography A</i> , 1983, 254, 45-52.	1.8	23
54	Observations on capacity factor determination for reversed-phase liquid chromatography with aqueous methanol eluents using the solubility parameter concept model and its derivatives. <i>Journal of Chromatography A</i> , 1983, 264, 47-62.	1.8	42
55	Application of local-composition theory to reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1983, 268, 277-280.	1.8	6
56	Isocratic chromatographic retention data for estimating aqueous solubilities of acidic, basic and neutral drugs. <i>International Journal of Pharmaceutics</i> , 1983, 17, 1-21.	2.6	29

#	ARTICLE	IF	CITATIONS
57	Behaviour of Macromolecular RNA in Reversed-Phase HPLC. Journal of Chromatographic Science, 1983, 21, 398-404.	0.7	29
58	Chapter 4 Column chromatography. Journal of Chromatography Library, 1983, , A137-A160.	0.1	0
59	THE COLUMN IN LIQUID CHROMATOGRAPHY. , 1984, , 213-351.		0
60	Automated optimisation of reversed-phase high-performance liquid chromatography separations. Journal of Chromatography A, 1984, 316, 69-79.	1.8	43
61	Examination of theoretical principles of gradient elution as applied to reversed-phase high-performance liquid chromatography separation of phenylthiohydantoin amino acids. Journal of Chromatography A, 1984, 316, 359-372.	1.8	17
62	Measurement and use of retention data from high-performance gradient elution. Journal of Chromatography A, 1984, 285, 19-51.	1.8	94
63	Phenyl-bonded phase liquid chromatography of styrene oligomers. Journal of Chromatography A, 1984, 312, 313-326.	1.8	19
64	Behaviour of biomolecules in water-organic solvent-in-organic salt two-phase ternary systems. Journal of Chromatography A, 1984, 288, 43-49.	1.8	1
65	Retention model for proteins in reversed-phase liquid chromatography. Journal of Chromatography A, 1984, 296, 15-30.	1.8	236
66	Reversed-phase liquid chromatography of homologous series. Journal of Chromatography A, 1984, 314, 13-36.	1.8	158
67	Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol/water and methanol/water. Journal of Chromatography A, 1984, 314, 141-153.	1.8	18
68	Determination of the lipophilicity of some aniline derivatives by reversed-phase thin-layer chromatography. The effect of the organic phase in the eluent. Chromatographia, 1984, 18, 318-322.	0.7	46
69	Mobile phase effects in reversed-phase chromatography VI. Thermodynamic models for retention and its dependence on mobile phase composition and temperature. Chromatographia, 1984, 18, 353-361.	0.7	35
70	Retention behaviour and molecular structure of the solute in thin-layer chromatography. Chromatographia, 1984, 18, 683-691.	0.7	7
72	Analysis of triglyceride species by high-performance liquid chromatography via a flame ionization detector. Lipids, 1984, 19, 142-150.	0.7	44
73	Gradient elution chromatography of polymers on reversed-phase columns with tetrahydrofuran as an eluent component. Chromatographia, 1984, 19, 55-61.	0.7	38
74	Optimal conditions for the reversed-phase chromatography of proteins. Chromatographia, 1984, 19, 77-84.	0.7	47
75	A method for characterization and optimization of reversed-phase liquid chromatographic separations based on the retention behaviour in homologous series. Chromatographia, 1984, 19, 101-112.	0.7	66

#	ARTICLE	IF	CITATIONS
76	Separations of water soluble polyethylene glycol oligomers on phenyl-bonded phase using low wavelength UV detection at 190 nm. Journal of High Resolution Chromatography, 1984, 7, 494-496.	2.0	12
77	Appendix 8 The practice of high-performance LC with four solvents. Journal of Chromatography Library, 1984, , 397-413.	0.1	0
78	Chapter 3 Theory and Practice of High-Speed Microbore HPLC. Journal of Chromatography Library, 1984, , 75-110.	0.1	4
79	Chapter 3: Fractionation. Journal of Chromatography Library, 1984, 29, 59-140.	0.1	0
80	Rotating arc direct current plasma as an emission excitation source. Analytical Chemistry, 1985, 57, 841-845.	3.2	7
81	Microbore reversed-phase chromatography of proteins with conventional gradient equipment for high-performance liquid chromatography. Journal of Chromatography A, 1985, 325, 187-194.	1.8	13
82	Mobile phase optimization for the reversed-phase liquid chromatography of complex hydroxyl aromatic mixtures. Journal of Chromatography A, 1985, 324, 1-18.	1.8	9
83	Stoichiometric displacement of solvent by non-polar solutes in reversed-phase liquid chromatography. Journal of Chromatography A, 1985, 332, 147-168.	1.8	90
84	Characterization of epoxy novolac resins using phenyl-bonded phase high-performance liquid chromatography. Journal of Chromatography A, 1985, 321, 325-333.	1.8	8
85	Lipophilicity measurements of protonated basic compounds by reversed-phase high-performance liquid chromatography. Journal of Chromatography A, 1985, 320, 293-304.	1.8	100
86	Effect of column temperature on high-performance liquid chromatographic behaviour of inorganic polyphosphates. Journal of Chromatography A, 1985, 348, 27-37.	1.8	56
87	Utilization of the functional group contribution concept in liquid chromatography on chemically bonded reversed phases. Journal of Chromatography A, 1985, 348, 49-65.	1.8	22
88	Physico-chemical characterization of cyclic nucleotides by reversed-phase high-performance liquid chromatography. Journal of Chromatography A, 1985, 350, 105-118.	1.8	61
89	Effect of column temperature on high-performance liquid chromatographic behaviour of inorganic polyphosphates. Journal of Chromatography A, 1985, 350, 119-125.	1.8	37
90	Lipophilicity determination of some new crown ethers by reversed-phase thin-layer chromatography. Chromatographia, 1985, 20, 253-260.	0.7	27
91	References to Part I. Journal of Chromatography Library, 1985, , 52-55.	0.1	0
92	References to Part II. Journal of Chromatography Library, 1985, 31, 180-182.	0.1	0
93	Mobile Phase Effects in the Reversed-Phase Liquid Chromatographic Separation of Coenzyme Q Homologs. Journal of Chromatographic Science, 1985, 23, 181-184.	0.7	3

#	ARTICLE	IF	CITATIONS
94	Relationship Between Log K _{av} Values of Benzodiazepines and Composition of the Mobile Phase. Journal of Liquid Chromatography and Related Technologies, 1985, 8, 1711-1729.	0.9	12
95	Effect of Modifier on the Retention of Solutes in Reversed-Phase Thin-Layer Chromatography. Journal of Liquid Chromatography and Related Technologies, 1985, 8, 1207-1224.	0.9	2
96	Optimization Methods For Hplc. Journal of Liquid Chromatography and Related Technologies, 1985, 8, 2725-2780.	0.9	31
97	Electrokinetic chromatography with micellar solution and open-tubular capillary. Analytical Chemistry, 1985, 57, 834-841.	3.2	1,215
98	Alkylation of amide linkages and cleavage of the C chain in the enzyme-activated-substrate inhibition of .alpha.-chymotrypsin with N-nitrosamides. Biochemistry, 1985, 24, 2447-2458.	1.2	9
99	Chapter 4 Optimization Criteria. Journal of Chromatography Library, 1986, 35, 116-169.	0.1	1
100	Chapter 5 Optimization Procedures. Journal of Chromatography Library, 1986, 35, 170-252.	0.1	2
101	Chapter 6 Programmed Analysis. Journal of Chromatography Library, 1986, 35, 253-295.	0.1	0
102	Prediction of precise isocratic retention data from two or more gradient elution runs. Analysis of some associated errors. Analytical Chemistry, 1986, 58, 907-917.	3.2	190
103	Chapter 3 Parameters Affecting Selectivity. Journal of Chromatography Library, 1986, , 37-115.	0.1	0
104	New mixing device for gradient elution. Journal of High Resolution Chromatography, 1986, 9, 537-538.	2.0	1
105	Solute and mobile phase contributions to retention in hydrophobic interaction chromatography of proteins. Journal of Chromatography A, 1986, 359, 131-146.	1.8	205
106	Stepwise gradient development in thin-layer chromatography. Journal of Chromatography A, 1986, 369, 11-17.	1.8	11
107	Prediction of retention of hydroxyl aromatics in reversed-phase liquid chromatography with slope-intercept relationships. Journal of Chromatography A, 1986, 360, 313-325.	1.8	19
108	Stepwise gradient development in thin-layer chromatography. Journal of Chromatography A, 1986, 370, 63-73.	1.8	19
109	Effects of modifier and molecular structure of some coumarins on retention in reversed-phase high-performance thin-layer and column chromatography. Journal of Chromatography A, 1986, 370, 281-292.	1.8	17
110	Method for chromaterization of selectivity in reversed-phase liquid chromatography. Journal of Chromatography A, 1986, 352, 91-110.	1.8	67
111	Temperature, methanol, and structure effects in the reversed-phase liquid chromatography of guanine and hypoxanthine and thier ribonucleosides and ribonucleotides. Journal of Chromatography A, 1986, 352, 221-230.	1.8	10

#	ARTICLE	IF	CITATIONS
112	Chromatographic analysis of 4-dimethylaminoazobenzene-4-sulphonyl and 4-naphthalene-1-azo-(4-dimethylaminobenzene)sulphonyl derivatives of aliphatic alcohols. <i>Journal of Chromatography A</i> , 1986, 362, 217-226.	1.8	6
113	Evaluation of the number of components in multi-component liquid chromatograms of plant extracts. <i>Analytica Chimica Acta</i> , 1986, 191, 261-273.	2.6	62
114	The preopt package for pre-optimization of gradient elutions in high-performance liquid chromatography. <i>Analytica Chimica Acta</i> , 1986, 191, 283-297.	2.6	31
115	Determination of hydrophobic parameters by reversed-phase liquid chromatography: theory, experimental techniques, and application in studies on quantitative structure-activity relationships. <i>Journal of Chromatography A</i> , 1986, 373, 191-225.	1.8	439
116	Stereoselective analysis of racemic psychotropic compounds by HPLC on chiral stationary phase. <i>Psychopharmacology</i> , 1986, 89, 392-3.	1.5	11
117	Isocratic and gradient-elution liquid chromatography of styrene oligomers on silica gel. <i>Journal of Chromatography A</i> , 1986, 362, 325-343.	1.8	27
118	Isolation, separation and quantification of metabolites of the carcinogenic polycyclic aromatic hydrocarbon dibenz(a,h)anthracene. <i>Fresenius Zeitschrift für Analytische Chemie</i> , 1986, 324, 357-357.	0.7	0
119	Effect of the eluent composition in HPLC on the retention of bi- and trifunctional aromatic solutes. <i>Chromatographia</i> , 1986, 22, 25-30.	0.7	24
120	The gradient volume principle for fast evaluation of optimum conditions for isocratic analysis. <i>Chromatographia</i> , 1986, 22, 31-39.	0.7	9
121	Effects of stationary-phase polarity on retention in reversed bonded phase HPLC columns. <i>Chromatographia</i> , 1986, 21, 335-341.	0.7	19
122	Selection of a Mobile Phase Optimization Technique in High Performance Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 1986, 24, 89-94.	0.7	6
123	Use of a Simple Solvent Optimization Program to Improve Separation of Limonin in Citrus Juices. <i>Journal of Chromatographic Science</i> , 1986, 24, 364-366.	0.7	11
124	Influences of Mobile Phase Methanol Content and Solute Character on Relationships Between Reversed-Phase Liquid Chromatographic Retention and Hydrophobic-Lipophilic Parameters of Aromatic Compounds. <i>Journal of Chromatographic Science</i> , 1986, 24, 307-316.	0.7	13
125	Quantitative Structure-Activity Relationships Using Hydrophobicity Constants Measured by High-Pressure Liquid Chromatography: A Comparison with Octanol-Water Partition Coefficients. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1987, 10, 2565-2589.	0.9	27
126	Planar Chromatography Application in Analytical Toxicology. <i>CRC Critical Reviews in Analytical Chemistry</i> , 1987, 17, 213-229.	2.3	0
127	Retention behavior of homologous series in reversed-phase liquid chromatography using micellar, hydro organic, and hybrid mobile phases. <i>Analytical Chemistry</i> , 1987, 59, 2738-2747.	3.2	84
128	Migration off Zones off Test Dyes in Preparative Thin-Layer Chromatography: Stepwise Gradient Elution. <i>Separation Science and Technology</i> , 1987, 22, 2101-2110.	1.3	5
129	Identification of New Zealand wheat cultivars by reversed-phase high-performance liquid chromatography. <i>New Zealand Journal of Agricultural Research</i> , 1987, 30, 125-130.	0.9	4

#	ARTICLE	IF	CITATIONS
130	Column liquid chromatography of integral membrane proteins. <i>Biomedical Applications</i> , 1987, 418, 223-243.	1.7	25
131	Gradient possibilities with the "PRISMA" model. HPLC separation of tobacco alkaloids. <i>Journal of High Resolution Chromatography</i> , 1987, 10, 208-210.	2.0	9
132	Improvement in the resolution of o-phthalaldehyde derivatized amino acids by applying gradient steepness optimization to five reversed-phase columns of different lengths and particle sizes. <i>Journal of Chromatography A</i> , 1987, 411, 121-138.	1.8	13
133	Isochronal optimization. <i>Journal of Chromatography A</i> , 1987, 400, 5-20.	1.8	4
134	Characterization of styrene-divinylbenzene column packings for liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 395, 159-169.	1.8	9
135	General method for the analysis of pharmaceutical dosage forms by high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 391, 233-242.	1.8	49
136	Eluotropic strength of solvents. <i>Journal of Chromatography A</i> , 1987, 389, 21-32.	1.8	39
137	Differences between retentions of various classes of aromatic hydrocarbons in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 388, 51-64.	1.8	44
138	Effect of model inaccuracy on selectivity optimization procedures in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 384, 117-133.	1.8	26
139	The determination of starting conditions for modifier optimization in reversed-phase HPLC. <i>Chromatographia</i> , 1987, 24, 108-114.	0.7	14
140	Computer simulation as a means of developing an optimized reversed-phase gradient-elution separation. <i>Chromatographia</i> , 1987, 24, 261-276.	0.7	83
141	Protein surface area and retention in hydrophobic interaction chromatography. <i>Chromatographia</i> , 1987, 24, 646-650.	0.7	40
142	Determination of hydrophobic parameters by reversed-phase high-performance liquid chromatography. Effect of stationary phase. <i>Chromatographia</i> , 1987, 24, 777-782.	0.7	14
143	Solute retention in the stationary phase of a liquid-solid chromatographic system. <i>Chromatographia</i> , 1987, 23, 915-924.	0.7	4
144	Comparison of the sequential simplex method and linear solvent strength theory in HPLC gradient elution optimization of multicomponent flavonoid mixtures. <i>Chromatographia</i> , 1987, 23, 844-849.	0.7	21
145	Retention behaviour of acidic, neutral and basic drugs on a CN column using phosphate buffers in the mobile phase. <i>Journal of Chromatography A</i> , 1987, 410, 77-94.	1.8	24
146	Solvent properties of liquid organic salts used as mobile phases in microcolumn reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1987, 411, 61-79.	1.8	110
147	Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. <i>Journal of Chromatography A</i> , 1988, 454, 95-107.	1.8	10

#	ARTICLE	IF	CITATIONS
148	Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. Journal of Chromatography A, 1988, 454, 83-94.	1.8	26
149	Prediction of high-performance liquid chromatography capacity factors following column calibration. Journal of Chromatography A, 1988, 437, 13-31.	1.8	7
150	Strategies for automated optimisation of high-performance liquid chromatographic separations incorporating diode-array detection. Journal of Chromatography A, 1988, 458, 335-353.	1.8	13
151	Computer-assisted retention prediction system for oligonucleotides in gradient anion-exchange chromatography. Journal of Chromatography A, 1988, 458, 385-394.	1.8	20
152	Methods for characterization of selectivity in reversed-phase liquid chromatography. Journal of Chromatography A, 1988, 449, 361-389.	1.8	57
153	Retention in reversed-phase liquid chromatography: solvatochromic investigation of homologous alcohol-water binary mobile phases. Journal of Chromatography A, 1988, 457, 85-98.	1.8	53
154	Modification of retention of some alkaloids in the system silanized silica/methanol + water + di(2-ethylhexyl)orthophosphoric acid. Journal of Chromatography A, 1988, 439, 257-266.	1.8	15
155	Quantitative structure-activity relationship of coumarin derivatives. Journal of Chromatography A, 1988, 445, 49-58.	1.8	6
156	Design and application of an expert system for mobile phase optimisation in reversed-phase liquid chromatography. Journal of Pharmaceutical and Biomedical Analysis, 1988, 6, 555-564.	1.4	15
157	Mechanism and prediction of retention of oligomers in normal-phase and reversed-phase HPLC. Chromatographia, 1988, 26, 417-422.	0.7	32
158	Molecular parameters and retention characteristics of unsubstituted polyaromatic hydrocarbons in HPLC. Chromatographia, 1988, 25, 272-278.	0.7	9
159	Olefin group selectivity in non-aqueous reversed-phase LC. Chromatographia, 1988, 25, 327-331.	0.7	0
160	Optimization of the mobile phase composition in gradient elution reversed-phase HPLC by stochastic prediction. Chromatographia, 1988, 25, 343-349.	0.7	9
161	Hydrophobic selectivity in micellar and hydro-organic reversed-phase liquid chromatography. Analytical Chemistry, 1988, 60, 876-887.	3.2	69
162	Determination of polycyclic aromatic hydrocarbons in a coal tar standard reference material. Analytical Chemistry, 1988, 60, 887-894.	3.2	225
163	On the Fundamental Retention Equation in Gradient Elution Liquid Chromatography. Journal of Liquid Chromatography and Related Technologies, 1988, 11, 1809-1826.	0.9	19
164	Optimization of the Mobile Phase for HPLC Separation of S-Alk(en)yl-L-Cysteine Derivatives and Their Corresponding Sulfoxide Isomers. Journal of Liquid Chromatography and Related Technologies, 1989, 12, 199-220.	0.9	11
165	HPLC Measurement of Cyclosporine in Blood Plasma and Urine and Simultaneous Measurement of its Four Metabolites in Blood. , 1989, 12, 1451-1462.		9

#	ARTICLE	IF	CITATIONS
166	Algorithms to correct gradient scan and transfer rule predictions of isocratic retention in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1989, 463, 1-20.	1.8	23
167	Predictive calculation methods for optimization of gradient elution using binary and ternary solvent gradients. <i>Journal of Chromatography A</i> , 1989, 485, 113-141.	1.8	56
168	Computer-assisted retention prediction for high-performance liquid chromatography in the ion-exchange mode. <i>Journal of Chromatography A</i> , 1989, 485, 143-168.	1.8	27
169	Comparison of optimization methods in reversed-phase high-performance liquid chromatography using mixture designs and multi-criteria decision making. <i>Journal of Chromatography A</i> , 1989, 485, 195-217.	1.8	53
170	Effects of different organic modifiers in optimization of reversed-phase high-performance liquid chromatographic gradient elution of a mixture of natural secoiridoid compounds. <i>Journal of Chromatography A</i> , 1989, 485, 631-645.	1.8	13
171	Synthetic peptide purification by application of linear solvent strength gradient theory. <i>Journal of Chromatography A</i> , 1989, 483, 131-143.	1.8	7
172	Systematic procedure for the determination of the nature of the solutes prior to the selection of the mobile phase parameters for optimization of reversed-phase ion-pair chromatographic separations. <i>Journal of Chromatography A</i> , 1989, 478, 21-38.	1.8	21
173	Comparison of reversed-phase thin-layer and high-performance liquid chromatography of flavonoid compounds. <i>Journal of Chromatography A</i> , 1989, 462, 205-217.	1.8	14
174	Solvent selectivity effects in reversed-phase high-performance liquid chromatography of flavonoid compounds. <i>Journal of Chromatography A</i> , 1989, 461, 281-292.	1.8	28
175	Computer-assisted selection of the optimum gradient programme in thin-layer chromatography. <i>Journal of Chromatography A</i> , 1989, 485, 517-532.	1.8	13
176	Practical approach for high-performance liquid chromatographic method development: assaying synthetic intermediates of a leukotriene inhibitor. <i>Journal of Chromatography A</i> , 1989, 485, 597-605.	1.8	5
177	Lipophilicity measurements of benzenesulfonamide inhibitors of carbonic anhydrase by reversed-phase HPLC. <i>International Journal of Pharmaceutics</i> , 1989, 56, 273-281.	2.6	13
178	Limitations of all empirical single-parameter solvent strength scales in reversed-phase liquid chromatography. <i>Analytical Chemistry</i> , 1989, 61, 1524-1529.	3.2	78
179	Influence of sample concentration and adsorption time on the yield of biomolecule ions in plasma desorption mass spectrometry. <i>Analytical Chemistry</i> , 1989, 61, 375-382.	3.2	4
180	Characterization and prediction of retention behavior in reversed-phase chromatography using factor analytical modeling. <i>Analytical Chemistry</i> , 1989, 61, 367-375.	3.2	27
181	Predictive Optimization of Gradient-Elution Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1989, 12, 117-137.	0.9	9
182	Reversed-phase high-performance liquid chromatography in the investigation of the hydrophobicity of selected ketones. <i>Journal of Chromatography A</i> , 1990, 506, 109-117.	1.8	11
183	Determination of hydrophobicity of organic bases on poly(butadiene)-coated alumina and on octadecylsilica columns. <i>Chromatographia</i> , 1990, 30, 277-282.	0.7	9

#	ARTICLE	IF	CITATIONS
184	Effects of stationary phase structure on retention and selectivity in reversed-phase liquid chromatography of flavonoid compounds with methanol as organic modifier. <i>Chromatographia</i> , 1990, 30, 447-452.	0.7	10
185	Optimization of conditions for the simultaneous separation of ten tryptophan metabolites using reversed-phase high-performance liquid chromatography. <i>Biomedical Applications</i> , 1990, 534, 13-21.	1.7	7
186	Possibilities of determination and prediction of solute capacity factors in reversed-phase systems with pure water as the mobile phase. <i>Journal of Chromatography A</i> , 1990, 500, 281-299.	1.8	51
187	Estimation of the reversed-phase liquid chromatographic lipophilicity parameter $\log k'_{\text{w}}$ using ET-30 solvatochromism. <i>Journal of Chromatography A</i> , 1990, 499, 435-451.	1.8	15
188	Effect of alcohol chain length, concentration and polarity on separations in high-performance liquid chromatography using bonded cyclodextrin columns. <i>Journal of Chromatography A</i> , 1990, 499, 477-488.	1.8	9
189	Optimization of the high-performance liquid chromatography of coumarins in <i>Angelica Archangelica</i> with reference to molecular structure. <i>Journal of Chromatography A</i> , 1990, 507, 367-380.	1.8	17
190	Tryptic fingerprinting on a poly(styrene- <i>co</i> -divinylbenzene) reversed-phase column. <i>Journal of Chromatography A</i> , 1990, 512, 315-323.	1.8	3
191	Chiral separations of atropine and homatropine on an $\hat{\pm}$ 1-acid glycoprotein-bonded stationary phase. <i>Journal of Chromatography A</i> , 1990, 506, 579-591.	1.8	27
192	Quantitative correlation of the parameters $\log k'_{\text{w}}$ and $\hat{\epsilon}^{\text{S}}$ in the retention equation in reversed-phase high-performance liquid chromatographic and solvatochromic parameters. <i>Journal of Chromatography A</i> , 1990, 522, 49-55.	1.8	25
193	Liquid Chromatographic Retention Behavior and Separation of Chlorophenols on a $\hat{\text{A}}$ -Cyclodextrin Bonded-Phase Column, Part I. Monoaromatic Chlorophenols: Retention Behavior. <i>Journal of Chromatographic Science</i> , 1990, 28, 311-318.	0.7	28
194	Reduction of reequilibration time following gradient elution reversed-phase liquid chromatography. <i>Analytical Chemistry</i> , 1990, 62, 16-21.	3.2	52
195	Determination of iodine in oyster tissue by isotope dilution laser resonance ionization mass spectrometry. <i>Analytical Chemistry</i> , 1990, 62, 386-389.	3.2	17
196	Adsorption coefficient (K_{oc}) and HPLC retention factors of aromatic hydrocarbons. <i>Chemosphere</i> , 1990, 21, 495-505.	4.2	25
197	Prediction of the adsorption coefficient (K_{oc}) for soil by a chemically immobilized humic acid column using RP-HPLC. <i>Chemosphere</i> , 1990, 21, 729-739.	4.2	41
198	Determination of the adsorption coefficient (K_{OC}) of some aromatics for soil by RP-HPLC on two immobilized humic acid phases. <i>Chemosphere</i> , 1990, 21, 777-788.	4.2	50
199	Optimization of separations in supercritical fluid chromatography using a modified simplex algorithm and short capillary columns. <i>Analytical Chemistry</i> , 1990, 62, 378-386.	3.2	25
200	Computerized optimization of the mobile phase composition in gradient elution reversed-phase HPLC. <i>Chromatographia</i> , 1990, 29, 482-488.	0.7	14
201	THE COLUMN IN LIQUID CHROMATOGRAPHY. , 1991, , 311-544.		3

#	ARTICLE	IF	CITATIONS
202	Predictive Optimization of the Separation of Phenylurea Pesticides using Ternary Mobile Phase Gradients in Reversed-Phase HPLC. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1991, 14, 3125-3151.	0.9	10
203	An evaluation of chemically immobilized humic acid: A new stationary phase for RP-HPLC prediction of uptake of organic pollutants by soils and sediments. , 1991, , 349-355.		3
204	Liquid chromatographic separation of phenothiazines and structurally-related drugs using a β -cyclodextrin bonded phase column. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1991, 9, 409-415.	1.4	16
205	Highly sensitive determination of photosynthetic pigments in marine in situ samples by high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1991, 558, 131-140.	1.8	26
206	Simplex optimization of liquid chromatography with the function of mutual information as a criterion. <i>Journal of Chromatography A</i> , 1991, 585, 187-193.	1.8	5
207	Influence of mobile phase composition on evaluation of lipophilicity by partition chromatography. <i>Journal of Chromatography A</i> , 1991, 557, 399-411.	1.8	21
208	Application of solute and mobile phase partition coefficient to describe solute retention in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1991, 555, 21-31.	1.8	7
209	Experimental model for predicting solute retention time as a function of mobile phase composition in ternary reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1991, 555, 33-51.	1.8	6
210	Multivariate characterization of solvent strength and solvent selectivity in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1991, 550, 397-410.	1.8	20
211	Gradient elution methods for predicting isocratic conditions. <i>Journal of Chromatography A</i> , 1991, 550, 425-447.	1.8	27
212	Chromatographic data for pharmacological classification of imidazol(in)e drugs. <i>Journal of Chromatography A</i> , 1991, 550, 573-584.	1.8	36
213	Description of retention mechanism by solvophobic theory. <i>Journal of Chromatography A</i> , 1991, 556, 219-234.	1.8	31
214	Mathematical structure of chromatographic optimization based on information theory: I. Effect of mobile phase composition on precision. <i>Chromatographia</i> , 1991, 31, 367-373.	0.7	9
215	Prediction of retention of O-ethyl, O-aryl and N-isopropyl phosphoramidothioates in RP-HPLC from molecular structure parameters. <i>Chromatographia</i> , 1991, 31, 143-146.	0.7	8
216	Influence of injection conditions in reversed-phase high-performance liquid chromatography of chlorophylls and carotenoids. <i>Chromatographia</i> , 1991, 31, 589-594.	0.7	91
217	Solvent strength, selectivity and retention mechanism studies on polybutadiene-coated alumina columns in reversed-phase liquid chromatography. <i>Analytica Chimica Acta</i> , 1991, 246, 113-130.	2.6	18
218	Effects of pH in reversed-phase liquid chromatography. <i>Analytica Chimica Acta</i> , 1991, 250, 1-19.	2.6	81
219	A different operational mode for addressing the general elution problem in rapid analysis gas chromatography. <i>Journal of High Resolution Chromatography</i> , 1991, 14, 542-548.	2.0	24

#	ARTICLE	IF	CITATIONS
220	Optimization of the gradient HPLC separation of selected phthalates using the overlapping resolution mapping technique. <i>Environmental Monitoring and Assessment</i> , 1991, 19, 83-91.	1.3	3
221	Application of Iso-Selective Gradient Elution for the Separation of Selected Phthalates. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1991, 14, 3153-3166.	0.9	4
222	Fourth Symposium on our Environment. , 1991, , .		0
223	Standardization of Different Octadecyl Bonded Phases Based on the Retention Equation in Reversed-Phase High-Performance Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 3141-3155.	0.9	1
224	The Selectivity Between Glycine and Taurine Conjugated Bile Acids in Reversed-Phase High-Performance Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 3157-3168.	0.9	0
225	Peak Identification of the Conjugated Bile Acids in Reversed-Phase Liquid Chromatography by Using the Conjugation Selectivity. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1992, 15, 1523-1531.	0.9	4
226	Soluteâ€“mobile phase and soluteâ€“stationary phase interactions in micellar liquid chromatography. A review. <i>Analyst, The</i> , 1992, 117, 831-837.	1.7	103
227	Evaluation of silica-humate and alumina-humate HPLC stationary phases for estimation of the adsorption coefficient, K _{oc} , of soil for some aromatics. <i>Chemosphere</i> , 1992, 24, 403-412.	4.2	30
228	Reversed-phase liquid chromatography of the opioid peptides â€” 2. Quantitative structureâ€“retention relationships and isocratic retention prediction. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1992, 10, 49-60.	1.4	6
229	The assay and resolution of the beta-blocker atenolol from its related impurities in a tablet pharmaceutical dosage form. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1992, 10, 329-334.	1.4	12
230	Study of amphiphilic behaviour of alkylglycoside surfactants using reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1992, 623, 277-287.	1.8	19
231	Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations. <i>Journal of Chromatography A</i> , 1992, 595, 53-61.	1.8	14
232	Effects of molecular structure on the S index in the retention equation in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1992, 606, 1-8.	1.8	23
233	Comparison and evaluation of HIC columns of different hydrophobicity. <i>Chromatographia</i> , 1992, 34, 391-397.	0.7	20
234	Behavior of large nucleic acids in reversed-phase high-performance liquid chromatography. <i>Biomedical Applications</i> , 1992, 584, 135-139.	1.7	3
235	The S index in the retention equation in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1992, 603, 35-42.	1.8	32
236	The retention behaviour of conjugated bile acids in reversed phase high performance liquid chromatography. <i>Biomedical Chromatography</i> , 1992, 6, 255-257.	0.8	5
237	An approximate experimental model for predicting solute retention times in RP-HPLC with quaternary solvent systems. <i>Journal of High Resolution Chromatography</i> , 1992, 15, 555-561.	2.0	0

#	ARTICLE	IF	CITATIONS
238	Effect of the dodecyl sulfate counter ion on selectivity and resolution in micellar electrokinetic capillary chromatography: SDSvs. Mg(DS) ₂ . Journal of Separation Science, 1993, 5, 347-360.	1.0	20
239	Optimization theory of chromatography. Chemometrics and Intelligent Laboratory Systems, 1993, 18, 1-16.	1.8	20
240	Retention behaviour of aromatic sulphonic acids in reversed-phase ion-pair liquid chromatography with methanol and acetonitrile as organic modifiers. Journal of Chromatography A, 1993, 644, 269-275.	1.8	9
241	UNIFAC model as a heuristic guide for estimating retention in reversed-phase liquid chromatography. Journal of Chromatography A, 1993, 656, 69-79.	1.8	15
242	Quantitative structure-retention relationships applied to reversed-phase high-performance liquid chromatography. Journal of Chromatography A, 1993, 656, 417-435.	1.8	195
243	Correlation of retention and selectivity of separation in reversed-phase high-performance liquid chromatography with interaction indices and with lipophilic and polar structural indices. Journal of Chromatography A, 1993, 656, 437-467.	1.8	40
244	Retention in reversed-phase liquid chromatography as a function of mobile-phase composition. Journal of Chromatography A, 1993, 656, 501-520.	1.8	230
245	Extra-thermodynamic relationships in chromatography—study of the relationship between the slopes and intercepts of plots of $\ln k'_{\text{R}}^2$ vs. mobile phase composition in reversed-phase chromatography. Journal of Chromatography A, 1993, 656, 521-535.	1.8	54
246	Variance of a zone migrating in a linear medium. Journal of Chromatography A, 1993, 637, 119-128.	1.8	40
247	Convergent solid-phase peptide synthesis. Tetrahedron, 1993, 49, 11065-11133.	1.0	205
248	Column-switching high-performance liquid chromatographic determination of a 2-pyridinone-based human immunodeficiency virus type 1 (HIV-1)-specific reverse transcriptase inhibitor in human plasma. Pharmaceutical Research, 1993, 10, 56-60.	1.7	8
249	Retention properties of cyclodextrins and modified cyclodextrins in reversed phase HPLC. Chromatographia, 1993, 36, 174-178.	0.7	20
250	Optimization of Mobile Phase Composition for HPLC Separations of Nitroaromatics Using Overlapping Resolution Mapping. Journal of Liquid Chromatography and Related Technologies, 1993, 16, 2223-2235.	0.9	7
251	A Separation Study of Tricyclic Antidepressant Drugs by HPLC with β -Cyclodextrin Bonded Stationary Phase. Journal of Liquid Chromatography and Related Technologies, 1993, 16, 3487-3508.	0.9	13
252	Effect of the Mobile and Stationary Phases on RP-HPLC Retention and Selectivity of Flavonoid Compounds. Journal of Liquid Chromatography and Related Technologies, 1994, 17, 3655-3670.	0.9	8
253	Comparison of Adsorption Coefficient (K_{OC}) for Soils and HPLC Retention Factors of Aromatic Hydrocarbons Using a Chemically Immobilized Humic Acid Column in RP-HPLC. Journal of Liquid Chromatography and Related Technologies, 1994, 17, 2593-2604.	0.9	19
254	Application of Nonlinear Regression Functions for the Modeling of Retention in Reversed-Phase LC. Journal of Chromatographic Science, 1994, 32, 144-152.	0.7	19
255	Determination of 2,4,6-trinitrotoluene and its biodegradation products by normal-phase liquid chromatography. Analytica Chimica Acta, 1994, 291, 205-210.	2.6	5

#	ARTICLE	IF	CITATIONS
256	Preparation of milligrams of pure enantiomers using analytical-scale high-performance liquid chromatography. <i>Chirality</i> , 1994, 6, 290-294.	1.3	20
257	Strategic approach for method selection in high-performance liquid chromatography. <i>Analytica Chimica Acta</i> , 1994, 298, 319-329.	2.6	2
258	Chromatographic behaviour of phenylurea pesticides in high-performance liquid chromatography with nitrile- and amino-bonded stationary phases. <i>Journal of Chromatography A</i> , 1994, 684, 77-92.	1.8	18
259	Gradient elution in micellar liquid chromatography I. Micelle concentration gradient. <i>Journal of Chromatography A</i> , 1994, 683, 321-334.	1.8	23
260	Gradient elution in micellar liquid chromatography II. Organic modifier gradients. <i>Journal of Chromatography A</i> , 1994, 683, 335-345.	1.8	30
261	Comparison of isocratic and gradient elution reversed-phase behaviour of high-molecular-mass polystyrenes in dichloromethane and acetonitrile. <i>Journal of Chromatography A</i> , 1994, 679, 105-114.	1.8	8
262	Determination of lipophilicity by means of reversed-phase thin-layer chromatography. <i>Journal of Chromatography A</i> , 1994, 662, 341-361.	1.8	129
263	Limitation of the ET(30) solvent strength scale in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1994, 677, 1-9.	1.8	18
264	Computational strategy for solvent strength optimization in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1994, 672, 67-85.	1.8	9
265	Computerized optimization of gradient elution conditions with ternary solvent mobile phases in RPLC. <i>Chromatographia</i> , 1994, 39, 216-223.	0.7	11
266	Limits of resolution and speed of analysis in linear chromatography with and without focusing. <i>Chromatographia</i> , 1994, 39, 719-728.	0.7	39
267	A method to determine the ability of drugs to diffuse through the blood-brain barrier.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 68-72.	3.3	190
268	Chapter 7 Characterization of retention and selectivity in reversed-phase LC using interaction indices. <i>Journal of Chromatography Library</i> , 1995, , 235-267.	0.1	4
269	Chromatographic evaluation of porous carbon-clad zirconia microparticles. <i>Analytical Chemistry</i> , 1995, 67, 3042-3050.	3.2	65
270	Temperature and solvation effects on homologous series selectivity in reversed phase liquid chromatography. <i>Analytica Chimica Acta</i> , 1995, 307, 203-215.	2.6	29
271	Retention characteristics of polypyrrole as a stationary phase for the electrochemically modulated liquid chromatographic (EMLC) separations of dansyl amino acids. <i>Journal of Electroanalytical Chemistry</i> , 1995, 387, 35-46.	1.9	41
272	Simultaneous determination of enalapril, felodipine and their degradation products in the dosage formulation by reversed-phase high-performance liquid chromatography using a Spherisorb C8 column. <i>Journal of Chromatography A</i> , 1995, 707, 245-254.	1.8	41
273	Bioavailability Estimation by Reversed-Phase Liquid Chromatography: High Bonding Density C-18 Phases for Modeling Biopartitioning Processes. <i>Analytical Chemistry</i> , 1995, 67, 48-57.	3.2	41

#	ARTICLE	IF	CITATIONS
274	Reverse-phase high performance liquid chromatography and linear solvation energy relationship for studies of alkyl (1-phenylsulfonyl) cycloalkane-carboxylates. <i>Bulletin of Environmental Contamination and Toxicology</i> , 1995, 55, 635-42.	1.3	1
275	Study of Pah's Separation and Phase-Solute Interaction by Micellar Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1995, 18, 2397-2425.	0.9	12
276	Prediction of Partition Coefficient and Toxicity for Phenylthio, Phenylsulfinyl and Phenylsulfonyl Acetates. <i>Environmental Science & Technology</i> , 1995, 29, 3044-3048.	4.6	11
277	Influence of Dodecyl Sulfate Counterion On Efficiency, Selectivity, Retention, Elution Range, and Resolution in Micellar Electrokinetic Chromatography. <i>Analytical Chemistry</i> , 1995, 67, 2315-2324.	3.2	48
278	Chapter 2 Reversed-Phase and Hydrophobic Interaction Chromatography of Carbohydrates and Glycoconjugates. <i>Journal of Chromatography Library</i> , 1995, 58, 41-101.	0.1	7
279	Determination and estimation of physicochemical properties for phenylsulfonyl acetates. <i>Chemosphere</i> , 1995, 30, 117-125.	4.2	16
280	Examination of silica-salicylic acid and silica-8-hydroxyquinoline HPLC stationary phases for estimation of the adsorption coefficient of soil for some aromatic hydrocarbons. <i>Chemosphere</i> , 1995, 30, 1717-1727.	4.2	17
281	Chapter 2 Retention prediction of pharmaceutical compounds. <i>Journal of Chromatography Library</i> , 1995, 57, 47-92.	0.1	4
282	Determination of Solvent Basicity Scale, β^2 , of Mixed Solvents for Three Chromatographic Solvent Systems: 2-Propanol/Hexane, Ethyl Acetate/Hexane, and Methanol/Water. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1996, 19, 277-291.	0.5	9
283	Characterization of polyaniline using chromatographic studies. <i>Chromatographia</i> , 1996, 42, 191-198.	0.7	15
284	Chromatographic behaviour in reversed-phase high-performance liquid chromatography with micellar and submicellar mobile phases: effects of the organic modifier. <i>Biomedical Applications</i> , 1996, 681, 3-19.	1.7	19
285	Recent progress in reversed-phase and hydrophobic interaction chromatography of carbohydrate species. <i>Journal of Chromatography A</i> , 1996, 720, 93-118.	1.8	67
286	Mobile phase effects on single-component and competitive adsorption isotherms in reversed-phase systems. <i>Journal of Chromatography A</i> , 1996, 734, 125-136.	1.8	22
287	Comparison of methods for extracting linear solvent strength gradient parameters from gradient chromatographic data. <i>Journal of Chromatography A</i> , 1996, 727, 1-11.	1.8	12
288	Estimation of Solvent-Modulated Linear Adsorption Parameters of Taxanes from Dilute Plant Tissue Culture Broth. <i>Biotechnology Progress</i> , 1996, 12, 810-821.	1.3	17
289	Correlations Between n-Octanol/Water Partition Coefficients and RP-HPLC Capacity Factors of 1,2-Dithiole-3-thiones and 1,2-Dithiol-3-ones. <i>Journal of Pharmaceutical Sciences</i> , 1996, 85, 990-998.	1.6	19
290	Retention in RP-HPLC: Lipophilicity Determination of Substituted Biphenyls by Reversed-Phase High Performance Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1997, 20, 3029-3037.	0.5	16
291	Mobile Phase Optimization in Thin Layer Chromatography (TLC). <i>Reviews in Analytical Chemistry</i> , 1997, 16, .	1.5	4

#	ARTICLE	IF	CITATIONS
292	Chromatographic Separation of Some Coumarins and Flavonoids on Diol-Modified Silica Gel Phase. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1997, 20, 2089-2098.	0.5	7
293	Determination of Lipophilicity by Gradient Elution High-Performance Liquid Chromatography. <i>Analytical Chemistry</i> , 1997, 69, 2575-2581.	3.2	74
294	Computerized optimization of RP-HPLC separation with nonaqueous or partially aqueous mobile phases. <i>Chromatographia</i> , 1997, 44, 529-537.	0.7	47
295	Chromatographic identification of phenolic compounds in human urine following oral administration of the herbal medicines Daisaiko-to and Shosaiko-to. <i>Biomedical Applications</i> , 1997, 693, 191-198.	1.7	18
297	Rate parameters and gradient correlations for gradient-elution chromatography. <i>AIChE Journal</i> , 1997, 43, 464-474.	1.8	10
298	Whey proteins eluted reversed phase gradients. <i>Journal of High Resolution Chromatography</i> , 1997, 20, 29-33.	2.0	1
299	Influence of differences between sample and mobile phase viscosities on the shape of chromatographic elution profiles. <i>Journal of Chromatography A</i> , 1997, 775, 73-79.	1.8	26
300	Determination of branched carbon number distribution of polyacrylate by liquid chromatography. <i>Analytica Chimica Acta</i> , 1997, 357, 219-224.	2.6	1
301	Peak distortion in reversed-phase liquid chromatography as a consequence of viscosity differences between sample solvent and mobile phase. <i>Journal of Chromatography A</i> , 1998, 805, 55-61.	1.8	51
302	Application of the phenomenological model to retention in reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1998, 829, 41-63.	1.8	11
303	Characterisation of retention in micellar high-performance liquid chromatography and in micellar electrokinetic chromatography using lipophilicity and polarity indices. <i>Journal of Chromatography A</i> , 1998, 807, 57-70.	1.8	22
304	Use of short columns and high flow rates for rapid gradient reversed-phase chromatography. <i>Chromatographia</i> , 1998, 47, 291-298.	0.7	38
305	Exothermodynamic Relationships in Liquid Chromatography. <i>Journal of Physical Chemistry B</i> , 1998, 102, 701-718.	1.2	71
306	Effect of the Chain Length and Concentration of the Alcohol Used in the Mobile Phase in Micellar Liquid Chromatography with a C ₁₈ Column. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1998, 21, 2117-2130.	0.5	1
307	Modelling of solute retention in the reversed-phase high-performance liquid chromatography system with the chemically bonded 3-cyanopropyl stationary phase. <i>Chemometrics and Intelligent Laboratory Systems</i> , 1999, 47, 253-266.	1.8	4
308	Structure and retention of 2,4-dihydroxythiobenzanilides in a reversed-phase system. <i>Journal of Chromatography A</i> , 1999, 830, 491-496.	1.8	24
309	Simultaneous optimisation of gradient time, gradient shape and initial composition of the mobile phase in the high-performance liquid chromatography of homologous and oligomeric series. <i>Journal of Chromatography A</i> , 1999, 845, 133-144.	1.8	38
310	Global linear solvation energy relationships for retention prediction in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1999, 848, 21-37.	1.8	88

#	ARTICLE	IF	CITATIONS
311	Analysis of commercial ceramides by non-aqueous reversed-phase liquid chromatography with evaporative light-scattering detection. <i>Chromatographia</i> , 1999, 49, 241-248.	0.7	23
312	Effects of organic modifiers on solute retention and electrokinetic migrations in micellar electrokinetic capillary chromatography. <i>Electrophoresis</i> , 1999, 20, 2898-2908.	1.3	34
313	Determination and prediction of partition coefficient and toxicity for sulfonylurea herbicides and their degradation products. <i>Chemosphere</i> , 1999, 38, 1713-1719.	4.2	6
314	Comparison of different HPLC stationary phases for determination of soil-water distribution coefficient, KOC values of organic chemicals in RP-HPLC system. <i>Chemosphere</i> , 1999, 39, 431-442.	4.2	26
315	REVERSED PHASE CHROMATOGRAPHIC STUDY OF THE INCLUSION SELECTIVITY OF TERPENE DERIVATIVES WITH β -CYCLODEXTRIN IN WATER/COSOLVENT MIXTURES. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1999, 22, 1267-1284.	0.5	11
316	Chapter 8 The development and industrial application of automated preparative HPLC. <i>Handbook of Analytical Separations</i> , 2000, 1, 293-336.	0.8	1
317	A CORRELATION BETWEEN Z AND S IN HIGH PERFORMANCE LIQUID CHROMATOGRAPHY. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2000, 23, 1267-1279.	0.5	2
318	New calixarene-bonded stationary phases in high-performance liquid chromatography: comparative studies on the retention behavior and on influences of the eluent. <i>Journal of Chromatography A</i> , 2000, 898, 35-52.	1.8	53
319	Chromatographic study of terpene derivatives on porous graphitic carbon stationary phase with β -cyclodextrin as mobile phase modifier. <i>Journal of Chromatography A</i> , 2000, 903, 67-76.	1.8	33
320	Adsorption/partition model of liquid chromatography for chemically bonded stationary phases of the aliphatic cyano, reversed-phase C8 and reversed-phase C18 types. <i>Journal of Chromatography A</i> , 2000, 869, 57-64.	1.8	39
321	Dependence of fungistatic activity of 2,4-dihydroxythiobenzanilides on the structure and lipophilic nature of the compounds. <i>European Journal of Medicinal Chemistry</i> , 2000, 35, 393-404.	2.6	42
322	Classification of structurally related compounds from <i>Astragalus</i> extract by correlation of the log k_w and S . <i>Chromatographia</i> , 2000, 51, 212-220.	0.7	14
323	Use of optimization software for comparing stationary phases in order to find HPLC conditions suitable for separating 16 PAHs. <i>Chromatographia</i> , 2000, 51, 536-544.	0.7	10
324	Chapter 1 Comparison of various modes and phase systems for analytical HPLC. <i>Handbook of Analytical Separations</i> , 2000, , 1-71.	0.8	6
325	Chapter 2 Fast generic HPLC methods. <i>Handbook of Analytical Separations</i> , 2000, 1, 73-85.	0.8	4
326	Reversed-Phase Chromatography. <i>Journal of Chromatography Library</i> , 2000, , 147-234.	0.1	8
327	Prediction of partition coefficient and toxicity for benzaldehyde compounds by their capacity factors and various molecular descriptors. <i>Chemosphere</i> , 2001, 42, 899-907.	4.2	37
328	THE HPLC SEPARATION OF TWO MIXTURES OF NEUTRAL PESTICIDES: EFFECTS OF TEMPERATURE, GRADIENT STEEPNESS, AND FLOW RATE. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2001, 24, 543-553.	0.5	6

#	ARTICLE	IF	CITATIONS
329	Capillary LC-MS at the Attomole Level for Monitoring and Discovering Endogenous Peptides in Microdialysis Samples Collected in Vivo. <i>Analytical Chemistry</i> , 2001, 73, 5005-5014.	3.2	136
330	Retention behavior of some flavonoids in 2D-TLC systems on cyano bonded polar phases. <i>Chromatographia</i> , 2001, 54, 789-794.	0.7	36
331	Automated capillary liquid chromatography for high sensitivity amino acid monitoring. <i>Journal of Separation Science</i> , 2001, 13, 24-32.	1.0	4
332	Ultrafast liquid chromatography/ultraviolet and liquid chromatography/tandem mass spectrometric analysis. <i>Rapid Communications in Mass Spectrometry</i> , 2001, 15, 141-151.	0.7	71
333	Determination of the Hydrophobicity of Local Anesthetic Agents. <i>Analytical Biochemistry</i> , 2001, 292, 102-106.	1.1	7
334	Enantioselectivity of 3-amino-2-oxazolidinone derivatives with potential psychotropic activity on cellulose tris(4-methylbenzoate) chiral selector. <i>Chirality</i> , 2001, 13, 148-152.	1.3	5
335	A theoretical study of the optimization of gradients at elevated temperature. <i>Journal of Separation Science</i> , 2001, 24, 921-929.	1.3	98
336	Comparison of selected retention models in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2001, 913, 3-13.	1.8	30
337	Purification of adipoyl-7-amino-3-deacetoxycephalosporanic acid from fermentation broth using stepwise elution with a synergistically adsorbed modulator. <i>Journal of Chromatography A</i> , 2001, 908, 273-291.	1.8	32
338	PREDICTION OF PARAMETERS IN REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY USING RETENTION PARAMETERS IN GAS LIQUID CHROMATOGRAPHY. <i>Analytical Letters</i> , 2001, 34, 785-802.	1.0	1
339	RETENTION OF SUBSTITUTED BENZALDEHYDES ON RP-HPLC. CORRELATION WITH PARTITION COEFFICIENTS AND MOLECULAR DESCRIPTORS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2001, 24, 937-945.	0.5	2
340	PARTITION PROPERTIES OF 18 POLYCHLORINATED ORGANIC COMPOUNDS (PCOCs): CORRELATION WITH MOLECULAR STRUCTURAL DESCRIPTORS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 627-637.	0.5	3
341	COMPARISON OF REVERSED-PHASE AND NORMAL-PHASE COLUMN LIQUID CHROMATOGRAPHIC TECHNIQUES FOR THE SEPARATION OF LOW AND HIGH MOLECULAR WEIGHT COMPOUNDS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 2901-2931.	0.5	10
342	Comparison of Different Retention Models in Normal and Reversed-Phase Liquid Chromatography with Binary Mobile Phases. <i>Journal of Chromatographic Science</i> , 2002, 40, 575-580.	0.7	8
343	Prediction of Soil Adsorption Coefficients from Retention Parameters on Three Reversed-Phase Liquid Chromatographic Columns. <i>Analytical Chemistry</i> , 2002, 74, 655-660.	3.2	9
344	Selective Enrichment of Low-Abundance Peptides in Complex Mixtures by Elution-Modified Displacement Chromatography and Their Identification by Electrospray Ionization Mass Spectrometry. <i>Analytical Chemistry</i> , 2002, 74, 3933-3941.	3.2	29
345	Chapter 2 Reversed-phase and hydrophobic interaction chromatography of carbohydrates and glycoconjugates. <i>Journal of Chromatography Library</i> , 2002, , 41-102.	0.1	16
346	Enantioseparation of fluoxetine on a new β -cyclodextrin bonded phase column by HPLC. <i>Separation Science and Technology</i> , 2002, 37, 1401-1415.	1.3	9

#	ARTICLE	IF	CITATIONS
347	Retention of ionizable compounds on HPLC. Modelling retention for neutral and ionizable compounds by linear solvation energy relationships. <i>Chromatographia</i> , 2002, 56, 431-437.	0.7	7
348	Synthesis and evaluation of a ferrocene-bonded silica phase for reversed-phase high-performance liquid chromatography. <i>Chromatographia</i> , 2002, 55, 549-553.	0.7	8
349	Retention of ionizable compounds on HPLC. Modelling retention in reversed-phase liquid chromatography as a function of pH and solvent composition with methanol-water mobile phases. <i>Chromatographia</i> , 2002, 55, 565-571.	0.7	27
350	Automated capillary liquid chromatography for simultaneous determination of neuroactive amines and amino acids. <i>Journal of Chromatography A</i> , 2002, 962, 105-115.	1.8	50
351	Mechanistic molecular model of liquid-liquid chromatography. <i>Journal of Chromatography A</i> , 2002, 965, 109-116.	1.8	66
352	Analysis of linear and cyclic oligomers in polyamide-6 without sample preparation by liquid chromatography using the sandwich injection method. <i>Journal of Chromatography A</i> , 2002, 949, 307-326.	1.8	14
353	Comparative study of the linear solvation energy relationship, linear solvent strength theory, and typical-conditions model for retention prediction in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2002, 965, 3-23.	1.8	58
354	Retention of ionizable compounds on high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2002, 945, 83-96.	1.8	52
355	Comparison of retention of aromatic hydrocarbons with polar groups in binary reversed-phase high-performance liquid chromatography systems. <i>Journal of Chromatography A</i> , 2002, 947, 167-183.	1.8	22
356	Combination of linear solvent strength model and quantitative structure-retention relationships as a comprehensive procedure of approximate prediction of retention in gradient liquid chromatography. <i>Journal of Chromatography A</i> , 2002, 962, 41-55.	1.8	73
357	Column selectivity in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2002, 961, 195-215.	1.8	227
358	Characterization and comparison of HPLC columns for gradient elution. <i>Chromatographia</i> , 2003, 57, S153-S161.	0.7	19
359	New approaches to chromatographic determination of lipophilicity of xenobiotics. <i>Analytical and Bioanalytical Chemistry</i> , 2003, 377, 803-811.	1.9	38
360	Monolithic silica columns with chemically bonded β -cyclodextrin as a stationary phase for enantiomer separations of chiral pharmaceuticals. <i>Analytical and Bioanalytical Chemistry</i> , 2003, 377, 892-901.	1.9	70
361	Prediction of gradient retention from the linear solvent strength (LSS) model, quantitative structure-retention relationships (QSRR), and artificial neural networks (ANN). <i>Journal of Separation Science</i> , 2003, 26, 271-282.	1.3	71
362	Simultaneous determination of neutral and acidic pharmaceuticals in wastewater by high-performance liquid chromatography-post-column photochemically induced fluorimetry. <i>Journal of Chromatography A</i> , 2003, 993, 29-37.	1.8	49
363	Preparative separation of quinolines by centrifugal partition chromatography with gradient elution. <i>Journal of Chromatography A</i> , 2003, 1011, 55-65.	1.8	22
364	Application of the reversed-phase liquid chromatographic model to describe the retention behaviour of polydisperse macromolecules in gradient and isocratic liquid chromatography. <i>Journal of Chromatography A</i> , 2003, 988, 53-67.	1.8	34

#	ARTICLE	IF	CITATIONS
365	Solvent Strength Parameters and Retention Factors in Pure Water Using UNIFAC-Predicted Activity Coefficients. <i>Industrial & Engineering Chemistry Research</i> , 2003, 42, 6320-6330.	1.8	3
366	The Column in Liquid Chromatography. , 2003, , 267-429.		4
367	Mechanistic Studies of the Separation of an HIV Protease Inhibitor from Its Piperazine Diastereomer by Reversed Phase High Performance Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2003, 26, 3343-3355.	0.5	0
368	QSPR Prediction of Three Partition Properties for Phenylacrylates. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2003, 26, 2065-2082.	0.5	1
370	Basic TLC Techniques, Materials, and Apparatus. , 2003, , .		3
371	HPLC-MS(MS) for bioanalysis in drug discovery and development. <i>Handbook of Analytical Separations</i> , 2003, , 271-292.	0.8	0
372	Method development in reversed-phase chromatography. <i>Handbook of Analytical Separations</i> , 2003, 4, 185-214.	0.8	12
373	Relationships Between LC Retention, Octanol-Water Partition Coefficient, and Fungistatic Properties of 2-(2,4-Dihydroxyphenyl)benzothiazoles. <i>Journal of AOAC INTERNATIONAL</i> , 2004, 87, 579-586.	0.7	12
374	Establishing a Selectivity Survey as Part of the Method Development Activity for an Isocratic Reversed-Phase Liquid Chromatographic Method. <i>Journal of Chromatographic Science</i> , 2004, 42, 184-190.	0.7	2
375	Using HPLC Retention Parameters to Estimate Fish Bioconcentration Factors of Organic Compounds. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2004, 27, 1861-1873.	0.5	4
376	Preparation and Evaluation of an End-Capped p-tert-Butyl-Calix[4]arene-Bonded-Silica Stationary Phase for Reversed-Phase High-Performance Liquid Chromatography. <i>Chromatographia</i> , 2004, 59, .	0.7	11
377	Phase System Selectivity and Peak Capacity in Liquid Column Chromatography ? the Impact on Two-Dimensional Separations. <i>Chromatographia</i> , 2004, 60, .	0.7	34
378	Demonstration of simultaneous anion-exchange and reversed-phase behavior on a strong anion-exchange column. <i>Journal of Chromatography A</i> , 2004, 1028, 239-245.	1.8	14
379	Accurate prediction of the retention behaviour of polydisperse macromolecules based on a minimum number of experiments. <i>Journal of Chromatography A</i> , 2004, 1041, 43-51.	1.8	5
380	High-performance liquid chromatography of di- and trisubstituted aromatic positional isomers on 1,3-alternate 25,27-dipropoxy-26,28-bis-[3-propyloxy]-calix[4]arene-bonded silica gel stationary phase. <i>Journal of Chromatography A</i> , 2004, 1055, 21-28.	1.8	30
381	Use of binary and ternary mobile phases in reversed-phase high performance liquid chromatographic separation of chiral compounds. <i>Journal of Separation Science</i> , 2004, 27, 304-310.	1.3	3
382	Controlling the retention in capillary LC with solvents, temperature, and electric fields. <i>Journal of Separation Science</i> , 2004, 27, 1402-1418.	1.3	25
383	Wheat digalactosyldiacylglycerol molecular species profiling using porous graphitic carbon stationary phase. <i>Journal of Separation Science</i> , 2004, 27, 1313-1322.	1.3	14

#	ARTICLE	IF	CITATIONS
384	Ghost peaks in reversed-phase gradient HPLC: a review and update. <i>Journal of Chromatography A</i> , 2004, 1052, 1-11.	1.8	64
385	Stationary-phase effects in gradient high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2004, 1030, 33-41.	1.8	22
386	Optimisation technique for stepwise gradient elution in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2004, 1033, 283-289.	1.8	50
387	Recursion equations in predicting band width under gradient elution. <i>Journal of Chromatography A</i> , 2004, 1040, 19-31.	1.8	15
388	Column selection for liquid chromatographic estimation of the $\log K_{ow}$ hydrophobicity parameter. <i>Journal of Chromatography A</i> , 2004, 1042, 37-48.	1.8	45
389	RP-HPLC Determination of Octanol-Water Partition Coefficients for Bioactive Compounds from Chinese Herbal Medicines. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2004, 27, 465-479.	0.5	6
390	Retention data for some carbonyl derivatives of imidazo[2,1-c][1,2,4]triazine in reversed-phase systems in TLC and HPLC and their use for determination of lipophilicity. Part 1. Lipophilicity of 8-aryl-3-phenyl-6,7-dihydro-4H-imidazo[2,1-c][1,2,4]triazin-4-ones. <i>Journal of Planar Chromatography - Modern TLC</i> , 2005, 18, 274-281.	0.6	19
391	Peak capacity in gradient ultra performance liquid chromatography (UPLC). <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 38, 337-343.	1.4	133
392	Use of reversed-phase liquid chromatography for determining the lipophilicity of β -aryl-N-cyclopropyl nitrones. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 39, 1057-1062.	1.4	14
393	New approach to linear gradient elution used for optimisation in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2005, 1068, 279-287.	1.8	42
394	Theory of peak capacity in gradient elution. <i>Journal of Chromatography A</i> , 2005, 1079, 153-161.	1.8	313
395	A practical approach to transferring linear gradient elution methods. <i>Journal of Chromatography A</i> , 2005, 1077, 110-119.	1.8	48
396	Characterization of immobilized artificial membrane (IAM) and XTerra columns by means of chromatographic models. <i>Journal of Chromatography A</i> , 2005, 1081, 163-173.	1.8	29
397	Fast Determination of Lipophilicity by HPLC. <i>Chromatographia</i> , 2005, 62, 251-255.	0.7	42
398	pH/Organic Solvent Double-Gradient Reversed-Phase HPLC. <i>Analytical Chemistry</i> , 2005, 77, 449-458.	3.2	44
399	The effect of stationary phase on lipophilicity determination of β -blockers using reverse-phase chromatographic systems. <i>Biomedical Chromatography</i> , 2005, 19, 725-736.	0.8	26
400	Retention behavior of phthalic, 3-nitrophthalic, and 4-nitrophthalic acids in ion-suppression reversed-phase high performance liquid chromatography using acids instead of buffers as ion-suppressors. <i>Journal of Separation Science</i> , 2005, 28, 1179-1187.	1.3	17
401	Retention process in reversed phase TLC systems with polar bonded stationary phases. <i>Journal of Separation Science</i> , 2005, 28, 566-574.	1.3	4

#	ARTICLE	IF	CITATIONS
402	Theoretical opportunities and actual limitations of pH gradient HPLC. <i>Analytical and Bioanalytical Chemistry</i> , 2005, 382, 718-727.	1.9	25
403	Fundamentals of Reversed Phase Chromatography: Thermodynamic and Exothermodynamic Treatment. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2005, 28, 965-1054.	0.5	56
404	Microfluidic routing of aqueous and organic flows at high pressures: fabrication and characterization of integrated polymer microvalve elements. <i>Lab on A Chip</i> , 2005, 5, 184.	3.1	27
405	Estimating Partition Coefficients for Fuel-Water Systems: Developing Linear Solvation Energy Relationships Using Linear Solvent Strength Theory To Handle Mixtures. <i>Environmental Science & Technology</i> , 2005, 39, 2702-2710.	4.6	15
406	Theory of Retention in Reversed Phase Liquid Chromatography with Ternary Mobile Phases. <i>Journal of Physical Chemistry B</i> , 2005, 109, 21056-21062.	1.2	7
407	Peak Capacity Optimization of Peptide Separations in Reversed-Phase Gradient Elution Chromatography: Fixed Column Format. <i>Analytical Chemistry</i> , 2006, 78, 3406-3416.	3.2	126
408	High-Performance Liquid Chromatography: Analytic and Preparative Applications in Protein-Structure Determination. <i>Methods of Biochemical Analysis</i> , 2006, 29, 59-136.	0.2	12
409	Analytical Solutions of the Ideal Model for Gradient Liquid Chromatography. <i>Analytical Chemistry</i> , 2006, 78, 7828-7840.	3.2	23
410	Chapter 15 High-pressure liquid chromatography. <i>Comprehensive Analytical Chemistry</i> , 2006, 47, 485-559.	0.7	6
411	Heteroscedasticity of retention factor and adequate modeling in micellar liquid chromatography. <i>Analytica Chimica Acta</i> , 2006, 576, 229-238.	2.6	16
412	Ultra-performance liquid chromatography-tandem mass spectrometry: A novel challenge in multiresidue pesticide analysis in food. <i>Analytica Chimica Acta</i> , 2006, 577, 8-17.	2.6	68
413	Multilinear gradient elution optimisation in reversed-phase liquid chromatography using genetic algorithms. <i>Journal of Chromatography A</i> , 2006, 1120, 299-307.	1.8	46
414	Peak compression in reversed-phase gradient elution. <i>Journal of Chromatography A</i> , 2006, 1111, 32-39.	1.8	107
415	Effect of temperature on the retention of ionizable compounds in reversed-phase liquid chromatography: Application to method development. <i>Journal of Chromatography A</i> , 2006, 1118, 234-243.	1.8	69
416	Can the theory of gradient liquid chromatography be useful in solving practical problems?. <i>Journal of Chromatography A</i> , 2006, 1126, 195-218.	1.8	85
417	Selectivity in reversed-phase separations. <i>Journal of Chromatography A</i> , 2006, 1127, 161-174.	1.8	110
418	Chromatographic behaviour of peptides on a mixed-mode stationary phase with an embedded charged group by capillary electrochromatography and high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2006, 1136, 221-225.	1.8	25
419	The chemical interpretation and practice of linear solvation energy relationships in chromatography. <i>Journal of Chromatography A</i> , 2006, 1126, 143-194.	1.8	461

#	ARTICLE	IF	CITATIONS
420	Chromatographic retention of adamantane derivatives in high-performance liquid chromatography. Russian Journal of Physical Chemistry A, 2006, 80, 435-439.	0.1	3
421	Nonlinear Retention Relationships in Reversed-Phase Chromatography. Chromatographia, 2006, 63, S45-S53.	0.7	42
422	The RP-HPLC measurement and QSPR analysis of logPo/w values of several Pt(II) complexes. Journal of Inorganic Biochemistry, 2006, 100, 1199-1207.	1.5	88
423	Column selectivity for two-dimensional liquid chromatography. Journal of Separation Science, 2006, 29, 1763-1783.	1.3	94
424	A generic stepwise optimization strategy for liquid chromatography electrospray ionization tandem mass spectrometry methods. Journal of Mass Spectrometry, 2006, 41, 1334-1345.	0.7	20
425	Reversed-Phase HPLC. , 2006, , 139-239.		13
426	Characterization of High-Pressure Liquid Chromatography Columns using Chromatographic Methods. Analytical Letters, 2006, 39, 2095-2152.	1.0	18
427	Mathematics of Optimization and Scaling for the Practicing Chromatographer. Journal of Liquid Chromatography and Related Technologies, 2007, 30, 605-647.	0.5	2
428	Effects of π - π Interactions on the Separation of PAHs on Phenyl-Type Stationary Phases. Journal of Liquid Chromatography and Related Technologies, 2007, 31, 324-347.	0.5	25
429	Analysis of the Ideal Model for a Single Component in Preparative Gradient Elution Chromatography. Analytical Chemistry, 2007, 79, 2507-2517.	3.2	4
430	Optimization of the separation of chlorophenols with stepwise gradient elution in reversed phase liquid chromatography. Journal of Separation Science, 2007, 30, 2687-2692.	1.3	6
431	Evaluation of electronic, lipophilic and membrane affinity effects on antiproliferative activity of 5-substituted-2-(2,4-dihydroxyphenyl)-1,3,4-thiadiazoles against various human cancer cells. European Journal of Medicinal Chemistry, 2007, 42, 940-947.	2.6	84
432	The bandwidth in gradient elution chromatography with a retained organic modifier. Journal of Chromatography A, 2007, 1145, 67-82.	1.8	43
433	Comprehensive two-dimensional liquid chromatography with parallel gradients for separation of phenolic and flavone antioxidants. Journal of Chromatography A, 2007, 1149, 73-87.	1.8	128
434	Simple algorithms for fitting and optimisation for multilinear gradient elution in reversed-phase liquid chromatography. Journal of Chromatography A, 2007, 1157, 178-186.	1.8	42
435	Gradient elution separation and peak capacity of columns packed with porous shell particles. Journal of Chromatography A, 2007, 1163, 203-211.	1.8	89
436	Could linear solvation energy relationships give insights into chiral recognition mechanisms?. Journal of Chromatography A, 2007, 1166, 70-78.	1.8	23
437	Could linear solvation energy relationships give insights into chiral recognition mechanisms?. Journal of Chromatography A, 2007, 1166, 61-69.	1.8	34

#	ARTICLE	IF	CITATIONS
438	QSAR of Antiproliferative Activity of <i>N</i> -Substituted 2-Amino-5-(2,4-dihydroxyphenyl)-1,3,4-thiadiazoles in Various Human Cancer Cells. QSAR and Combinatorial Science, 2008, 27, 607-617.	1.3	9
439	Preparation and HPLC evaluation of a new 1,3-alternate 25,27-bis(4-chlorobenzoyloxy)-26,28-bis(3-propyloxy)calix[4]arene silica bonded stationary phase. Journal of Separation Science, 2008, 31, 926-934.	1.3	16
440	Stationary phases for hydrophilic interaction chromatography, their characterization and implementation into multidimensional chromatography concepts. Journal of Separation Science, 2008, 31, 1421-1437.	1.3	219
441	Brief analysis of the retention process in RP-HPLC systems with a C ₃₀ bonded stationary phase. Journal of Separation Science, 2008, 31, 2417-2423.	1.3	2
442	Response surface methodology and support vector machine for the optimization of separation in linear gradient elution. Journal of Separation Science, 2008, 31, 3864-3870.	1.3	16
443	The ultimate band compression factor in gradient elution chromatography. Journal of Chromatography A, 2008, 1178, 79-91.	1.8	38
444	Optimization of separation in two-dimensional high-performance liquid chromatography by adjusting phase system selectivity and using programmed elution techniques. Journal of Chromatography A, 2008, 1189, 207-220.	1.8	70
445	Hydrophilic interaction chromatography of nucleotides and their pathway intermediates on titania. Journal of Chromatography A, 2008, 1187, 87-93.	1.8	77
446	Methacrylate monolithic capillary columns for gradient peptide separations. Journal of Chromatography A, 2008, 1208, 109-115.	1.8	17
447	Effect of n-octanol in the mobile phase on lipophilicity determination by reversed-phase high-performance liquid chromatography on a modified silica column. Journal of Chromatography A, 2008, 1209, 111-119.	1.8	37
448	The application of gradient reversed-phase high-performance liquid chromatography to the pKa and determination of polyprotic analytes. Journal of Chromatography A, 2008, 1214, 109-114.	1.8	32
449	Application of Multilinear Gradient Elution for Optimization of Separation of Chlorophenols Using Derringer's Desirability Function. Chromatographia, 2008, 67, 169-172.	0.7	6
450	Influence of pH on Retention in Linear Organic Modifier Gradient RP HPLC. Analytical Chemistry, 2008, 80, 7855-7861.	3.2	24
451	Method transfer for fast liquid chromatography in pharmaceutical analysis: Application to short columns packed with small particle. Part II: Gradient experiments. European Journal of Pharmaceutics and Biopharmaceutics, 2008, 68, 430-440.	2.0	191
452	Prediction of Analyte Retention for Ion Chromatography Separations Performed Using Elution Profiles Comprising Multiple Isocratic and Gradient Steps. Analytical Chemistry, 2008, 80, 2474-2482.	3.2	38
453	Effect of Mobile Phase Composition on Henry's Constants of 2-Amino-3-phenyl-propanoic Acid, 2-Amino-3-(3-indolyl)-propanoic Acid, and 2-Amino-3-(4-hydroxyphenyl)-propanoic Acid in a Capcell Pak C ₁₈ Chromatography. Journal of Chemical & Engineering Data, 2008, 53, 2613-2621.	1.0	16
454	Optimization of Gradient Separation for Chromatographic Fingerprint of Herbal Medicine: A Predictive Model Combined with Grid Search. Journal of Chromatographic Science, 2008, 46, 722-729.	0.7	2
455	Characterization of the GEMINI C18 _μ Column: Lipophilicity Measurement and LSER. Journal of Liquid Chromatography and Related Technologies, 2008, 31, 2198-2218.	0.5	10

#	ARTICLE	IF	CITATIONS
456	Selectivity of Calixarene-bonded Silica-phases in HPLC: Description of Special Characteristics with a Multiple Term Linear Equation at Two Different pH-Values. <i>Analytical Sciences</i> , 2008, 24, 1157-1164.	0.8	9
458	RP-HPLC measurement and quantitative structure - property relationship analysis of the n-octanol - water partitioning coefficients of selected metabolites of polybrominated diphenyl ethers. <i>Environmental Chemistry</i> , 2008, 5, 332.	0.7	23
460	Synthesis and separation of protected tripeptide epimers by RP-HPLC. <i>International Journal of Peptide and Protein Research</i> , 1995, 45, 217-224.	0.1	9
462	Preparation and characterization of polymethacrylate monolithic capillary columns with dual hydrophilic interaction reversed-phase retention mechanism for polar compounds. <i>Journal of Separation Science</i> , 2009, 32, 2530-2543.	1.3	65
463	Utilization of dual retention mechanism on columns with bonded PEG and diol stationary phases for adjusting the separation selectivity of phenolic and flavone natural antioxidants. <i>Journal of Separation Science</i> , 2009, 32, 3603-3619.	1.3	93
464	Derived chromatographic indices as effective tools to study the self-aggregation process of bile acids. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 50, 613-621.	1.4	23
465	Peak compression technique in high-performance liquid chromatography. <i>Science Bulletin</i> , 2009, 54, 2805-2809.	4.3	5
466	Analytical tools for the physicochemical profiling of drug candidates to predict absorption/distribution. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 394, 707-729.	1.9	68
467	Fast log P determination by ultra-high-pressure liquid chromatography coupled with UV and mass spectrometry detections. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 394, 1919-1930.	1.9	41
468	Theoretical models in LC based bioanalytical method development. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 49, 1-10.	1.4	8
469	Optimization of two-dimensional gradient liquid chromatography separations. <i>Journal of Chromatography A</i> , 2009, 1216, 3443-3457.	1.8	89
470	Retention models for isocratic and gradient elution in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 1737-1755.	1.8	134
471	Lipophilicity data for some preservatives estimated by reversed-phase liquid chromatography and different computation methods. <i>Journal of Chromatography A</i> , 2009, 1216, 2456-2465.	1.8	48
472	Preparation and evaluation of 1,3-alternate 25,27-bis-(pentafluorobenzoyloxy)-26,28-bis-(3-propyloxy)-calix[4]arene-bonded silica gel high performance liquid chromatography stationary phase. <i>Journal of Chromatography A</i> , 2009, 1216, 3954-3960.	1.8	30
473	Critical comparison of performances of superficially porous particles and sub-2 μ m particles under optimized ultra-high pressure conditions. <i>Journal of Chromatography A</i> , 2009, 1216, 4597-4605.	1.8	82
474	Description of retention characteristics of calixarene-bonded stationary phases in dependence of the methanol content in the mobile phase. <i>Journal of Chromatography A</i> , 2009, 1216, 6285-6294.	1.8	21
475	Detection issues in two-dimensional on-line chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 7785-7792.	1.8	32
476	Interconversion of gradient and isocratic retention data in reversed-phase liquid chromatography: Effect of the uptake of eluent modifier on the retention of analytes. <i>Journal of Chromatography A</i> , 2009, 1216, 8630-8635.	1.8	5

#	ARTICLE	IF	CITATIONS
477	The influence of the eluent composition on the retention of derivatives of some aromatic heterocyclic compounds in reversed-phase high-performance liquid chromatography. <i>Russian Journal of Physical Chemistry A</i> , 2009, 83, 1223-1229.	0.1	7
478	Generation and Limitations of Peak Capacity in Online Two-Dimensional Liquid Chromatography. <i>Analytical Chemistry</i> , 2009, 81, 3879-3888.	3.2	38
479	New Approaches to Linear Gradient Elution Used for Optimization in Reversed-Phase Liquid Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 1527-1576.	0.5	23
480	Optimization of Chromatographic Systems for Determination of Lipophilicity for Selected Isoquinoline Alkaloids. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 2265-2280.	0.5	6
481	Chromatographic retention prediction and octanol/water partition coefficient determination of monobasic weak acidic compounds in ion-suppression reversed-phase liquid chromatography using acids as ion-suppressors. <i>Talanta</i> , 2009, 79, 752-761.	2.9	17
482	Simultaneous Determination of Medicinal Ingredients in So-called Health-promoting Food Using Liquid Chromatography Tandem Mass Spectrometry with a Pentafluorophenyl Stationary Phase. <i>Journal of Health Science</i> , 2009, 55, 183-191.	0.9	7
483	Chromatographic and computational studies of the physicochemical properties of cholinesterase inhibitors – alkyl- and arylcarbamate derivatives of <i>N</i> -benzylpiperidine and <i>N</i> -benzylpiperazine. <i>Journal of Planar Chromatography - Modern TLC</i> , 2010, 23, 359-364.	0.6	2
484	The prediction of the peak width at half height in HPLC. <i>Chinese Journal of Chemistry</i> , 2010, 9, 237-244.	2.6	1
485	Retention prediction and hydrophobicity estimation of weak acidic compounds by reversed-phase liquid chromatography using acetic and perchloric acids as ion suppressors. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 398, 2731-2743.	1.9	12
486	Two-column Sequential Injection Chromatography – New approach for fast and effective analysis and its comparison with gradient elution chromatography. <i>Analytica Chimica Acta</i> , 2010, 668, 61-66.	2.6	24
487	Quantitative structure-retention relationship for photosystem II inhibitors in RP-HPLC. <i>Chinese Journal of Chemistry</i> , 1998, 16, 514-520.	2.6	0
490	Characterization of stationary phases for reversed-phase chromatography. <i>Journal of Separation Science</i> , 2010, 33, 453-463.	1.3	30
491	Comparison of various second-dimension gradient types in comprehensive two-dimensional liquid chromatography. <i>Journal of Separation Science</i> , 2010, 33, 1382-1397.	1.3	51
492	Chromatographic behavior of a new hybrid type RP material containing silica bonded 1,3-bis(2,2,7,7-tetrakis(cyanopropoxy)hexafluoroethyl)hexane. <i>Journal of Separation Science</i> , 2010, 33, 2956-2964.	1.3	10
493	Characterization of calixarene-bonded stationary phases. <i>Journal of Separation Science</i> , 2010, 33, 2930-2942.	1.3	10
494	Selectivity of calixarene-bonded silica phases in HPLC: Description of special characteristics with a multiple term linear equation at different methanol concentrations. <i>Journal of Separation Science</i> , 2010, 33, 2943-2955.	1.3	6
495	Scorpion venom complexity fractal analysis. Its relevance for comparing venoms. <i>Journal of Theoretical Biology</i> , 2010, 267, 405-416.	0.8	8
496	Improved reversed-phase gradient retention modeling. <i>Journal of Chromatography A</i> , 2010, 1217, 3794-3803.	1.8	138

#	ARTICLE	IF	CITATIONS
497	Comparison of hypercrosslinked polystyrene columns for the separation of nitrogen group-types in petroleum using High Performance Liquid Chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 6178-6185.	1.8	22
498	Characterization of Ascentis RP-Amide column: Lipophilicity measurement and linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2010, 1217, 65-74.	1.8	20
499	Characterization of adsorption processes in analytical liquid–solid chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 792-812.	1.8	92
500	3D structure-based protein retention prediction for ion-exchange chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 1343-1353.	1.8	62
501	The kinetic plot method applied to gradient chromatography: Theoretical framework and experimental validation. <i>Journal of Chromatography A</i> , 2010, 1217, 2787-2795.	1.8	90
502	Ƴ-Selective stationary phases: (III) Influence of the propyl phenyl ligand density on the aromatic and methylene selectivity of aromatic compounds in reversed phase liquid chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 5377-5383.	1.8	19
503	Automated instrumental method for on-line fraction analysis and peak deconvolution in gradient multicomponent overloaded high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 4919-4924.	1.8	5
504	Evaluation of 1.0mm i.d. column performances on ultra high pressure liquid chromatography instrumentation. <i>Journal of Chromatography A</i> , 2010, 1217, 4925-4933.	1.8	36
505	Combined effects of mobile phase composition and temperature on the retention of homologous and polar test compounds on polydentate C8 column. <i>Journal of Chromatography A</i> , 2010, 1217, 6052-6060.	1.8	18
506	Protein- versus peptide fractionation in the first dimension of two-dimensional high-performance liquid chromatography-matrix-assisted laser desorption/ionization tandem mass spectrometry for qualitative proteome analysis of tissue samples. <i>Journal of Chromatography A</i> , 2010, 1217, 6159-6168.	1.8	16
507	Study of the retention and selectivity of cholesterol bonded phases with different linkage spacers. <i>Journal of Chromatography A</i> , 2010, 1217, 6891-6897.	1.8	27
508	Selectivity optimization in green chromatography by gradient stationary phase optimized selectivity liquid chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 7222-7230.	1.8	42
509	A parallel pore and surface diffusion model for predicting the adsorption and elution profiles of lispro insulin and two impurities in gradient-elution reversed phase chromatography. <i>Journal of Chromatography A</i> , 2010, 1217, 8103-8120.	1.8	8
510	Lipophilicity of novel antitumour and analgesic active 8-aryl-2,6,7,8-tetrahydroimidazo[2,1-c][1,2,4]triazine-3,4-dione derivatives determined by reversed-phase HPLC and computational methods. <i>European Journal of Medicinal Chemistry</i> , 2010, 45, 2644-2649.	2.6	24
512	Adsorption Isotherms of Substituted Benzophenones on a Reverse-Phase Liquid Chromatography System: Effect of the Mobile-Phase Composition. <i>Journal of Chemical & Engineering Data</i> , 2010, 55, 4768-4774.	1.0	5
513	Effect of Ethanol Content on Adsorption Equilibria of Some Useful Amino Acids in Poly-4-vinylpyridine Chromatography. <i>Journal of Chemical & Engineering Data</i> , 2010, 55, 3327-3333.	1.0	7
515	General Strategy for Performing Temperature Programming in High Performance Liquid Chromatography: Prediction of Linear Temperature Gradients. <i>Analytical Chemistry</i> , 2011, 83, 2227-2233.	3.2	18
516	Correlation between n-octanol/water partition coefficient and liquid chromatographic retention for caffeine and its metabolites, and some structure-pharmacokinetic considerations. <i>Journal of Pharmacy and Pharmacology</i> , 2011, 39, 252-260.	1.2	26

#	ARTICLE	IF	CITATIONS
517	Selectivity Tests of Stationary Phases for Reversed-Phase HPLC. <i>Analytical Letters</i> , 2011, 44, 1640-1662.	1.0	10
518	RP-HPLC Method for Simultaneous Determination of Butenafine Hydrochloride and Betamethasone Dipropionate in a Cream Formulation. <i>Journal of AOAC INTERNATIONAL</i> , 2011, 94, 106-109.	0.7	10
519	Gradient reversed-phase high-performance chromatography of ionogenic analytes. <i>TrAC - Trends in Analytical Chemistry</i> , 2011, 30, 1372-1381.	5.8	20
520	Synthesis, structure elucidation, determination of the lipophilicity and identification of antitumour activities in vitro of novel 3-(2-furanyl)-8-aryl-7,8-dihydroimidazo[2,1-c][1,2,4]triazin-4(6H)-ones with a low cytotoxicity towards normal human skin fibroblast cells. <i>Bioorganic and Medicinal Chemistry</i> , 2011, 19, 5103-5116.	1.4	22
521	Determination of n-octanol/water partition coefficient for DDT-related compounds by RP-HPLC with a novel dual-point retention time correction. <i>Chemosphere</i> , 2011, 83, 131-136.	4.2	43
522	Retention times and bandwidths in reversed-phase gradient liquid chromatography of peptides and proteins. <i>Journal of Chromatography A</i> , 2011, 1218, 8874-8889.	1.8	24
523	Determination of thiamine (vitamin B1) in pharmaceutical tablets and human urine by titania-based ligand-exchange hydrophilic interaction chromatography. <i>Analytical Methods</i> , 2011, 3, 1568.	1.3	24
524	Comparison of LC Columns Packed with 2.6 μ m Core-Shell and Sub-2 μ m Porous Particles for Gradient Separation of Antibiotics. <i>Chromatographia</i> , 2011, 74, 19-27.	0.7	13
525	Theory-guided efficient strategy to maximize speed and resolution in rapid gradient LC-MS/MS bioanalysis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2011, 879, 1917-1926.	1.2	9
526	Comparison of theoretical and experimental models for characterizing solvent properties using reversed phase liquid chromatography. <i>Analytica Chimica Acta</i> , 2011, 702, 37-44.	2.6	10
527	Preparation and chromatographic performance of calix[4]crown \AA 5 macrocycle \AA bonded silica stationary phase. <i>Journal of Separation Science</i> , 2011, 34, 393-401.	1.3	7
528	Novel criteria for fast searching for optimal method in gradient ion chromatography: An integrated approach. <i>Journal of Separation Science</i> , 2011, 34, 780-788.	1.3	4
529	Effects of functional monomers on retention behavior of small and large molecules in monolithic capillary columns at isocratic and gradient conditions. <i>Journal of Separation Science</i> , 2011, 34, 2054-2062.	1.3	21
530	Peak capacity optimization in comprehensive two dimensional liquid chromatography: A practical approach. <i>Journal of Chromatography A</i> , 2011, 1218, 64-73.	1.8	53
531	Kinetic performance limits of constant pressure versus constant flow rate gradient elution separations. Part II: Experimental. <i>Journal of Chromatography A</i> , 2011, 1218, 1170-1184.	1.8	20
532	Innovative high-performance liquid chromatography method development for the screening of 19 antimalarial drugs based on a generic approach, using design of experiments, independent component analysis and design space. <i>Journal of Chromatography A</i> , 2011, 1218, 5205-5215.	1.8	86
533	Multilinear gradient elution optimization in reversed-phase liquid chromatography based on logarithmic retention models: Application to separation of a set of purines, pyrimidines and nucleosides. <i>Journal of Chromatography A</i> , 2011, 1218, 5658-5663.	1.8	17
534	Stationary and mobile phases in hydrophilic interaction chromatography: a review. <i>Analytica Chimica Acta</i> , 2011, 692, 1-25.	2.6	610

#	ARTICLE	IF	CITATIONS
535	Transfer-volume effects in two-dimensional chromatography: Adsorption-phenomena in second-dimension size-exclusion chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 1147-1152.	1.8	9
536	Kinetic performance limits of constant pressure versus constant flow rate gradient elution separations. Part I: Theory. <i>Journal of Chromatography A</i> , 2011, 1218, 1153-1169.	1.8	23
537	EFFECT OF ETHANOL CONTENT ON MASS-TRANSFER PARAMETERS AND HETP INDEXES OF AMINO ACIDS IN A POLY-4-VINYLPYRIDINE CHROMATOGRAPHY. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2011, 34, 456-475.	0.5	4
538	Evaluation of the Chromatographic Performance of di-and tri-amide Calix[4]arene Derivatives Bonded Silica Stationary Phases. <i>Journal of Macromolecular Science - Pure and Applied Chemistry</i> , 2012, 49, 1022-1029.	1.2	2
539	Optimum separation condition of five bioactive compounds by HCl program in HPLC. <i>Acta Chromatographica</i> , 2012, 24, 65-74.	0.7	0
540	Optimization of comprehensive two-dimensional gradient chromatography coupling in-line hydrophilic interaction and reversed phase liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1268, 91-101.	1.8	65
541	On-line comprehensive two-dimensional separations of charged compounds using reversed-phase high performance liquid chromatography and hydrophilic interaction chromatography. Part I: Orthogonality and practical peak capacity considerations. <i>Journal of Chromatography A</i> , 2012, 1262, 148-159.	1.8	58
542	Maximizing the peak capacity using coupled columns packed with 2.6 μ m core-shell particles operated at 1200bar. <i>Journal of Chromatography A</i> , 2012, 1256, 72-79.	1.8	23
543	Determination of n-octanol/water partition coefficients of weak ionizable solutes by RP-HPLC with neutral model compounds. <i>Talanta</i> , 2012, 97, 355-361.	2.9	24
544	Determination of the lipophilicity of some new thiosemicarbaside derivatives by reversed-phase thin-layer chromatography. <i>Acta Chromatographica</i> , 2012, 24, 271-290.	0.7	3
545	A novel evaluation method for extrapolated retention factor in determination of n-octanol/water partition coefficient of halogenated organic pollutants by reversed-phase high performance liquid chromatography. <i>Analytica Chimica Acta</i> , 2012, 713, 130-135.	2.6	18
546	The influence of the organic modifier in hydro-organic mobile phase on separation selectivity of steroid hormones separation using cholesterol-bonded stationary phases. <i>Journal of Chromatography A</i> , 2012, 1245, 90-97.	1.8	17
547	Inversion of chromatographic elution orders of peptides and its importance for proteomics. <i>Journal of Analytical Chemistry</i> , 2012, 67, 1014-1025.	0.4	10
548	Methods to Isolate and Identify New Plant Signaling Peptides. <i>Signaling and Communication in Plants</i> , 2012, , 217-239.	0.5	3
549	Influence of variation in mobile phase pH and solute pKa with the change of organic modifier fraction on QSRRs of hydrophobicity and RP-HPLC retention of weakly acidic compounds. <i>Talanta</i> , 2012, 101, 64-70.	2.9	16
550	Determination of Melamine in Liquid Milk and Milk Powder by Titania-Based Ligand-Exchange Hydrophilic Interaction Liquid Chromatography. <i>Food Analytical Methods</i> , 2012, 5, 1062-1069.	1.3	23
551	Plant Signaling Peptides. <i>Signaling and Communication in Plants</i> , 2012, , .	0.5	2
552	Novel tetracarboxylatoplatinum($\text{Pt}(\text{CO}_2)_4$) complexes as carboplatin prodrugs. <i>Dalton Transactions</i> , 2012, 41, 14404-14415.	1.6	76

#	ARTICLE	IF	CITATIONS
553	Programmed elution in comprehensive two-dimensional liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1255, 112-129.	1.8	61
554	Columns and optimum gradient conditions for fast secondâ€dimension separations in comprehensive twoâ€dimensional liquid chromatography. <i>Journal of Separation Science</i> , 2012, 35, 1712-1722.	1.3	17
555	Chromatographic Separation of Synthesized Phenolic Lipids from Krill Oil and Dihydroxyphenyl Acetic Acid. <i>JAACS, Journal of the American Oil Chemists' Society</i> , 2012, 89, 597-608.	0.8	4
556	Synthesis and RP HPLC studies of biologically active semicarbazides and their cyclic analogues 1,2,4-triazol-3-ones. <i>Monatshefte FÃ¼r Chemie</i> , 2012, 143, 657-667.	0.9	7
557	Kinetic plot based comparison of the efficiency and peak capacity of high-performance liquid chromatography columns: Theoretical background and selected examples. <i>Journal of Chromatography A</i> , 2012, 1228, 20-30.	1.8	49
558	Effect of pH additive and column temperature on kinetic performance of two different sub-2µm stationary phases for ultrafast separation of charged analytes. <i>Journal of Chromatography A</i> , 2012, 1228, 135-147.	1.8	19
559	Comprehensive two-dimensional liquid chromatography â€” practical impacts of theoretical considerations. A review. <i>Open Chemistry</i> , 2012, 10, 844-875.	1.0	24
560	Hydrophilic interaction liquid chromatography (HILIC)â€”a powerful separation technique. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 231-247.	1.9	950
561	Thorough investigation of the retention mechanisms and retention behavior of amides and sulfonamides on amino column in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1301, 27-37.	1.8	15
563	On-line comprehensive two dimensional separations of charged compounds using reversed-phase high performance liquid chromatography and hydrophilic interaction chromatography. Part II: Application to the separation of peptides. <i>Journal of Chromatography A</i> , 2013, 1306, 27-36.	1.8	61
564	Combined effects of mobile phase composition and temperature on the retention of phenolic antioxidants on an octylsilica polydentate column. <i>Journal of Chromatography A</i> , 2013, 1317, 49-58.	1.8	6
565	Separation and identification of phenolic compounds in canned artichoke by LC/DAD/ESI-MS using coreâ€shell C18 column: A comparative study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 927, 173-180.	1.2	12
566	EFFECT OF LIQUID-PHASE COMPOSITION ON THE HENRY'S CONSTANTS OF VALINE, ISOLEUCINE, AND LEUCINE IN A CAPCELL-PACK C₁₈ CHROMATOGRAPHY. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2013, 36, 180-196.	0.5	1
567	Design Spaces for analytical methods. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 42, 157-167.	5.8	188
568	Chemometric-assisted method development in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1287, 2-22.	1.8	62
569	Approaches to model the retention and peak profile in linear gradient reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1284, 28-35.	1.8	28
570	A novel fractionized sampling and stacking strategy for online hyphenation of solid-phase-based extraction to ultra-high performance liquid chromatography for ultrasensitive analysis. <i>Journal of Chromatography A</i> , 2013, 1316, 29-36.	1.8	25
571	Possibilities of retention prediction in fast gradient liquid chromatography. Part 1: Comparison of separation on packed fully porous, nonporous and monolithic columns. <i>Journal of Chromatography A</i> , 2013, 1278, 37-45.	1.8	7

#	ARTICLE	IF	CITATIONS
572	Fast estimation of adsorption isotherm parameters in gradient elution preparative liquid chromatography II: The competitive case. <i>Journal of Chromatography A</i> , 2013, 1314, 70-76.	1.8	15
573	Fast estimation of adsorption isotherm parameters in gradient elution preparative liquid chromatography. I: The single component case. <i>Journal of Chromatography A</i> , 2013, 1299, 64-70.	1.8	23
574	Study of the peak variance in isocratic and gradient liquid chromatography using the transport model. <i>Journal of Chromatography A</i> , 2013, 1295, 67-81.	1.8	12
575	Quantification aspects of constant pressure (ultra) high pressure liquid chromatography using mass-sensitive detectors with a nebulizing interface. <i>Journal of Chromatography A</i> , 2013, 1274, 118-128.	1.8	5
576	Influence of a strong sample solvent on analyte dispersion in chromatographic columns. <i>Journal of Chromatography A</i> , 2013, 1297, 46-55.	1.8	17
577	Synthesis, <i>In Vitro</i> Antimicrobial and Antioxidant Activities of Some New 4,5-Dihydro-1 <i>H</i> -1,2,4-triazol-5-one Derivatives. <i>Archiv Der Pharmazie</i> , 2013, 346, 470-480.	2.1	18
578	Quantitative Structure Retention Relationship Models in an Analytical Quality by Design Framework: Simultaneously Accounting for Compound Properties, Mobile-Phase Conditions, and Stationary-Phase Properties. <i>Industrial & Engineering Chemistry Research</i> , 2013, 52, 12269-12284.	1.8	28
579	Five different columns in the analysis of basic drugs in hydrophilic interaction liquid chromatography. <i>Open Chemistry</i> , 2013, 11, 1150-1162.	1.0	5
580	Characterization of new N,O-dialkyl phosphoramidate-bonded stationary phases for reversed-phase HPLC – retention and selectivity. <i>Analyst</i> , 2013, 138, 5221.	1.7	22
581	Synthesis and Structure-Activity Relationship Study of Organometallic Bioconjugates of the Cyclic Octapeptide Octreotate. <i>ChemBioChem</i> , 2013, 14, 2472-2479.	1.3	14
582	Optimization of IC Separation Based on Isocratic-to-Gradient Retention Modeling in Combination with Sequential Searching or Evolutionary Algorithm. <i>Journal of Analytical Methods in Chemistry</i> , 2013, 2013, 1-11.	0.7	5
583	The retention behaviour of amino acids in hydrophilic interaction liquid chromatography on zwitterionic stationary phases. <i>Journal of Separation Science</i> , 2013, 36, 2217-2222.	1.3	12
585	Retention Prediction and Hydrophobicity Measurement of Weakly Basic Compounds in Reversed-phase Liquid Chromatography Using Ammonia and Triethylamine as Ion-suppressors. <i>Current Analytical Chemistry</i> , 2013, 10, 172-181.	0.6	8
586	Partial Least Square and Hierarchical Clustering in ADMET Modeling: Prediction of Blood – Brain Barrier Permeation of \pm -Adrenergic and Imidazoline Receptor Ligands. <i>Journal of Pharmacy and Pharmaceutical Sciences</i> , 2013, 16, 622.	0.9	24
587	Evolution and Current Trends in Liquid and Supercritical Fluid Chromatography. <i>Current Chromatography</i> , 2014, 1, 15-40.	0.1	18
588	Fingering dynamics on the adsorbed solute with influence of less viscous and strong sample solvent. <i>Journal of Chemical Physics</i> , 2014, 141, 214701.	1.2	11
589	Theoretical and practical interest in UHPLC technology for 2D-LC. <i>TrAC - Trends in Analytical Chemistry</i> , 2014, 63, 104-112.	5.8	37
590	Evaluation and comparison of <i>n</i> -alkyl chain and polar ligand bonded stationary phases for protein separation in reversed-phase liquid chromatography. <i>Journal of Separation Science</i> , 2014, 37, 2467-2473.	1.3	8

#	ARTICLE	IF	CITATIONS
591	Retention behavior of a homologous series and positional isomers of aliphatic amino acids in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2014, 37, 739-747.	1.3	20
592	Polycyclic aromatic hydrocarbons as test probes to investigate the retention behavior of 1,3-alternate-calix[4]arene silica-bonded stationary phases. <i>Journal of Separation Science</i> , 2014, 37, 543-550.	1.3	17
593	Development of on-line comprehensive two-dimensional liquid chromatography method for the separation of biomass compounds. <i>Journal of Chromatography A</i> , 2014, 1340, 90-98.	1.8	54
594	Retention mechanism assessment and method development for the analysis of iohexol and its related compounds in hydrophilic interaction liquid chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4217-4232.	1.9	7
595	Evaluation of the peak variance in gradient liquid chromatography. <i>Analyst, The</i> , 2014, 139, 1504.	1.7	2
596	Equivalence between the equilibrium dispersive and the transport model in describing band broadening in analytical gradient liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1369, 191-195.	1.8	3
597	Effect of injection matrix concentration on peak shape and separation efficiency in ion chromatography. <i>Journal of Chromatography A</i> , 2014, 1371, 177-183.	1.8	8
598	THEORETICAL AND EMPIRICAL MODELS IN HYDROPHILIC INTERACTION LIQUID CHROMATOGRAPHY. <i>Instrumentation Science and Technology</i> , 2014, 42, 230-266.	0.9	16
599	Combined influences of viscous fingering and solvent effect on the distribution of adsorbed solutes in porous media. <i>RSC Advances</i> , 2014, 4, 34369.	1.7	20
600	A novel method for characterization and comparison of reversed-phase column selectivity. <i>Journal of Chromatography A</i> , 2014, 1361, 153-161.	1.8	9
601	RELATIONSHIP BETWEEN HYDROPHOBICITY AND RPLC RETENTION BEHAVIOR OF AMPHOTERIC COMPOUNDS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 2711-2724.	0.5	5
602	A Novel Class of Bis- and Tris-Chelate Diam(m)inebis(dicarboxylato)platinum(IV) Complexes as Potential Anticancer Prodrugs. <i>Journal of Medicinal Chemistry</i> , 2014, 57, 6751-6764.	2.9	49
603	Some insights on the description of gradient elution in reversed-phase liquid chromatography. <i>Journal of Separation Science</i> , 2014, 37, 2269-2277.	1.3	16
604	Retention and bandwidths prediction in fast gradient liquid chromatography. Part 2 – Core-shell columns. <i>Journal of Chromatography A</i> , 2014, 1337, 57-66.	1.8	16
605	Implementation of gradients of organic solvent in micellar liquid chromatography using DryLab®: Separation of basic compounds in urine samples. <i>Journal of Chromatography A</i> , 2014, 1344, 31-41.	1.8	21
606	Solvent selectivity and strength in reversed-phase liquid chromatography separation of peptides. <i>Journal of Chromatography A</i> , 2014, 1337, 140-146.	1.8	13
607	Gradient-elution parameters in capillary liquid chromatography for high-speed separations of peptides and intact proteins. <i>Journal of Chromatography A</i> , 2014, 1355, 149-157.	1.8	34
608	Investigation of plateau methods for adsorption isotherm determination in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2014, 1354, 129-138.	1.8	19

#	ARTICLE	IF	CITATIONS
609	Importance of instrumentation for fast liquid chromatography in pharmaceutical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 87, 105-119.	1.4	113
610	Determination of the <i>n</i> -octanol/water partition coefficients of weakly ionizable basic compounds by reversed-phase high-performance liquid chromatography with neutral model compounds. <i>Journal of Separation Science</i> , 2014, 37, 3226-3234.	1.3	4
612	Determination of lipophilicity of some new 1,2,4-triazole derivatives by RP-HPLC and RP-TLC and calculated methods. <i>Current Chemistry Letters</i> , 2015, , 101-110.	0.5	5
613	A preliminary study on the selectivity of linear polynuclear aromatic hydrocarbons in SFC using phenyl-type stationary phases. <i>Microchemical Journal</i> , 2015, 121, 136-140.	2.3	9
614	Alternative solvents can make preparative liquid chromatography greener. <i>Green Chemistry</i> , 2015, 17, 4073-4081.	4.6	42
615	Retention modelling in hydrophilic interaction chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 9135-9152.	1.9	34
616	Structure-retention behaviour of biologically active fused 1,2,4-triazinones – Correlation with in silico molecular properties. <i>European Journal of Pharmaceutical Sciences</i> , 2015, 68, 114-126.	1.9	12
617	Instrument contributions to resolution and sensitivity in ultra high performance liquid chromatography using small bore columns: Comparison of diode array and triple quadrupole mass spectrometry detection. <i>Journal of Chromatography A</i> , 2015, 1377, 64-74.	1.8	42
618	Repetitive injection method: A tool for investigation of injection zone formation and its compression in microfluidic liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1381, 110-117.	1.8	16
619	Model-based description of peptide retention on doped reversed-phase media. <i>Journal of Chromatography A</i> , 2015, 1407, 169-175.	1.8	17
620	QuEChERS: a sample preparation for extraction of carbaryl from rat feces. <i>Toxicological and Environmental Chemistry</i> , 2015, 97, 687-699.	0.6	4
621	Effect of Strong Sample Solvent on the Solute Dynamics for More or Less Viscous Sample: A Comparative Study. <i>Procedia IUTAM</i> , 2015, 15, 249-255.	1.2	0
622	Influence of <i>n</i> -octanol in mobile phase on QSRRs of lipophilicity and retention mechanism of acidic and basic compounds in RP-HPLC. <i>RSC Advances</i> , 2015, 5, 28840-28847.	1.7	5
623	Mechanism and Dynamics of Molecular Exchange at the Silica/Binary Solvent Mixtures Interface. <i>Journal of Physical Chemistry A</i> , 2015, 119, 12073-12081.	1.1	5
624	Chromatographic Evaluation of the Lipophilic Properties of Some 1,2,4-Triazole with Potential Antitumour Activity. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 1199-1206.	0.5	4
625	Recent advances in lipophilicity measurement by reversed-phase high-performance liquid chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2015, 68, 28-36.	5.8	58
626	Wide injection zone compression in gradient reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1390, 86-94.	1.8	20
627	A new methodology to determine the isoeluotropic conditions on ultra-performance flash purification stationary phases from analytical reversed liquid chromatography stationary phase. <i>Journal of Chromatography A</i> , 2015, 1397, 59-72.	1.8	2

#	ARTICLE	IF	CITATIONS
628	Comparison of a C30 Bonded Silica Column and Columns with Shorter Bonded Ligands in Reversed-Phase LC. <i>Chromatographia</i> , 2015, 78, 861-871.	0.7	8
629	Synthesis, in vitro antioxidant activity, and physicochemical properties of novel 4,5-dihydro-1H-1,2,4-triazol-5-one derivatives. <i>Journal of Molecular Liquids</i> , 2015, 206, 359-366.	2.3	13
630	Retention prediction and separation optimization under multilinear gradient elution in liquid chromatography with Microsoft Excel macros. <i>Journal of Chromatography A</i> , 2015, 1395, 109-115.	1.8	17
631	Prediction of retention characteristics of heterocyclic compounds. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 9185-9189.	1.9	2
632	Rapid sequential separation of sedimentary lipid biomarkers via selective accelerated solvent extraction. <i>Organic Geochemistry</i> , 2015, 88, 29-34.	0.9	13
633	Possibilities of retention prediction in fast gradient liquid chromatography. Part 3: Short silica monolithic columns. <i>Journal of Chromatography A</i> , 2015, 1410, 76-89.	1.8	15
634	Optimization of conditions in on-line comprehensive two-dimensional reversed phase liquid chromatography. Experimental comparison with one-dimensional reversed phase liquid chromatography for the separation of peptides. <i>Journal of Chromatography A</i> , 2015, 1421, 48-59.	1.8	59
635	Effect of gradient steepness on the kinetic performance limits and peak compression for reversed-phase gradient separations of small molecules. <i>Journal of Chromatography A</i> , 2015, 1409, 152-158.	1.8	8
636	Optimization of reversed-phase chromatography methods for peptide analytics. <i>Journal of Chromatography A</i> , 2015, 1425, 198-203.	1.8	7
637	Determination of Lipophilicity of New Thiosemicarbazide and 1,2,4-triazole-3-thione Derivatives Using Reversed-Phase HPLC Method and Theoretical Calculations. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 430-437.	0.5	6
638	Contrasting selectivity between HPLC and SFC using phenyl-type stationary phases: A study on linear polynuclear aromatic hydrocarbons. <i>Microchemical Journal</i> , 2015, 119, 40-43.	2.3	17
639	Analysis of antibody-drug conjugates by comprehensive on-line two-dimensional hydrophobic interaction chromatography x reversed phase liquid chromatography hyphenated to high resolution mass spectrometry. I. Optimization of separation conditions. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016, 1032, 103-111.	1.2	51
640	Central composite design with the help of multivariate curve resolution in loadability optimization of RP-HPLC to scale-up a binary mixture. <i>Journal of Separation Science</i> , 2016, 39, 1031-1040.	1.3	9
641	Determination of the lipophilicity of quinolinesulfonamides by reversed-phase HPLC and theoretical calculations. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 702-709.	0.5	10
642	Applicability of the critical chromatography concept to proteomic problems. II. Effect of mobile phase on the separation of peptides and proteins taking into account the amino acid sequence. <i>Journal of Analytical Chemistry</i> , 2016, 71, 1188-1197.	0.4	0
643	Chromatographic methods enabling the characterization of stationary phases and retention prediction in high-performance liquid chromatography and supercritical fluid chromatography. <i>Journal of Separation Science</i> , 2016, 39, 115-131.	1.3	21
644	Combined solvent- and non-uniform temperature-programmed gradient liquid chromatography. I. A theoretical investigation. <i>Journal of Chromatography A</i> , 2016, 1473, 38-47.	1.8	4
645	Retention in Porous Layer Pillar Array Planar Separation Platforms. <i>Analytical Chemistry</i> , 2016, 88, 8741-8748.	3.2	14

#	ARTICLE	IF	CITATIONS
646	Comparison of isocratic retention models for hydrophilic interaction liquid chromatographic separation of native and fluorescently labeled oligosaccharides. <i>Journal of Chromatography A</i> , 2016, 1438, 179-188.	1.8	36
647	General theory of peak compression in liquid chromatography. <i>Journal of Chromatography A</i> , 2016, 1433, 114-122.	1.8	20
648	Chromatographic study of the conformational behavior of graft copolymers with a broad distribution of grafting densities in dilute solutions in selective solvents for grafts. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 50-58.	0.5	11
649	Theoretical and experimental comparison of one dimensional versus on-line comprehensive two dimensional liquid chromatography for optimized sub-hour separations of complex peptide samples. <i>Journal of Chromatography A</i> , 2017, 1498, 183-195.	1.8	36
650	Mixed-mode reversed phase/positively charged repulsion chromatography for intact protein separation. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 138, 63-69.	1.4	8
651	Recent advances in stationary phases and understanding of retention in hydrophilic interaction chromatography. A review. <i>Analytica Chimica Acta</i> , 2017, 967, 12-32.	2.6	226
652	Advances in Hydrophilic Interaction Liquid Chromatography. , 2017, , 39-87.		3
654	Applicability of retention modelling in hydrophilic-interaction liquid chromatography for algorithmic optimization programs with gradient-scanning techniques. <i>Journal of Chromatography A</i> , 2017, 1530, 104-111.	1.8	36
655	Imidazolium Ionic Liquids as Mobile Phase Additives in Reversed Phase Liquid Chromatography for the Analysis of Anions. <i>Chromatographia</i> , 2017, 80, 1615-1622.	0.7	10
656	Modeling of preparative liquid chromatography. , 2017, , 573-592.		2
657	Determination of reversed-phase high performance liquid chromatography based octanol-water partition coefficients for neutral and ionizable compounds: Methodology evaluation. <i>Journal of Chromatography A</i> , 2017, 1528, 25-34.	1.8	36
658	Chromatographic and Computational Assessment of Lipophilicity of New Anticancer Acetylenequinoline Derivatives. <i>Journal of Chromatographic Science</i> , 2017, 55, 934-939.	0.7	4
659	Applicability of the critical-chromatography concept to proteomics problems: Separation of peptides modeled by a heterogeneous rod. <i>Polymer Science - Series A</i> , 2017, 59, 451-464.	0.4	1
660	Measuring and tracking vitamin B12: A review of current methods with a focus on optical spectroscopy. <i>Applied Spectroscopy Reviews</i> , 2017, 52, 439-455.	3.4	21
661	Separation of structurally related primary aliphatic amines using hydrophilic interaction chromatography with fluorescence detection after postcolumn derivatization with 2-mercaptoethanol. <i>Journal of Separation Science</i> , 2017, 40, 4689-4699.	1.3	6
662	pH-Gradient Liquid Chromatography: Fundamentals and Examples. <i>Journal of AOAC INTERNATIONAL</i> , 2017, 100, 1590-1598.	0.7	3
663	Synthesis and In Vitro Antimycobacterial Activity of Novel N-Arylpiperazines Containing an Ethane-1,2-diyl Connecting Chain. <i>Molecules</i> , 2017, 22, 2100.	1.7	9
664	A QuEChERS-Based Sample Preparation Method for the Analysis of 5-Nitroimidazoles in Bovine Milk by HPLC-DAD. <i>Journal of AOAC INTERNATIONAL</i> , 2017, 100, 1671-1680.	0.7	12

#	ARTICLE	IF	CITATIONS
665	Chromatographic retention behaviour, modelling and optimization of a UHPLC-UV separation of the regioisomers of the Novel Psychoactive Substance (NPS) methoxphenidine (MXP). <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 153, 238-247.	1.4	11
666	Mobile phase effects in reversed-phase and hydrophilic interaction liquid chromatography revisited. <i>Journal of Chromatography A</i> , 2018, 1543, 48-57.	1.8	19
667	Comparison between centrifugal partition chromatography and preparative liquid chromatography as first dimensions in off-line two-dimensional separation: Application to the isolation of multi-targeted compounds from Edelweiss plant. <i>Electrophoresis</i> , 2018, 39, 2011-2019.	1.3	6
668	Chromatographic performance of microfluidic liquid chromatography devices: Experimental evaluation of straight versus serpentine packed channels. <i>Journal of Chromatography A</i> , 2018, 1533, 127-135.	1.8	5
669	Enhancing detectability of anabolic-steroid residues in bovine urine by actively modulated online comprehensive two-dimensional liquid chromatography " high-resolution mass spectrometry. <i>Analytica Chimica Acta</i> , 2018, 1013, 87-97.	2.6	31
670	Mobile phase effects on the retention on polar columns with special attention to the dual hydrophilic interaction "reversed-phase liquid chromatography mechanism, a review. <i>Journal of Separation Science</i> , 2018, 41, 145-162.	1.3	45
671	Optimizing separations in online comprehensive two-dimensional liquid chromatography. <i>Journal of Separation Science</i> , 2018, 41, 68-98.	1.3	176
672	Comprehensive two dimensional liquid chromatography as analytical strategy for pharmaceutical analysis. <i>Journal of Chromatography A</i> , 2018, 1536, 195-204.	1.8	28
673	Studies on the retention mechanism of solutes in hydrophilic interaction chromatography using stoichiometric displacement theory I. The linear relationship of $\lg k'$ vs. $\lg[H_2O]$. <i>Talanta</i> , 2018, 176, 499-508.	2.9	10
674	1H-Tetrazole-5-amine Immobilized on Substituted Polymer Gel/Silica as a New Stationary Phase for Hydrophilic Interaction Chromatography. <i>Chromatographia</i> , 2018, 81, 349-357.	0.7	6
675	Development and Validation of Liquid Chromatography-Based Methods to Assess the Lipophilicity of Cytotoxic Platinum(IV) Complexes. <i>Inorganics</i> , 2018, 6, 130.	1.2	30
676	Investigation of the retention mechanisms of amlodipine besylate, bisoprolol fumarate, and their impurities on three different HILIC columns. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018, 41, 523-531.	0.5	9
677	Imidazolium ionic liquids as mobile phase additives in reversed phase liquid chromatography for the determination of iodide and iodate. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 7347-7355.	1.9	19
678	Separation of Triacylglycerols from Edible Oil Using a Liquid Chromatography "Mass Spectrometry System with a Porous Graphitic Carbon Column and a Toluene "Isopropanol Gradient Mobile Phase. <i>JAACS, Journal of the American Oil Chemists' Society</i> , 2018, 95, 1253-1266.	0.8	11
679	Suitable interface for coupling liquid chromatography to inductively coupled plasma-mass spectrometry for the analysis of organic matrices. 1 Theoretical and experimental considerations on solute dispersion. <i>Journal of Chromatography A</i> , 2018, 1565, 68-80.	1.8	10
680	Visualizing Events Under Solvent-Gradient and Insights Thereof. <i>Chromatographia</i> , 2018, 81, 575-584.	0.7	1
681	A novel strategy for retention prediction of nucleic acids with their sequence information in ion-pair reversed phase liquid chromatography. <i>Talanta</i> , 2018, 185, 592-601.	2.9	9
682	Measuring log Kow coefficients of neutral species of perfluoroalkyl carboxylic acids using reversed-phase high-performance liquid chromatography. <i>Environmental Pollution</i> , 2018, 242, 1283-1290.	3.7	16

#	ARTICLE	IF	CITATIONS
683	Controlled Fabrication of Silica@Covalent Triazine Polymer Core@Shell Spheres as a Reversed-Phase/Hydrophilic Interaction Mixed-Mode Chromatographic Stationary Phase. <i>ACS Applied Materials & Interfaces</i> , 2019, 11, 46149-46156.	4.0	40
684	Synthesis and structure-activity relationship of nitrile-based cruzain inhibitors incorporating a trifluoroethylamine-based P2 amide replacement. <i>Bioorganic and Medicinal Chemistry</i> , 2019, 27, 115083.	1.4	18
685	Evaluation of three temperature- and mobile phase-dependent retention models for reversed-phase liquid chromatographic retention and apparent retention enthalpy. <i>Journal of Chromatography A</i> , 2019, 1589, 73-82.	1.8	6
686	Solvent-induced modulation of the chemical sensing performance of gold nanoparticle film chemiresistors. <i>Sensors and Actuators B: Chemical</i> , 2019, 284, 316-322.	4.0	7
687	Gradient method transfer after changing the average pore diameter of the chromatographic stationary phase I @ One-dimensional sample mixture. <i>Journal of Chromatography A</i> , 2019, 1597, 119-131.	1.8	6
688	The comparison of retention behaviour of imidazoline and serotonin receptor ligands in non-aqueous hydrophilic interaction chromatography and supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2019, 1603, 371-379.	1.8	4
689	Computer-aided gradient optimization of hydrophilic interaction liquid chromatographic separations of intact proteins and protein glycoforms. <i>Journal of Chromatography A</i> , 2019, 1598, 67-76.	1.8	16
690	Retention Modeling of Gradient Elutions: Application of Iso-to-Grad Approach for LC Systems with Dual-Species Eluent. <i>Chromatographia</i> , 2019, 82, 749-755.	0.7	2
691	Column Characterization and Selection Systems in Reversed-Phase High-Performance Liquid Chromatography. <i>Chemical Reviews</i> , 2019, 119, 3674-3729.	23.0	191
692	Modeling Retention Behavior on Analysis of Hallucinogenic Mushrooms Using Hydrophilic Interaction Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 2019, 57, 230-237.	0.7	11
693	Accurate modelling of the retention behaviour of peptides in gradient-elution hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1614, 460650.	1.8	19
694	Utility of linear and nonlinear models for retention prediction in liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1613, 460690.	1.8	14
695	Comprehensive two-dimensional liquid chromatography with inductively coupled plasma mass spectrometry detection for the characterization of sulfur, vanadium and nickel compounds in petroleum products. <i>Journal of Chromatography A</i> , 2020, 1611, 460605.	1.8	10
696	Modeling the nonlinear behavior of a bioactive peptide in reversed-phase gradient elution chromatography. <i>Journal of Chromatography A</i> , 2020, 1616, 460789.	1.8	14
697	Separation optimization in HPLC analysis implemented in R programming language. <i>Journal of Chromatography A</i> , 2020, 1617, 460823.	1.8	7
698	Pushing the limits of resolving power and analysis time in on-line comprehensive hydrophilic interaction x reversed phase liquid chromatography for the analysis of complex peptide samples. <i>Journal of Chromatography A</i> , 2020, 1615, 460753.	1.8	19
699	Extension of the linear solvent strength retention model including a parameter that describes the elution strength changes in liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1615, 460757.	1.8	15
700	Assessing effects of ultra-high-pressure liquid chromatography instrument configuration on dispersion, system pressure, and retention. <i>Journal of Chromatography A</i> , 2020, 1634, 461660.	1.8	4

#	ARTICLE	IF	CITATIONS
701	Comparison of various modes and phase systems for analytical HPLC. Handbook of Analytical Separations, 2020, 8, 1-91.	0.8	1
702	Fast-generic HPLC methods. Handbook of Analytical Separations, 2020, 8, 93-112.	0.8	0
703	History of liquid chromatographyâ€”mass spectrometry couplings. , 2020, , 279-295.		5
704	Influence of the pre-elution of solute in initial mobile phase on retention time and peak compression under linear gradient elution. Journal of Chromatography A, 2020, 1618, 460858.	1.8	4
705	A simple, versatile, and automated pulse-diffusion-focusing strategy for sensitive milliliter-level-injection HILIC-MS/MS analysis of hydrophilic toxins. Journal of Hazardous Materials, 2020, 392, 122318.	6.5	5
706	Improving selectivity and performing online on-column fractioning in liquid chromatography for the separation of therapeutic biopharmaceutical products. Journal of Chromatography A, 2020, 1618, 460901.	1.8	13
707	Reducing the influence of geometry-induced gradient deformation in liquid chromatographic retention modelling. Journal of Chromatography A, 2021, 1635, 461714.	1.8	14
708	Computer aided optimization of multilinear gradient elution in liquid chromatography. Journal of Chromatography A, 2021, 1635, 461754.	1.8	5
709	Recent applications of retention modelling in liquid chromatography. Journal of Separation Science, 2021, 44, 88-114.	1.3	57
710	Novel computer-assisted separation prediction strategy for online-enrichment-HPLC-FLD in simultaneous monitoring of bisphenols in childrenâ€™s water bottles. Food Chemistry, 2021, 339, 127766.	4.2	10
711	Reversed-phase and hydrophobic interaction chromatography of carbohydrates and glycoconjugates. , 2021, , 35-124.		0
712	Deep Q-learning for the selection of optimal isocratic scouting runs in liquid chromatography. Journal of Chromatography A, 2021, 1638, 461900.	1.8	10
713	New Succinimides with Potent Anticancer Activity: Synthesis, Activation of Stress Signaling Pathways and Characterization of Apoptosis in Leukemia and Cervical Cancer Cells. International Journal of Molecular Sciences, 2021, 22, 4318.	1.8	9
714	Perspective on the Future Approaches to Predict Retention in Liquid Chromatography. Analytical Chemistry, 2021, 93, 5653-5664.	3.2	43
715	The development of retention time prediction model using multilinear gradient profiles of seven pharmaceuticals. Journal of Pharmaceutical and Biomedical Analysis, 2021, 198, 114024.	1.4	2
716	Recent advances in modelling and control of liquid chromatography. Current Opinion in Chemical Engineering, 2021, 32, 100685.	3.8	7
717	Numerical approximation of nonlinear and non-equilibrium model of gradient elution chromatography. Journal of Liquid Chromatography and Related Technologies, 2021, 44, 382-394.	0.5	3
718	CRB-SWATH: A Method for Enhancing Untargeted Precursor Ion Extraction and Automatically Constructing Their Tandem Mass Spectra from SWATH Datasets by Chromatographic Retention Behaviors. Analytical Chemistry, 2021, 93, 12273-12280.	3.2	6

#	ARTICLE	IF	CITATIONS
719	Application of retention modeling in chiral method development. I. Selection of isocratic composition for preparative separation with SFC. <i>Journal of Chromatography A</i> , 2021, 1651, 462308.	1.8	3
720	Measuring and using scanning-gradient data for use in method optimization for liquid chromatography. <i>Journal of Chromatography A</i> , 2021, 1636, 461780.	1.8	22
724	Das Elutionsmittel. , 1997, , 83-98.		1
725	Novel Approaches to the Liquid Chromatographic Analysis of Primary Amines, Amino Acids and Peptides. <i>Progress in Pharmaceutical and Biomedical Analysis</i> , 1994, , 41-89.	0.1	2
726	HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY SEPARATIONS OF LARGE MOLECULES: A GENERAL MODEL. , 1986, , 195-312.		49
728	Use of Planar Chromatography in Pesticide Residue Analysis. , 2009, , 187-264.		5
729	HILIC Retention Behavior and Method Development for Highly Polar Basic Compounds Used in Pharmaceutical Synthesis. <i>Chromatographic Science</i> , 2011, , 345-372.	0.1	2
730	A comparative study of the molecular lipophilicity indices of vitamins A and E, and of some precursors of vitamin A, estimated by HPLC and by different computation methods. <i>Acta Chromatographica</i> , 2009, 21, 237-250.	0.7	12
731	RP-HPTLC determination of the lipophilicity of some new derivatives of thiosemicarbazide and 1,2,4-triazole of sulphonylacetic acid. <i>Acta Chromatographica</i> , 2010, 22, 37-55.	0.7	7
732	Determination of the lipophilicity of bioactive 2-phenylbenzothiazoles by RPTLC. <i>Journal of Planar Chromatography - Modern TLC</i> , 2002, 15, 380-383.	0.6	5
734	Ultrahigh Sensitivity Analysis of Amino Acids and Peptides by Capillary Liquid Chromatography with Electrochemical Detection. , 2002, , 52-82.		0
735	Gradient HPLC. , 2005, , 725-732.		0
736	Eluotropic Series of Solvents for TLC. , 2005, , 547-553.		0
737	Tlc. , 2005, , 1700-1704.		0
738	Gradient Elution. , 2005, , 718-719.		0
739	Retention Behavior of Fluorobenzenes on Fluoro-Derivatized Poly(<i>p</i> -phenylene) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 144 <i>Chromatography</i> , 2011, 32, 127-133.	0.8	10
740	Multi-Solvent Optimization of HPLC Separations. , 1984, , 71-80.		0
742	Das Elutionsmittel. , 1988, , 101-114.		0

#	ARTICLE	IF	CITATIONS
746	Programmierte Elution. , 1997, , 154-174.		0
747	Separation of Macromolecules. , 1998, , 202-216.		0
748	Method Development of Chromatography: Retentionâ€“Eluent Composition Relationships and Application to Analysis of Pesticides. <i>Chromatographic Science</i> , 2015, , 77-98.	0.1	0
749	Optimization of Normal-Phase and Reversed-Phase Systems for Analysis of Pesticides: Choice of the Mode of Elutionâ€“Isocratic and Gradient Elution. <i>Chromatographic Science</i> , 2015, , 215-242.	0.1	0
750	Application of the Horner Method for a Well Produced at a Constant Bottom-hole Pressure. , 2015, , 103-107.		0
753	On-line 2D-RPLC x RPLC â€“ HRMS to assess wastewater treatment in a pharmaceutical plant. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2022, 208, 114465.	1.4	7
754	Mixed-acidic cation-exchange material for the separation of underivatized amino acids. <i>Journal of Chromatography A</i> , 2022, 1664, 462790.	1.8	5
755	Impact of UPLC-MS in Food and Drug/Metabolite Analysis. <i>Current Pharmaceutical Analysis</i> , 2021, 17, 10-30.	0.3	4
756	Solvent strength of aqueous phase for two typical biphasic solvent systems in high-speed countercurrent chromatography. <i>Journal of Chromatography A</i> , 2022, 1663, 462767.	1.8	5
757	Separation of ampicillin on polar-endcapped phase: Development of the HPLC method to achieve its correct dosage in cardiac surgery. <i>Acta Chromatographica</i> , 2023, 35, 1-9.	0.7	4
758	Protein Purification by Reversed Phase Chromatography and Hydrophobic Interaction Chromatography. , 2022, , 221-247.		1
759	Model of retention time and density of gradient peak capacity for improved LC-MS method optimization: Application to metabolomics. <i>Analytica Chimica Acta</i> , 2022, 1197, 339492.	2.6	1
760	The evaluation of drug-plasma protein binding interaction on immobilized human serum albumin stationary phase, aided by different computational approaches. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2022, 211, 114593.	1.4	8
761	Development of a retention prediction model in ion-pair reversed-phase HPLC for nucleoside triphosphates used as mRNA vaccine raw materials. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2022, 1193, 123168.	1.2	2
762	Analytical tools for antibodyâ€“drug conjugates: From inÂvitro to inÂvivo. <i>TrAC - Trends in Analytical Chemistry</i> , 2022, 152, 116621.	5.8	4
763	Improving retention-time prediction in supercritical-fluid chromatography by multivariate modelling. <i>Journal of Chromatography A</i> , 2022, 1668, 462909.	1.8	4
764	Detection of organic hydrates in reversed phase high performance liquid chromatography using recurrent approximation of their retention times. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2021, 44, 588-598.	0.5	5
765	On the Application of the Critical Chromatography Concept for Investigation of the Nucleotide Sequence of Single-Stranded DNA. <i>Polymer Science - Series A</i> , 2022, 64, 63.	0.4	0

#	ARTICLE	IF	CITATIONS
766	The development of a generic analysis method for natural and synthetic dyes by ultra-high-pressure liquid chromatography with photo-diode-array detection and triethylamine as an ion-pairing agent. <i>Journal of Chromatography A</i> , 2022, 1673, 463038.	1.8	9
770	A Novel Numerical Treatment of Nonlinear and Nonequilibrium Model of Gradient Elution Chromatography considering Core-Shell Particles in the Column. <i>Mathematical Problems in Engineering</i> , 2022, 2022, 1-14.	0.6	1
771	A simple mathematical treatment for predicting linear solvent strength behaviour in gradient elution: application to biomolecules. <i>Journal of Separation Science</i> , 2022, , .	1.3	2
773	Solvent Strength of Organic Phase for Two Biphasic Solvent Systems in High Speed Countercurrent Chromatography. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
774	Retention prediction of monoamine neurotransmitters in gradient liquid chromatography. <i>Journal of Separation Science</i> , 2022, 45, 3319-3327.	1.3	2
775	Liquid chromatography at the University of Pardubice: A tribute to Professor Pavel Jandera. <i>Journal of Separation Science</i> , 2022, 45, 3214-3231.	1.3	0
776	Solvent strength of organic phase for two biphasic solvent systems in high speed countercurrent chromatography. <i>Journal of Chromatography A</i> , 2022, 1680, 463422.	1.8	4
777	Comparing Two Versions of a Separation Map in Reversed Phase Liquid Chromatography. <i>Russian Journal of Physical Chemistry A</i> , 2022, 96, 1768-1772.	0.1	0
778	Numerical Approximation of the Nonequilibrium Model of Gradient Elution Chromatography Considering Linear and Nonlinear Solvent Strength Models. <i>ACS Omega</i> , 2022, 7, 31905-31915.	1.6	1
779	Theoretical and practical comparison of RPLC and RPLC—RPLC: how to consider dilution effects and sensitivity in addition to separation power?. <i>Analytical and Bioanalytical Chemistry</i> , 2023, 415, 2357-2369.	1.9	6
780	Reversed-Phase Liquid Chromatography of Peptides for Bottom-Up Proteomics: A Tutorial. <i>Journal of Proteome Research</i> , 2022, 21, 2846-2892.	1.8	17
781	An online internal standard technique for high-performance liquid chromatography-inductively coupled plasma mass spectrometry (HPLC-ICP-MS). <i>Analytical Methods</i> , 2023, 15, 240-246.	1.3	1
782	User-Driven Strategy for In Silico Screening of Reversed-Phase Liquid Chromatography Conditions for Known Pharmaceutical-Related Small Molecules. <i>Molecules</i> , 2022, 27, 8306.	1.7	2
783	Graph Neural Networks for Improved Retention Time Predictions. <i>Lc-gc Europe</i> , 2022, , 410-413.	0.2	0
784	Magnetism-Assisted Density Gradient Separation of Microplastics. <i>Analytical Chemistry</i> , 2022, 94, 17947-17955.	3.2	4
785	Novel computer-assisted approach to quick prediction and optimization of gradient separation for online enrichment-reversed phase liquid chromatography tandem system. <i>Chinese Chemical Letters</i> , 2023, 34, 108139.	4.8	1
786	Use of ultrasound in gradient elution of Polycyclic Aromatic Hydrocarbon mixtures by HPLC. <i>Bulletin of the Korean Chemical Society</i> , 2023, 44, 613-618.	1.0	0
787	Practical Approaches to Overcome the Challenges of Comprehensive Two-Dimensional Liquid Chromatography. <i>Lc-gc Europe</i> , 2018, , 242-249.	0.2	12

#	ARTICLE	IF	CITATIONS
792	Modeling of preparative liquid chromatography. , 2023, , 603-624.		0
793	Åšvod do Å¡portovej antropolÅ³gie. , 0, , .		0