The freshwater Crustacea of the island of Rhum (Inner Lecological survey

Biological Journal of the Linnean Society 11, 333-367 DOI: 10.1111/j.1095-8312.1979.tb00043.x

Citation Report

#	Article	IF	CITATIONS
1	Acidity and species diversity in freshwater crustacean faunas. Freshwater Biology, 1980, 10, 41-45.	2.4	92
2	The Distribution of Some Freshwater Copepods and its Bearing on the History of the Fauna and Flora of the British Isles. Journal of Biogeography, 1981, 8, 281.	3.0	13
3	The Crustacea of some chalk streams in southern England. Hydrobiologia, 1982, 97, 193-201.	2.0	12
4	The Entomology of the Isle of Rhum National Nature Reserve. Biological Journal of the Linnean Society, 1982, 18, 291-401.	1.6	14
5	Inland waters in the Inner Hebrides. Proceedings of the Royal Society of Edinburgh Section B Biological Sciences, 1983, 83, 229-244.	0.2	0
6	Freshwater science. Proceedings of the Royal Society of Edinburgh Section B Biological Sciences, 1983, 84, 171-210.	0.2	0
7	Aquatic angiosperm communities from lochs on Rhum. Transactions of the Botanical Society of Edinburgh, 1984, 44, 229-236.	0.1	0
8	An ecological validation of a taxonomic distinction: the ecology of Acanthocyclops vernalis and A. robustus (Crustacea: Copepoda). Zoological Journal of the Linnean Society, 1985, 84, 165-180.	2.3	19
9	The ecology and distribution of the genusDaphnia(Crustacea: Cladocera) in restricted areas: the pattern in Yorkshire. Journal of Natural History, 1985, 19, 97-128.	0.5	33
10	New Biostratigraphic Evidence of the Post-Glacial Colonization of Ireland and for Mesolithic Forest Disturbance. Journal of Biogeography, 1986, 13, 487.	3.0	74
11	Cladoceran remains as indicators of lake acidification. Hydrobiologia, 1986, 143, 129-134.	2.0	46
12	Guidelines and limitations to cladoceran paleoecological interpretations. Palaeogeography, Palaeoclimatology, Palaeoecology, 1988, 62, 405-412.	2.3	50
13	The ecology of some microcrustacea from standing waters in Tayside, Scotland. Journal of Natural History, 1989, 23, 375-406.	0.5	5
14	A Study of the Distribution and Ecology of Littoral Freshwater Chydorid (Crustacea, Cladocera) Communities in Ireland Using Multivariate Analyses. Journal of Biogeography, 1991, 18, 267.	3.0	35
15	The ecology and distribution of the littoral freshwater Chydoridae (Branchiopoda, Anomopoda) of Ireland, with taxonomic comments on some species. Hydrobiologia, 1992, 241, 1-70.	2.0	80
16	Variation in acid tolerance of certain freshwater crustaceans in different natural waters. Hydrobiologia, 1993, 250, 119-125.	2.0	20
17	Relationships between littoral microcrustacea and aquatic macrophyte communities on the Isle of Skye (Scotland), with implications for the conservation of standing waters. Aquatic Conservation: Marine and Freshwater Ecosystems, 1994, 4, 307-331.	2.0	9
18	The response of three chydorid species to temperature, pH and food. , 2001, 459, 165-172.		44

ATION RED

#	Article	IF	CITATIONS
19	Cladocera and Other Branchiopod Crustaceans. Developments in Paleoenvironmental Research, 2001, , 5-41.	8.0	200
20	Ecological distribution of pelagic copepods and species relationship to acidification, liming and natural recovery in a boreal area. Journal of Limnology, 2003, 62, 97.	1.1	15
21	The effect of habitat complexity on the contribution of some littoral–benthic Cladocera to the pelagic food web. Marine and Freshwater Research, 2013, 64, 1049.	1.3	9
22	Diversity, abundance, and life histories of littoral chydorids (Cladocera: Chydoridae) in a subarctic European lake. Journal of Crustacean Biology, 2020, 40, 534-543.	0.8	5
23	Inland waters in the Inner Hebrides. Proceedings of the Royal Society of Edinburgh Section B: Biology, 1983, 83, 229-244.	0.0	1
24	The use of Members of the Family Chydoridae (Anomopoda, Branchiopoda) as an Indicator of Lake Ecological Quality in Ireland. Biology and Environment, 2002, 102, 81-91.	0.3	10