

Plasma Cholesterol, Triglyceride and Uric Acid in Urban New Guinea

Australian and New Zealand Journal of Medicine

10, 491-495

DOI: [10.1111/j.1445-5994.1980.tb04964.x](https://doi.org/10.1111/j.1445-5994.1980.tb04964.x)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Mortality and morbidity of diabetes in Papua New Guinea. <i>Diabetologia</i> , 1982, 23, 136-137.	6.3	14
2	Nutritional status of some Papua New Guinea highlanders as assessed by physical measurements and blood analysis. <i>Ecology of Food and Nutrition</i> , 1988, 20, 185-196.	1.6	3
3	Levels of serum cholesterol, triglyceride, HDL-cholesterol, apoproteins A-I and B, and plasma glucose, and prevalence of diastolic hypertension and cigarette smoking in Papua New Guinea highlanders. <i>Pathology</i> , 1989, 21, 46-50.	0.6	11
4	Diet in an urban Papua New Guinea population with high levels of cardiovascular risk factors. <i>Ecology of Food and Nutrition</i> , 1996, 35, 311-324.	1.6	8
5	Serum Lipids and Modernization in Coastal and Highland Papua New Guinea. <i>American Journal of Epidemiology</i> , 1996, 144, 1129-1142.	3.4	20
6	Adult lipids associated with early life growth in traditional Melanesian societies undergoing rapid modernization: A longitudinal study of the mid-20th century. <i>American Journal of Physical Anthropology</i> , 2014, 153, 551-558.	2.1	4
7	Hyperuricaemia in the Pacific: why the elevated serum urate levels?. <i>Rheumatology International</i> , 2014, 34, 743-757.	3.0	37
8	Urban-rural differences in the association between blood lipids and characteristics of the built environment: a systematic review and meta-analysis. <i>BMJ Global Health</i> , 2019, 4, e001017.	4.7	27
9	Epidemiology of Gout and Hyperuricemia. , 2019, , 59-72.		0
10	Prevalence of non-communicable diseases and their risk factors in Papua New Guinea: A systematic review. <i>SAGE Open Medicine</i> , 2020, 8, 205031212097384.	1.8	7