

CITATION REPORT

List of articles citing

Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei

DOI: 10.1093/nar/11.5.1475

Nucleic Acids Research, 1983, 11, 1475-89.

Source: <https://exaly.com/paper-pdf/16160988/citation-report.pdf>

Version: 2024-04-10

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
1930	NQO1-Selective Activated Prodrug of Triptolide: Synthesis and Antihepatocellular Carcinoma Activity Evaluation.		
1929	Abstract.		2
1928	Identification of trans-acting factors by electrophoretic mobility shift assay. 2004 , 249, 7-20		2
1927	REVIEWS. 1929 , os-XXXI, 93-96		1
1926	Is actin a transcription initiation factor for RNA polymerase B?. 1984 , 3, 2363-2371		164
1925	Stimulation of in vitro transcription from the SV40 early promoter by the enhancer involves a specific trans-acting factor.. 1984 , 3, 3129-3133		106
1924	Human genes and pseudogenes for the 7SL RNA component of signal recognition particle.. 1984 , 3, 3303-3310		77
1923	Accurate transcription of homologous 5S rRNA and tRNA genes and splicing of tRNA in vitro by soluble extracts of Neurospora. <i>Nucleic Acids Research</i> , 1984 , 12, 5737-55	20.1	14
1922	Stimulation of in vitro transcription by the upstream element of the adenovirus-2 major late promoter involves a specific factor. <i>Nucleic Acids Research</i> , 1984 , 12, 8779-99	20.1	125
1921	Unwinding of the DNA helix in simian virus 40 chromosome templates by RNA polymerase. 1984 , 143, 39-45		8
1920	The enhancement of specific gene transcription in isolated nuclei by added HeLa whole cell extract. 1984 , 16, 963-9		2
1919	Transcription of adenovirus cores in vitro. 1984 , 139, 152-63		2
1918	Normal and mutant human beta-globin pre-mRNAs are faithfully and efficiently spliced in vitro. 1984 , 36, 993-1005		684
1917	Interrelations between the maturation of a 100 kDa nucleolar protein and pre rRNA synthesis in CHO cells. <i>Nucleic Acids Research</i> , 1984 , 12, 3025-35	20.1	123
1916	Purification and characterization of ternary complexes containing accurately initiated RNA polymerase II and less than 20 nucleotides of RNA. 1984 , 178, 415-37		33
1915	In vivo splicing products of the rabbit beta-globin pre-mRNA. 1984 , 39, 589-602		228
1914	Efficient coupled transcription and mRNA splicing in vitro using plasmids derived from early region 3 of adenovirus 2 and a nondefective adenovirus-simian virus 40 hybrid. 1984 , 81, 2679-83		2

1913	Transcription of human histone genes in extracts from synchronized HeLa cells. 1984 , 81, 2713-7	90
1912	Homologous globin cell-free transcription system with comparison of heterologous factors. 1984 , 4, 17-22	24
1911	Isolation and characterization of two fractions from HeLa cells required for mRNA splicing in vitro. 1985 , 82, 4351-5	52
1910	Factors involved in specific transcription by human RNA polymerase II: analysis by a rapid and quantitative in vitro assay. 1985 , 82, 4394-8	566
1909	Sequences upstream of c-mos(rat) that block RNA accumulation in mouse cells do not inhibit in vitro transcription. 1985 , 5, 406-9	13
1908	Accurate transcription of cloned Neurospora RNA polymerase II-dependent genes in vitro by homologous soluble extracts. 1985 , 82, 5450-4	27
1907	Cap-dependent RNA splicing in a HeLa nuclear extract. 1985 , 82, 7590-4	142
1906	Accurate initiation of rat ribosomal RNA gene transcription using a fractionated nuclear extract from normal liver and a hepatoma. 1985 , 82, 1059-63	21
1905	Characterization of the branch site in lariat RNAs produced by splicing of mRNA precursors. 1985 , 313, 552-7	213
1904	Nuclear extracts from globin-synthesizing cells enhance globin transcription in vitro. 1985 , 317, 824-8	27
1903	Upstream sequences modulate the internal promoter of the human 7SL RNA gene. 1985 , 318, 371-4	183
1902	The two intervening sequences of human beta- and gamma-globin pre-mRNAs are excised in a preferred temporal order in vitro.. 1985 , 4, 1991-1996	17
1901	Specific interaction between a transcription factor and the upstream element of the adenovirus-2 major late promoter.. 1985 , 4, 3563-3570	136
1900	Purification of a protein required for the splicing of pre-mRNA and its separation from the lariat debranching enzyme.. 1985 , 4, 3571-3581	45
1899	Functional domains of the Xenopus laevis 5S gene promoter.. 1985 , 4, 3751-3756	40
1898	Specific switching on of silent egg protein genes in vitro by an S-100 fraction in isolated nuclei from male Xenopus.. 1985 , 4, 3253-3258	10
1897	An RNA processing activity that debranches RNA lariats. 1985 , 229, 135-40	274
1896	Eukaryotic RNA polymerases. 1985 , 18, 31-90	410

1895	Structure of a Neurospora RNA polymerase I promoter defined by transcription in vitro with homologous extracts. <i>Nucleic Acids Research</i> , 1985 , 13, 4311-32	20.1	31
1894	In vitro transcription with extracts of nuclei of Drosophila embryos. <i>Nucleic Acids Research</i> , 1985 , 13, 2709-30	20.1	62
1893	Characterization of an unique RNA initiated immediately upstream from human alpha 1 globin gene in vivo and in vitro: polymerase II-dependence, tissue specificity, and subcellular location. <i>Nucleic Acids Research</i> , 1985 , 13, 6059-74	20.1	11
1892	Spacer sequences downstream of the 28S RNA coding region are part of the mouse rDNA transcription unit. <i>Nucleic Acids Research</i> , 1985 , 13, 2293-304	20.1	30
1891	STEM Footprints and Bound Mass Distributions for DNA Control Proteins. 1985 , 3, 549-553		
1890	Poly(A) site cleavage in a HeLa nuclear extract is dependent on downstream sequences. 1985 , 43, 677-83		154
1889	Formation of a rate-limiting intermediate in 5S RNA gene transcription. 1985 , 40, 119-27		208
1888	Specific and stable intron-factor interactions are established early during in vitro pre-mRNA splicing. 1985 , 43, 131-42		196
1887	Multiple factors including the small nuclear ribonucleoproteins U1 and U2 are necessary for pre-mRNA splicing in vitro. 1985 , 42, 725-36		466
1886	Enhancement of RNA polymerase III transcription by the E1A gene product of adenovirus. 1985 , 41, 955-63		190
1885	U2 as well as U1 small nuclear ribonucleoproteins are involved in premessenger RNA splicing. 1985 , 42, 737-50		599
1884	Trans splicing of mRNA precursors. 1985 , 42, 157-64		190
1883	Transcription of mouse rDNA terminates downstream of the 3' end of 28S RNA and involves interaction of factors with repeated sequences in the 3' spacer. 1985 , 43, 801-10		194
1882	A multicomponent complex is involved in the splicing of messenger RNA precursors. 1985 , 42, 345-53		465
1881	Alternative splicing caused by RNA secondary structure. 1985 , 43, 667-76		268
1880	Cloning specific complete polyadenylylated 3'-terminal cDNA segments. 1985 , 33, 191-6		17
1879	In vitro stimulation of specific RNA polymerase II-mediated transcription by the pseudorabies virus immediate early protein. 1985 , 43, 821-9		45
1878	Accurate cleavage and polyadenylation of exogenous RNA substrate. 1985 , 41, 845-55		343

1877	Growth-dependent regulation of rRNA synthesis is mediated by a transcription initiation factor (TIF-IA). <i>Nucleic Acids Research</i> , 1985 , 13, 8165-80	20.1	109
1876	The U2 RNA analogue of <i>Trypanosoma brucei gambiense</i> : implications for a splicing mechanism in trypanosomes. <i>Nucleic Acids Research</i> , 1986 , 14, 8893-903	20.1	92
1875	Electrophoretic separation of complexes involved in the splicing of precursors to mRNAs. 1986 , 46, 845-55		501
1874	Identification of a protein-binding site that mediates transcriptional response of the c-fos gene to serum factors. 1986 , 46, 567-74		878
1873	A transcription terminator located upstream of the mouse rDNA initiation site affects rRNA synthesis. 1986 , 47, 901-11		161
1872	Identification of a cellular transcription factor involved in E1A trans-activation. 1986 , 45, 219-28		538
1871	Sequence requirements for splicing of higher eukaryotic nuclear pre-mRNA. 1986 , 47, 555-65		379
1870	A protein that specifically recognizes the 3' splice site of mammalian pre-mRNA introns is associated with a small nuclear ribonucleoprotein. 1986 , 47, 755-66		287
1869	Purification and biochemical characterization of the promoter-specific transcription factor, Sp1. 1986 , 234, 47-52		1435
1868	A soluble biotinyl-alpha-amanitin affinity probe for the isolation of RNA polymerase B. 1986 , 64, 923-9		2
1867	Identification of DNA binding proteins from human cells. 1986 , 3, 96-101		5
1866	Pre-mRNA splicing in vitro requires intact U4/U6 small nuclear ribonucleoprotein. 1986 , 46, 697-704		233
1865	A protein associated with small nuclear ribonucleoprotein particles recognizes the 3' splice site of premessenger RNA. 1986 , 47, 973-84		284
1864	Inducible binding of a factor to the c-fos enhancer. 1986 , 47, 777-84		313
1863	A repeated 18 bp sequence motif in the mouse rDNA spacer mediates binding of a nuclear factor and transcription termination. 1986 , 45, 837-46		138
1862	U1, U2, and U4/U6 small nuclear ribonucleoproteins are required for in vitro splicing but not polyadenylation. 1986 , 46, 691-6		198
1861	The human tRNA ^{Val} gene family: organization, nucleotide sequences and homologous transcription of three single-copy genes. 1986 , 44, 287-97		47
1860	Inducibility of kappa immunoglobulin enhancer-binding protein Nf-kappa B by a posttranslational mechanism. 1986 , 47, 921-8		1850

1859	A small nuclear ribonucleoprotein associates with the AAUAAA polyadenylation signal in vitro. 1986 , 45, 581-91	143
1858	Transcriptional selectivity of viral genes in mammalian cells. 1986 , 46, 795-805	737
1857	A Drosophila RNA polymerase II cell-free DNA-dependent transcription system. 1986 , 866, 83-8	
1856	The presence of nucleosomes on a DNA template prevents initiation by RNA polymerase II in vitro. 1986 , 45, 95-104	308
1855	Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. 1986 , 46, 417-28	705
1854	Multiple nuclear factors interact with the immunoglobulin enhancer sequences. 1986 , 46, 705-16	2389
1853	Accurate in vitro splicing of two pre-mRNA plant introns in a HeLa cell nuclear extract.. 1986 , 5, 2749-2758	58
1852	In vitro transcription by Xenopus oocytes RNA polymerase III requires a DNA topoisomerase II activity.. 1986 , 5, 151-155	16
1851	Proteins that bind to RNA polymerase II are required for accurate initiation of transcription at the adenovirus 2 major late promoter.. 1986 , 5, 2923-2930	37
1850	Generation of histone mRNA 3' ends by endonucleolytic cleavage of the pre-mRNA in a snRNP-dependent in vitro reaction.. 1986 , 5, 1319-1326	87
1849	DNA topoisomerase II cleaves at specific sites in the 5' flanking region of c-fos proto-oncogenes in vitro.. 1986 , 5, 2257-2265	30
1848	Regulation of adenovirus mRNA formation. 1986 , 31, 169-228	52
1847	In vitro cleavage of the simian virus 40 early polyadenylation site adjacent to a required downstream TG sequence. 1986 , 6, 4734-41	42
1846	Identification of proteins that bind tightly to pre-mRNA during in vitro splicing. 1986 , 83, 3718-22	17
1845	Inhibition of splicing but not cleavage at the 5' splice site by truncating human beta-globin pre-mRNA. 1986 , 83, 927-31	51
1844	Alternative branch points are selected during splicing of a yeast pre-mRNA in mammalian and yeast extracts. 1986 , 83, 2022-6	50
1843	Identification of a factor in HeLa cells specific for an upstream transcriptional control sequence of an E1A-inducible adenovirus promoter and its relative abundance in infected and uninfected cells. 1986 , 83, 5914-8	120
1842	Cell-type-specific transcription of an immunoglobulin kappa light chain gene in vitro. 1986 , 83, 8511-5	104

1841	The AAUAAA sequence is required both for cleavage and for polyadenylation of simian virus 40 pre-mRNA in vitro. 1986 , 6, 2317-23	163
1840	Transcriptional role for the nontranscribed spacer of rat ribosomal DNA. 1986 , 6, 2766-73	52
1839	Herpes simplex virus immediate early infected-cell polypeptide 4 binds to DNA and promotes transcription. 1986 , 83, 4016-20	66
1838	In vitro transcription and delimitation of promoter elements of the murine dihydrofolate reductase gene. 1986 , 6, 2392-401	74
1837	A purified transcription factor (TIF-IB) binds to essential sequences of the mouse rDNA promoter. 1986 , 83, 604-8	102
1836	Specific protein binding to the simian virus 40 enhancer in vitro. 1986 , 6, 2098-105	156
1835	Alpha 4, the major regulatory protein of herpes simplex virus type 1, is stably and specifically associated with promoter-regulatory domains of alpha genes and of selected other viral genes. 1986 , 83, 3218-22	137
1834	DNA-binding site of major regulatory protein alpha 4 specifically associated with promoter-regulatory domains of alpha genes of herpes simplex virus type 1. 1986 , 83, 4700-4	91
1833	In vitro splicing of influenza viral NS1 mRNA and NS1-beta-globin chimeras: possible mechanisms for the control of viral mRNA splicing. 1986 , 83, 5444-8	49
1832	Transcription antitermination during influenza viral template RNA synthesis requires the nucleocapsid protein and the absence of a 5' capped end. 1986 , 83, 6282-6	133
1831	Interaction of a common factor with conserved promoter and enhancer sequences in histone H2B, immunoglobulin, and U2 small nuclear RNA (snRNA) genes. 1986 , 83, 6382-6	211
1830	Interferon-stimulated transcription: isolation of an inducible gene and identification of its regulatory region. 1986 , 83, 8929-33	165
1829	Mutational analysis of the immunoglobulin heavy chain promoter region. 1986 , 83, 9626-30	98
1828	RNA splicing products formed with isolated fractions from HeLa cells are associated with fast-sedimenting complexes. 1986 , 83, 887-91	74
1827	Polymorphism in an androgen-regulated mouse gene is the result of the insertion of a B1 repetitive element into the transcription unit. 1986 , 6, 209-17	27
1826	Control of transcription initiation in vitro requires binding of a transcription factor to the distal promoter of the ovalbumin gene. 1986 , 6, 2784-91	101
1825	Nuclear factors bind to regulatory regions of the mouse kappa immunoglobulin gene. <i>Nucleic Acids Research</i> , 1986 , 14, 4837-48	20.1 43
1824	Specific transcription of a 15-kilodalton zene gene in HeLa cell extracts. 1986 , 7, 71-9	10

1823	The expression of a nopaline synthase - human growth hormone chimaeric gene in transformed tobacco and sunflower callus tissue. 1986 , 6, 347-57	212
1822	Effects of DNA methylation on specific transcription by RNA polymerase II in vitro. 1986 , 11, 13-7	6
1821	A nuclear factor that binds to a conserved sequence motif in transcriptional control elements of immunoglobulin genes. 1986 , 319, 154-8	1167
1820	Transcription factor Sp1 recognizes a DNA sequence in the mouse dihydrofolate reductase promoter. 1986 , 319, 246-8	272
1819	Distinct factors bind to apparently homologous sequences in the immunoglobulin heavy-chain enhancer. 1986 , 322, 846-8	177
1818	Interaction of cell-type-specific nuclear proteins with immunoglobulin VH promoter region sequences. 1986 , 323, 548-51	317
1817	A lymphoid-specific protein binding to the octamer motif of immunoglobulin genes. 1986 , 323, 640-3	728
1816	Effect of mutations at the lariat branch acceptor site on beta-globin pre-mRNA splicing in vitro. 1986 , 324, 589-91	115
1815	Multiple protein-binding sites in the 5'-flanking region regulate c-fos expression. 1986 , 6, 4305-16	535
1814	Heterogeneous nuclear ribonucleoproteins: role in RNA splicing. 1986 , 231, 1534-9	395
1813	Upstream regulatory sequences of immunoglobulin genes are recognized by nuclear proteins which also bind to other gene regions. <i>Nucleic Acids Research</i> , 1986 , 14, 8829-44	20.1 48
1812	In vitro processing of a plant pre-mRNA in a HeLa cell nuclear extract. <i>Nucleic Acids Research</i> , 1986 , 14, 7513-28	20.1 45
1811	Antibodies to hnRNP core proteins inhibit in vitro splicing of human beta-globin pre-mRNA. <i>Nucleic Acids Research</i> , 1986 , 14, 5241-54	20.1 124
1810	In vitro transcription of a human hsp 70 heat shock gene by extracts prepared from heat-shocked and non-heat-shocked human cells. <i>Nucleic Acids Research</i> , 1986 , 14, 8933-48	20.1 20
1809	Endonucleolytic activity that cleaves immunoglobulin recombination sequences. 1986 , 231, 1141-5	56
1808	A sequence upstream from the coding region is required for the transcription of the 7SK RNA genes. <i>Nucleic Acids Research</i> , 1986 , 14, 9243-60	20.1 63
1807	Alternative splicing of SV40 early pre-mRNA in vitro. <i>Nucleic Acids Research</i> , 1986 , 14, 9911-26	20.1 32
1806	The different intron 2 species excised in vivo from the E2A premRNA of adenovirus-2: an approach to analyse alternative splicing. <i>Nucleic Acids Research</i> , 1986 , 14, 5207-27	20.1 11

1805	Cell proliferation and expression of the transferrin receptor gene: promoter sequence homologies and protein interactions. 1986 , 103, 1781-8		65
1804	In vitro splicing of simian virus 40 early pre mRNA. <i>Nucleic Acids Research</i> , 1986 , 14, 1219-35	20.1	51
1803	A 100-kD HeLa cell octamer binding protein (OBP100) interacts differently with two separate octamer-related sequences within the SV40 enhancer. 1987 , 1, 1147-60		232
1802	Factors that interact with the rat albumin promoter are present both in hepatocytes and other cell types. 1987 , 1, 256-67		81
1801	Tissue-specific expression of the human alpha 1-antitrypsin gene is controlled by multiple cis-regulatory elements. <i>Nucleic Acids Research</i> , 1987 , 15, 8399-415	20.1	62
1800	Characterization of xenobiotic responsive elements upstream from the drug-metabolizing cytochrome P-450c gene: a similarity to glucocorticoid regulatory elements. <i>Nucleic Acids Research</i> , 1987 , 15, 4179-91	20.1	328
1799	The HLA-DQ beta gene upstream region contains an immunoglobulin-like octamer motif that binds cell-type specific nuclear factors. <i>Nucleic Acids Research</i> , 1987 , 15, 8057-67	20.1	21
1798	Purification of intercalator-released p67, a polypeptide that interacts specifically with the c-fos serum response element. <i>Nucleic Acids Research</i> , 1987 , 15, 10145-58	20.1	101
1797	Identification of the human U7 snRNP as one of several factors involved in the 3' end maturation of histone premessenger RNA's. 1987 , 238, 1682-7		206
1796	Specific factors binding to the late E2A promoter region of adenovirus type 2 DNA: no apparent effects of 5'-CCGG-3' methylation. 1987 , 6, 449-60		25
1795	A mutational analysis of spliceosome assembly: evidence for splice site collaboration during spliceosome formation. 1987 , 1, 532-43		160
1794	Characterization of human DNA sequences synthesized at the onset of S-phase. <i>Nucleic Acids Research</i> , 1987 , 15, 10211-32	20.1	41
1793	Nuclear proteins from lactating mammary glands bind to the promoter of a milk protein gene. <i>Nucleic Acids Research</i> , 1987 , 15, 2103-21	20.1	73
1792	Two forms of transcription factor TFIIIC in extracts from HeLa cells. <i>Nucleic Acids Research</i> , 1987 , 15, 5031-9	20.1	19
1791	In vitro processing of the human growth hormone primary transcript. <i>Nucleic Acids Research</i> , 1987 , 15, 7005-25	20.1	14
1790	Only two of the four sites of interaction with nuclear factors within the Xenopus U2 gene promoter are necessary for efficient transcription. <i>Nucleic Acids Research</i> , 1987 , 15, 6437-53	20.1	21
1789	A protein factor binding to an octamer motif in the gamma-globin promoter disappears upon induction of differentiation and hemoglobin synthesis in K562 cells. <i>Nucleic Acids Research</i> , 1987 , 15, 9349-64	20.1	51
1788	OVEC, a versatile system to study transcription in mammalian cells and cell-free extracts. <i>Nucleic Acids Research</i> , 1987 , 15, 6787-98	20.1	276

1787	Purification of an octamer sequence (ATGCAAAT)-binding protein from human B cells. <i>Nucleic Acids Research</i> , 1987 , 15, 10105-16	20.1	41
1786	Assembly of pre-mRNA splicing complex is cap dependent. <i>Nucleic Acids Research</i> , 1987 , 15, 1387-99	20.1	47
1785	Nuclear factors binding to the human immunoglobulin heavy-chain gene enhancer. <i>Nucleic Acids Research</i> , 1987 , 15, 2851-69	20.1	35
1784	Novobiocin interferes with the binding of transcription factors TFIIIA and TFIIIC to the promoters of class III genes. <i>Nucleic Acids Research</i> , 1987 , 15, 4365-74	20.1	11
1783	Multiple transcription factors interact with the adenovirus-2 EII-late promoter: evidence for a novel CCAAT recognition factor. <i>Nucleic Acids Research</i> , 1987 , 15, 7761-80	20.1	41
1782	Natural point mutations within rat rDNA transcription terminator elements reveal the functional importance of single bases for factor binding and termination. <i>Nucleic Acids Research</i> , 1987 , 15, 4131-43	20.1	20
1781	Requirements for accurate and efficient mRNA 3' end cleavage and polyadenylation of a simian virus 40 early pre-RNA in vitro. 1987 , 7, 495-503		65
1780	Properties of a U1 RNA enhancer-like sequence. <i>Nucleic Acids Research</i> , 1987 , 15, 2403-16	20.1	49
1779	In vitro binding of several cell-specific and ubiquitous nuclear proteins to the GT-I motif of the SV40 enhancer. 1987 , 1, 794-807		122
1778	Identification of factors that interact with the E1A-inducible adenovirus E3 promoter. 1987 , 1, 1132-46		175
1777	Identification of nuclear cap specific proteins in HeLa cells. <i>Nucleic Acids Research</i> , 1987 , 15, 6489-500	20.1	45
1776	Upstream regulatory regions required to stabilize binding to the TATA sequence in an adenovirus early promoter. <i>Nucleic Acids Research</i> , 1987 , 15, 8367-85	20.1	61
1775	Linker scanner mutagenesis of the <i>Xenopus laevis</i> ribosomal gene promoter. <i>Nucleic Acids Research</i> , 1987 , 15, 7429-41	20.1	44
1774	A purine-rich DNA sequence motif present in SV40 and lymphotropic papovavirus binds a lymphoid-specific factor and contributes to enhancer activity in lymphoid cells. 1987 , 1, 962-72		102
1773	Specific cellular proteins bind to critical promoter sequences of the adenovirus early EIIa promoter. <i>Nucleic Acids Research</i> , 1987 , 15, 509-27	20.1	42
1772	Regulation in vitro of metallothionein gene binding factors. 1987 , 235, 1383-7		101
1771	A mammalian mitochondrial RNA processing activity contains nucleus-encoded RNA. 1987 , 235, 1178-84		279
1770	A 5'----3' exoribonuclease of human placental nuclei: purification and substrate specificity. <i>Nucleic Acids Research</i> , 1987 , 15, 695-708	20.1	32

1769	Products of in vitro cleavage and polyadenylation of simian virus 40 late pre-mRNAs. 1987 , 7, 1518-29	71
1768	Mutation of the c-fos gene dyad symmetry element inhibits serum inducibility of transcription in vivo and the nuclear regulatory factor binding in vitro. 1987 , 7, 1217-25	257
1767	Evolutionary changes of sequences and factors that direct transcription termination of human and mouse ribosomal genes. 1987 , 7, 2521-9	37
1766	Selective transcription and DNase I protection of the rat prolactin gene by GH3 pituitary cell-free extracts. 1987 , 84, 5211-5	79
1765	Drosophila Krüppel gene product produced in a baculovirus expression system is a nuclear phosphoprotein that binds to DNA. 1987 , 84, 5700-4	63
1764	Binding of the virion protein mediating alpha gene induction in herpes simplex virus 1-infected cells to its cis site requires cellular proteins. 1987 , 84, 7061-5	200
1763	Accurate 5' splice-site selection in mouse kappa immunoglobulin light chain premessenger RNAs is not cell-type-specific. 1987 , 84, 7928-32	22
1762	Cell-cycle-specific interaction of nuclear DNA-binding proteins with a CCAAT sequence from the human thymidine kinase gene. 1987 , 84, 8350-4	76
1761	Regulatory elements in the first intron contribute to transcriptional control of the human alpha 1(I) collagen gene. 1987 , 84, 8869-73	174
1760	Complex protein binding within the mouse immunoglobulin heavy-chain enhancer. 1987 , 7, 4194-203	70
1759	Recognition of mutant and cryptic 5' splice sites by the U1 small nuclear ribonucleoprotein in vitro. 1987 , 7, 698-707	38
1758	Identification of a complex associated with processing and polyadenylation in vitro of herpes simplex virus type 1 thymidine kinase precursor RNA. 1987 , 7, 3277-86	76
1757	c-fos sequence necessary for basal expression and induction by epidermal growth factor, 12-O-tetradecanoyl phorbol-13-acetate and the calcium ionophore. 1987 , 7, 3490-502	316
1756	An embryonic DNA-binding protein specific for the promoter of the retrovirus long terminal repeat. 1987 , 7, 3548-53	21
1755	Multiple basal elements of a human hsp70 promoter function differently in human and rodent cell lines. 1987 , 7, 3646-55	128
1754	In vitro transcription of immunoglobulin genes in a B-cell extract: effects of enhancer and promoter sequences. 1987 , 7, 1989-94	28
1753	Binding of transcription factors and creation of a large nucleoprotein complex on the human cytomegalovirus enhancer. 1987 , 84, 3658-62	114
1752	A plasmacytoma-specific factor binds the c-myc promoter region. 1987 , 84, 7031-5	47

1751	Reconstitution of the U1 small nuclear ribonucleoprotein particle. 1987 , 7, 4030-7	82
1750	Upstream DNA sequences required for tissue-specific expression of the HLA-DR alpha gene. 1987 , 84, 4254-8	120
1749	Sequence-specific interactions of nuclear factors with conserved sequences of human class II major histocompatibility complex genes. 1987 , 84, 4939-43	100
1748	Conserved major histocompatibility complex class II boxes--X and Y--are transcriptional control elements and specifically bind nuclear proteins. 1987 , 84, 6249-53	192
1747	CArG boxes in the human cardiac alpha-actin gene are core binding sites for positive trans-acting regulatory factors. 1987 , 84, 6702-6	87
1746	Fractionation of HeLa cell nuclear extracts reveals minor small nuclear ribonucleoprotein particles. 1987 , 84, 8408-12	33
1745	Protein-binding site at the immunoglobulin mu membrane polyadenylation signal: possible role in transcription termination. 1987 , 84, 9160-4	33
1744	Detection of three protein binding sites in the serum-regulated promoter of the human gene encoding the 70-kDa heat shock protein. 1987 , 84, 2203-7	100
1743	Pre-mRNA splicing and the nuclear matrix. 1987 , 7, 111-20	202
1742	Purification of the c-fos enhancer-binding protein. 1987 , 7, 3482-9	162
1741	The abundance and in vitro DNA binding of three cellular proteins interacting with the adenovirus Ela early promoter are not modified by the Ela gene products. 1987 , 7, 3806-17	47
1740	Binding of multiple nuclear factors to the 5' upstream regulatory element of the murine major histocompatibility class I gene. 1987 , 7, 4542-8	96
1739	In vitro splicing of kappa immunoglobulin precursor mRNA. 1987 , 7, 1346-51	16
1738	Heat-inducible human factor that binds to a human hsp70 promoter. 1987 , 7, 1530-4	247
1737	Multiple interactions between the splicing substrate and small nuclear ribonucleoproteins in spliceosomes. 1987 , 7, 281-93	164
1736	Variations in template protection by the RNA polymerase II transcription complex during the initiation process. 1987 , 7, 3371-9	30
1735	Reconstitution of cell-type-specific transcription of the rat prolactin gene in vitro. 1987 , 7, 3402-8	56
1734	In vitro splicing pathways of pre-mRNAs containing multiple intervening sequences?. 1987 , 7, 3428-37	38

1733	Multiple proteins bind to VA RNA genes of adenovirus type 2. 1987 , 7, 1021-31	36
1732	Host cell proteins bind to the cis-acting site required for virion-mediated induction of herpes simplex virus 1 alpha genes. 1987 , 84, 71-5	144
1731	Preferential excision of the 5' proximal intron from mRNA precursors with two introns as mediated by the cap structure. 1987 , 84, 5187-91	95
1730	Two promoter-specific host factors interact with adjacent sequences in an E1A-inducible adenovirus promoter. 1987 , 84, 6112-6	78
1729	A cellular protein, activating transcription factor, activates transcription of multiple E1A-inducible adenovirus early promoters. 1987 , 84, 8355-9	261
1728	Role of an adenovirus E2 promoter binding factor in E1A-mediated coordinate gene control. 1987 , 84, 2180-4	249
1727	Protein-nucleotide contacts in the immunoglobulin heavy-chain promoter region. 1987 , 84, 3851-5	43
1726	B lymphocyte-specific protein binding near an immunoglobulin kappa-chain gene J segment. 1987 , 84, 1516-20	34
1725	Two transcriptional activators, CCAAT-box-binding transcription factor and heat shock transcription factor, interact with a human hsp70 gene promoter. 1987 , 7, 1129-38	183
1724	Both conserved signals on mammalian histone pre-mRNAs associate with small nuclear ribonucleoproteins during 3' end formation in vitro. 1987 , 7, 1663-72	78
1723	Nucleoprotein complexes that regulate gene expression in adipocyte differentiation: direct participation of c-fos. 1987 , 49, 835-44	510
1722	Splicing of SV40 early pre-mRNA to large T and small t mRNAs utilizes different patterns of lariet branch sites. 1987 , 50, 227-36	96
1721	Posttranscriptional Processing (Splicing and 3' End Formation) of Nuclear Messenger RNA Precursors in Vitro. 1987 , 99-111	
1720	Unrelated leader sequences can efficiently promote human tRNA gene transcription. 1987 , 51, 237-46	39
1719	Splicing of plant pre-mRNAs in animal systems and vice versa. 1987 , 56, 253-65	73
1718	The in vitro transcription of the 7SK RNA gene by RNA polymerase III is dependent only on the presence of an upstream promoter. 1987 , 51, 81-7	194
1717	Steroid-dependent interaction of transcription factors with the inducible promoter of mouse mammary tumor virus in vivo. 1987 , 48, 261-70	427
1716	Interaction of protein with DNA in vitro. 1987 , 152, 721-35	160

1715	Tissue-specific binding of a nuclear factor to the insulin gene promoter. 1987 , 222, 332-6	3
1714	A mouse renin promoter containing the conserved decanucleotide element binds the same B-cell factors as an authentic immunoglobulin heavy chain promoter. 1987 , 222, 337-40	8
1713	In vitro identification of a B19 parvovirus promoter. 1987 , 157, 534-8	93
1712	A transcriptional promoter of the human parvovirus B19 active in vitro and in vivo. 1987 , 157, 539-42	31
1711	Priming of influenza mRNA transcription is inhibited in CHO cells treated with the methylation inhibitor, neplanocin A. 1987 , 7, 317-27	34
1710	Developmental control of a promoter-specific factor that is also regulated by the E1A gene product. 1987 , 48, 501-6	118
1709	Identification and purification of a human lymphoid-specific octamer-binding protein (OTF-2) that activates transcription of an immunoglobulin promoter in vitro. 1987 , 51, 783-93	400
1708	A common octamer motif binding protein is involved in the transcription of U6 snRNA by RNA polymerase III and U2 snRNA by RNA polymerase II. 1987 , 51, 71-9	242
1707	Factors involved in control of tissue-specific expression of albumin gene. 1987 , 50, 627-38	363
1706	Genomic footprinting reveals cell type-specific DNA binding of ubiquitous factors. 1987 , 51, 435-43	336
1705	Interactions between small nuclear ribonucleoprotein particles in formation of spliceosomes. 1987 , 49, 763-74	516
1704	The role of the kappa enhancer and its binding factor NF-kappa B in the developmental regulation of kappa gene transcription. 1987 , 48, 121-8	261
1703	A variant gene and a pseudogene for human 5S RNA are transcriptionally active in vitro. 1987 , 60, 137-44	10
1702	A multiplicity of CCAAT box-binding proteins. 1987 , 50, 863-72	734
1701	A pre-B cell nuclear protein that specifically interacts with the immunoglobulin V-J recombination sequences. 1987 , 51, 909-17	58
1700	5' cleavage site in eukaryotic pre-mRNA splicing is determined by the overall 5' splice region, not by the conserved 5' GU. 1987 , 50, 237-46	200
1699	Binding of transcription factor TFIID to the major late promoter during in vitro nucleosome assembly potentiates subsequent initiation by RNA polymerase II. 1987 , 51, 613-22	375
1698	42 bp element from LDL receptor gene confers end-product repression by sterols when inserted into viral TK promoter. 1987 , 48, 1061-9	219

1697	Purification and characterization of OTF-1, a transcription factor regulating cell cycle expression of a human histone H2b gene. 1987 , 51, 773-81	514
1696	Alternative splicing of E1A transcripts of adenovirus requires appropriate ionic conditions in vitro. 1987 , 50, 31-9	86
1695	Analysis of RNase-A-resistant regions of adenovirus 2 major late precursor-mRNA in splicing extracts reveals an ordered interaction of nuclear components with the substrate RNA. 1987 , 196, 559-73	56
1694	Novobiocin inhibits initiation of RNA polymerase II-directed transcription of the mouse metallothionein-I gene independent of its effect on DNA topoisomerase II. <i>Nucleic Acids Research</i> , 1987 , 15, 8547-60	20.1 23
1693	Formation of mRNA 3' termini: stability and dissociation of a complex involving the AAUAAA sequence.. 1987 , 6, 177-186	48
1692	Cell type-specificity elements of the immunoglobulin heavy chain gene enhancer.. 1987 , 6, 1323-1330	153
1691	RNA 3' processing regulates histone mRNA levels in a mammalian cell cycle mutant. A processing factor becomes limiting in G1-arrested cells.. 1987 , 6, 1721-1726	77
1690	One cell-specific and three ubiquitous nuclear proteins bind in vitro to overlapping motifs in the domain B1 of the SV40 enhancer.. 1987 , 6, 3005-3013	81
1689	In vitro binding of cell-specific and ubiquitous nuclear proteins to the octamer motif of the SV40 enhancer and related motifs present in other promoters and enhancers.. 1987 , 6, 3015-3025	95
1688	Functional characterization of X. laevis U5 snRNA genes.. 1987 , 6, 3071-3078	34
1687	Interactions of cellular proteins involved in the transcriptional regulation of the human immunodeficiency virus.. 1987 , 6, 3761-3770	214
1686	Six distinct nuclear factors interact with the 75-base-pair repeat of the Moloney murine leukemia virus enhancer. 1987 , 7, 1101-10	262
1685	Two spliceosomes can form simultaneously and independently on synthetic double-intron messenger RNA precursors.. 1987 , 6, 1747-1755	10
1684	Promoter interaction of the E1A-inducible factor E2F and its potential role in the formation of a multi-component complex.. 1987 , 6, 2061-2068	92
1683	A cell-free plant extract for accurate pre-tRNA processing, splicing and modification.. 1987 , 6, 2811-2818	47
1682	Cleavage and polyadenylation of messenger RNA precursors in vitro occurs within large and specific 3' processing complexes.. 1987 , 6, 4159-4168	51
1681	Analysis of mRNA 3' end formation by modification interference: the only modifications which prevent processing lie in AAUAAA and the poly(A) site.. 1987 , 6, 4177-4184	48
1680	A cellular transcription factor E4F1 interacts with an E1a-inducible enhancer and mediates constitutive enhancer function in vitro.. 1987 , 6, 1345-1353	129

1679	Identification and purification of a polypeptide that binds to the c-fos serum response element.. 1987 , 6, 2711-2717	266
1678	Specific pre-cleavage and post-cleavage complexes involved in the formation of SV40 late mRNA 3' termini in vitro.. 1987 , 6, 4185-4192	34
1677	Xenopus cytoskeletal actin and human c-fos gene promoters share a conserved protein-binding site.. 1987 , 6, 667-673	114
1676	Linear oligomeric potato spindle tuber viroid (PSTV) RNAs are accurately processed in vitro to the monomeric circular viroid proper when incubated with a nuclear extract from healthy potato cells. 1987 , 6, 2173-2183	35
1675	A human and a plant intron-containing tRNA ^{Tyr} gene are both transcribed in a HeLa cell extract but spliced along different pathways.. 1987 , 6, 35-41	55
1674	Processing of human CALC-I RNA. 1987 , 12, 194-195	1
1673	Isolation, fractionation and reconstitution of a nuclear extract capable of transcribing ribosomal DNA. 1987 , 73, 11-20	34
1672	Partial purification of plant transcription factors. I. Initiation. 1987 , 9, 147-58	14
1671	Partial purification of plant transcription factors. II. An in vitro transcription system is inefficient. 1987 , 9, 159-69	21
1670	A rapidly inducible DNA-binding activity which binds upstream of the c-fos proto-oncogene. 1987 , Suppl 5, 63-8	6
1669	A transcription factor which binds to the enhancers of SV40, immunoglobulin heavy chain and U2 snRNA genes. 1987 , 325, 268-72	168
1668	An inducible transcription factor activates expression of human immunodeficiency virus in T cells. 1987 , 326, 711-3	1896
1667	Heat shock factor is regulated differently in yeast and HeLa cells. 1987 , 329, 81-4	471
1666	A c-erb-A binding site in rat growth hormone gene mediates trans-activation by thyroid hormone. 1987 , 329, 738-41	344
1665	A homologous in vitro system to analyze transcription of a mouse immunoglobulin mu heavy-chain gene. 1988 , 172, 679-85	6
1664	Interferon and growth factor modulation of nuclear factors binding to 5' upstream elements of the 2-5A synthetase gene. 1988 , 38, 261-7	2
1663	Reversible chemical cross-linking and ribonuclease digestion analysis of the organization of proteins in ribonucleoprotein particles. 1988 , 84, 17-28	11
1662	Cooperativity of the glucocorticoid receptor and the CACCC-box binding factor. 1988 , 332, 87-90	382

1661	HTLV-1 transactivator induces interleukin-2 receptor expression through an NF-kappa B-like factor. 1988 , 333, 776-8	587
1660	Spliceosomal RNA U6 is remarkably conserved from yeast to mammals. 1988 , 334, 213-8	385
1659	Transcriptional autoregulation of the proto-oncogene fos. 1988 , 334, 314-9	590
1658	Phosphorylation-induced binding and transcriptional efficacy of nuclear factor CREB. 1988 , 334, 494-8	1253
1657	Thyroid hormone receptor alpha isoforms generated by alternative splicing differentially activate myosin HC gene transcription. 1988 , 334, 539-42	406
1656	Activation in vitro of sequence-specific DNA binding by a human regulatory factor. 1988 , 335, 372-5	276
1655	Trans splicing involves a novel form of small nuclear ribonucleoprotein particles. 1988 , 335, 559-62	205
1654	DNA binding factors for the CpG-rich island containing the promoter of the human X-linked PGK gene. 1988 , 14, 461-72	14
1653	DNA bending induced by specific interaction of decamer binding proteins with immunoglobulin gene control sequences. 1988 , 101, 145-58	3
1652	The alpha- and beta-tubulins of <i>Toxoplasma gondii</i> are encoded by single copy genes containing multiple introns. 1988 , 29, 261-73	136
1651	Transcription by RNA polymerase III. 1988 , 57, 873-914	644
1650	Activation of cell-specific expression of rat growth hormone and prolactin genes by a common transcription factor. 1988 , 239, 1400-5	477
1649	Chemical cross-linking of Sm and RNP antigenic proteins. 1988 , 27, 4595-600	16
1648	Reconstituted U1 small nuclear ribonucleoprotein complex restores 5' splice site cleavage activity. 1988 , 154, 1010-7	1
1647	Differential enzymatic accessibilities of the 5' and 3' splice sites of beta-globin pre-messenger RNA in splicing competent HeLa cell nuclear extract. 1988 , 150, 517-25	
1646	I kappa B: a specific inhibitor of the NF-kappa B transcription factor. 1988 , 242, 540-6	2056
1645	An in vitro system for accurate methylation of internal adenosine residues in messenger RNA. 1988 , 242, 1159-62	113
1644	Sequence-specific DNA binding protein(s) that bind(s) to a putative human DNA replication origin. 1988 , 37, 1807-8	3

1643	Binding of tissue-specific factors to the enhancer sequence of hepatitis B virus. 1988 , 229, 349-54	3
1642	A 64 kd nuclear protein binds to RNA segments that include the AAUAAA polyadenylation motif. 1988 , 52, 221-8	181
1641	Presence of transcription signals in two putative DNA replication origins of human cells. 1988 , 951, 430-42	22
1640	Induction of kappa transcription by interferon-gamma without activation of NF-kappa B. 1988 , 242, 1036-7	70
1639	A small-scale procedure for preparation of nuclear extracts that support efficient transcription and pre-mRNA splicing. 1988 , 5, 22-31	433
1638	3' cleavage and polyadenylation of mRNA precursors in vitro requires a poly(A) polymerase, a cleavage factor, and a snRNP. 1988 , 54, 875-89	202
1637	Effects of RNA secondary structure on alternative splicing of pre-mRNA: is folding limited to a region behind the transcribing RNA polymerase?. 1988 , 54, 393-401	248
1636	A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. 1988 , 52, 207-19	497
1635	Transcription factor ATF interacts with the TATA factor to facilitate establishment of a preinitiation complex. 1988 , 54, 1033-42	466
1634	A promoter of the rat insulin-like growth factor II gene consists of minimal control elements. 1988 , 199, 61-81	84
1633	Regulation of SV40 early gene expression. 1988 , 66, 567-77	29
1632	DNA-stimulated protein phosphorylation in HeLa whole cell and nuclear extracts. 1988 , 157, 535-40	20
1631	Inhibition of c-H-ras oncogene induced transformation of rat embryo fibroblasts by cotransfected polynucleotides containing alternating purine-pyrimidine sequences. 1988 , 152, 449-55	
1630	Selective transcription of an insulin-regulated gene in nuclear extracts of rat hepatoma cells. 1988 , 153, 116-27	4
1629	Separation and characterization of a poly(A) polymerase and a cleavage/specificity factor required for pre-mRNA polyadenylation. 1988 , 52, 731-42	188
1628	Cellular factor affecting the stability of beta-globin mRNA. 1988 , 62, 65-74	26
1627	Congenital immunodeficiency with a regulatory defect in MHC class II gene expression lacks a specific HLA-DR promoter binding protein, RF-X. 1988 , 53, 897-906	242
1626	Herpes simplex virus regulatory elements and the immunoglobulin octamer domain bind a common factor and are both targets for virion transactivation. 1988 , 52, 435-45	388

1625	Activation of DNA-binding activity in an apparently cytoplasmic precursor of the NF-kappa B transcription factor. 1988 , 53, 211-7	1152
1624	Activation of transcription factor IIIc by the adenovirus E1A protein. 1988 , 53, 907-20	210
1623	fos-associated cellular p39 is related to nuclear transcription factor AP-1. 1988 , 54, 553-60	491
1622	Analysis of the role of the transcription factor ATF in the assembly of a functional preinitiation complex. 1988 , 54, 1043-51	183
1621	An adenovirus E1A protein domain activates transcription in vivo and in vitro in the absence of protein synthesis. 1988 , 53, 921-6	74
1620	GAL4 derivatives function alone and synergistically with mammalian activators in vitro. 1988 , 54, 659-64	295
1619	Hepatocyte-specific promoter element HP1 of the Xenopus albumin gene interacts with transcriptional factors of mammalian hepatocytes. 1988 , 202, 307-20	100
1618	Synergism of closely adjacent estrogen-responsive elements increases their regulatory potential. 1988 , 201, 537-44	105
1617	A complex formed between cell components and an HSV structural polypeptide binds to a viral immediate early gene regulatory DNA sequence. 1988 , 52, 425-34	358
1616	The HeLa cell protein TEF-1 binds specifically and cooperatively to two SV40 enhancer motifs of unrelated sequence. 1988 , 54, 931-42	302
1615	Nucleotide sequence and transcription of a rat tRNA(Phe) gene and a neighboring Alu-like element. 1988 , 69, 275-85	4
1614	Factors involved in specific transcription by mammalian RNA polymerase II: purification, genetic specificity, and TATA box-promoter interactions of TFIID. 1988 , 8, 4028-40	509
1613	Cell-specific activity of the human immunodeficiency virus enhancer repeat in vitro. 1988 , 4, 381-91	17
1612	The ubiquitous octamer-binding protein Oct-1 contains a POU domain with a homeo box subdomain. 1988 , 2, 1582-99	594
1611	cAMP acts in synergy with E1A protein to activate transcription of the adenovirus early genes E4 and E1A. 1988 , 2, 1517-28	61
1610	Multiple factors are required for poly(A) addition to a mRNA 3' end. 1988 , 2, 588-97	63
1609	The regions of the herpes simplex virus type 1 immediate early protein Vmw175 required for site specific DNA binding closely correspond to those involved in transcriptional regulation. <i>Nucleic Acids Research</i> , 1988 , 16, 11005-25	20.1 74
1608	Functional dissection of 5' and 3' extragenic control regions of human tRNA(Val) genes reveals two different regulatory effects. 1988 , 7, 87-97	17

1607	A high-efficiency HeLa cell nuclear transcription extract. 1988 , 7, 47-55		642
1606	Multiple factors are required for specific RNA cleavage at a poly(A) addition site. 1988 , 2, 578-87		102
1605	Sp1 transcription factor binds DNA and activates transcription even when the binding site is CpG methylated. 1988 , 2, 1127-35		265
1604	Components required for in vitro cleavage and polyadenylation of eukaryotic mRNA. <i>Nucleic Acids Research</i> , 1988 , 16, 5323-44	20.1	23
1603	The pseudorabies immediate early protein stimulates in vitro transcription by facilitating TFIID: promoter interactions. 1988 , 2, 542-53		157
1602	A 5S rRNA/L5 complex is a precursor to ribosome assembly in mammalian cells. 1988 , 106, 545-56		159
1601	Alternative splicing of SV40 early pre-mRNA is determined by branch site selection. 1988 , 2, 1460-75		71
1600	The 5' flanking region of the gene for the Epstein-Barr virus-encoded nuclear antigen 2 contains a cell type specific cis-acting regulatory element that activates transcription in transfected B-cells. <i>Nucleic Acids Research</i> , 1988 , 16, 8391-410	20.1	51
1599	A DNA binding protein regulated by IL-4 and by differentiation in B cells. 1988 , 242, 1559-62		63
1598	The DNA-binding properties of the major regulatory protein alpha 4 of herpes simplex viruses. 1988 , 239, 1531-4		131
1597	Selective activation of transcription by a novel CCAAT binding factor. 1988 , 241, 582-5		203
1596	Cloning and expression of AP-2, a cell-type-specific transcription factor that activates inducible enhancer elements. 1988 , 2, 1557-69		503
1595	Functional cooperativity between transcription factors UBF1 and SL1 mediates human ribosomal RNA synthesis. 1988 , 241, 1192-7		325
1594	Gel electrophoretic isolation of splicing complexes containing U1 small nuclear ribonucleoprotein particles. 1988 , 8, 814-21		206
1593	Detection of two tissue-specific DNA-binding proteins with affinity for sites in the mouse beta-globin intervening sequence 2. 1988 , 8, 381-92		71
1592	A highly conserved, 5' untranslated, inverted repeat sequence is ineffective in translational control of the alpha 1(I) collagen gene. <i>Nucleic Acids Research</i> , 1988 , 16, 9721-36	20.1	19
1591	An erythroid specific nuclear factor binding to the proximal CACCC box of the beta-globin gene promoter. <i>Nucleic Acids Research</i> , 1988 , 16, 4299-313	20.1	76
1590	3' processing of pre-mRNA plays a major role in proliferation-dependent regulation of histone gene expression. <i>Nucleic Acids Research</i> , 1988 , 16, 9399-414	20.1	70

1589	Pre-tRNA splicing in a nuclear extract from human leukaemia cells: separation of endonuclease and ligase activities. <i>Nucleic Acids Research</i> , 1988 , 16, 7713-4	20.1	1
1588	Sigma region located between C mu and C delta genes of human immunoglobulin heavy chain: possible involvement of tRNA-like structure in RNA splicing. <i>Nucleic Acids Research</i> , 1988 , 16, 9497-511	20.1	20
1587	A 5' exo-ribonuclease and RNA ligase of <i>T. brucei</i> . <i>Nucleic Acids Research</i> , 1988 , 16, 9737-59	20.1	8
1586	An embryonic DNA-binding protein specific for a region of the human IFN beta 1 promoter. <i>Nucleic Acids Research</i> , 1988 , 16, 10575-92	20.1	10
1585	An ACCC-containing protein-binding sequence in the neighbourhood of the decanucleotide recognition site of the immunoglobulin gene promoter. <i>Nucleic Acids Research</i> , 1988 , 16, 3693-704	20.1	7
1584	An in vitro interaction between the human U3 snRNP and 28S rRNA sequences near the alpha-sarcin site. <i>Nucleic Acids Research</i> , 1988 , 16, 10493-509	20.1	40
1583	Transcription of tRNA genes in a S100 extract from human leukaemia cells. <i>Nucleic Acids Research</i> , 1988 , 16, 7715	20.1	
1582	Binding of a cytosolic protein to the iron-responsive element of human ferritin messenger RNA. 1988 , 241, 1207-10		346
1581	The mouse albumin promoter and a distal upstream site are simultaneously DNase I hypersensitive in liver chromatin and bind similar liver-abundant factors in vitro. 1988 , 2, 528-41		66
1580	Unusual branch point selection involved in splicing of the alternatively processed Calcitonin/CGRP-I pre-mRNA. <i>Nucleic Acids Research</i> , 1988 , 16, 9513-26	20.1	48
1579	Rapid detection of DNA-binding factors using protein-blotting and digoxigenin-dUTP marked probes. <i>Nucleic Acids Research</i> , 1988 , 16, 11839	20.1	30
1578	Induction of proto-oncogene fos transcription through the adenylate cyclase pathway: characterization of a cAMP-responsive element. 1988 , 2, 1529-38		327
1577	In vitro splicing of adenovirus E1A transcripts: characterization of novel reactions and of multiple branch points abnormally far from the 3' splice site. <i>Nucleic Acids Research</i> , 1988 , 16, 2389-409	20.1	45
1576	All human tRNA ^{Tyr} genes contain introns as a prerequisite for pseudouridine biosynthesis in the anticodon. <i>Nucleic Acids Research</i> , 1988 , 16, 1951-66	20.1	58
1575	Heavy metal ions in transcription factors from HeLa cells: Sp1, but not octamer transcription factor requires zinc for DNA binding and for activator function. <i>Nucleic Acids Research</i> , 1988 , 16, 5771-81	20.1	57
1574	A family of immunologically related transcription factors that includes multiple forms of ATF and AP-1. 1988 , 2, 1216-26		234
1573	Inducible DNA-protein interactions of the murine kappa immunoglobulin enhancer in intact cells: comparison with in vitro interactions. <i>Nucleic Acids Research</i> , 1988 , 16, 953-67	20.1	24
1572	Model for tissue specific Calcitonin/CGRP-I RNA processing from in vitro experiments. <i>Nucleic Acids Research</i> , 1988 , 16, 7867-83	20.1	23

1571	Interferon-induced nuclear factors that bind a shared promoter element correlate with positive and negative transcriptional control. 1988 , 2, 383-93		479
1570	Alternative splicing of tropomyosin pre-mRNAs in vitro and in vivo. 1988 , 2, 1627-38		105
1569	The effects of HPFH mutations in the human gamma-globin promoter on binding of ubiquitous and erythroid specific nuclear factors. <i>Nucleic Acids Research</i> , 1988 , 16, 7783-97	20.1	88
1568	Estrogen-dependent in vitro transcription from the vitellogenin promoter in liver nuclear extracts. 1988 , 239, 1137-9		79
1567	Xeroderma pigmentosum group E cells lack a nuclear factor that binds to damaged DNA. 1988 , 242, 564-7		369
1566	Octamer-binding proteins from B or HeLa cells stimulate transcription of the immunoglobulin heavy-chain promoter in vitro. 1988 , 2, 1227-37		117
1565	Multiple nuclear factors interact with promoter sequences of the urokinase-type plasminogen activator gene. <i>Nucleic Acids Research</i> , 1988 , 16, 7527-44	20.1	23
1564	EivF, a factor required for transcription of the adenovirus EIV promoter, binds to an element involved in Ela-dependent activation and cAMP induction. 1988 , 2, 975-90		59
1563	Presplicing complex formation requires two proteins and U2 snRNP. 1988 , 2, 1155-67		105
1562	The human beta-globin gene 3' enhancer contains multiple binding sites for an erythroid-specific protein. 1988 , 2, 1089-100		318
1561	In vitro transcription of the Drosophila engrailed gene. 1988 , 2, 68-81		266
1560	Two adjacent promoter elements mediate nerve growth factor activation of the c-fos gene and bind distinct nuclear complexes. 1988 , 85, 9474-8		79
1559	B-cell control region at the 5' end of a major histocompatibility complex class II gene: sequences and factors. 1988 , 8, 3975-87		72
1558	A common factor regulates skeletal and cardiac alpha-actin gene transcription in muscle. 1988 , 8, 4120-33		108
1557	A common trans-acting factor is involved in transcriptional regulation of neurotransmitter genes by cyclic AMP. 1988 , 8, 4225-33		155
1556	Prolactin upstream factor I mediates cell-specific transcription. 1988 , 8, 5432-8		27
1555	A herpesvirus trans-activating protein interacts with transcription factor OTF-1 and other cellular proteins. 1988 , 85, 6347-51		357
1554	Several distinct "CCAAT" box binding proteins coexist in eukaryotic cells. 1988 , 85, 757-61		233

1553	Interaction of nuclear proteins with muscle-specific regulatory sequences of the human cardiac alpha-actin promoter. 1988 , 8, 4110-9	94
1552	Inducible expression and cytogenetic effects of the EcoRI restriction endonuclease in Chinese hamster ovary cells. 1988 , 8, 4204-11	43
1551	Multiple hepatic trans-acting factors are required for in vitro transcription of the human alpha-1-antitrypsin gene. 1988 , 8, 4362-9	60
1550	Sequence-specific interactions between a cellular DNA-binding protein and the simian virus 40 origin of DNA replication. 1988 , 8, 903-11	52
1549	Identification of two transcription factors that bind to specific elements in the promoter of the adenovirus early-region 4. 1988 , 8, 1290-300	67
1548	Unusual branch point selection in processing of human growth hormone pre-mRNA. 1988 , 8, 2011-20	68
1547	A DNA-binding factor specific for xenobiotic responsive elements of P-450c gene exists as a cryptic form in cytoplasm: its possible translocation to nucleus. 1988 , 85, 5859-63	195
1546	An erythrocyte-specific DNA-binding factor recognizes a regulatory sequence common to all chicken globin genes. 1988 , 85, 5976-80	463
1545	Identification of a functional mammalian spliceosome containing unspliced pre-mRNA. 1988 , 85, 7216-20	74
1544	Splicing mutation in human hereditary analbuminemia. 1988 , 85, 2125-9	46
1543	Short donor site sequences inserted within the intron of beta-globin pre-mRNA serve for splicing in vitro. 1988 , 8, 4484-91	46
1542	Identification of protein-binding sites in the hepatitis B virus enhancer and core promoter domains. 1988 , 8, 5159-65	49
1541	Nuclear proteins that bind the human gamma-globin gene promoter: alterations in binding produced by point mutations associated with hereditary persistence of fetal hemoglobin. 1988 , 8, 5310-22	120
1540	Identification and characterization of multiple erythroid cell proteins that interact with the promoter of the murine alpha-globin gene. 1988 , 8, 3215-26	58
1539	Identification of an inducible factor that binds to a positive regulatory element of the human beta-interferon gene. 1988 , 85, 3309-13	101
1538	Association of U2, U4, U5, and U6 small nuclear ribonucleoproteins in a spliceosome-type complex in absence of precursor RNA. 1988 , 85, 5459-62	64
1537	NF-kappa B protein purification from bovine spleen: nucleotide stimulation and binding site specificity. 1988 , 85, 8825-9	141
1536	Nuclear proteins interacting with the promoter region of the human granulocyte/macrophage colony-stimulating factor gene. 1988 , 85, 674-8	114

1535	Accurate and efficient pre-mRNA splicing in Drosophila cell-free extracts. 1988 , 85, 2904-8	73
1534	Cytoplasmic protein binds in vitro to a highly conserved sequence in the 5' untranslated region of ferritin heavy- and light-subunit mRNAs. 1988 , 85, 2171-5	667
1533	Two distinct nuclear factors bind the conserved regulatory sequences of a rabbit major histocompatibility complex class II gene. 1988 , 8, 2034-41	17
1532	DNA affinity labeling of adenovirus type 2 upstream promoter sequence-binding factors identifies two distinct proteins. 1988 , 8, 105-13	44
1531	A cell-specific enhancer of the mouse alpha 1-antitrypsin gene has multiple functional regions and corresponding protein-binding sites. 1988 , 8, 1055-66	83
1530	Delineation of DNA sequences that are important for in vitro transcription from the adenovirus Ella late promoter. 1988 , 8, 1906-14	9
1529	trans splicing in Leishmania enriettii and identification of ribonucleoprotein complexes containing the spliced leader and U2 equivalent RNAs. 1988 , 8, 2597-603	26
1528	Negative regulation of the human polyomavirus BK enhancer involves cell-specific interaction with a nuclear repressor. 1988 , 8, 3448-57	38
1527	Translation of globin RNA spliced in vitro. 1988 , 254, 21-6	2
1526	Conserved region of the rat alpha-lactalbumin promoter is a target site for protein binding in vitro. 1988 , 256, 391-6	42
1525	Assembly of a polyadenylation-specific 25S ribonucleoprotein complex in vitro. 1988 , 8, 2052-62	47
1524	Interferon-induced transcription of a major histocompatibility class I gene accompanies binding of inducible nuclear factors to the interferon consensus sequence. 1988 , 85, 5884-8	138
1523	Cyclic AMP induction of early adenovirus promoters involves sequences required for E1A trans-activation. 1988 , 85, 7192-6	131
1522	The cell-specific enhancer of the mouse transthyretin (prealbumin) gene binds a common factor at one site and a liver-specific factor(s) at two other sites. 1988 , 8, 81-90	116
1521	Purification of multiple erythroid cell proteins that bind the promoter of the alpha-globin gene. 1988 , 8, 4270-81	54
1520	Differential protein binding in lymphocytes to a sequence in the enhancer of the mouse retrovirus SL3-3. 1988 , 8, 1625-37	110
1519	DNA-binding factors of B lymphoid cells are susceptible to limited proteolytic cleavage during nuclear extract preparation. 1988 , 8, 1812-5	13
1518	Transcription of HLA class II genes in the absence of B-cell-specific octamer-binding factor. 1988 , 8, 3734-9	30

1517	Identification of a novel TA-rich DNA binding protein that recognizes a TATA sequence within the brain creatine kinase promoter. <i>Nucleic Acids Research</i> , 1988 , 16, 8925-44	20.1	54
1516	Identification of murine nuclear proteins that bind to the conserved octamer sequence of the immunoglobulin promoter region. 1988 , 85, 3560-4		17
1515	Adenoviral control regions activated by E1A and the cAMP response element bind to the same factor. 1988 , 85, 4171-5		112
1514	Phosphorylation of serum response factor, a factor that binds to the serum response element of the c-FOS enhancer. 1988 , 85, 7206-10		180
1513	Spliceosome assembly involves the binding and release of U4 small nuclear ribonucleoprotein. 1988 , 85, 411-5		189
1512	T-cell-specific expression of interleukin 2: evidence for a negative regulatory site. 1988 , 85, 2934-8		58
1511	Identification and characterization of two functional domains within the murine heavy-chain enhancer. 1988 , 8, 145-52		96
1510	The C proteins of heterogeneous nuclear ribonucleoprotein complexes interact with RNA sequences downstream of polyadenylation cleavage sites. 1988 , 8, 4477-83		88
1509	Concentration dependence of transcriptional transactivation in inducible E1A-containing human cells. 1988 , 8, 4799-807		33
1508	DNA-binding site for two skeletal actin promoter factors is important for expression in muscle cells. 1988 , 8, 1800-2		79
1507	Mutations in poly(A) site downstream elements affect in vitro cleavage activity. 1988 , 8, 1839-41		29
1506	Three sequence-specific DNA-protein complexes are formed with the same promoter element essential for expression of the rat somatostatin gene. 1988 , 8, 1947-56		67
1505	Multiple cis-acting sequence elements are required for efficient splicing of simian virus 40 small-t antigen pre-mRNA. 1988 , 8, 3582-90		57
1504	Adenovirus transcriptional regulatory regions are conserved in mammalian cells and <i>Saccharomyces cerevisiae</i> . 1988 , 8, 3717-25		12
1503	Purification and characterization of TTFI, a factor that mediates termination of mouse ribosomal DNA transcription. 1988 , 8, 3891-7		69
1502	Processing at immunoglobulin polyadenylation sites in lymphoid cell extracts.. 1988 , 7, 1421-1429		14
1501	Nuclear factor NF-kappa B can interact functionally with its cognate binding site to provide lymphoid-specific promoter function.. 1988 , 7, 3109-3113		53
1500	Two transcription factors, NF-kappa B and H2TF1, interact with a single regulatory sequence in the class I major histocompatibility complex promoter. 1988 , 85, 723-7		315

1499	The human beta-globin promoter; nuclear protein factors and erythroid specific induction of transcription.. 1988 , 7, 4203-4212	135
1498	Interferon-induced binding of nuclear factors to promoter elements of the 2-5A synthetase gene.. 1988 , 7, 751-759	128
1497	Trypanosome trans-splicing utilizes 2'-5' branches and a corresponding debranching activity.. 1988 , 7, 1431-1437	25
1496	Tissue specific trans-acting factor interaction with proximal rat prolactin gene promoter sequences.. 1988 , 7, 1721-1733	20
1495	Purification of the human immunodeficiency virus type 1 enhancer and TAR binding proteins EBP-1 and UBP-1.. 1988 , 7, 2117-2130	152
1494	Antisense RNA inhibits splicing of pre-mRNA in vitro.. 1988 , 7, 2523-2532	57
1493	Physical identification of branched intron side-products of splicing in <i>Trypanosoma brucei</i> .. 1988 , 7, 2539-2545	16
1492	RNA binding specificity of hnRNP proteins: a subset bind to the 3' end of introns.. 1988 , 7, 3519-3529	153
1491	A zinc-responsive factor interacts with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I gene.. 1988 , 7, 3763-3770	204
1490	Proteins bound at adjacent DNA elements act synergistically to regulate human proenkephalin cAMP inducible transcription.. 1988 , 7, 3793-3805	256
1489	Wheat germ splicing endonuclease is highly specific for plant pre-tRNAs.. 1988 , 7, 3823-3828	26
1488	Identification of a novel lymphoid specific octamer binding protein (OTF-2B) by proteolytic clipping bandshift assay (PCBA).. 1988 , 7, 4221-4229	188
1487	Comparison of in vitro and in vivo splice site selection in kappa-immunoglobulin precursor mRNA. 1988 , 8, 2610-9	22
1486	Simian virus 40 T antigen alters the binding characteristics of specific simian DNA-binding factors. 1988 , 8, 1648-56	38
1485	Negative regulation of transcription in vitro by a glucocorticoid response element is mediated by a trans-acting factor. 1988 , 8, 3872-81	66
1484	The role of the polypyrimidine stretch at the SV40 early pre-mRNA 3' splice site in alternative splicing.. 1988 , 7, 809-817	67
1483	Purification of a NF1-like DNA-binding protein from rat liver and cloning of the corresponding cDNA.. 1988 , 7, 3115-3123	172
1482	Complementation between two cell lines lacking kappa enhancer activity: implications for the developmental control of immunoglobulin transcription.. 1988 , 7, 4213-4220	24

1481	Enhancer-like interferon responsive sequences of the human and murine (2?-5?) oligoadenylate synthetase gene promoters.. 1988 , 7, 1411-1419	135
1480	The promoter of Alzheimer's disease amyloid A4 precursor gene.. 1988 , 7, 2807-2813	231
1479	Heat shock but not other stress inducers leads to the disruption of a sub-set of snRNPs and inhibition of in vitro splicing in HeLa cells.. 1988 , 7, 3509-3518	126
1478	Interaction of a nodule specific, trans -acting factor with distinct DNA elements in the soybean leghaemoglobin lbc 3 5' upstream region. 1988 , 7, 1265-1271	100
1477	Two different liver-specific factors stimulate in vitro transcription from the human alpha 1-antitrypsin promoter.. 1988 , 7, 2075-2087	141
1476	A cell-specific activator in the Xenopus A2 vitellogenin gene: promoter elements functioning with rat liver nuclear extracts.. 1988 , 7, 2495-2501	23
1475	The -17 mutation in Greek HPFH affects the binding of three nuclear factors to the CCAAT region of the gamma-globin gene.. 1988 , 7, 3099-3107	89
1474	Direct combinatorial interaction between a herpes simplex virus regulatory protein and a cellular octamer-binding factor mediates specific induction of virus immediate-early gene expression.. 1988 , 7, 4231-4238	103
1473	Cells resistant to interferon are defective in activation of a promoter-binding factor.. 1988 , 7, 3779-3783	57
1472	The first intron of the 4F2 heavy-chain gene contains a transcriptional enhancer element that binds multiple nuclear proteins. 1989 , 9, 2588-97	70
1471	Mapping of branch sites in trans-spliced pre-mRNAs of Trypanosoma brucei. 1989 , 9, 4291-7	46
1470	An abundant U6 snRNP found in germ cells and embryos of Xenopus laevis.. 1989 , 8, 4179-4187	51
1469	TPA can overcome the requirement for Ela and together act synergistically in stimulating expression of the adenovirus E111 promoter.. 1989 , 8, 4239-4250	18
1468	Independent removal of two identical or different introns from mRNA precursors in vitro. 1989 , 105, 1014-23	
1467	Accurate processing of human pre-rRNA in vitro. 1989 , 9, 4422-31	51
1466	The Xenopus ribosomal gene enhancers bind an essential polymerase I transcription factor, xUBF. 1989 , 3, 1779-88	122
1465	Double-stranded RNA activates binding of NF-kappa B to an inducible element in the human beta-interferon promoter.. 1989 , 8, 1129-1138	220
1464	Suppression of MHC class I gene expression by N-myc through enhancer inactivation.. 1989 , 8, 3351-3355	57

1463	Electron microscopy of U1 small nuclear ribonucleoprotein particles: shape of the particle and position of the 5' RNA terminus.. 1989 , 8, 277-286	34
1462	Non-enzymatic excision of pre-tRNA introns?. 1989 , 8, 293-300	11
1461	Interactions of alpha- and gamma-interferon in the transcriptional regulation of the gene encoding a guanylate-binding protein.. 1989 , 8, 2009-2014	113
1460	The SV40 TC-II(kappa B) enhanson binds ubiquitous and cell type specifically inducible nuclear proteins from lymphoid and non-lymphoid cell lines.. 1989 , 8, 4215-4227	44
1459	NF-kappa B activation of the cytomegalovirus enhancer is mediated by a viral transactivator and by T cell stimulation.. 1989 , 8, 4251-4258	252
1458	DNA replication origin and transcriptional enhancer in c-myc gene share the c-myc protein binding sequences.. 1989 , 8, 4273-4279	71
1457	Phosphorylation-dependent activation of the adenovirus-inducible E2F transcription factor in a cell-free system. 1989 , 86, 4352-6	99
1456	Identification of an octamer-binding site in the mouse kappa light-chain immunoglobulin enhancer. 1989 , 9, 4239-47	39
1455	Human immunodeficiency virus type 1 LTR TATA and TAR region sequences required for transcriptional regulation.. 1989 , 8, 765-778	197
1454	A thyroid-specific nuclear protein essential for tissue-specific expression of the thyroglobulin promoter.. 1989 , 8, 2537-2542	245
1453	Repression of c-fos transcription is mediated through p67SRF bound to the SRE.. 1989 , 8, 2567-2574	98
1452	U3, U8 and U13 comprise a new class of mammalian snRNPs localized in the cell nucleolus.. 1989 , 8, 3113-3119	235
1451	Purification and analysis of RNA polymerase II transcription factors by using wheat germ agglutinin affinity chromatography. 1989 , 86, 1781-5	170
1450	Two distinct cellular phosphoproteins bind to the c-fos serum response element.. 1989 , 8, 1785-1792	87
1449	A family of octamer-specific proteins present during mouse embryogenesis: evidence for germline-specific expression of an Oct factor.. 1989 , 8, 2543-2550	452
1448	Octamer binding proteins confer transcriptional activity in early mouse embryogenesis.. 1989 , 8, 2551-2557	224
1447	Identification of the RNA binding segment of human U1 A protein and definition of its binding site on U1 snRNA.. 1989 , 8, 4163-4170	202
1446	In vitro synthesis of vertebrate U1 snRNA.. 1989 , 8, 287-292	27

1445	Transcriptional enhancement by upstream activators is brought about by different molecular mechanisms for class I and II RNA polymerase genes.. 1989 , 8, 3011-3017		7
1444	Mapping U2 snRNP--pre-mRNA interactions using biotinylated oligonucleotides made of 2'-OMe RNA.. 1989 , 8, 4171-4178		41
1443	The octamer-binding proteins form multi-protein--DNA complexes with the HSV alpha TIF regulatory protein.. 1989 , 8, 4229-4238		171
1442	Evidence from complementation assays in vitro that U5 snRNP is required for both steps of mRNA splicing.. 1989 , 8, 3105-3112		23
1441	A common protein binds to two silencers 5' to the human beta-globin gene. <i>Nucleic Acids Research</i> , 1989 , 17, 8833-52	20.1	88
1440	Nuclear factor E2F mediates basic transcription and trans-activation by E1a of the human MYC promoter. 1989 , 3, 527-36		218
1439	Cis-acting sequences from mouse rDNA promote plasmid DNA amplification and persistence in mouse cells: implication of HMG-I in their function. <i>Nucleic Acids Research</i> , 1989 , 17, 9909-32	20.1	61
1438	Cytoplasmic activation of ISGF3, the positive regulator of interferon-alpha-stimulated transcription, reconstituted in vitro. 1989 , 3, 1362-71		421
1437	In vitro reconstitution of snRNPs: a reconstituted U4/U6 snRNP participates in splicing complex formation. 1989 , 3, 479-87		36
1436	Identification of a functional NF-kappa B binding site in the murine T cell receptor beta 2 locus. 1989 , 170, 1737-43		28
1435	An ordered pathway of assembly of components required for polyadenylation site recognition and processing. 1989 , 3, 2180-90		176
1434	Induction of transcription factor AP-1 by adenovirus E1A protein and cAMP. 1989 , 3, 1991-2002		63
1433	The Hox-1.3 homeo box protein is a sequence-specific DNA-binding phosphoprotein. 1989 , 3, 158-72		123
1432	Four factors are required for 3'-end cleavage of pre-mRNAs. 1989 , 3, 1711-24		190
1431	In vitro 3' end processing and poly(A) tailing of RNA in <i>Trypanosoma cruzi</i> . <i>Nucleic Acids Research</i> , 1989 , 17, 4647-60	20.1	21
1430	Effect of the cap structure on pre-mRNA splicing in <i>Xenopus</i> oocyte nuclei. 1989 , 3, 1472-9		68
1429	3'-end formation of mouse pre-rRNA involves both transcription termination and a specific processing reaction. 1989 , 3, 224-31		62
1428	Identification and purification of a 62,000-dalton protein that binds specifically to the polypyrimidine tract of introns. 1989 , 3, 1874-86		295

1427	Two tissue-specific factors bind the erythroid promoter of the human porphobilinogen deaminase gene. <i>Nucleic Acids Research</i> , 1989 , 17, 37-54	20.1	297
1426	DNA curvature in front of the human mitochondrial L-strand replication origin with specific protein binding. <i>Nucleic Acids Research</i> , 1989 , 17, 6077-86	20.1	19
1425	A secondary structure at the 3' splice site affects the in vitro splicing reaction of mouse immunoglobulin mu chain pre-mRNAs. <i>Nucleic Acids Research</i> , 1989 , 17, 8159-69	20.1	76
1424	Purification and characterization of a protein that binds to the recombination signal sequence of the immunoglobulin J kappa segment. <i>Nucleic Acids Research</i> , 1989 , 17, 9015-26	20.1	64
1423	Binding of the herpes simplex virus type 1 UL9 gene product to an origin of viral DNA replication. <i>Nucleic Acids Research</i> , 1989 , 17, 1409-25	20.1	77
1422	Enhancer functions and in vitro protein binding of native and mutated interferon-responsive sequences. <i>Nucleic Acids Research</i> , 1989 , 17, 1679-95	20.1	29
1421	Construction and characterization of herpes simplex virus type 1 mutants with defined lesions in immediate early gene 1. 1989 , 70 (Pt 5), 1185-202		158
1420	Nuclear proteins TREF1 and TREF2 bind to the transcriptional control element of the transferrin receptor gene and appear to be associated as a heterodimer. 1989 , 1, 151-64		51
1419	Branch point selection in alternative splicing of tropomyosin pre-mRNAs. <i>Nucleic Acids Research</i> , 1989 , 17, 5633-50	20.1	90
1418	Unique sequence organization and erythroid cell-specific nuclear factor-binding of mammalian theta 1 globin promoters. <i>Nucleic Acids Research</i> , 1989 , 17, 5687-700	20.1	36
1417	CpG methylation of the cAMP-responsive enhancer/promoter sequence TGACGTCA abolishes specific factor binding as well as transcriptional activation. 1989 , 3, 612-9		435
1416	Increased Sp1 binding mediates erythroid-specific overexpression of a mutated (HPFH) gamma-globulin promoter. <i>Nucleic Acids Research</i> , 1989 , 17, 10231-41	20.1	57
1415	Identification of a novel site specific endonuclease produced by Mycoplasma fermentans: discovery while characterizing DNA binding proteins in T lymphocyte cell lines. <i>Nucleic Acids Research</i> , 1989 , 17, 3491-9	20.1	5
1414	Differential binding of interferon-induced factors to an oligonucleotide that mediates transcriptional activation. <i>Nucleic Acids Research</i> , 1989 , 17, 3415-24	20.1	72
1413	Advanced Research on Animal Cell Technology. 1989 ,		0
1412	RNA editing: a novel mechanism for regulating lipid transport from the intestine. 1989 , 30 Spec No, 35-43		8
1411	A transcription factor, TFIS, interacts with both the promoter and enhancer of the Xenopus rRNA genes. 1989 , 3, 1768-78		41
1410	Efficiency of utilization of the simian virus 40 late polyadenylation site: effects of upstream sequences. 1989 , 9, 4248-58		195

1409	Evidence for nuclear factors involved in recognition of 5' splice sites. <i>Nucleic Acids Research</i> , 1989 , 17, 2655-74	20.1	40
1408	Genetic analysis of the role of human U1 snRNA in mRNA splicing: I. Effect of mutations in the highly conserved stem-loop I of U1. 1989 , 3, 697-707		27
1407	CCAAT/enhancer binding protein activates the promoter of the serum albumin gene in cultured hepatoma cells. 1989 , 3, 1314-22		446
1406	Specific recognition of cruciform DNA by nuclear protein HMG1. 1989 , 243, 1056-9		569
1405	Regulatory elements controlling the basal and drug-inducible expression of glutathione S-transferase Ya subunit gene. 1989 , 8, 399-408		30
1404	Analysis of possible repressor elements in the 5'-flanking region of the human beta-globin gene. 1989 , 8, 715-21		7
1403	Specific binding of estrogen receptor to the estrogen response element. 1989 , 9, 43-9		194
1402	The adenovirus early region 4 open reading frame 6/7 protein regulates the DNA binding activity of the cellular transcription factor, E2F, through a direct complex. 1989 , 3, 1699-710		114
1401	Developmental and tissue-specific expression of nuclear proteins that bind the regulatory element of the major histocompatibility complex class I gene. 1989 , 169, 1309-21		90
1400	The cell-specific transcription factor PTF1 contains two different subunits that interact with the DNA. 1989 , 3, 1613-24		71
1399	An adenovirus early region 4 gene product is required for induction of the infection-specific form of cellular E2F activity. 1989 , 3, 1062-74		92
1398	Nuclear protein factors and erythroid transcription of the human A gamma-globin gene. <i>Nucleic Acids Research</i> , 1989 , 17, 3811-27	20.1	49
1397	Anti-OTF-1 antibodies inhibit NFIII stimulation of in vitro adenovirus DNA replication. <i>Nucleic Acids Research</i> , 1989 , 17, 1845-63	20.1	29
1396	Clusters of nuclear factor I binding sites identify enhancers of several papillomaviruses but alone are not sufficient for enhancer function. <i>Nucleic Acids Research</i> , 1989 , 17, 3519-33	20.1	82
1395	Dual enhancer activities of the cyclic-AMP responsive element with cell type and promoter specificity. <i>Nucleic Acids Research</i> , 1989 , 17, 1521-36	20.1	32
1394	The role of a repetitive palindromic sequence element in the human cytomegalovirus major immediate early enhancer. 1989 , 70 (Pt 1), 107-23		47
1393	Distamycins inhibit the binding of OTF-1 and NFE-1 transactors to their conserved DNA elements. <i>Nucleic Acids Research</i> , 1989 , 17, 1051-9	20.1	99
1392	Point mutational analysis of the human c-fos serum response factor binding site. <i>Nucleic Acids Research</i> , 1989 , 17, 1177-95	20.1	97

1391	Nuclear factors interacting with an interleukin-6 responsive element of rat alpha 2-macroglobulin gene. <i>Nucleic Acids Research</i> , 1989 , 17, 9425-35	20.1	62
1390	Identification of an amplification promoting DNA sequence from the hypotrichous ciliate <i>Stylonychia lemnae</i> . <i>Nucleic Acids Research</i> , 1989 , 17, 8783-802	20.1	18
1389	Purification of DNA-binding transcription factors by their selective adsorption on the affinity latex particles. <i>Nucleic Acids Research</i> , 1989 , 17, 6229-40	20.1	59
1388	Mutations in two regions upstream of the A gamma globin gene canonical promoter affect gene expression. <i>Nucleic Acids Research</i> , 1989 , 17, 4339-52	20.1	35
1387	Developmental modulation of protein binding to beta-globin gene regulatory sites within chicken erythrocyte nuclei. 1989 , 3, 1860-73		77
1386	Purified U5 small nuclear ribonucleoprotein can relieve the inhibition of spliceosome assembly and splicing by snRNP-free nuclear proteins. <i>Nucleic Acids Research</i> , 1989 , 17, 5223-43	20.1	11
1385	DNA recognition element required for PUF-I mediated cell-type-specific transcription of the rat prolactin gene. <i>Nucleic Acids Research</i> , 1989 , 17, 2705-22	20.1	17
1384	Determinant differences between the rabbit and mouse immunoglobulin kappa enhancers impair the activity of the rabbit enhancer in mouse myeloma cells. <i>Nucleic Acids Research</i> , 1989 , 17, 4745-55	20.1	6
1383	Cooperative interaction of chicken lysozyme enhancer sub-domains partially overlapping with a steroid receptor binding site. <i>Nucleic Acids Research</i> , 1989 , 17, 4975-91	20.1	36
1382	Binding of a pancreatic nuclear protein is correlated with amylase enhancer activity. <i>Nucleic Acids Research</i> , 1989 , 17, 8185-95	20.1	26
1381	Autonomous splicing and complementation of in vivo-assembled spliceosomes. 1989 , 108, 765-77		61
1380	The deletion of the distal CCAAT box region of the A gamma-globin gene in black HPFH abolishes the binding of the erythroid specific protein NFE3 and of the CCAAT displacement protein. <i>Nucleic Acids Research</i> , 1989 , 17, 6681-91	20.1	64
1379	Regulation of mouse CYP1A1 gene expression by dioxin: requirement of two cis-acting elements during induction. 1989 , 9, 2378-86		141
1378	AP-1/jun binding sites mediate serum inducibility of the human vimentin promoter. <i>Nucleic Acids Research</i> , 1989 , 17, 1619-33	20.1	76
1377	Functional analysis of GC element binding and transcription in the hamster dihydrofolate reductase gene promoter. <i>Nucleic Acids Research</i> , 1989 , 17, 9291-304	20.1	121
1376	Interleukin 1 and cyclic AMP induce kappa immunoglobulin light-chain expression via activation of an NF-kappa B-like DNA-binding protein. 1989 , 9, 959-64		192
1375	The upstream muscle-specific enhancer of the rat muscle creatine kinase gene is composed of multiple elements. 1989 , 9, 2396-413		128
1374	Identification of a myocyte nuclear factor that binds to the muscle-specific enhancer of the mouse muscle creatine kinase gene. 1989 , 9, 2627-40		366

1373	Proximal rat prolactin promoter sequences direct optimal, pituitary cell-specific transcription. 1989 , 3, 559-66		29
1372	Regulation of beta-glucuronidase expression in transgenic tobacco plants by an A/T-rich, cis-acting sequence found upstream of a French bean beta-phaseolin gene. 1989 , 1, 839-53		196
1371	Sequence-specific functions of the early palindrome domain within the SV40 core origin of replication. <i>Nucleic Acids Research</i> , 1989 , 17, 9279-9289	20.1	3
1370	Identification of thyrotroph-specific factors and cis-acting sequences of the murine thyrotropin beta subunit gene. 1989 , 3, 1037-45		23
1369	Regulation of b-Glucuronidase Expression in Transgenic Tobacco Plants by an A/T-Rich, cis-Acting Sequence Found Upstream of a French Bean b-Phaseolin Gene. 1989 , 1, 839		101
1368	Binding of thyroid hormone receptors to the rat thyrotropin-beta gene. 1989 , 3, 1359-68		64
1367	Transcription factor E2F is required for efficient expression of the hamster dihydrofolate reductase gene in vitro and in vivo. 1989 , 9, 4994-5002		335
1366	A nuclease-hypersensitive element of the human c-myc promoter interacts with a transcription initiation factor. 1989 , 9, 5123-33		145
1365	Two phases in the addition of a poly(A) tail. 1989 , 3, 1401-12		81
1364	Binding sites of HeLa cell nuclear proteins on the upstream region of adenovirus type 5 E1A gene. <i>Nucleic Acids Research</i> , 1989 , 17, 10015-34	20.1	23
1363	Accurate and efficient RNA polymerase III transcription in a cell-free extract prepared from <i>Ascaris suum</i> embryos. 1989 , 35, 277-83		7
1362	Inhibitory effects of vesicular stomatitis virus on cellular and influenza viral RNA metabolism and protein synthesis. 1989 , 172, 274-84		6
1361	The minimal transforming fragment of HSV-2 mtrIII can function as a complex promoter element. 1989 , 169, 346-53		22
1360	Nuclear proteins binding to an enhancer element of the major histocompatibility class I promoter: differences between highly oncogenic and nononcogenic adenovirus-transformed rat cells. 1989 , 172, 643-6		28
1359	Identification of DNA-protein interactions and enhancer activity at the 5' end of the upstream regulatory region in human papillomavirus type 11. 1989 , 170, 123-30		13
1358	Requirement of DNA topoisomerases for in vitro chromatin assembly by 3T6 mouse cell extracts. 1989 , 181, 531-7		5
1357	Heat-shock protein 83 of <i>Leishmania mexicana amazonensis</i> is an abundant cytoplasmic protein with a tandemly repeated genomic arrangement. 1989 , 185, 231-6		35
1356	Molecular control of tissue-specific expression at the mouse TNF locus. 1989 , 19, 549-52		30

1355	Abolishment of c-fos inducibility in ras-transformed mouse osteoblast cell lines. 1989 , 2, 208-16	15
1354	South western blot mapping: a procedure for simultaneous characterization of DNA binding proteins and their specific genomic DNA target sites. 1989 , 179, 299-303	17
1353	Demonstration of an amino terminal La epitope recognized by human anti-La sera. 1989 , 22, 65-71	23
1352	Binding of cellular protein(s) to U3 region of human T-cell leukemia virus type-I long terminal repeat. 1989 , 2, 307-11	
1351	5'-flanking DNA of the human insulin receptor gene and long terminal repeat of mouse mammary tumour virus bind to the same nuclear protein(s). 1989 , 32, 877-80	2
1350	Functional cooperativity between protein molecules bound at two distinct sequence elements of the immunoglobulin heavy-chain promoter. 1989 , 337, 573-6	129
1349	In vitro correction of G.T mispairs to G.C pairs in nuclear extracts from human cells. 1989 , 339, 234-6	190
1348	Occupation of the c-fos serum response element in vivo by a multi-protein complex is unaltered by growth factor induction. 1989 , 340, 68-70	364
1347	COUP transcription factor is a member of the steroid receptor superfamily. 1989 , 340, 163-6	444
1346	Muscle-specific gene expression controlled by a regulatory element lacking a MyoD1-binding site. 1989 , 341, 716-20	47
1345	Conservation between yeast and man of a protein associated with U5 small nuclear ribonucleoprotein. 1989 , 342, 819-21	107
1344	Five intermediate complexes in transcription initiation by RNA polymerase II. 1989 , 56, 549-61	987
1343	Role of the octamer motif in hybrid cell extinction of immunoglobulin gene expression: extinction is dominant in a two enhancer system. 1989 , 58, 441-8	91
1342	Antisense probing of the human U4/U6 snRNP with biotinylated 2'-OMe RNA oligonucleotides. 1989 , 59, 531-9	166
1341	Congenital immunodeficiencies associated with absence of HLA class II antigens on lymphocytes result from distinct mutations in trans-acting factors. 1989 , 26, 288-309	79
1340	Proteins binding to the liver-specific pyruvate kinase gene promoter. A unique combination of known factors. 1989 , 209, 205-19	147
1339	The 4F2 heavy chain gene: a molecular model of inducible gene expression in human T cells. 1989 , 2 Suppl, 67-79	3
1338	Ontogeny of the opiate phenotype: an approach to defining transsynaptic mechanisms at the molecular level in the rat adrenal medulla. 1989 , 7, 499-511	18

1337	Cloning and analysis of a human 86-kDa heat-shock-protein-encoding gene. 1989 , 83, 105-15	6
1336	Structure, chromosome mapping and regulation of the mouse zinc-finger gene Krox-24; evidence for a common regulatory pathway for immediate-early serum-response genes. 1989 , 80, 325-36	62
1335	Preproenkephalin DNA-binding proteins in the rat: 5' flanking region. 1989 , 5, 131-40	19
1334	Nuclear factor I can functionally replace transcription factor Sp1 in a U2 small nuclear RNA gene enhancer. 1989 , 205, 387-96	25
1333	Epstein-Barr virus small RNA (EBER) genes: unique transcription units that combine RNA polymerase II and III promoter elements. 1989 , 57, 825-34	153
1332	Probing the structure and function of U2 snRNP with antisense oligonucleotides made of 2'-OMe RNA. 1989 , 58, 383-90	121
1331	Complementation of in vitro-assembled spliceosomes. 1989 , 209, 379-92	2
1330	Sequence-specific methylation in a downstream region of the late E2A promoter of adenovirus type 2 DNA prevents protein binding. 1989 , 210, 411-5	40
1329	A 7 bp mutation converts a human RNA polymerase II snRNA promoter into an RNA polymerase III promoter. 1989 , 58, 55-67	226
1328	The human telomere terminal transferase enzyme is a ribonucleoprotein that synthesizes TTAGGG repeats. 1989 , 59, 521-9	1331
1327	A recombination hotspot in the LTR of a mouse retrotransposon identified in an in vitro system. 1989 , 57, 937-46	68
1326	Biochemical analysis of transcriptional activation by Jun: differential activity of c- and v-Jun. 1989 , 59, 709-17	210
1325	Erythroid-specific activation and derepression of the chick beta-globin promoter in vitro. 1989 , 57, 1189-200	80
1324	An in vitro system for the editing of apolipoprotein B mRNA. 1989 , 58, 519-25	162
1323	Protein-DNA interactions in the 5' region of the mouse alcohol dehydrogenase gene Adh-1. 1989 , 78, 277-85	25
1322	A brain-specific transcription activator. 1989 , 3, 563-72	139
1321	The liver-specific transcription factor LF-B1 contains a highly diverged homeobox DNA binding domain. 1989 , 59, 145-57	489
1320	An improved nuclear extract preparation method. 1989 , 6, 101-9	29

1319	Definition of T-cell specific DNA-binding factors that interact with a 3'-silencer in the CD4+ T-cell gene Rpt-1. 1989 , 85, 461-9	2
1318	Mutually exclusive splicing of alpha-tropomyosin exons enforced by an unusual lariat branch point location: implications for constitutive splicing. 1989 , 56, 749-58	221
1317	The ability of a ternary complex to form over the serum response element correlates with serum inducibility of the human c-fos promoter. 1989 , 56, 563-72	493
1316	The protein CDP, but not CP1, footprints on the CCAAT region of the gamma-globin gene in unfractionated B-cell extracts. 1989 , 1007, 237-42	48
1315	Initiation of transcription by RNA polymerase II. 1989 , 1009, 1-10	56
1314	Down-regulation of histone H3 and H4 gene transcription in differentiated L6 myotubes. 1989 , 1009, 177-83	10
1313	Cloning and in vitro transcription of the Sarcophaga lectin gene. 1989 , 1009, 244-50	12
1312	Transcription of rat aldolase B gene: minor RNAs are transcribed by RNA polymerase III. 1989 , 1007, 91-8	4
1311	Stage-specific detection of a DNA-binding protein for the storage protein gene of Sarcophaga peregrina. 1989 , 1008, 79-84	4
1310	Identification of an inhibitor of transcription in extracts prepared from wheat shoot nuclei. 1989 , 64, 91-98	3
1309	Enhancer and promoter element interactions dictate cyclic adenosine monophosphate mediated and cell-specific expression of the glycoprotein hormone alpha-gene. 1989 , 3, 763-72	76
1308	Pre-mRNA with a trimethylguanosine cap structure can be spliced efficiently in vitro. 1989 , 162, 977-83	3
1307	Tissue-specific expression of human terminal deoxynucleotidyl transferase is regulated at the transcriptional level. 1989 , 164, 750-7	10
1306	The same nuclear proteins bind the proximal CACCC box of the human beta-globin promoter and a similar sequence in the enhancer. 1989 , 164, 149-55	22
1305	Nuclear factor I represses the reverse-oriented transcription from the adenovirus type 5 DNA terminus. 1989 , 164, 1212-9	15
1304	Interferon-induced early changes in nuclear protein interactions with the interferon consensus sequence. 1989 , 163, 370-7	2
1303	RNase H cleavage of RNA hybridized to oligonucleotides containing methylphosphonate, phosphorothioate and phosphodiester bonds. <i>Nucleic Acids Research</i> , 1989 , 17, 9193-204	20.1 148
1302	Ligand-dependent interaction of the dioxin receptor with target DNA. 1989 , 34, 375-7	5

1301	Immunological evidence for the role of phosphoprotein p68/pi = 7.3 in premessenger RNA splicing. 1989 , 248, 101-4	6
1300	Site-selective 8-Cl-cAMP which causes growth inhibition and differentiation increases DNA (CRE)-binding activity in cancer cells. 1989 , 254, 83-8	12
1299	In vitro transcription of the mouse whey acidic protein promoter is affected by upstream sequences. 1989 , 251, 173-6	8
1298	Constitutive in vitro binding of nuclear proteins to the 5'-flanking region of 6-16, a human gene inducible by alpha, beta-interferons. 1989 , 258, 323-30	6
1297	RNA polymerase II-directed gene transcription by rat skeletal muscle nuclear extracts. 1989 , 185, 8-20	35
1296	Evidence that a nuclear matrix protein participates in premessenger RNA splicing. 1989 , 182, 521-33	43
1295	The "initiator" as a transcription control element. 1989 , 57, 103-13	1496
1294	An enhancer stimulates transcription in trans when attached to the promoter via a protein bridge. 1989 , 58, 767-77	180
1293	Analysis of splicing complexes and small nuclear ribonucleoprotein particles by native gel electrophoresis. 1989 , 180, 442-53	54
1292	Rapid detection of octamer binding proteins with 'mini-extracts', prepared from a small number of cells. <i>Nucleic Acids Research</i> , 1989 , 17, 6419	20.1 3646
1291	Genes, variant genes and pseudogenes of the human tRNA(Val) gene family. Expression and pre-tRNA maturation in vitro. 1989 , 209, 505-23	36
1290	Functional analysis of point mutations in the AAUAAA motif of the SV40 late polyadenylation signal. <i>Nucleic Acids Research</i> , 1989 , 17, 3899-908	20.1 75
1289	Mouse RNAase MRP RNA is encoded by a nuclear gene and contains a decamer sequence complementary to a conserved region of mitochondrial RNA substrate. 1989 , 56, 131-9	195
1288	The involvement of NF-kappa B in beta-interferon gene regulation reveals its role as widely inducible mediator of signal transduction. 1989 , 57, 287-94	484
1287	Characterization of the spliced leader genes and transcripts in <i>Trypanosoma cruzi</i> . 1989 , 82, 177-89	49
1286	Determination of RNA-protein and RNA-ribonucleoprotein interactions by nuclease probing. 1989 , 180, 454-68	11
1285	Identification of a zinc finger protein that binds to the sterol regulatory element. 1989 , 245, 640-3	231
1284	Identification and purification of a <i>Saccharomyces cerevisiae</i> protein with the DNA binding specificity of mammalian activating transcription factor. 1989 , 86, 109-13	46

1283	Human immunodeficiency virus 1 tat protein binds trans-activation-responsive region (TAR) RNA in vitro. 1989 , 86, 6925-9	443
1282	Differential binding of nuclear factors to the intron 1 sequences containing the transcriptional pause site correlates with c-myb expression. 1989 , 86, 7326-30	51
1281	T-cell nuclei contain a protein that binds upstream of the murine granulocyte-macrophage colony-stimulating factor gene. 1989 , 86, 7392-6	4
1280	Ribozyme, antisense RNA, and antisense DNA inhibition of U7 small nuclear ribonucleoprotein-mediated histone pre-mRNA processing in vitro. 1989 , 9, 4479-87	74
1279	Functional serum response elements upstream of the growth factor-inducible gene zif268. 1989 , 9, 4889-95	181
1278	Evidence that xeroderma pigmentosum cells from complementation group E are deficient in a homolog of yeast photolyase. 1989 , 9, 5105-12	57
1277	Multiple splicing factors are released from endogenous complexes during in vitro pre-mRNA splicing. 1989 , 9, 5273-80	10
1276	Transcription factor MBF-I interacts with metal regulatory elements of higher eucaryotic metallothionein genes. 1989 , 9, 5315-23	55
1275	A B-cell-specific nuclear protein that binds to DNA sites 5' to immunoglobulin S alpha tandem repeats is regulated during differentiation. 1989 , 9, 5594-601	61
1274	The sarcomeric actin CArG-binding factor is indistinguishable from the c-fos serum response factor. 1989 , 9, 515-22	187
1273	Octamer transcription factors 1 and 2 each bind to two different functional elements in the immunoglobulin heavy-chain promoter. 1989 , 9, 747-56	98
1272	Forskolin inducibility and tissue-specific expression of the fibronectin promoter. 1989 , 9, 1498-506	88
1271	Purification and characterization of multiple nuclear factors that bind to the TAX-inducible enhancer within the human T-cell leukemia virus type 1 long terminal repeat. 1989 , 9, 1733-45	88
1270	Interaction of nuclear proteins with a positive cis-acting element of rat embryonic myosin heavy-chain promoter: identification of a new transcriptional factor. 1989 , 9, 1839-49	32
1269	Remarkable intron and exon sequence conservation in human and mouse homeobox Hox 1.3 genes. 1989 , 9, 2273-8	31
1268	Identification of a cell-specific DNA-binding activity that interacts with a transcriptional activator of genes expressed in the acinar pancreas. 1989 , 9, 2464-76	135
1267	Regulation of the visna virus long terminal repeat in macrophages involves cellular factors that bind sequences containing AP-1 sites. 1989 , 9, 2728-33	68
1266	Identification of multiple proteins that interact with functional regions of the human cardiac alpha-actin promoter. 1989 , 9, 3269-83	112

1265	Novel regulation of transcription initiation of the peptide IX gene of adenovirus 2. 1989 , 9, 4265-71		6
1264	A multiple cytokine- and second messenger-responsive element in the enhancer of the human interleukin-6 gene: similarities with c-fos gene regulation. 1989 , 9, 5537-47		175
1263	The C-terminal domain of the largest subunit of RNA polymerase II and transcription initiation. 1989 , 9, 5750-3		26
1262	Factor interactions with the simian virus 40 early pre-mRNA influence branch site selection and alternative splicing. 1989 , 9, 2007-17		10
1261	U5 small nuclear ribonucleoprotein: RNA structure analysis and ATP-dependent interaction with U4/U6. 1989 , 9, 3350-9		138
1260	Specific protein-DNA interactions at a xenobiotic-responsive element: copurification of dioxin receptor and DNA-binding activity. 1989 , 86, 60-4		112
1259	The octamer motif is a B-lymphocyte-specific regulatory element of the HLA-DR alpha gene promoter. 1989 , 86, 6739-43		41
1258	UACUAAC is the preferred branch site for mammalian mRNA splicing. 1989 , 86, 2752-6		166
1257	A characterization of the elements comprising the promoter of the mouse ribosomal protein gene RPS16. <i>Nucleic Acids Research</i> , 1989 , 17, 5323-37	20.1	55
1256	Infection with the intracellular protozoan parasite <i>Theileria parva</i> induces constitutively high levels of NF-kappa B in bovine T lymphocytes. 1989 , 9, 4677-86		81
1255	A block in mammalian splicing occurring after formation of large complexes containing U1, U2, U4, U5, and U6 small nuclear ribonucleoproteins. 1989 , 9, 259-67		31
1254	A new myocyte-specific enhancer-binding factor that recognizes a conserved element associated with multiple muscle-specific genes. 1989 , 9, 5022-33		577
1253	New methods to investigate ATP requirement for pre-mRNA splicing: inhibition by hexokinase/glucose or an ATP-binding site blocker. 1989 , 106, 372-5		8
1252	Assembly of transcriptionally inactive chromatin in vitro. 1989 , 106, 29-33		3
1251	Effects of anti-IgM suppression on polyclonally activated murine B cells: analysis of immunoglobulin mRNA, gene specific nuclear factors and cell cycle distribution. <i>Nucleic Acids Research</i> , 1989 , 17, 10455-72	20.1	11
1250	Poly(A) polymerase purified from HeLa cell nuclear extract is required for both cleavage and polyadenylation of pre-mRNA in vitro. 1989 , 9, 193-203		99
1249	Transcriptional promiscuity of the human alpha-globin gene. 1989 , 9, 241-51		40
1248	An AP-2 element acts synergistically with the cyclic AMP- and phorbol ester-inducible enhancer of the human proenkephalin gene. 1989 , 9, 321-4		174

1247	Indirect binding of human T-cell leukemia virus type I tax1 to a responsive element in the viral long terminal repeat. 1989 , 9, 4152-60	92
1246	Identification of a new promoter upstream of the murine dihydrofolate reductase gene. 1989 , 9, 4568-70	20
1245	Dual functions of a cis-acting element within the rat prolactin gene promoter. 1989 , 9, 817-9	12
1244	Regulation of 2',5'-oligoadenylate synthetase gene expression by interferons and platelet-derived growth factor. 1989 , 9, 1060-8	33
1243	Sea urchin early and late H4 histone genes bind a specific transcription factor in a stable preinitiation complex. 1989 , 9, 1476-87	12
1242	Cellular factors that interact with the hepatitis B virus enhancer. 1989 , 9, 1804-9	101
1241	In vitro activation and nuclear translocation of NF-kappa B catalyzed by cyclic AMP-dependent protein kinase and protein kinase C. 1989 , 9, 2424-30	416
1240	Unusual levels of heat shock element-binding activity in embryonal carcinoma cells. 1989 , 9, 3888-96	47
1239	Negative transcriptional regulatory element that functions in embryonal carcinoma cells. 1989 , 9, 4032-7	11
1238	The adenovirus-inducible factor E2F stimulates transcription after specific DNA binding. 1989 , 9, 578-85	145
1237	Structure, chromosome location, and expression of the mouse zinc finger gene Krox-20: multiple gene products and coregulation with the proto-oncogene c-fos. 1989 , 9, 787-97	160
1236	NF-kappa B binds within a region required for B-cell-specific expression of major histocompatibility complex class II gene E alpha d. 1989 , 9, 844-6	42
1235	Cell-specific activity of the modulator region in the human cytomegalovirus major immediate-early gene. 1989 , 9, 1342-5	59
1234	Role of poly(A) polymerase in the cleavage and polyadenylation of mRNA precursor. 1989 , 9, 1435-44	42
1233	In vivo evidence of interaction between interferon-stimulated gene factors and the interferon-stimulated response element. 1989 , 9, 3533-7	20
1232	Alteration of a single nucleotide allows efficient binding of H2TF1/KBF1 to the immunoglobulin kappa enhancer B motif. 1989 , 9, 3548-52	29
1231	Conserved terminal hairpin sequences of histone mRNA precursors are not involved in duplex formation with the U7 RNA but act as a target site for a distinct processing factor. 1989 , 86, 4345-9	81
1230	Hydroxyl radical "footprinting" of RNA: application to pre-mRNA splicing complexes. 1989 , 86, 7795-9	31

1229	Identification of cellular proteins that bind to the human immunodeficiency virus type 1 trans-activation-responsive TAR element RNA. 1989 , 86, 7828-32	121
1228	A lipopolysaccharide-induced DNA-binding protein for a class II gene in B cells is distinct from NF-kappa B. 1989 , 9, 3184-92	27
1227	RNA polymerase II is capable of pausing and prematurely terminating transcription at a precise location in vivo and in vitro. 1989 , 86, 12-6	83
1226	Altered binding of human histone gene transcription factors during the shutdown of proliferation and onset of differentiation in HL-60 cells. 1989 , 86, 1865-9	54
1225	DNA bending is induced by a transcription factor that interacts with the human c-FOS and alpha-actin promoters. 1989 , 86, 2162-6	72
1224	Tissue-specific transcription of the mouse alpha-fetoprotein gene promoter is dependent on HNF-1. 1989 , 9, 4204-12	134
1223	Multiple forms of poly(A) polymerases purified from HeLa cells function in specific mRNA 3'-end formation. 1989 , 9, 4229-38	66
1222	Human transcription factor TFIIC2 specifically interacts with a unique sequence in the <i>Xenopus laevis</i> 5S rRNA gene. 1989 , 9, 4941-50	28
1221	Expression of the glycoprotein hormone alpha-subunit gene in the placenta requires a functional cyclic AMP response element, whereas a different cis-acting element mediates pituitary-specific expression. 1989 , 9, 5113-22	133
1220	E1A-induced enhancer activity of the poly(dG-dT).poly(dA-dC) element (GT element) and interactions with a GT-specific nuclear factor. 1989 , 9, 5248-53	33
1219	Identification of cis-acting sequences responsible for phorbol ester induction of human serum amyloid A gene expression via a nuclear factor kappaB-like transcription factor. 1989 , 9, 1908-16	140
1218	Localization of transcriptional regulatory elements and nuclear factor binding sites in mouse ribosomal protein gene rpl32. 1989 , 9, 2067-74	93
1217	Cross-binding of factors to functionally different promoter elements in c-fos and skeletal actin genes. 1989 , 9, 2191-201	147
1216	Purification of a yeast centromere-binding protein that is able to distinguish single base-pair mutations in its recognition site. 1989 , 9, 2544-50	88
1215	ras-induced neuronal differentiation of PC12 cells: possible involvement of fos and jun. 1989 , 9, 3174-83	141
1214	Identification of HeLa cell nuclear factors that bind to and activate the early promoter of human polyomavirus BK in vitro. 1989 , 9, 3821-8	30
1213	A cytosolic protein binds to structural elements within the iron regulatory region of the transferrin receptor mRNA. 1989 , 86, 3574-8	215
1212	E1A-dependent trans-activation of the human MYC promoter is mediated by the E2F factor. 1989 , 86, 3594-8	275

1211	Identification, purification, and biochemical characterization of U2 small nuclear ribonucleoprotein auxiliary factor. 1989 , 86, 9243-7	339
1210	Estrogen receptor selectively binds the "coding strand" of an estrogen responsive element. 1989 , 86, 863-7	89
1209	Tumor necrosis factor alpha and interleukin 1 stimulate the human immunodeficiency virus enhancer by activation of the nuclear factor kappa B. 1989 , 86, 2336-40	1466
1208	Cloning of the major histocompatibility complex class II promoter binding protein affected in a hereditary defect in class II gene regulation. 1989 , 86, 4200-4	118
1207	Purification of a yeast TATA box-binding protein that exhibits human transcription factor IID activity. 1989 , 86, 4843-7	111
1206	Specific binding of a HeLa cell nuclear protein to RNA sequences in the human immunodeficiency virus transactivating region. 1989 , 86, 4858-62	121
1205	Dysfunction of chromosomal loop attachment sites: illegitimate recombination linked to matrix association regions and topoisomerase II. 1989 , 86, 5497-501	189
1204	20S small nuclear ribonucleoprotein U5 shows a surprisingly complex protein composition. 1989 , 86, 6038-42	131
1203	A T-cell-specific transcriptional enhancer element 3' of C alpha in the human T-cell receptor alpha locus. 1989 , 86, 6714-8	148
1202	The mammalian analogue of the yeast PRP8 splicing protein is present in the U4/5/6 small nuclear ribonucleoprotein particle and the spliceosome. 1989 , 86, 8742-6	77
1201	Positive and negative regulatory elements in the mouse albumin enhancer. 1989 , 86, 1553-7	109
1200	In vitro transcription directed from the somatostatin promoter is dependent upon a purified 43-kDa DNA-binding protein. 1989 , 86, 2181-5	42
1199	New B-lymphocyte-specific enhancer-binding protein. 1989 , 9, 312-20	35
1198	E2F from adenovirus-infected cells binds cooperatively to DNA containing two properly oriented and spaced recognition sites. 1989 , 9, 4495-506	59
1197	A gamma interferon-unresponsive variant of cell line 70Z/3, IFN-4, can be partially rescued by phorbol myristate acetate. 1989 , 9, 5231-3	14
1196	Functional identification of regulatory elements within the promoter region of platelet-derived growth factor 2. 1989 , 9, 396-405	61
1195	Major histocompatibility complex class I genes in murine fibrosarcoma IC9 are down regulated at the level of the chromatin structure. 1989 , 9, 3136-42	23
1194	Isolation and characterization of branched oligonucleotides from RNA. 1989 , 180, 177-91	6

1193	Modification interference analysis of reactions using RNA substrates. 1989 , 180, 369-79	33
1192	Immunoprecipitation of ribonucleoproteins using autoantibodies. 1989 , 180, 468-81	67
1191	A 65-kappaD subunit of active NF-kappaB is required for inhibition of NF-kappaB by I kappaB. 1989 , 3, 1689-98	470
1190	A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family.. 1990 , 9, 1897-1906	1108
1189	????????HBV DNA?????????. 1990 , 31, 1137-1137	
1188	Binding of a liver-specific factor to the human albumin gene promoter and enhancer. 1990 , 10, 991-9	34
1187	Characterization of the rat transforming growth factor alpha gene and identification of promoter sequences. 1990 , 10, 2111-21	101
1186	Myosin light-chain 1/3 gene alternative splicing: cis regulation is based upon a hierarchical compatibility between splice sites. 1990 , 10, 2133-44	37
1185	A novel tumor necrosis factor-responsive transcription factor which recognizes a regulatory element in hemopoietic growth factor genes. 1990 , 10, 2950-9	61
1184	Brain and muscle creatine kinase genes contain common TA-rich recognition protein-binding regulatory elements. 1990 , 10, 4826-36	92
1183	A cellular factor binds to the herpes simplex virus type 1 transactivator Vmw65 and is required for Vmw65-dependent protein-DNA complex assembly with Oct-1. 1990 , 10, 4974-7	107
1182	Identification and characterization of a HeLa nuclear protein that specifically binds to the trans-activation-response (TAR) element of human immunodeficiency virus. 1990 , 87, 3624-8	205
1181	Wound-inducible nuclear protein binds DNA fragments that regulate a proteinase inhibitor II gene from potato. 1990 , 87, 603-7	76
1180	Transcriptional activation and repression by Fos are independent functions: the C terminus represses immediate-early gene expression via CArG elements. 1990 , 10, 4243-55	203
1179	Selection and analysis of galactose metabolic pathway variants of a mouse liver cell line. 1990 , 10, 4582-9	8
1178	A naturally occurring gamma globin gene mutation enhances SP1 binding activity. 1990 , 10, 95-102	47
1177	A transcriptionally active form of TFIIC is modified in poliovirus-infected HeLa cells. 1990 , 10, 5106-13	58
1176	DNA-binding protein activated by gamma radiation in human cells. 1990 , 10, 5279-85	56

1175	Inhibition of c-fos transcription and phosphorylation of the serum response factor by an inhibitor of phospholipase C-type reactions. 1990 , 10, 5558-61	32
1174	Activation of octamer-containing promoters by either octamer-binding transcription factor 1 (OTF-1) or OTF-2 and requirement of an additional B-cell-specific component for optimal transcription of immunoglobulin promoters. 1990 , 10, 6204-15	127
1173	Characterization of a thyroid-specific enhancer located 5.5 kilobase pairs upstream of the human thyroid peroxidase gene. 1990 , 10, 6216-24	69
1172	Sequence elements required for activity of a murine major histocompatibility complex class II promoter bind common and cell-type-specific nuclear factors. 1990 , 10, 593-604	72
1171	Mutational analysis of the DRA promoter: cis-acting sequences and trans-acting factors. 1990 , 10, 711-9	107
1170	NF-kappa B as inducible transcriptional activator of the granulocyte-macrophage colony-stimulating factor gene. 1990 , 10, 1281-6	182
1169	Sequences downstream of the transcription initiation site modulate the activity of the murine dihydrofolate reductase promoter. 1990 , 10, 1390-8	95
1168	Purification and cloning of interferon-stimulated gene factor 2 (ISGF2): ISGF2 (IRF-1) can bind to the promoters of both beta interferon- and interferon-stimulated genes but is not a primary transcriptional activator of either. 1990 , 10, 2448-57	319
1167	Constitutive and interleukin-1 (IL-1)-inducible factors interact with the IL-1-responsive element in the IL-6 gene. 1990 , 10, 2757-64	269
1166	Developmental regulation of topoisomerase II sites and DNase I-hypersensitive sites in the chicken beta-globin locus. 1990 , 10, 2774-86	113
1165	Complex regulation of the immunoglobulin mu heavy-chain gene enhancer: microB, a new determinant of enhancer function. 1990 , 10, 3145-54	71
1164	Synthesis and processing of small B2 transcripts in mouse embryonal carcinoma cells. 1990 , 10, 4058-67	31
1163	Tumor necrosis factor alpha regulates expression of the major histocompatibility complex class II-associated invariant chain by binding of an NF-kappa B-like factor to a promoter element. 1990 , 10, 4146-54	71
1162	Factors involved in specific transcription by mammalian RNA polymerase II: identification of general transcription factor TFIIG. 1990 , 87, 9158-62	83
1161	Identification of a DNA sequence responsible for binding of the 1,25-dihydroxyvitamin D3 receptor and 1,25-dihydroxyvitamin D3 enhancement of mouse secreted phosphoprotein 1 (SPP-1 or osteopontin) gene expression. 1990 , 87, 9995-9	399
1160	The promoter of the human interleukin-2 gene contains two octamer-binding sites and is partially activated by the expression of Oct-2. 1990 , 10, 5464-72	96
1159	Analysis of the upstream elements of the xenobiotic compound-inducible and positionally regulated glutathione S-transferase Ya gene. 1990 , 10, 1841-52	114
1158	Regulation of the mouse alpha A-crystallin gene: isolation of a cDNA encoding a protein that binds to a cis sequence motif shared with the major histocompatibility complex class I gene and other genes. 1990 , 10, 3700-8	113

1157	Interleukin-6 induction by tumor necrosis factor and interleukin-1 in human fibroblasts involves activation of a nuclear factor binding to a kappa B-like sequence. 1990 , 10, 3818-23	265
1156	The mouse albumin enhancer contains a negative regulatory element that interacts with a novel DNA-binding protein. 1990 , 10, 3896-905	54
1155	A transcription factor interacting with the class I gene enhancer is inactive in tumorigenic cell lines which suppress major histocompatibility complex class I genes. 1990 , 10, 4100-9	41
1154	Regulation of the human T-cell receptor alpha gene enhancer: multiple ubiquitous and T-cell-specific nuclear proteins interact with four hypomethylated enhancer elements. 1990 , 10, 4720-7	53
1153	Gene induction by interferons: functional complementation between trans-acting factors induced by alpha interferon and gamma interferon. 1990 , 10, 5055-63	70
1152	Analysis of the mouse Dhfr/Rep-3 major promoter region by using linker-scanning and internal deletion mutations and DNase I footprinting. 1990 , 10, 6003-12	25
1151	A uridylate tract mediates efficient heterogeneous nuclear ribonucleoprotein C protein-RNA cross-linking and functionally substitutes for the downstream element of the polyadenylation signal. 1990 , 10, 6397-407	113
1150	A cloned human CCAAT-box-binding factor stimulates transcription from the human hsp70 promoter. 1990 , 10, 6709-17	69
1149	Transcriptional and posttranscriptional regulation of the rat prolactin gene by calcium. 1990 , 10, 442-8	28
1148	In vitro analysis of promoter elements regulating transcription of the phosphoenolpyruvate carboxykinase (GTP) gene. 1990 , 10, 480-5	39
1147	Involvement of a NF-kappa B-like transcription factor in the activation of the interleukin-6 gene by inflammatory lymphokines. 1990 , 10, 561-8	353
1146	An inducible 50-kilodalton NF kappa B-like protein and a constitutive protein both bind the acute-phase response element of the angiotensinogen gene. 1990 , 10, 1023-32	121
1145	Two DNA-binding proteins discriminate between the promoters of different members of the major histocompatibility complex class II multigene family. 1990 , 10, 965-71	66
1144	Initiation and processing of two kappa immunoglobulin germ line transcripts in mouse B cells. 1990 , 10, 1950-8	56
1143	A negative element involved in vimentin gene expression. 1990 , 10, 2349-58	70
1142	Functional organization of the hepatitis B virus enhancer. 1990 , 10, 3683-9	70
1141	A nonerythroid GATA-binding protein is required for function of the human preproendothelin-1 promoter in endothelial cells. 1990 , 10, 4854-62	155
1140	Sequence requirements in different steps of the pre-mRNA splicing reaction: analysis by the RNA modification-exclusion technique. 1990 , 10, 4942-7	10

1139	Multiple liver-specific factors bind to the hepatitis B virus core/pregenomic promoter: trans-activation and repression by CCAAT/enhancer binding protein. 1990 , 87, 5069-73	133
1138	NF-kappa B: a family of inducible and differentially expressed enhancer-binding proteins in human T cells. 1990 , 87, 10028-32	244
1137	Molecular mechanisms in down-regulation of tumor necrosis factor expression. 1990 , 87, 9563-7	99
1136	Vitamin D-mediated modifications in protein-DNA interactions at two promoter elements of the osteocalcin gene. 1990 , 87, 1701-5	133
1135	Cisplatin-resistant cells express increased levels of a factor that recognizes damaged DNA. 1990 , 87, 3324-7	110
1134	Identification of a mammalian nuclear factor and human cDNA-encoded proteins that recognize DNA containing apurinic sites. 1990 , 87, 3396-400	42
1133	Cell type-specific protein-DNA interactions in the human zeta-globin upstream promoter region: displacement of Sp1 by the erythroid cell-specific factor NF-E1. 1990 , 10, 282-94	45
1132	Interleukin-1 synergy with phosphoinositide pathway agonists for induction of interleukin-2 gene expression: molecular basis of costimulation. 1990 , 10, 6325-34	47
1131	The ubiquitous octamer-binding protein(s) is sufficient for transcription of immunoglobulin genes. 1990 , 10, 982-90	51
1130	Transcriptional regulation of interleukin 3 gene expression in T lymphocytes. 1990 , 87, 9650-4	78
1129	Coordinate occupancy of AP-1 sites in the vitamin D-responsive and CCAAT box elements by Fos-Jun in the osteocalcin gene: model for phenotype suppression of transcription. 1990 , 87, 9990-4	173
1128	Transcriptional stimulation of the delta 1-crystallin gene by insulin-like growth factor I and insulin requires DNA cis elements in chicken. 1990 , 87, 3353-7	40
1127	Lipopolysaccharide-unresponsive mutant pre-B-cell lines blocked in NF-kappa B activation. 1990 , 10, 422-5	12
1126	Fos and jun cooperate in transcriptional regulation via heterologous activation domains. 1990 , 10, 5532-5	79
1125	Increased expression and DNA-binding activity of transcription factor Sp1 in doxorubicin-resistant HL-60 leukemia cells. 1990 , 10, 5541-7	44
1124	Interaction of a nuclear protein with a palindromic sequence of the mouse immunoglobulin lambda 2-chain gene promoter is important for its transcription. 1990 , 10, 5894-902	2
1123	A 36-kilodalton cellular transcription factor mediates an indirect interaction of human T-cell leukemia/lymphoma virus type I TAX1 with a responsive element in the viral long terminal repeat. 1990 , 10, 4192-201	85
1122	Double-stranded RNA unwinding and modifying activity is detected ubiquitously in primary tissues and cell lines. 1990 , 10, 5586-90	97

1121	Positive and negative transcriptional elements of the human type IV collagenase gene. 1990 , 10, 6524-32	104
1120	Transcription initiation from the dihydrofolate reductase promoter is positioned by HIP1 binding at the initiation site. 1990 , 10, 653-61	254
1119	Octamer-binding proteins in diverse hemopoietic cells. 1990 , 10, 1293-6	33
1118	Capping of mammalian U6 small nuclear RNA in vitro is directed by a conserved stem-loop and AUAUAC sequence: conversion of a noncapped RNA into a capped RNA. 1990 , 10, 939-46	50
1117	A novel cis-acting DNA element required for a high level of inducible expression of the rat P-450c gene. 1990 , 10, 1470-5	98
1116	Detection of an immunoglobulin switch region-specific DNA-binding protein in mitogen-stimulated mouse splenic B cells. 1990 , 10, 1714-8	44
1115	Identification of a novel factor that interacts with an immunoglobulin heavy-chain promoter and stimulates transcription in conjunction with the lymphoid cell-specific factor OTF2. 1990 , 10, 2145-53	22
1114	Transcriptional potentiation of the vitellogenin B1 promoter by a combination of both nucleosome assembly and transcription factors: an in vitro dissection. 1990 , 10, 3926-33	31
1113	Positive and negative elements regulate human interleukin 3 expression. 1990 , 87, 5046-50	72
1112	Polyoma virus small tumor antigen pre-mRNA splicing requires cooperation between two 3' splice sites. 1990 , 87, 3338-42	23
1111	Multiple cis- and trans-acting elements involved in regulation of the neu gene. 1990 , 10, 6306-15	53
1110	Complex requirements for RNA polymerase III transcription of the Xenopus U6 promoter. <i>Nucleic Acids Research</i> , 1990 , 18, 5649-57	20.1 23
1109	Characterization of promoter elements of an interferon-inducible Ly-6E/A differentiation antigen, which is expressed on activated T cells and hematopoietic stem cells. 1990 , 10, 5150-9	41
1108	Surveying cis-acting sequences of pre-mRNA by adding antisense 2'-O-methyl oligoribonucleotides to a splicing reaction. 1990 , 108, 399-405	26
1107	Specific Binding of Nuclear Protein from Tobacco Hairy Roots Cultured in Vitro to a 5'-Upstream Region of the rolC Gene of the Ri Plasmid. 1990 ,	
1106	A T cell-specific transcriptional enhancer within the human T cell receptor delta locus. 1990 , 247, 1225-9	127
1105	Evidence for three distinct D proteins, which react differentially with anti-Sm autoantibodies, in the cores of the major snRNPs U1, U2, U4/U6 and U5. <i>Nucleic Acids Research</i> , 1990 , 18, 6475-84	20.1 110
1104	Differential expression of human placental growth-hormone variant and chorionic somatomammotropin in culture. 1990 , 267, 653-8	41

1103	Detection in non-erythroid cells of a factor with the binding characteristics of the erythroid cell transcription factor EF1. 1990 , 269, 543-5	2
1102	The role of the 5'-flanking sequence of a human tRNA(Glu) gene in modulation of its transcriptional activity in vitro. 1990 , 272, 797-803	6
1101	Glucocorticoid inhibition of transcription from adenovirus major late promoter. 1990 , 4, 1515-21	2
1100	Preparation and fractionation of mammalian extracts active in pre-mRNA splicing. 1990 , 181, 3-19	24
1099	Preparation of mammalian extracts active in polyadenylation. 1990 , 181, 49-74	15
1098	Histone-specific RNA 3' processing in nuclear extracts from mammalian cells. 1990 , 181, 74-89	14
1097	Nuclear cap binding protein from HeLa cells. 1990 , 181, 209-15	3
1096	Purification of small nuclear ribonucleoprotein particles with antibodies against modified nucleosides of small nuclear RNAs. 1990 , 181, 232-57	58
1095	Purification and characterization of U small nuclear ribonucleoproteins in cesium chloride gradients. 1990 , 181, 264-73	3
1094	Structural analysis of U small nuclear ribonucleoproteins by in vitro assembly. 1990 , 181, 273-84	6
1093	Site-specific degradation of RNA of small nuclear ribonucleoprotein particles with complementary oligodeoxynucleotides and RNase H. 1990 , 181, 284-92	5
1092	Immunoselective cell separation. 1990 , 184, 314-9	5
1091	Cell-specific expression of the mouse glial fibrillary acidic protein gene: identification of the cis- and trans-acting promoter elements for astrocyte-specific expression. 1990 , 55, 1180-8	131
1090	High levels of c-fos proto-oncogene expression in normal human adult skin. 1990 , 94, 418-22	31
1089	Progesterone receptor stimulates transcription of mouse mammary tumour virus in a cell-free system. 1990 , 344, 360-2	69
1088	Nucleolar transcription factor hUBF contains a DNA-binding motif with homology to HMG proteins. 1990 , 344, 830-6	645
1087	Disruption of a C/EBP binding site in the factor IX promoter is associated with haemophilia B. 1990 , 345, 444-6	143
1086	Major determinants of the specificity of interaction between small nuclear ribonucleoproteins U1A and U2B'' and their cognate RNAs. 1990 , 345, 502-6	299

1085	Functional analysis of cis-linked regulatory sequences in the HLA DRA promoter by transcription in vitro. 1990 , 36, 108-15	12
1084	A factor known to bind to endogenous Ig heavy chain enhancer only in lymphocytes is a ubiquitously active transcription factor. 1990 , 187, 507-13	4
1083	Interaction of upstream stimulatory factor with the human heme oxygenase gene promoter. 1990 , 188, 231-7	51
1082	Characterisation of tissue-specific trans-acting factor binding to a proximal element in the rat growth hormone gene promoter. 1990 , 191, 605-15	5
1081	The tissue-specific expression of the thyroglobulin gene requires interaction between thyroid-specific and ubiquitous factors. 1990 , 193, 311-8	71
1080	Identification of a recombinational signal sequence-specific DNA-binding protein(s) of Mr 115,000 in the nuclear extracts from immature lymphoid cell lines. 1990 , 17, 67-75	5
1079	The gene family encoding eggshell proteins of <i>Schistosoma japonicum</i> . 1990 , 42, 69-82	24
1078	Localization by mutational analysis of transcription factor binding sequences in the U3 region of Rous sarcoma virus LTR. 1990 , 176, 483-93	15
1077	Interactions between cellular regulatory proteins and a unique sequence region in the human cytomegalovirus major immediate-early promoter. 1990 , 174, 18-25	30
1076	Pseudorabies virus immediate-early gene overlaps with an oppositely oriented open reading frame: characterization of their promoter and enhancer regions. 1990 , 179, 365-77	51
1075	Antibodies specific for branched ribonucleic acids. 1990 , 185, 125-30	4
1074	Hepadnavirus enhancer and its binding proteins. 1990 , 25 Suppl 2, 11-9	8
1073	Anti-Sm-RNP activity in sera of patients with rheumatic and autoimmune diseases. 1990 , 9, 346-55	5
1072	Slow response variant of the B lymphoma 70Z/3 defective in LPS activation of NF-kappa B. 1990 , 31, 65-72	10
1071	1.3E2, a variant of the B lymphoma 70Z/3, defective in activation of NF-kappa B and OTF-2. 1990 , 31, 73-8	17
1070	In vitro transcription from cauliflower mosaic virus promoters by a cell-free extract from tobacco cells. 1990 , 14, 391-405	25
1069	Identification of a 55 Kdal nuclear protein sharing homology with hnRNP type L. 1990 , 14, 89-90	1
1068	The X boxes from promoters of HLA class II B genes at different loci do not compete for nuclear protein-specific binding. 1990 , 32, 117-28	16

1067	Synthesis of the herpes simplex virus type 1 trans-activator Vmw65 in insect cells using a baculovirus vector. 1990 , 94, 45-52	3
1066	Differential utilization of the SV40 enhancer and early promoter in various human lung carcinoma cell lines. 1990 , 45, 911-5	1
1065	Structural and functional analysis of the rat testis-specific histone H1t gene. 1990 , 44, 1-17	40
1064	Glial cells of the human developing brain and B cells of the immune system share a common DNA binding factor for recognition of the regulatory sequences of the human polyomavirus, JCV. 1990 , 27, 461-71	76
1063	Expression of chimeric human transferrin genes in vitro. 1990 , 27, 633-41	4
1062	Splicing factor SF4 is dispensable for the assembly of a functional splicing complex and participates in the subsequent steps of the splicing reaction.. 1990 , 9, 4119-4126	20
1061	Serum and v-src increase the level of a CCAAT-binding factor required for transcription from a retroviral long terminal repeat. 1990 , 4, 243-54	43
1060	A novel upstream element compensates for an ineffectual octamer motif in an immunoglobulin V kappa promoter.. 1990 , 9, 3109-3117	30
1059	A 3? transcriptional enhancer regulates tissue-specific expression of the human CD2 gene.. 1990 , 9, 3129-3136	50
1058	Lymphoid specific gene expression of the adenovirus early region 3 promoter is mediated by NF-kappa B binding motifs.. 1990 , 9, 4435-4442	36
1057	Expression of a human 7S K RNA gene in vivo requires a novel pol III upstream element.. 1990 , 9, 711-718	21
1056	Tax1 induction of the HTLV-I 21 bp enhancer requires cooperation between two cellular DNA-binding proteins.. 1990 , 9, 957-964	53
1055	Amino-terminal domain of NF1 binds to DNA as a dimer and activates adenovirus DNA replication.. 1990 , 9, 559-566	95
1054	A role for the adenovirus inducible E2F transcription factor in a proliferation dependent signal transduction pathway.. 1990 , 9, 2179-2184	170
1053	The specific DNA binding activity of the dioxin receptor is modulated by the 90 kd heat shock protein.. 1990 , 9, 69-76	167
1052	Protein recognition sites in polyomavirus enhancer: formation of a novel site for NF-1 factor in an enhancer mutant and characterization of a site in the enhancer D domain.. 1990 , 9, 947-955	18
1051	Synergistic interaction between interferon-alpha and interferon-gamma through induced synthesis of one subunit of the transcription factor ISGF3.. 1990 , 9, 1105-1111	154
1050	The beta-globin dominant control region: hypersensitive site 2.. 1990 , 9, 2159-2167	182

1049	Detailed analysis of the site 3 region of the human beta-globin dominant control region.. 1990 , 9, 2169-2177	200
1048	Tissue-specific nuclear factors mediate expression of the CD3 delta gene during T cell development.. 1990 , 9, 109-115	18
1047	Discrete domains of human U6 snRNA required for the assembly of U4/U6 snRNP and splicing complexes.. 1990 , 9, 251-255	29
1046	Transcription factor E4TF1 contains two subunits with different functions.. 1990 , 9, 841-847	56
1045	A mitigator sequence in the downstream region of the major late promoter of adenovirus type 12 DNA.. 1990 , 9, 1615-1623	30
1044	The presence of a DNA binding complex correlates with E beta class II MHC gene expression.. 1990 , 9, 1543-1549	14
1043	A growth-dependent transcription initiation factor (TIF-IA) interacting with RNA polymerase I regulates mouse ribosomal RNA synthesis.. 1990 , 9, 2857-2863	76
1042	Identification of sequences responsible for positive and negative regulation by E1A in the promoter of H-2Kbm1 class I MHC gene.. 1990 , 9, 127-135	38
1041	In vitro splicing of mutually exclusive exons from the chicken beta-tropomyosin gene: role of the branch point location and very long pyrimidine stretch.. 1990 , 9, 241-249	69
1040	Extinction of Ig genes expression in myeloma x fibroblast somatic cell hybrids is accompanied by repression of the oct-2 gene encoding a B-cell specific transcription factor.. 1990 , 9, 849-855	27
1039	In vitro analysis of a transcription termination site for RNA polymerase II. 1990 , 10, 5782-95	47
1038	The SIF binding element confers sis/PDGF inducibility onto the c-fos promoter.. 1990 , 9, 4477-4484	456
1037	DNA sequences involved in the transcriptional activation of a human placental lactogen gene. 1990 , 4, 1815-26	32
1036	Thyrotropin and adenosine 3',5'-monophosphate stimulate the activity of the ferritin-H promoter. 1990 , 4, 1117-24	16
1035	A DNA-activated protein kinase from HeLa cell nuclei. 1990 , 10, 6460-71	257
1034	Interplay between U2 snRNP and 3' splice factor(s) for branch point selection on human beta-globin pre-mRNA. <i>Nucleic Acids Research</i> , 1990 , 18, 235-45	20.1 5
1033	Lipopolysaccharide is a potent monocyte/macrophage-specific stimulator of human immunodeficiency virus type 1 expression. 1990 , 172, 253-61	176
1032	Characterization of a cellular factor which interacts functionally with Oct-1 in the assembly of a multicomponent transcription complex. <i>Nucleic Acids Research</i> , 1990 , 18, 6871-80	20.1 106

1031	A novel B-cell lineage-specific transcription factor present at early but not late stages of differentiation. 1990 , 4, 849-59		197
1030	Transformed and nontransformed cells differ in stability and cell cycle regulation of a binding activity to the murine thymidine kinase promoter. 1990 , 87, 9310-4		27
1029	Tandem AP-1-binding sites within the human beta-globin dominant control region function as an inducible enhancer in erythroid cells. 1990 , 4, 993-1006		267
1028	A nuclear cap binding protein from HeLa cells. <i>Nucleic Acids Research</i> , 1990 , 18, 6989-95	20.1	69
1027	Regulation of 'haploid expressed genes' in male germ cells. 1990 , 88, 679-93		136
1026	A Sequential splicing mechanism promotes selection of an optimal exon by repositioning a downstream 5' splice site in preprotachykinin pre-mRNA. 1990 , 4, 1172-84		61
1025	TFE3: a helix-loop-helix protein that activates transcription through the immunoglobulin enhancer muE3 motif. 1990 , 4, 167-79		425
1024	Sarkosyl defines three intermediate steps in transcription initiation by RNA polymerase III: application to stimulation of transcription by E1A. 1990 , 4, 646-58		44
1023	Crosslinking of hnRNP proteins to pre-mRNA requires U1 and U2 snRNPs. <i>Nucleic Acids Research</i> , 1990 , 18, 3307-18	20.1	56
1022	Identification of a second inducible DNA-protein interaction in the kappa immunoglobulin enhancer. <i>Nucleic Acids Research</i> , 1990 , 18, 1037-43	20.1	35
1021	A functional splice site in the 5' untranslated region of a zein gene. <i>Nucleic Acids Research</i> , 1990 , 18, 111-20	20.1	16
1020	Purification and characterization of pre-mRNA splicing factor SF2 from HeLa cells. 1990 , 4, 1158-71		308
1019	Compilation of DNA sequences of Escherichia coli (update 1990). <i>Nucleic Acids Research</i> , 1990 , 18 Suppl, 2549-87	20.1	27
1018	In vitro splicing of fibronectin pre-mRNAs. <i>Nucleic Acids Research</i> , 1990 , 18, 4089-97	20.1	32
1017	Muscle-specific expression of the cardiac alpha-actin gene requires MyoD1, CARG-box binding factor, and Sp1. 1990 , 4, 1811-22		317
1016	Site-specific excision from RNA by RNase H and mixed-phosphate-backbone oligodeoxynucleotides. 1990 , 87, 1401-5		139
1015	X-box-binding proteins positively and negatively regulate transcription of the HLA-DRA gene through interaction with discrete upstream W and V elements. 1990 , 87, 7703-7		39
1014	Regulation of a human epidermal keratin gene: sequences and nuclear factors involved in keratinocyte-specific transcription. 1990 , 4, 1985-98		134

1013	Cooperative interactions between transcription factors Sp1 and OTF-1. 1990 , 87, 4732-6	106
1012	Myogenin resides in the nucleus and acquires high affinity for a conserved enhancer element on heterodimerization. 1990 , 4, 582-95	261
1011	Regulation of the class II-associated invariant chain gene in normal and mutant B lymphocytes. 1990 , 87, 4590-4	20
1010	Coupled and uncoupled induction of fos and jun transcription by different second messengers in cells of hematopoietic origin. <i>Nucleic Acids Research</i> , 1990 , 18, 221-8	20.1 69
1009	Genetic mechanisms that determine oxidative capacity of striated muscles. Control of gene transcription. 1990 , 82, 319-31	11
1008	The effect of immunosuppressive agents on the induction of nuclear factors that bind to sites on the interleukin 2 promoter. 1990 , 172, 1869-72	119
1007	Identification of two distinct intron elements involved in alternative splicing of beta-tropomyosin pre-mRNA. 1990 , 4, 98-110	98
1006	NRF-1: a trans-activator of nuclear-encoded respiratory genes in animal cells. 1990 , 4, 1023-34	309
1005	Assembly of alternative multiprotein complexes directs rRNA promoter selectivity. 1990 , 4, 943-54	141
1004	Casein kinase II enhances the DNA binding activity of serum response factor. 1990 , 4, 955-67	172
1003	In vivo monitoring of a cAMP-stimulated DNA-binding activity. 1990 , 4, 1437-49	82
1002	Uridine branch acceptor is a cis-acting element involved in regulation of the alternative processing of calcitonin/CGRP-I pre-mRNA. <i>Nucleic Acids Research</i> , 1990 , 18, 5365-73	20.1 52
1001	Inducibility of the HS II enhancer depends on binding of an erythroid specific nuclear protein. <i>Nucleic Acids Research</i> , 1990 , 18, 6011-7	20.1 137
1000	Conserved domains of human U4 snRNA required for snRNP and spliceosome assembly. <i>Nucleic Acids Research</i> , 1990 , 18, 6223-9	20.1 33
999	Macromolecular interaction on a cAMP responsive region in the urokinase-type plasminogen activator gene: a role of protein phosphorylation. <i>Nucleic Acids Research</i> , 1990 , 18, 1991-9	20.1 33
998	In vitro activation of heat shock transcription factor DNA-binding by calcium and biochemical conditions that affect protein conformation. 1990 , 87, 3748-52	274
997	Mutational analysis of the promoter region of the alpha 27 gene of herpes simplex virus 1 within the context of the viral genome. 1990 , 87, 5268-72	22
996	Light regulates expression of a Fos-related protein in rat suprachiasmatic nuclei. 1990 , 87, 5959-62	241

995	Transcription of the testis-specific mouse protamine 2 gene in a homologous in vitro transcription system. 1990 , 87, 891-5	75
994	Interferon gamma regulates binding of two nuclear protein complexes in a macrophage cell line. 1990 , 87, 914-8	35
993	A mammalian protein of 220 kDa binds pre-mRNAs in the spliceosome: a potential homologue of the yeast PRP8 protein. 1990 , 87, 3082-6	57
992	An insulin response element in the glyceraldehyde-3-phosphate dehydrogenase gene binds a nuclear protein induced by insulin in cultured cells and by nutritional manipulations in vivo. 1990 , 87, 5273-7	130
991	A mutant of herpes simplex virus type 1 immediate early polypeptide Vmw175 binds to the cap site of its own promoter in vitro but fails to autoregulate in vivo. 1990 , 71 (Pt 4), 851-61	19
990	A novel enhancer in the immunoglobulin lambda locus is duplicated and functionally independent of NF kappa B. 1990 , 4, 978-92	91
989	The specific binding of nuclear protein(s) to the cAMP responsive element (CRE) sequence (TGACGTCA) is reduced by the misincorporation of U and increased by the deamination of C. <i>Nucleic Acids Research</i> , 1990 , 18, 5775-80	20.1 62
988	Small-scale preparation of extracts from radiolabeled cells efficient in pre-mRNA splicing. 1990 , 181, 20-30	72
987	Reversal of creatine kinase translational repression by 3' untranslated sequences. 1990 , 248, 1003-6	73
986	RNA lariat debranching enzyme as tool for analyzing RNA structure. 1990 , 181, 180-8	19
985	RNA polymerase II transcription blocked by Escherichia coli lac repressor. 1990 , 248, 480-3	85
984	Different combinations of regulatory elements may account for expression of the glycoprotein hormone alpha-subunit gene in primate and horse placenta. 1990 , 4, 1480-7	41
983	Thyroid hormone receptors form distinct nuclear protein-dependent and independent complexes with a thyroid hormone response element. 1990 , 4, 1627-35	73
982	Calcitonin exon sequences influence alternative RNA processing. 1990 , 4, 1744-9	22
981	Transcription of the human calcitonin gene is mediated by a C cell-specific enhancer containing E-box-like elements. 1990 , 4, 1750-7	37
980	Similar nuclear factors mediate stimulation of rat thyroglobulin gene transcription by thyrotropin and insulin-like growth factor-I. 1990 , 4, 2021-9	9
979	The homeodomain protein, Pit-1/GHF-1, is capable of binding to and activating cell-specific elements of both the growth hormone and prolactin gene promoters. 1990 , 4, 1069-80	109
978	Xenobiotic-inducible expression of murine glutathione S-transferase Ya subunit gene is controlled by an electrophile-responsive element. 1990 , 87, 6258-62	421

977	Loss of a DNA-protein complex correlates with extinguished major histocompatibility complex class II expression in a human B cell. 1990 , 171, 2159-64		9
976	Transcription factor PEA3 participates in the induction of urokinase plasminogen activator transcription in murine keratinocytes stimulated with epidermal growth factor or phorbol-ester. <i>Nucleic Acids Research</i> , 1990 , 18, 5009-17	20.1	132
975	Transcriptional activation by Sp1 as directed through TATA or initiator: specific requirement for mammalian transcription factor IID. 1990 , 87, 4509-13		477
974	Structural and functional basis of the developmental regulation of human coagulation factor IX gene: factor IX Leyden. 1990 , 87, 4421-5		45
973	Differential regulation of human immunodeficiency viruses (HIVs): a specific regulatory element in HIV-2 responds to stimulation of the T-cell antigen receptor. 1990 , 87, 9098-102		61
972	A TATA-like sequence located downstream of the transcription initiation site is required for expression of an RNA polymerase II transcribed gene. 1990 , 4, 1611-22		76
971	Regulation of gene expression with double-stranded phosphorothioate oligonucleotides. 1990 , 250, 997-1000		355
970	Protein binding elements in the human beta-polymerase promoter. <i>Nucleic Acids Research</i> , 1990 , 18, 919-28	20.1	28
969	Site-directed mutagenesis reveals a liver transcription factor essential for the albumin transcriptional enhancer. 1990 , 87, 5469-73		102
968	Activation of the heat shock transcription factor by hypoxia in mammalian cells. 1990 , 87, 6263-7		151
967	Dioxin-inducible, Ah receptor-dependent transcription in vitro. 1990 , 87, 8545-9		26
966	Characterization of chicken octamer-binding proteins demonstrates that POU domain-containing homeobox transcription factors have been highly conserved during vertebrate evolution. 1990 , 87, 1099-103		42
965	A heteromeric transcription factor required for mammalian RNA polymerase II. <i>Nucleic Acids Research</i> , 1990 , 18, 4843-9	20.1	27
964	Initiation of transcription of the erythroid promoter of the porphobilinogen deaminase gene is regulated by a cis-acting sequence around the cap site. <i>Nucleic Acids Research</i> , 1990 , 18, 6509-15	20.1	58
963	Transcriptional initiation is controlled by upstream GC-box interactions in a TATAA-less promoter. 1990 , 10, 6632-41		256
962	Transcription vectors that facilitate the identification and mapping of RNA splice sites in genomic DNA. 1990 , 9, 535-42		2
961	Structure and transcription of a human gene for H1 RNA, the RNA component of human RNase P. <i>Nucleic Acids Research</i> , 1990 , 18, 97-103	20.1	139
960	Sequence specificity of the human mRNA N6-adenosine methylase in vitro. <i>Nucleic Acids Research</i> , 1990 , 18, 5735-41	20.1	157

959	A sensitive method for the determination of protein-DNA binding specificities. <i>Nucleic Acids Research</i> , 1990 , 18, 6197-204	20.1	346
958	HrpF, a human sequence-specific DNA-binding protein homologous to XrpFI, a <i>Xenopus laevis</i> oocyte transcription factor. <i>Nucleic Acids Research</i> , 1990 , 18, 5811-6	20.1	12
957	Structure and function of the enhancer 3' to the human A gamma globin gene. <i>Nucleic Acids Research</i> , 1990 , 18, 7407-15	20.1	46
956	A plant DNA-binding protein increases the number of active preinitiation complexes in a human in vitro transcription system. 1990 , 4, 1899-909		66
955	Detection of a novel minisatellite-specific DNA-binding protein. <i>Nucleic Acids Research</i> , 1990 , 18, 625-9	20.1	61
954	Unusual c-fos induction upon chromaffin PC12 differentiation by sodium butyrate: loss of fos autoregulatory function. <i>Nucleic Acids Research</i> , 1990 , 18, 3605-10	20.1	35
953	Splicing of a <i>C. elegans</i> myosin pre-mRNA in a human nuclear extract. <i>Nucleic Acids Research</i> , 1990 , 18, 143-9	20.1	17
952	The T----C substitution at -198 of the A gamma-globin gene associated with the British form of HPFH generates overlapping recognition sites for two DNA-binding proteins. <i>Nucleic Acids Research</i> , 1990 , 18, 5685-93	20.1	32
951	A multisubunit factor, CstF, is required for polyadenylation of mammalian pre-mRNAs. 1990 , 4, 2112-20		214
950	Albumin improves formation and detection of some specific protein-DNA complexes in the mobility shift assay. <i>Nucleic Acids Research</i> , 1990 , 18, 2198	20.1	12
949	The promoter of the human parathyroid hormone gene contains a functional cyclic AMP-response element. <i>Nucleic Acids Research</i> , 1990 , 18, 5677-83	20.1	26
948	Point mutations in AAUAAA and the poly (A) addition site: effects on the accuracy and efficiency of cleavage and polyadenylation in vitro. <i>Nucleic Acids Research</i> , 1990 , 18, 5799-805	20.1	451
947	A 62,000 molecular weight spliceosome protein crosslinks to the intron polypyrimidine tract. <i>Nucleic Acids Research</i> , 1990 , 18, 5995-6001	20.1	29
946	<i>Drosophila melanogaster</i> genes for U1 snRNA variants and their expression during development. <i>Nucleic Acids Research</i> , 1990 , 18, 6971-9	20.1	55
945	Beta-globin transcripts carrying a single intron with three adjacent nucleotides of 5' exon are efficiently spliced in vitro irrespective of intron position or surrounding exon sequences. <i>Nucleic Acids Research</i> , 1990 , 18, 4671-6	20.1	19
944	Analysis and in vitro localization of internal methylated adenine residues in dihydrofolate reductase mRNA. <i>Nucleic Acids Research</i> , 1990 , 18, 4803-8	20.1	19
943	A factor binding GATAAG confers tissue specificity on the promoter of the human zeta-globin gene. <i>Nucleic Acids Research</i> , 1990 , 18, 1339-50	20.1	45
942	Isolation and functional characterization of TIF-IB, a factor that confers promoter specificity to mouse RNA polymerase I. <i>Nucleic Acids Research</i> , 1990 , 18, 1385-93	20.1	69

941	Structure-probing of U1 snRNPs gradually depleted of the U1-specific proteins A, C and 70k. Evidence that A interacts differentially with developmentally regulated mouse U1 snRNA variants. <i>Nucleic Acids Research</i> , 1990 , 18, 449-57	20.1	60
940	The enhancer factor R of Epstein-Barr virus (EBV) is a sequence-specific DNA binding protein. <i>Nucleic Acids Research</i> , 1990 , 18, 6835-43	20.1	103
939	Identification of tissue specific nuclear proteins: DNA sequence and protein binding regions in the T cell receptor beta J-C intron. <i>Nucleic Acids Research</i> , 1990 , 18, 3027-33	20.1	7
938	A DNA helicase from human cells. <i>Nucleic Acids Research</i> , 1990 , 18, 6785-92	20.1	54
937	In vitro capping in <i>Trypanosoma cruzi</i> identifies and shows specificity for the spliced leader RNA and U-RNAs. <i>Nucleic Acids Research</i> , 1990 , 18, 4197-206	20.1	10
936	Protein-DNA interactions upstream from the human A gamma globin gene. <i>Nucleic Acids Research</i> , 1990 , 18, 1977-82	20.1	23
935	Binding sites in mammalian genes and viral gene regulatory regions recognized by methylated DNA-binding protein. <i>Nucleic Acids Research</i> , 1990 , 18, 6253-60	20.1	57
934	The major transcriptional regulatory protein of herpes simplex virus type 1 includes a protease resistant DNA binding domain. <i>Nucleic Acids Research</i> , 1990 , 18, 4579-85	20.1	12
933	A novel C-T transition within the distal CCAAT motif of the G gamma-globin gene in the Japanese HPFH: implication of factor binding in elevated fetal globin expression. <i>Nucleic Acids Research</i> , 1990 , 18, 5245-53	20.1	86
932	DNA-protein interactions at the interferon-responsive promoter: evidences for an involvement of phosphorylation. <i>Nucleic Acids Research</i> , 1990 , 18, 2125-31	20.1	12
931	Structural requirements of iron-responsive elements for binding of the protein involved in both transferrin receptor and ferritin mRNA post-transcriptional regulation. <i>Nucleic Acids Research</i> , 1990 , 18, 1819-24	20.1	116
930	A novel transcriptional activator (PO-B) binds between the TATA box and cap site of the pro-opiomelanocortin gene. <i>Nucleic Acids Research</i> , 1990 , 18, 4513-21	20.1	31
929	Regulated splicing of the <i>Drosophila</i> P transposable element third intron in vitro: somatic repression. 1990 , 248, 1200-8		152
928	An amplification-promoting sequence from mouse genomic DNA: interaction with a trans-acting factor that also affects gene expression. 1990 , 9, 311-21		3
927	Transcriptional control of the human Harvey ras proto-oncogene: role of multiple elements in the promoter region. 1990 , 94, 249-53		11
926	CpG methylation inhibits proenkephalin gene expression and binding of the transcription factor AP-2. <i>Nucleic Acids Research</i> , 1990 , 18, 3975-82	20.1	355
925	Monoclonal anti-La antibody derived from a mouse with experimental SLE is similar to human anti-La antibodies. 1990 , 3, 701-13		6
924	Single-strand-binding factor(s) which interact with ARS1 of <i>Saccharomyces cerevisiae</i> . 1990 , 95, 73-7		30

923	Different pathways mediate virus inducibility of the human IFN-alpha 1 and IFN-beta genes. 1990 , 60, 767-79	166
922	A novel transcriptional regulatory factor that binds to the polyoma virus enhancer in a developmental stage-specific manner. 1990 , 72, 327-36	6
921	Conserved sequences in the U2 snRNA-encoding genes of Kinetoplastida do not include the putative branchpoint recognition region. 1990 , 91, 71-7	38
920	Generating highly labeled oligonucleotides for DNA-protein interaction. 1990 , 7, 133-7	4
919	Multiple bandshift assay: rapid identification and cloning of DNA fragments containing specific protein-binding sites. 1990 , 90, 287-91	13
918	Two nuclear proteins bind to the major positive element of the apolipoprotein B gene promoter. 1990 , 94, 295-301	12
917	The synthesis of influenza virus negative-strand RNA takes place in insoluble complexes present in the nuclear matrix fraction. 1990 , 16, 325-37	37
916	The progesterone receptor stimulates cell-free transcription by enhancing the formation of a stable preinitiation complex. 1990 , 60, 247-57	210
915	Regulation of the Human α (I) Procollagen Gene by a Trans-Acting Factor. 1990 , 580, 451-453	4
914	Localization and characterization of the DNase I-hypersensitive site II (HS II) enhancer. A critical regulatory element within the beta-globin locus-activating region. 1990 , 612, 141-51	5
913	The regulation of gamma-globin gene expression. 1990 , 612, 160-6	1
912	Novel Brain Specific Transcription Factors Form DNA-Protein Complexes with SP1 and TATA Regions of the Rat Neuropeptide Y Gene Promoter. 1990 , 611, 382-387	
911	The U3 small nucleolar ribonucleoprotein functions in the first step of preribosomal RNA processing. 1990 , 60, 897-908	446
910	RNA polymerase B (II) and general transcription factors. 1990 , 59, 711-54	471
909	Preparation of extracts from higher eukaryotes. 1990 , 182, 194-203	201
908	Synergistic activation of transcription by multiple binding sites for NF-kappa B even in absence of co-operative factor binding to DNA. 1990 , 214, 373-80	42
907	Strategy for statistical-mapping of potential regulatory regions in the human genome. 1990 , 216, 485-90	4
906	Multiple cell type-specific proteins differentially regulate target sequence recognition by the alpha retinoic acid receptor. 1990 , 63, 729-38	207

905	HIV-1 Tat protein trans-activates transcription in vitro. 1990 , 63, 791-802	255
904	The v-rel oncogene encodes a kappa B enhancer binding protein that inhibits NF-kappa B function. 1990 , 63, 803-14	320
903	Targeted snRNP depletion reveals an additional role for mammalian U1 snRNP in spliceosome assembly. 1990 , 63, 293-302	150
902	A functional ribonucleoprotein complex forms around the 5' end of poliovirus RNA. 1990 , 63, 369-80	434
901	The essential pre-mRNA splicing factor SF2 influences 5' splice site selection by activating proximal sites. 1990 , 62, 35-42	443
900	Spliced leader RNA sequences can substitute for the essential 5' end of U1 RNA during splicing in a mammalian in vitro system. 1990 , 62, 889-99	91
899	GC box binding induces phosphorylation of Sp1 by a DNA-dependent protein kinase. 1990 , 63, 155-65	621
898	Mechanism of transcriptional activation by Sp1: evidence for coactivators. 1990 , 61, 1187-97	1024
897	A myosin-like dimerization helix and an extra-large homeodomain are essential elements of the tripartite DNA binding structure of LFB1. 1990 , 61, 1225-36	171
896	Trans splicing of nematode pre-messenger RNA in vitro. 1990 , 61, 1247-55	89
895	Purified human I kappa B can rapidly dissociate the complex of the NF-kappa B transcription factor with its cognate DNA. 1990 , 61, 255-65	412
894	Structure of spliceosomal snRNPs and their role in pre-mRNA splicing. 1990 , 1087, 265-92	344
893	Nuclear factor I stimulates transcription of the adenovirus 12 E1A gene in a cell-free system. 1990 , 1048, 85-92	11
892	Functional analysis of multiple promoters of the rat insulin-like factor II gene. 1990 , 1048, 165-70	28
891	In vitro transcription of the plasma protein genes of Bombyx mori. 1990 , 1087, 18-24	9
890	Class II ribonuclease H comigrates with, but is distinct from, the third largest subunit of calf thymus RNA polymerase I. 1990 , 1087, 31-8	2
889	Tumor necrosis factor beta (TNF-beta) induces binding of the NF-kappa B transcription factor to a high-affinity kappa B element in the TNF-beta promoter. 1990 , 2, 389-97	54
888	Activation of interleukin-6 gene expression through the NF-kappa B transcription factor. 1990 , 10, 2327-34	1146

887	Analysis of a CR1 (chicken repeat) sequence flanking the 5' end of the gene encoding alpha-skeletal actin. 1990 , 88, 173-80	3
886	A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. 1990 , 60, 461-72	670
885	Cell type specific and viral regulation of visna virus gene expression. 1990 , 16, 175-83	5
884	Primary structure and assignment to chromosome 6 of three related rat genes encoding liver serine protease inhibitors. 1990 , 94, 273-82	28
883	Sequence analysis of the promoter region of the rat vasopressin gene. 1990 , 260, 305-8	68
882	Analysis of U1snRNP-specific A protein cross-linked complexes. 1990 , 272, 213-6	3
881	A putative AP-2 binding site in the 5' flanking region of the mouse POMC gene. 1990 , 264, 125-9	9
880	Nuclear factor-1 (NF-1) binds to multiple sites within the transcriptional enhancer of Moloney murine leukemia virus. 1990 , 277, 209-11	7
879	Human HLA-DR alpha gene: a rare oligonucleotide (GTATA) identifies an upstream sequence required for nuclear protein binding. 1990 , 268, 51-4	13
878	Tissue and species distribution of liver type and tumor type nuclear poly(A) polymerases. 1990 , 169, 325-31	1
877	Transcription factor levels in medullary thyroid carcinoma cells differentiated by Harvey ras oncogene: c-jun is increased. 1990 , 170, 140-6	12
876	Coordinated induction of two unrelated glucose-regulated protein genes by a calcium ionophore: human BiP/GRP78 and GAPDH. 1990 , 171, 431-8	30
875	The interaction of a c-Jun/Fos related protein factor with the U3 sequences of the mouse mammary tumor virus LTR. 1990 , 172, 348-56	4
874	A simplified method for the preparation of transcriptionally active liver nuclear extracts. 1990 , 9, 777-81	120
873	Inhibition of in vitro transcription by specific double-stranded oligodeoxyribonucleotides. 1990 , 89, 203-9	16
872	Early events of RNA polymerase II transcription initiation. 1990 , 46, 729-35	3
871	Effects of the DNA topoisomerase II inhibitor, VM26, on transcriptional initiation in vitro. 1990 , 46, 1309-18	1
870	Purification of a group of HeLa nuclear proteins that bind to a regulatory element (-1430/-1327) of the human proliferating cell nucleolar protein P120 gene. 1991 , 180, 805-12	5

869	Expression of enhancer binding factors associated with various cell types of lung cancer. 1991 , 180, 1064-9	2
868	DNase I hypersensitive sites of the 5' region of the fibronectin gene of the liver of the rat. 1991 , 181, 131-7	8
867	Universality of c-fos transcriptional regulation: the Dyad Symmetry Element mediates activation by PMA in T lymphocytes. 1991 , 181, 353-60	3
866	Interactions of nuclear protein from cultured rat hepatocytes with the cyclic AMP responsive elements and the NF1-CTF site in the promoter of the rat phosphoenolpyruvate carboxykinase gene. 1991 , 181, 367-74	11
865	Stability of antisense DNA oligodeoxynucleotide analogs in cellular extracts and sera. 1991 , 49, 1793-801	183
864	Control of class II gene transcription during in vitro nucleosome assembly. 1991 , 35, 419-47	53
863	The orientation and spacing of core DNA-binding motifs dictate selective transcriptional responses to three nuclear receptors. 1991 , 65, 1267-79	559
862	Constitutive expression of zif268 in neocortex is regulated by synaptic activity. 1991 , 88, 5106-10	250
861	A general method for purification of H1 histones that are active for repression of basal RNA polymerase II transcription. 1991 , 2, 162-9	32
860	Three individual regulatory elements of the promoter positively activate the transcription of the murine gene encoding granulocyte colony-stimulating factor. 1991 , 107, 241-6	10
859	Transcription stimulation of the adenovirus type-12 E1a gene in vitro by a novel factor bound to a region adjacent to a TATA box. 1991 , 109, 171-6	1
858	Physiology and cell biology update: mechanisms of type IV collagen gene regulation. 1991 , 18, 134-9	2
857	Molecular pathways of pain: Fos/Jun-mediated activation of a noncanonical AP-1 site in the prodynorphin gene. 1991 , 6, 607-17	258
856	Inducible preproenkephalin DNA binding proteins in the rat striatum. 1991 , 2, 427-39	6
855	Identification of cis-acting regulatory elements in the human oxytocin gene promoter. 1991 , 2, 501-10	9
854	In vivo stimulation of H-2Dd expression following RadLV infection of thymocytes: increased transcription and DNA-binding activity to sequences 5' of the Dd gene. 1991 , 142, 431-40	5
853	Importance of the CArG box in regulation of beta-actin-encoding genes. 1991 , 108, 211-7	23
852	Primary structure of the rat gene encoding an inhibitor of the insulin receptor tyrosine kinase. 1991 , 98, 209-16	34

851	Protein-binding A + T-rich motifs flank the duck beta A-globin enhancer. 1991 , 103, 253-7	8
850	Neither the endogenous nor a functional steroid hormone receptor binding site transactivate the ribosomal RNA gene promoter in vitro. 1991 , 39, 409-18	7
849	Splicing of the Drosophila P element ORF2-ORF3 intron is inhibited in a human cell extract. 1991 , 35, 65-72	22
848	Characterization of the murine Hox-2.3 promoter: involvement of the transcription factor USF (MLTF). 1991 , 33, 179-90	21
847	Evidence that interleukin-1 and phorbol esters activate NF-kappa B by different pathways: role of protein kinase C. 1991 , 2, 329-35	65
846	Role of nucleosomal cores and histone H1 in regulation of transcription by RNA polymerase II. 1991 , 254, 238-245	266
845	Nucleotides within both proximal and distal parts of the consensus sequence are important for specific DNA recognition by the herpes simplex virus regulatory protein ICP4. <i>Nucleic Acids Research</i> , 1991 , 19, 477-83	20.1 12
844	Hypoxia-inducible nuclear factors bind to an enhancer element located 3' to the human erythropoietin gene. 1991 , 88, 5680-4	677
843	Purification of sequence-specific binding proteins by DNA affinity chromatography. 1991 , 208, 10-23	114
842	Transcription factor OTF-1 interacts with two distinct DNA elements in the A gamma-globin gene promoter. 1991 , 30, 2961-7	14
841	Apolipoprotein B upstream suppressor site: identification of an element which can decrease apolipoprotein B transcription. 1991 , 176, 1116-22	3
840	Effects of chronic exposure of NG108-15 cells to morphine or ethanol on binding of nuclear factors to cAMP-response element. 1991 , 174, 25-31	14
839	Binding of cytosolic protein from cultured rat hepatocytes to the 3'-end of phosphoenolpyruvate carboxykinase mRNA--significance for protein-mediated mRNA stabilization. 1991 , 177, 1273-82	14
838	Purification of a novel 55 kDa HeLa cell nuclear DNA-binding protein. 1991 , 174, 542-8	5
837	Transactivation of the human insulin receptor gene by the CAAT/enhancer binding protein. 1991 , 174, 721-8	32
836	UV damage-specific DNA-binding protein in xeroderma pigmentosum complementation group E. 1991 , 175, 1139-43	74
835	A Pit-1 binding site 3' to the transcription start site inhibits transcription elongation in vitro. 1991 , 177, 790-6	12
834	Interference with protein binding at AP2 sites by sequence-specific methylation in the late E2A promoter of adenovirus type 2 DNA. 1991 , 281, 191-5	26

833	A 53 kDa protein binds to the negative regulatory region of JC virus early promoter. 1991 , 281, 272-4	12
832	Identification of a polypeptide bound to the beta region of the mouse r protein L32 promoter. 1991 , 286, 163-6	6
831	Functional properties of human thyroid hormone receptor beta 1 overexpressed using baculovirus. 1991 , 291, 315-8	3
830	Characterization of an enhancer-like structure in the promoter region of the proliferating cell nuclear antigen (PCNA) gene. 1991 , 193, 283-90	16
829	A cellular factor binding to the TAATGARAT DNA sequence prevents the expression of the HSV immediate-early genes following infection of nonpermissive cell lines derived from dorsal root ganglion neurons. 1991 , 194, 78-82	15
828	Genes for collagen types I, IV, and V are transcribed in HeLa cells but a postinitiation block prevents the accumulation of type I mRNA. 1991 , 192, 118-21	12
827	The 165-kDa DNA topoisomerase I from <i>Xenopus laevis</i> oocytes is a tissue-specific variant. 1991 , 146, 4-11	9
826	Distamycin inhibits the binding of a nuclear factor to the -278/-256 upstream sequence of the human HLA-DR alpha gene. 1991 , 41, 497-502	22
825	Developmentally and regionally regulated alterations of octamer- and GC-box-binding activities during the postnatal development of mouse cerebellum. 1991 , 61, 161-8	6
824	Transcriptional repression by YY1, a human GLI-Krüppel-related protein, and relief of repression by adenovirus E1A protein. 1991 , 67, 377-88	914
823	Activation domains of stably bound GAL4 derivatives alleviate repression of promoters by nucleosomes. 1991 , 64, 533-44	250
822	Ubiquitous transcription factor OTF-1 mediates induction of the MMTV promoter through synergistic interaction with hormone receptors. 1991 , 64, 565-72	219
821	DNA methylation inhibits transcription indirectly via a methyl-CpG binding protein. 1991 , 64, 1123-34	644
820	Mechanism of action of an acidic transcriptional activator in vitro. 1991 , 64, 971-81	554
819	IP-1: a dominant inhibitor of Fos/Jun whose activity is modulated by phosphorylation. 1991 , 64, 983-93	215
818	Nuclear targeting of the transcription factor PTF1 is mediated by a protein subunit that does not bind to the PTF1 cognate sequence. 1991 , 67, 987-94	68
817	Promoter melting by a stage-specific vaccinia virus transcription factor is independent of the presence of RNA polymerase. 1991 , 65, 105-13	49
816	G1/S phosphorylation of the retinoblastoma protein is associated with an altered affinity for the nuclear compartment. 1991 , 65, 381-93	298

815	Specific regulation of mRNA splicing in vitro by a peptide from HIV-1 Rev. 1991 , 67, 169-78	143
814	Primary structure of the human splicing factor ASF reveals similarities with Drosophila regulators. 1991 , 66, 373-82	340
813	Functional expression of cloned human splicing factor SF2: homology to RNA-binding proteins, U1 70K, and Drosophila splicing regulators. 1991 , 66, 383-94	475
812	Purification of a stage-specific and sequence-specific DNA-binding protein for the arylphorin gene of <i>Sarcophaga peregrina</i> . 1991 , 1089, 21-6	4
811	Regulatory elements mediating transcription of the human Ha-ras gene. 1991 , 220, 599-611	18
810	Influences of separation and adjacent sequences on the use of alternative 5' splice sites. 1991 , 217, 265-81	30
809	During <i>Drosophila</i> embryogenesis the beta 1 tubulin gene is specifically expressed in the nervous system and the apodemes. 1991 , 33, 107-18	32
808	Cholinergic agonist-induced binding of adrenomedullary nuclear proteins to the rat preproenkephalin promoter. 1991 , 2, 517-25	10
807	Constitutive and NF-kappa B-like proteins in the regulation of the serum amyloid A gene by interleukin 1. 1991 , 3, 380-8	54
806	The octamer-binding protein Oct-2 represses HSV immediate-early genes in cell lines derived from latently infectable sensory neurons. 1991 , 7, 381-90	114
805	Role of nucleosomal cores and histone H1 in regulation of transcription by RNA polymerase II. 1991 , 254, 238-45	310
804	Affinity purification of transcription factor IIA from HeLa cell nuclear extracts.. 1991 , 10, 2305-2310	29
803	Protein-DNA interactions in vivo of an erythroid-specific, human beta-globin locus enhancer. 1991 , 88, 8676-80	68
802	Transcription factor requirement for multiple rounds of initiation by human RNA polymerase II. 1991 , 88, 10691-5	18
801	The interaction of ICP4 with cell/infected-cell factors and its state of phosphorylation modulate differential recognition of leader sequences in herpes simplex virus DNA.. 1991 , 10, 397-406	24
800	VITAMIN D REGULATION OF THE BONE SPECIFIC OSTEOCALCIN GENE IS FUNCTIONALLY RELATED TO OSTEOBLAST GROWTH AND DIFFERENTIATION. 1991 , 12-20	
799	Pax-3, a novel murine DNA binding protein expressed during early neurogenesis.. 1991 , 10, 1135-1147	667
798	Changes in NF-kappa B and ISGF3 DNA binding activities are responsible for differences in MHC and beta-IFN gene expression in Ad5- versus Ad12-transformed cells.. 1991 , 10, 4169-4175	36

797	The 5'HS2 of the globin locus control region enhances transcription through the interaction of a multimeric complex binding at two functionally distinct NF-E2 binding sites.. 1991 , 10, 1391-1398	153
796	Three protein factors (SF1, SF3 and U2AF) function in pre-splicing complex formation in addition to snRNPs.. 1991 , 10, 1503-1509	65
795	TFIID is required for in vitro transcription of the human U6 gene by RNA polymerase III.. 1991 , 10, 1853-1862	82
794	Poliovirus proteinase 3C converts an active form of transcription factor III _C to an inactive form: a mechanism for inhibition of host cell polymerase III transcription by poliovirus.. 1991 , 10, 2941-2947	127
793	In vitro deadenylation of mammalian mRNA by a HeLa cell 3' exonuclease.. 1991 , 10, 3067-3071	84
792	An intact Box C sequence in the U3 snRNA is required for binding of fibrillarin, the protein common to the major family of nucleolar snRNPs.. 1991 , 10, 2645-2651	102
791	Novel downstream elements upregulate transcription initiated from an Epstein-Barr virus latent promoter.. 1991 , 10, 143-151	18
790	Mammalian nuclei contain foci which are highly enriched in components of the pre-mRNA splicing machinery.. 1991 , 10, 195-206	147
789	Biochemical characterization of U2 snRNP auxiliary factor: an essential pre-mRNA splicing factor with a novel intranuclear distribution.. 1991 , 10, 207-214	164
788	A novel splicing factor is an integral component of 200S large nuclear ribonucleoprotein (InRNP) particles.. 1991 , 10, 425-432	24
787	N-myc disrupts protein kinase C-mediated signal transduction in neuroblastoma.. 1991 , 10, 1119-1125	38
786	Binding of thymic factors to the conserved decanucleotide promoter element of the T-cell receptor V beta gene is developmentally regulated and is absent in SCID mice. 1991 , 88, 8131-5	10
785	Interaction of the v-rel protein with an NF-kappa B DNA binding site. 1991 , 88, 1783-7	36
784	A novel lineage-specific nuclear factor regulates mb-1 gene transcription at the early stages of B cell differentiation.. 1991 , 10, 3409-3417	104
783	Novel cyclic adenosine 3',5'-monophosphate response element in the human chorionic gonadotropin beta-subunit gene. 1991 , 5, 693-702	50
782	Cytoplasmic activation of GAF, an IFN-gamma-regulated DNA-binding factor.. 1991 , 10, 927-932	261
781	Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1.. 1991 , 10, 2247-2258	2809
780	Nuclear RNase MRP processes RNA at multiple discrete sites: interaction with an upstream G box is required for subsequent downstream cleavages. 1991 , 5, 1264-76	70

779	Muscle-specific expression of the troponin I gene requires interactions between helix-loop-helix muscle regulatory factors and ubiquitous transcription factors. 1991 , 11, 267-80	180
778	Activation and repression of transcription by the gap proteins hunchback and Krüppel in cultured <i>Drosophila</i> cells. 1991 , 5, 254-64	98
777	Recombinant 43-kDa USF binds to DNA and activates transcription in a manner indistinguishable from that of natural 43/44-kDa USF. 1991 , 11, 5125-36	103
776	Two distinct alpha-interferon-dependent signal transduction pathways may contribute to activation of transcription of the guanylate-binding protein gene. 1991 , 11, 5147-53	133
775	Analysis of small nuclear ribonucleoproteins (RNPs) in <i>Trypanosoma brucei</i> : structural organization and protein components of the spliced leader RNP. 1991 , 11, 5516-26	59
774	Natural and synthetic DNA elements with the CArG motif differ in expression and protein-binding properties. 1991 , 11, 6296-305	44
773	Role of the mammalian transcription factors IIF, IIS, and IIX during elongation by RNA polymerase II. 1991 , 11, 1195-206	163
772	A stem cell-specific silencer in the primer-binding site of a retrovirus. 1991 , 11, 1214-21	84
771	Cross-resistance to UV radiation of a cisplatin-resistant human cell line: overexpression of cellular factors that recognize UV-modified DNA. 1991 , 11, 2075-80	60
770	Stereospecific alignment of the X and Y elements is required for major histocompatibility complex class II DRA promoter function. 1991 , 11, 2406-15	73
769	Transcriptional role of a conserved GATA-1 site in the human epsilon-globin gene promoter. 1991 , 11, 2558-66	61
768	Identification of a multiprotein complex interacting with the c-fos serum response element. 1991 , 11, 2752-9	19
767	A 32-kilodalton protein binds to AU-rich domains in the 3' untranslated regions of rapidly degraded mRNAs. 1991 , 11, 3355-64	267
766	Structural analyses of the 7SK ribonucleoprotein (RNP), the most abundant human small RNP of unknown function. 1991 , 11, 3432-45	133
765	The B-cell and neuronal forms of the octamer-binding protein Oct-2 differ in DNA-binding specificity and functional activity. 1991 , 11, 3925-30	51
764	Characterization of a protein that regulates the DNA-binding activity of NF-AT, the nuclear factor of activated T cells. 1991 , 88, 11431-4	15
763	Affinity-purified CCAAT-box-binding protein (YEBP) functionally regulates expression of a human class II major histocompatibility complex gene and the herpes simplex virus thymidine kinase gene. 1991 , 88, 1873-7	70
762	Differential DNA binding by monomeric, homodimeric, and potentially heteromeric forms of the thyroid hormone receptor. 1991 , 11, 5005-15	145

761	A novel protein factor is required for use of distal alternative 5' splice sites in vitro. 1991 , 11, 5945-53	50
760	The <i>Drosophila</i> suppressor of sable gene encodes a polypeptide with regions similar to those of RNA-binding proteins. 1991 , 11, 894-905	42
759	RNA polymerase II elongation complexes paused after the synthesis of 15- or 35-base transcripts have different structures. 1991 , 11, 1508-22	51
758	Interaction of a lens cell transcription factor with the proximal domain of the mouse gamma F-crystallin promoter. 1991 , 11, 1531-7	31
757	Identification of a cis-acting sequence required for germ line-specific splicing of the P element ORF2-ORF3 intron. 1991 , 11, 1538-46	45
756	Sequence requirements for premature transcription arrest within the first intron of the mouse c-fos gene. 1991 , 11, 2832-41	84
755	Liver cells contain constitutive DNase I-hypersensitive sites at the xenobiotic response elements 1 and 2 (XRE1 and -2) of the rat cytochrome P-450IA1 gene and a constitutive, nuclear XRE-binding factor that is distinct from the dioxin receptor. 1991 , 11, 4314-23	25
754	Transactivation of pancreas-specific gene sequences in somatic cell hybrids. 1991 , 11, 4423-30	8
753	cAMP response element-binding protein is activated by Ca ²⁺ /calmodulin- as well as cAMP-dependent protein kinase. 1991 , 88, 5061-5	453
752	Cell-free transcription directed by the 422 adipose P2 gene promoter: activation by the CCAAT/enhancer binding protein. 1991 , 88, 8465-9	51
751	Possible involvement of (2'5')oligoadenylate synthetase activity in pre-mRNA splicing. 1991 , 88, 10377-81	37
750	Accurate and efficient RNA polymerase II transcription with a soluble nuclear fraction derived from <i>Drosophila</i> embryos. 1991 , 88, 1024-8	92
749	In vitro and in vivo analysis of the c-myc RNA polymerase III promoter. <i>Nucleic Acids Research</i> , 1991 , 19, 5045-52	20.1 22
748	The serum response factor is extensively modified by phosphorylation following its synthesis in serum-stimulated fibroblasts. 1991 , 11, 4545-54	90
747	Maintenance of NF-kappa B activity is dependent on protein synthesis and the continuous presence of external stimuli. 1991 , 11, 259-66	102
746	c-myc reverses neu-induced transformed morphology by transcriptional repression. 1991 , 11, 354-62	58
745	Novel protein-DNA interactions associated with increased immunoglobulin transcription in response to antigen plus interleukin-5. 1991 , 11, 5197-205	43
744	A heat-labile factor promotes premature 3' end formation in exon 1 of the murine adenosine deaminase gene in a cell-free transcription system. 1991 , 11, 5398-409	22

743	Extracellular signals that regulate liver transcription factors during hepatic differentiation in vitro. 1991 , 11, 773-84	168
742	Multiple basal promoter elements determine the level of human c-fos transcription. 1991 , 11, 1270-80	45
741	Dissection of the mouse N-ras gene upstream regulatory sequences and identification of the promoter and a negative regulatory element. 1991 , 11, 1334-43	23
740	Leucine periodicity of U2 small nuclear ribonucleoprotein particle (snRNP) A' protein is implicated in snRNP assembly via protein-protein interactions. 1991 , 11, 1578-89	44
739	Identification of single-stranded-DNA-binding proteins that interact with muscle gene elements. 1991 , 11, 1944-53	66
738	Detection and characterization of a factor which rescues spliceosome assembly from a heat-inactivated HeLa cell nuclear extract. 1991 , 11, 3425-31	15
737	Striking similarities between the regulatory mechanisms governing yeast mating-type genes and mammalian major histocompatibility complex genes. 1991 , 11, 4228-34	17
736	Identification of a specific exon sequence that is a major determinant in the selection between a natural and a cryptic 5' splice site. 1991 , 11, 4581-90	50
735	Primary structure of a human arginine-rich nuclear protein that colocalizes with spliceosome components. 1991 , 88, 8189-93	39
734	Identification of a fungal cutinase promoter that is inducible by a plant signal via a phosphorylated trans-acting factor. 1991 , 88, 8208-12	46
733	Human beta-globin locus control region: analysis of the 5' DNase I hypersensitive site HS 2 in transgenic mice. 1991 , 88, 1626-30	120
732	B-cell- and myocyte-specific E2-box-binding factors contain E12/E47-like subunits. 1991 , 11, 1156-60	162
731	Synergistic activation of a human promoter in vivo by transcription factor Sp1. 1991 , 11, 1935-43	108
730	The functional domains of the murine Thy-1 gene promoter. 1991 , 11, 2216-28	72
729	A GC-rich element confers epidermal growth factor responsiveness to transcription from the gastrin promoter. 1991 , 11, 2686-96	75
728	In vitro analysis of the tissue plasminogen activator promoter reveals a GC box-binding activity present in murine brain but undetectable in kidney and liver. 1991 , 11, 3139-47	32
727	The myosin light chain enhancer and the skeletal actin promoter share a binding site for factors involved in muscle-specific gene expression. 1991 , 11, 3735-44	47
726	The small subunit of transcription factor IIF recruits RNA polymerase II into the preinitiation complex. 1991 , 88, 9999-10003	158

725	Cloning of the DNA-binding subunit of human nuclear factor kappa B: the level of its mRNA is strongly regulated by phorbol ester or tumor necrosis factor alpha. 1991 , 88, 966-70	233
724	The immunosuppressant FK-506 specifically inhibits mitogen-induced activation of the interleukin-2 promoter and the isolated enhancer elements NFIL-2A and NF-AT1. 1991 , 11, 4074-87	42
723	Overlapping elements in the guanylate-binding protein gene promoter mediate transcriptional induction by alpha and gamma interferons. 1991 , 11, 182-91	271
722	The same CCAAT box-binding factor binds to the promoter of two coordinately regulated major histocompatibility complex class II genes. 1991 , 11, 578-81	6
721	Conserved mechanism of tRNA splicing in eukaryotes. 1991 , 11, 5410-6	79
720	Identification of an essential site for transcriptional activation within the human T-cell receptor delta enhancer. 1991 , 11, 5671-80	59
719	trans-Activation of a globin promoter in nonerythroid cells. 1991 , 11, 843-53	96
718	Tissue-specific transcription of the cardiac myosin light-chain 2 gene is regulated by an upstream repressor element. 1991 , 11, 1676-85	54
717	DNase I-hypersensitive sites and transcription factor-binding motifs within the mouse E beta meiotic recombination hot spot. 1991 , 11, 1813-9	64
716	Recognition of U1 and U2 small nuclear RNAs can be altered by a 5-amino-acid segment in the U2 small nuclear ribonucleoprotein particle (snRNP) B' protein and through interactions with U2 snRNP-A' protein. 1991 , 11, 1829-39	104
715	An in vitro transcription analysis of early responses of the human immunodeficiency virus type 1 long terminal repeat to different transcriptional activators. 1991 , 11, 1883-93	76
714	A new U6 small nuclear ribonucleoprotein-specific protein conserved between cis- and trans-splicing systems. 1991 , 11, 2026-34	23
713	Ecdysterone receptor is a sequence-specific transcription factor involved in the developmental regulation of heat shock genes. 1991 , 11, 3660-75	80
712	High expression of a 3'----5' exonuclease activity is specific to B lymphocytes. 1991 , 11, 4398-404	22
711	Identification of nuclear proteins that specifically bind to RNAs containing 5' splice sites. 1991 , 88, 320-4	22
710	A triplex DNA-binding protein from human cells: purification and characterization. 1991 , 88, 10450-4	84
709	Positive and negative regulation of immunoglobulin gene expression by a novel B-cell-specific enhancer element. 1991 , 11, 75-83	16
708	An RNA-binding protein specifically interacts with a functionally important domain of the downstream element of the simian virus 40 late polyadenylation signal. 1991 , 11, 5312-20	38

707	Transcriptional activation through ETS domain binding sites in the cytochrome c oxidase subunit IV gene. 1991 , 11, 5631-8	143
706	A position-dependent silencer plays a major role in repressing alpha-fetoprotein expression in human hepatoma. 1991 , 11, 5885-93	125
705	Characterization of the mouse granulocyte-macrophage colony-stimulating factor (GM-CSF) gene promoter: nuclear factors that interact with an element shared by three lymphokine genes--those for GM-CSF, interleukin-4 (IL-4), and IL-5. 1991 , 11, 5894-901	79
704	Pseudouridine modification of U5 RNA in ribonucleoprotein particles assembled in vitro. 1991 , 11, 5998-6006	31
703	Repression of immunoglobulin enhancers by the helix-loop-helix protein Id: implications for B-lymphoid-cell development. 1991 , 11, 6185-91	137
702	Nuclear and cytoplasmic location of the FER tyrosine kinase. 1991 , 11, 1180-3	53
701	Synergistic interactions between transcription factors control expression of the apolipoprotein AI gene in liver cells. 1991 , 11, 677-87	128
700	Cyclic AMP-independent activation of transcription factor NF-kappa B in HL60 cells by tumor necrosis factors alpha and beta. 1991 , 11, 2315-8	29
699	A complex regulatory DNA element associated with a major histocompatibility complex class I gene consists of both a silencer and an enhancer. 1991 , 11, 4217-27	69
698	Proximal regulatory domains of rat atrial natriuretic factor gene. 1991 , 84, 1256-65	47
697	Identification and characterization of a T-cell-specific enhancer adjacent to the murine CD4 gene. 1991 , 11, 5506-15	117
696	Characterization of the major regulatory element upstream of the human alpha-globin gene cluster. 1991 , 11, 4679-89	168
695	Transcription factors nuclear factor I and Sp1 interact with the murine collagen alpha 1 (I) promoter. 1991 , 11, 4065-73	124
694	Regulatory elements involved in constitutive and phorbol ester-inducible expression of the plasminogen activator inhibitor type 2 gene promoter. <i>Nucleic Acids Research</i> , 1991 , 19, 3881-6	20.1 53
693	Transcriptional regulation by Fos and Jun in vitro: interaction among multiple activator and regulatory domains. 1991 , 11, 3624-32	182
692	Molecular analyses of two poly(A) site-processing factors that determine the recognition and efficiency of cleavage of the pre-mRNA. 1991 , 11, 2432-8	112
691	The overlapping octamer/TAATGARAT motif is a high-affinity binding site for the cellular transcription factors Oct-1 and Oct-2. 1991 , 277 (Pt 2), 541-5	14
690	Differential expression of the mouse U1a and U1b SnRNA genes is not dependent on sequence differences in the octamer motif. 1991 , 277 (Pt 3), 719-22	5

689	Identification of inducible damage-recognition proteins that are overexpressed in HeLa cells resistant to cis-diamminedichloroplatinum (II). 1991 , 277 (Pt 3), 875-8	57
688	A heat-inducible nuclear factor that binds to the heat-shock element of the human haem oxygenase gene. 1991 , 277 (Pt 3), 895-7	26
687	Transcription in vitro of a napin gene, napA, from Brassica napus with a HeLa cell nuclear extract. 1991 , 115, 191-3	2
686	Isolation of multiple protein factors involved in ribosomal DNA transcription. 1991 , 587, 25-32	5
685	A rapid micropreparation technique for extraction of DNA-binding proteins from limiting numbers of mammalian cells. <i>Nucleic Acids Research</i> , 1991 , 19, 2499	20.1 2056
684	A nuclear factor that interacts with metal responsive elements of a human metallothionein gene. 1991 , 80, 145-57	11
683	Regulation of JCvL promoter function: evidence that a pentanucleotide "silencer" repeat sequence AGGGAAGGGA down-regulates transcription of the JC virus late promoter. 1991 , 180, 327-38	43
682	The Vmw175 binding site in the IE-1 promoter has no apparent role in the expression of Vmw110 during herpes simplex virus type 1 infection. 1991 , 180, 509-17	13
681	A key DNA-protein interaction determines the function of the 5'URR enhancer in human papillomavirus type 11. 1991 , 181, 132-8	15
680	In vitro transcription of the hepatitis B virus gene by nuclear extracts of human hepatoma cells. 1991 , 182, 545-52	6
679	Potential negative regulation of damage-recognition proteins in cisplatin-resistant HeLa cells in response to DNA damage. 1991 , 264, 59-66	7
678	Multiple sets of adjacent mu E1 and oct-1 binding sites upstream of the pseudorabies virus immediate-early gene promoter. 1991 , 182, 239-49	9
677	Identification of a major regulatory sequence in the latency associated transcript (LAT) promoter of herpes simplex virus type 1 (HSV-1). 1991 , 182, 287-97	75
676	DNA replication of human polyomavirus JC is stimulated by NF-I in vivo. 1991 , 182, 298-308	32
675	Upstream sequences within the terminal hairpin positively regulate the P6 promoter of B19 parvovirus. 1991 , 185, 39-47	16
674	Identification of a novel transcription factor, ACF, in cultured avian fibroblast cells that interacts with a Marek's disease virus late gene promoter. 1991 , 185, 80-9	2
673	Transactivation by herpes simplex virus proteins ICP4 and ICP0 in vaccinia virus infected cells. 1991 , 184, 67-78	8
672	TAPP (tetra-p-amidinophenoxyneopentane) inhibits the binding of nuclear factors to target DNA sequences. 1991 , 72, 251-258	7

671	Two related helix-loop-helix proteins participate in separate cell-specific complexes that bind the insulin enhancer. 1991 , 5, 292-9	142
670	Oncogenic germ-line mutations in Sp1 and ATF sites in the human retinoblastoma gene. 1991 , 353, 83-6	195
669	Binding of general transcription factor TFIIB to an acidic activating region. 1991 , 353, 569-71	375
668	YY1 is an initiator sequence-binding protein that directs and activates transcription in vitro. 1991 , 354, 241-5	419
667	Mapping of the bovine oxytocin gene control region: identification of binding sites for luteal nuclear proteins in the 5' non-coding region of the gene. 1991 , 3, 539-49	25
666	High-level expression of human c-jun gene causes cellular transformation of chicken embryo fibroblasts. 1991 , 82, 58-64	34
665	Organization of mouse DNA polymerase beta gene silencer elements and identification of the silencer-binding factor(s). 1991 , 82, 72-81	7
664	Identification and characterization of three DNA-binding proteins on the promoter of the human MDR1 gene in drug-sensitive and -resistant cells. 1991 , 82, 1151-9	17
663	Inhibitory effect of 8-chloro-cyclic adenosine 3',5'-monophosphate on cell growth of gastric carcinoma cell lines. 1991 , 82, 325-31	13
662	Activation of class I HLA expression by TNF-alpha and gamma-interferon is mediated through protein kinase C-dependent pathway in CML cell lines. 1991 , 78, 359-67	4
661	Structures of regulatory regions in the human cytochrome P-450sc (desmolase) gene. 1991 , 195, 563-9	40
660	ADP is a substrate for the AAUAAA-directed poly(A) addition reaction catalyzed by HeLa cell nuclear extracts. 1991 , 195, 685-9	
659	Differences in 2,3,7,8-tetrachlorodibenzo-p-dioxin-inducible CYP1A1 expression in human breast carcinoma cell lines involve altered trans-acting factors. 1991 , 197, 577-82	18
658	Baculovirus-expressed myogenic determination factors require E12 complex formation for binding to the myosin-light-chain enhancer. 1991 , 198, 187-93	8
657	A simple procedure for isolation of eukaryotic mRNA polyadenylation factors. 1991 , 202, 765-73	8
656	Binding of proteins of HeLa S3 cell extract to oligonucleotides containing the consensus interferon-response sequence (IRS) and to the IRS-containing fragment of the human c-myc gene. 1991 , 200, 107-11	3
655	3' processing of histone H4 precursor mRNA requires the presence of a small nuclear RNP particle. 1991 , 23, 1043-7	3
654	U7 snRNP particles bind H4 pre-mRNA in vitro. 1991 , 23, 1049-52	2

653	Transcriptional element H4-site II of cell cycle regulated human H4 histone genes is a multipartite protein/DNA interaction site for factors HiNF-D, HiNF-M, and HiNF-P: involvement of phosphorylation. 1991 , 46, 174-89	50
652	Influence of dexamethasone on the vitamin D-mediated regulation of osteocalcin gene expression. 1991 , 47, 184-96	44
651	Analysis of the c-myc P2 promoter. 1991 , 148, 75-84	15
650	Cross-linking of surface IgM activates NF-kappa B in B lymphocyte. 1991 , 21, 2993-8	33
649	Nuclear and DNA-binding proteins in human brain tumors. 1991 , 12, 515-23	4
648	Identification of locus-specific DNA-binding factors for the regulation of HLA class-I genes in human colorectal cancer. 1991 , 6, 131-7	7
647	Sequence-specific DNA-binding proteins within the Mbcr on the Ph1 chromosome. 1991 , 3, 308-12	3
646	Steroid hormone receptors and in vitro transcription. 1991 , 13, 73-8	16
645	Increased protein binding to a -530 mutation of the human beta-globin gene associated with decreased beta-globin synthesis. 1991 , 36, 42-7	48
644	Analysis of CYP1A1 promoter function by transcription in vitro. 1991 , 4, 93-6	8
643	Properties of the promoter region of the T18d (T13c) Tla gene. 1991 , 33, 171-7	4
642	E1A gene products stimulate in vitro transcription from the adenovirus early region 4 promoter by enhancing a stable preinitiation complex. 1991 , 5, 47-55	2
641	Differential binding of rat pituitary-specific nuclear factors to the 5'-flanking region of pituitary and placental members of the human growth hormone gene family. 1991 , 106, 181-7	16
640	Generation of monoclonal antibodies to the zinc finger domain of the eukaryotic transcription factor Sp1. 1991 , 101, 73-81	2
639	Accurate in vitro transcription of plant promoters with nuclear extracts prepared from cultured plant cells. 1991 , 16, 771-86	17
638	Expression of nuclear tRNA(Tyr) genes from Arabidopsis thaliana in HeLa cell and wheat germ extracts. 1991 , 16, 865-75	32
637	Silencer region of a chalcone synthase promoter contains multiple binding sites for a factor, SBF-1, closely related to GT-1. 1991 , 16, 235-49	88
636	A role for the interferon response DNA sequence in directing transcription of the T18d Tla gene. 1991 , 34, 293-8	1

635	Developmental and tissue-specific expression of the Q5k gene. 1991 , 34, 28-38		14
634	A set of 4 nuclear proteins binds to a DNA sequence within the FOS promoter region. 1991 , 15, 87-92		2
633	Identification of novel single-stranded d(TC)n binding proteins in several mammalian species. <i>Nucleic Acids Research</i> , 1991 , 19, 949-53	20.1	58
632	NF-I proteins from brain interact with the proenkephalin cAMP inducible enhancer. <i>Nucleic Acids Research</i> , 1991 , 19, 2721-8	20.1	43
631	A rapid microscale procedure for the simultaneous preparation of cytoplasmic RNA, nuclear DNA binding proteins and enzymatically active luciferase extracts. <i>Nucleic Acids Research</i> , 1991 , 19, 5080	20.1	64
630	The plasmacytoma J558L lacks constitutively active NF-kappa B and is deficient in early response gene activation. <i>Nucleic Acids Research</i> , 1991 , 19, 7235-41	20.1	5
629	Order of probe and nuclear protein extract addition can determine specificity of protein-DNA complexes in tested mobility shift assays. <i>Nucleic Acids Research</i> , 1991 , 19, 677-8	20.1	10
628	Polypeptide components of Drosophila small nuclear ribonucleoprotein particles. <i>Nucleic Acids Research</i> , 1991 , 19, 5877-82	20.1	31
627	Identification of transcriptional suppressor proteins that bind to the negative regulatory element of the human immunodeficiency virus type 1. <i>Nucleic Acids Research</i> , 1991 , 19, 6107-12	20.1	14
626	The 5' end domain of U2 snRNA is required to establish the interaction of U2 snRNP with U2 auxiliary factor(s) during mammalian spliceosome assembly. <i>Nucleic Acids Research</i> , 1991 , 19, 877-84	20.1	9
625	An NF kappa B-like factor is essential but not sufficient for cytokine induction of endothelial leukocyte adhesion molecule 1 (ELAM-1) gene transcription. <i>Nucleic Acids Research</i> , 1991 , 19, 2645-53	20.1	237
624	Transcription factors binding to the mouse HTF9 housekeeping promoter differ between cell types. <i>Nucleic Acids Research</i> , 1991 , 19, 4451-8	20.1	16
623	Multiple interdependent regulatory sites in the mouse c-fos promoter determine basal level transcription: cell type-specific effects. <i>Nucleic Acids Research</i> , 1991 , 19, 3583-91	20.1	21
622	Apolipoprotein B mRNA sequences 3' of the editing site are necessary and sufficient for editing and editosome assembly. <i>Nucleic Acids Research</i> , 1991 , 19, 6781-6	20.1	69
621	Requirement of protein factors and ATP for the disassembly of the spliceosome after mRNA splicing reaction. <i>Nucleic Acids Research</i> , 1991 , 19, 6819-21	20.1	6
620	Characterization of a third major SINE family of repetitive sequences in the galago genome. <i>Nucleic Acids Research</i> , 1991 , 19, 1649-56	20.1	41
619	Sequences involved in the control of adenovirus L1 alternative RNA splicing. <i>Nucleic Acids Research</i> , 1991 , 19, 2379-86	20.1	29
618	The housekeeping promoter from the mouse CpG island HTF9 contains multiple protein-binding elements that are functionally redundant. <i>Nucleic Acids Research</i> , 1991 , 19, 2817-24	20.1	32

617	Two hypervariable minisatellite DNA binding proteins. <i>Nucleic Acids Research</i> , 1991 , 19, 3269-74	20.1	46
616	The amdR product and a CCAAT-binding factor bind to adjacent, possibly overlapping DNA sequences in the promoter region of the <i>Aspergillus nidulans</i> amdS gene. <i>Nucleic Acids Research</i> , 1991 , 19, 2655-60	20.1	62
615	Validation of an in vitro RNA processing system for CT/CGRP precursor mRNA. <i>Nucleic Acids Research</i> , 1991 , 19, 3601-6	20.1	26
614	Alterations of RNase H sensitivity of the 3' splice site region during the in vitro splicing reaction. <i>Nucleic Acids Research</i> , 1991 , 19, 3953-8	20.1	18
613	Ara-ATP impairs 3'-end processing of pre-mRNAs by inhibiting both cleavage and polyadenylation. <i>Nucleic Acids Research</i> , 1991 , 19, 5871-5	20.1	9
612	Characterization of the apolipoprotein B mRNA editing enzyme: no similarity to the proposed mechanism of RNA editing in kinetoplastid protozoa. <i>Nucleic Acids Research</i> , 1991 , 19, 3569-76	20.1	56
611	A nuclear micrococcal-sensitive, ATP-dependent exoribonuclease degrades uncapped but not capped RNA substrates. <i>Nucleic Acids Research</i> , 1991 , 19, 2685-92	20.1	58
610	The C-terminal 70 amino acids of the adenovirus E4-ORF6/7 protein are essential and sufficient for E2F complex formation. <i>Nucleic Acids Research</i> , 1991 , 19, 6579-86	20.1	20
609	GATA-1 transactivates erythropoietin receptor gene, and erythropoietin receptor-mediated signals enhance GATA-1 gene expression. <i>Nucleic Acids Research</i> , 1991 , 19, 3843-8	20.1	132
608	LR1, a lipopolysaccharide-responsive factor with binding sites in the immunoglobulin switch regions and heavy-chain enhancer. 1991 , 5, 2353-61		57
607	DNA protein interactions at the interferon-responsive promoter elements: potential for an H-DNA conformation. <i>Nucleic Acids Research</i> , 1991 , 19, 517-24	20.1	9
606	Characterization of human 5S rRNA genes. <i>Nucleic Acids Research</i> , 1991 , 19, 4147-51	20.1	66
605	Alpha-tropomyosin mutually exclusive exon selection: competition between branchpoint/polypyrimidine tracts determines default exon choice. 1991 , 5, 642-55		132
604	The constitutively expressed octamer binding protein OTF-1 and a novel octamer binding protein expressed specifically in cervical cells bind to an octamer-related sequence in the human papillomavirus 16 enhancer. <i>Nucleic Acids Research</i> , 1991 , 19, 4531-5	20.1	22
603	Thyrotropin-releasing hormone regulation of human TSHB expression: role of a pituitary-specific transcription factor (Pit-1/GHF-1) and potential interaction with a thyroid hormone-inhibitory element. 1991 , 88, 3130-4		111
602	The cloned RNA polymerase II transcription factor IID selects RNA polymerase III to transcribe the human U6 gene in vitro. 1991 , 5, 1477-89		136
601	cAMP stimulates the C/EBP-related transcription factor rNFIL-6 to trans-locate to the nucleus and induce c-fos transcription. 1991 , 5, 1754-66		339
600	tat regulates binding of the human immunodeficiency virus trans-activating region RNA loop-binding protein TRP-185. 1991 , 5, 2128-40		125

599	DNA helicase IV from HeLa cells. <i>Nucleic Acids Research</i> , 1991 , 19, 3613-8	20.1	49
598	An intron binding protein is required for transformation ability of p53. <i>Nucleic Acids Research</i> , 1991 , 19, 4747-52	20.1	39
597	Intracellular distribution of microinjected antisense oligonucleotides. 1991 , 88, 2702-6		344
596	Modular recognition of 5-base-pair DNA sequence motifs by human heat shock transcription factor. 1991 , 11, 3504-14		27
595	Novel upstream elements and the TATA-box region mediate preferential transcription from the uteroglobin promoter in endometrial cells. <i>Nucleic Acids Research</i> , 1991 , 19, 2849-59	20.1	32
594	The herpes simplex virus trans-activator VP16 recognizes the Oct-1 homeo domain: evidence for a homeo domain recognition subdomain. 1991 , 5, 2555-66		115
593	The mouse proline-rich protein MP6 promoter binds isoprenaline-inducible parotid nuclear proteins via a highly conserved NFkB/rel-like site. <i>Nucleic Acids Research</i> , 1991 , 19, 5205-11	20.1	6
592	Constitutive expression of the urokinase plasminogen activator gene in murine RAW264 macrophages involves distal and 5' non-coding sequences that are conserved between mouse and pig. <i>Nucleic Acids Research</i> , 1991 , 19, 6839-47	20.1	47
591	The Zta trans-activator protein stabilizes TFIIID association with promoter DNA by direct protein-protein interaction. 1991 , 5, 2441-54		231
590	Interaction of DNA-binding proteins with a milk protein gene promoter in vitro: identification of a mammary gland-specific factor. <i>Nucleic Acids Research</i> , 1991 , 19, 6603-10	20.1	122
589	Isolation of intact nuclei from hematopoietic cell types. <i>Nucleic Acids Research</i> , 1991 , 19, 4301	20.1	45
588	Site-specific creation of uridine from cytidine in apolipoprotein B mRNA editing. <i>Nucleic Acids Research</i> , 1991 , 19, 1197-201	20.1	54
587	A new fast and easy method for the oligonucleotide affinity purification of DNA binding proteins. <i>Nucleic Acids Research</i> , 1991 , 19, 3156	20.1	2
586	Plant nonsense suppressor tRNA(Tyr) genes are expressed at very low levels in vitro due to inefficient splicing of the intron-containing pre-tRNAs. <i>Nucleic Acids Research</i> , 1991 , 19, 707-12	20.1	21
585	Isolation of the murine ribonuclease gene Rib-1: structure and tissue specific expression in pancreas and parotid gland. <i>Nucleic Acids Research</i> , 1991 , 19, 6935-41	20.1	14
584	SV40 activates transcription from the transferrin receptor promoter by inducing a factor which binds to the CRE/AP-1 recognition sequence. <i>Nucleic Acids Research</i> , 1991 , 19, 7117-23	20.1	17
583	LyF-1, a transcriptional regulator that interacts with a novel class of promoters for lymphocyte-specific genes. 1991 , 11, 5229-43		196
582	Functional characterization of the developmentally controlled immunoglobulin kappa 3' enhancer: regulation by Id, a repressor of helix-loop-helix transcription factors. 1991 , 11, 1040-7		119

581	Ca(2+)-mediated inhibition of a nuclear protein that recognizes UV-damaged DNA and is constitutively overexpressed in resistant human cells: DNA-binding assay. <i>Nucleic Acids Research</i> , 1991 , 19, 6413-8	20.1	5
580	2'-O-methyl, 2'-O-ethyl oligoribonucleotides and phosphorothioate oligodeoxyribonucleotides as inhibitors of the in vitro U7 snRNP-dependent mRNA processing event. <i>Nucleic Acids Research</i> , 1991 , 19, 2629-35	20.1	54
579	Cytoplasmic protein binding to highly conserved sequences in the 3' untranslated region of mouse protamine 2 mRNA, a translationally regulated transcript of male germ cells. 1991 , 88, 3584-8		140
578	Activation of endothelial-leukocyte adhesion molecule 1 (ELAM-1) gene transcription. 1991 , 88, 6523-7		245
577	An erythroid specific enhancer upstream to the gene encoding the cell-type specific transcription factor GATA-1. <i>Nucleic Acids Research</i> , 1991 , 19, 5285-91	20.1	81
576	Modulation of transcription factor NF-kappa B binding activity by oxidation-reduction in vitro. 1991 , 88, 4328-32		573
575	Role for the Epstein-Barr virus nuclear antigen 2 in viral promoter switching during initial stages of infection. 1991 , 88, 3942-6		115
574	Transcription from a TATA-less promoter requires a multisubunit TFIID complex. 1991 , 5, 1935-45		533
573	Insulin regulation of the glucagon gene is mediated by an insulin-responsive DNA element. 1991 , 88, 7224-7		110
572	In vitro apolipoprotein B mRNA editing: identification of a 27S editing complex. 1991 , 88, 1489-93		100
571	Two-step affinity purification of U7 small nuclear ribonucleoprotein particles using complementary biotinylated 2'-O-methyl oligoribonucleotides. 1991 , 88, 9784-8		59
570	Isolation of a cDNA for HSF2: evidence for two heat shock factor genes in humans. 1991 , 88, 6911-5		310
569	Inducible proteins binding to the murine thymidine kinase promoter in late G1/S phase. 1991 , 88, 1157-61		74
568	LEF-1, a gene encoding a lymphoid-specific protein with an HMG domain, regulates T-cell receptor alpha enhancer function [corrected]. 1991 , 5, 880-94		524
567	c-rel activates but v-rel suppresses transcription from kappa B sites. 1991 , 88, 3715-9		152
566	Footprinting analysis of mammalian RNA polymerase II along its transcript: an alternative view of transcription elongation. 1991 , 88, 4245-9		87
565	Potential role of poly(A) polymerase in the assembly of polyadenylation-specific RNP complexes. <i>Nucleic Acids Research</i> , 1991 , 19, 343-51	20.1	5
564	Modulation of alternative splicing of adenoviral E1A transcripts: factors involved in the early-to-late transition. 1991 , 5, 1847-58		26

563	Characterization of a nuclear protein that binds to three elements within the silencer region of a bean chalcone synthase gene promoter. 1991 , 88, 2515-9		61
562	Different forms of soluble cytoplasmic mRNA binding proteins and particles in <i>Xenopus laevis</i> oocytes and embryos. 1991 , 112, 1-11		105
561	Sp1 is essential and its position is important for p120 gene transcription: a 35 bp juxtaposed positive regulatory element enhances transcription 2.5 fold. <i>Nucleic Acids Research</i> , 1991 , 19, 6559-63	20.1	10
560	The mouse Pkg-1 gene promoter contains an upstream activator sequence. <i>Nucleic Acids Research</i> , 1991 , 19, 5755-61	20.1	189
559	Prolactin-deficient GH3B3 cells are defective in the utilization of the endogenous prolactin promoter yet are fully competent to initiate transcription from a transfected prolactin promoter. 1991 , 10, 105-12		4
558	Interleukin-1 or phorbol induction of the stromelysin promoter requires an element that cooperates with AP-1. <i>Nucleic Acids Research</i> , 1991 , 19, 335-41	20.1	84
557	Human SRF-related proteins: DNA-binding properties and potential regulatory targets. 1991 , 5, 2327-41		388
556	Immunoaffinity purification of a [U4/U6.U5] tri-snRNP from human cells. 1991 , 5, 1439-52		108
555	Electron microscopy of U4/U6 snRNP reveals a Y-shaped U4 and U6 RNA containing domain protruding from the U4 core RNP. 1991 , 112, 1065-72		10
554	Proteins of the nuclear factor-1 family act as an activator of the late promoter in human polyomavirus BK in vitro. 1991 , 72 (Pt 8), 1935-42		20
553	O6-methylguanine inhibits the binding of transcription factors to DNA. <i>Nucleic Acids Research</i> , 1991 , 19, 5739-42	20.1	23
552	High level expression and purification of herpes simplex virus type 1 immediate early polypeptide Vmw110. <i>Nucleic Acids Research</i> , 1991 , 19, 6155-61	20.1	44
551	In vitro transcription of baculovirus immediate early genes: accurate mRNA initiation by nuclear extracts from both insect and human cells. 1991 , 88, 4513-7		76
550	Identification and characterization of a cis-acting element that interferes with glucocorticoid-inducible activation of the mouse mammary tumor virus promoter. 1991 , 88, 5393-7		23
549	RNA-protein complexes mediate in vitro capping of the spliced-leader primary transcript and U-RNAs in <i>Trypanosoma cruzi</i> . 1991 , 88, 5626-30		4
548	Inhibition by interferon of herpes simplex virus type 1-activated transcription of tat-defective provirus. 1991 , 88, 9573-7		18
547	Identification of nuclear factor delta EF1 and its binding site essential for lens-specific activity of the delta 1-crystallin enhancer. <i>Nucleic Acids Research</i> , 1991 , 19, 3543-7	20.1	61
546	Activation of IFN-beta element by IRF-1 requires a posttranslational event in addition to IRF-1 synthesis. <i>Nucleic Acids Research</i> , 1991 , 19, 4421-8	20.1	190

545	Proteolysis of splicing factors during rat and monkey cell fractionation. <i>Nucleic Acids Research</i> , 1991 , 19, 4509-14	20.1	6
544	Unique topography of DNA-protein interactions in the non-coding region of epidermodysplasia verruciformis-associated human papillomaviruses. 1991 , 72 (Pt 12), 2989-97		11
543	Activation of the cellular transcription factor AP-1 in herpes simplex virus infected cells is dependent on the viral immediate-early protein ICPO. <i>Nucleic Acids Research</i> , 1991 , 19, 4879-83	20.1	38
542	Retinoic acid response element in the human alcohol dehydrogenase gene ADH3: implications for regulation of retinoic acid synthesis. 1991 , 11, 1638-46		173
541	3,5,3'-triiodothyronine receptor auxiliary protein (TRAP) enhances receptor binding by interactions within the thyroid hormone response element. 1991 , 5, 85-93		74
540	The pancreatic islet-specific glucagon G3 transcription factors recognize control elements in the rat somatostatin and insulin-I genes. 1991 , 5, 1457-66		51
539	Binding specificity of cyclic adenosine 3',5'-monophosphate-responsive element (CRE)-binding proteins and activating transcription factors to naturally occurring CRE sequence variants. 1991 , 5, 1541-51		37
538	Structure-function analysis of the rat prolactin promoter: phasing requirements of proximal cell-specific elements. 1991 , 5, 836-43		27
537	Overexpression of the human vitamin D3 receptor in mammalian cells using recombinant adenovirus vectors. 1991 , 5, 867-78		18
536	Regulation of interleukin-2 gene enhancer activity by the T cell accessory molecule CD28. 1991 , 251, 313-6		545
535	Multiple Oct2 isoforms are generated by alternative splicing. <i>Nucleic Acids Research</i> , 1991 , 19, 43-51	20.1	138
534	Sequence-specific antirepression of histone H1-mediated inhibition of basal RNA polymerase II transcription. 1991 , 251, 643-9		379
533	Signal transduction by interferon-alpha through arachidonic acid metabolism. 1991 , 251, 204-7		146
532	Functional dissection of the lck proximal promoter. 1992 , 12, 2758-68		113
531	MAZ, a zinc finger protein, binds to c-MYC and C2 gene sequences regulating transcriptional initiation and termination. 1992 , 89, 7452-6		251
530	Binding of transcription factor sp1 to exon-1 of C-myc is altered in burkitts-lymphoma. 1992 , 1, 175-80		1
529	Inhibition of the activation of heat shock factor in vivo and in vitro by flavonoids. 1992 , 12, 3490-8		188
528	Analysis of the mechanism of steroid hormone receptor-dependent gene activation in cell-free systems. 1992 , 13, 525-35		52

527	Thyrotropin (TSH)-releasing hormone-responsive elements in the rat TSH beta gene have distinct biological and nuclear protein-binding properties. 1992 , 6, 43-52		9
526	Phosphorothioate substitution identifies phosphate groups important for pre-mRNA splicing. <i>Nucleic Acids Research</i> , 1992 , 20, 1949-57	20.1	16
525	Thioredoxin regulates the DNA binding activity of NF-kappa B by reduction of a disulphide bond involving cysteine 62. <i>Nucleic Acids Research</i> , 1992 , 20, 3821-30	20.1	727
524	A single-stranded DNA binding protein from mouse tumor cells specifically recognizes the C-rich strand of the (AGG:CCT) _n repeats that can alter DNA conformation. <i>Nucleic Acids Research</i> , 1992 , 20, 6631-5	20.1	26
523	Independent modes of transcriptional activation by the p50 and p65 subunits of NF-kappa B. 1992 , 6, 775-87		329
522	A combination of upstream and proximal elements is required for efficient expression of the mouse renin promoter in cultured cells. <i>Nucleic Acids Research</i> , 1992 , 20, 3617-23	20.1	24
521	Multiple octamer binding sites in the promoter region of the bovine alpha s2-casein gene. <i>Nucleic Acids Research</i> , 1992 , 20, 4311-8	20.1	48
520	Different binding site requirements for binding and activation for the bipartite enhancer factor EF-1A. <i>Nucleic Acids Research</i> , 1992 , 20, 6555-64	20.1	25
519	Identification of novel factors that bind to the PRD I region of the human beta-interferon promoter. <i>Nucleic Acids Research</i> , 1992 , 20, 1531-8	20.1	61
518	Functional characterization of the promoter of pp63, a gene encoding a natural inhibitor of the insulin receptor tyrosine kinase. <i>Nucleic Acids Research</i> , 1992 , 20, 1983-90	20.1	19
517	Specific transcription from the adenovirus E2E promoter by RNA polymerase III requires a subpopulation of TFIID. <i>Nucleic Acids Research</i> , 1992 , 20, 5705-12	20.1	24
516	The cooperative interaction between two motifs of an enhancer element of the chicken alpha A-crystallin gene, alpha CE1 and alpha CE2, confers lens-specific expression. <i>Nucleic Acids Research</i> , 1992 , 20, 3701-12	20.1	61
515	The abolition of collagen gene expression in SV40-transformed fibroblasts is associated with trans-acting factor switching. <i>Nucleic Acids Research</i> , 1992 , 20, 5825-30	20.1	13
514	E2F mediates dihydrofolate reductase promoter activation and multiprotein complex formation in human cytomegalovirus infection. 1992 , 12, 4364-74		109
513	Two upstream elements activate transcription of a major histocompatibility complex class I gene in vitro. <i>Nucleic Acids Research</i> , 1992 , 20, 2533-40	20.1	30
512	Two distinct pathways of interferon induction as revealed by 2-aminopurine. <i>Nucleic Acids Research</i> , 1992 , 20, 2749-54	20.1	9
511	Characterization of the spermidine-dependent, sequence-specific endoribonuclease that requires transfer RNA for its activity. <i>Nucleic Acids Research</i> , 1992 , 20, 3737-42	20.1	22
510	Multiple promoter elements govern expression of the human ornithine decarboxylase gene in colon carcinoma cells. <i>Nucleic Acids Research</i> , 1992 , 20, 2581-90	20.1	21

509	DNA helicase III from HeLa cells: an enzyme that acts preferentially on partially unwound DNA duplexes. <i>Nucleic Acids Research</i> , 1992 , 20, 5329-37	20.1	34
508	A direct-repeat sequence of the human BiP gene is required for A23187-mediated inducibility and an inducible nuclear factor binding. <i>Nucleic Acids Research</i> , 1992 , 20, 6481-5	20.1	9
507	An additional long-range interaction in human U1 snRNA. <i>Nucleic Acids Research</i> , 1992 , 20, 1215-21	20.1	4
506	Transcriptional induction of IFN-gamma-responsive genes is modulated by DNA surrounding the interferon stimulation response element. <i>Nucleic Acids Research</i> , 1992 , 20, 3865-72	20.1	17
505	Interaction of hnRNP A1 with snRNPs and pre-mRNAs: evidence for a possible role of A1 RNA annealing activity in the first steps of spliceosome assembly. <i>Nucleic Acids Research</i> , 1992 , 20, 5017-25	20.1	53
504	Cofractionation of the TATA-binding protein with the RNA polymerase III transcription factor TFIIIB. <i>Nucleic Acids Research</i> , 1992 , 20, 5889-98	20.1	57
503	U1-U2 snRNPs interaction induced by an RNA complementary to the 5' end sequence of U1 snRNA. <i>Nucleic Acids Research</i> , 1992 , 20, 3625-30	20.1	15
502	The AT-rich tract of the SV40 ori core: negative synergism and specific recognition by single stranded and duplex DNA binding proteins. <i>Nucleic Acids Research</i> , 1992 , 20, 3333-9	20.1	12
501	Characterization of a retinoic acid responsive element isolated by whole genome PCR. <i>Nucleic Acids Research</i> , 1992 , 20, 3223-32	20.1	19
500	In vitro and in vivo protein-DNA interactions on the rat erythroid-specific L' pyruvate kinase gene promoter. <i>Nucleic Acids Research</i> , 1992 , 20, 5669-76	20.1	19
499	An ultraviolet light-damaged DNA recognition protein absent in xeroderma pigmentosum group E cells binds selectively to pyrimidine (6-4) pyrimidone photoproducts. <i>Nucleic Acids Research</i> , 1992 , 20, 5805-10	20.1	65
498	Human calcitonin gene regulation by helix-loop-helix recognition sequences. <i>Nucleic Acids Research</i> , 1992 , 20, 117-23	20.1	37
497	Identification of exon sequences and an exon binding protein involved in alternative RNA splicing of calcitonin/CGRP. <i>Nucleic Acids Research</i> , 1992 , 20, 2361-6	20.1	62
496	Unusual DNA binding characteristics of an in vitro translation product of the CCAAT binding protein mYB-1. <i>Nucleic Acids Research</i> , 1992 , 20, 601-6	20.1	33
495	A role for the TATA-box-binding protein component of the transcription factor IID complex as a general RNA polymerase III transcription factor. 1992 , 89, 1949-53		132
494	The rat poly pyrimidine tract binding protein (PTB) interacts with a single-stranded DNA motif in a liver-specific enhancer. <i>Nucleic Acids Research</i> , 1992 , 20, 1243-9	20.1	44
493	Mechanism of DNase I hypersensitive site formation within the human globin locus control region. 1992 , 89, 1143-7		90
492	Pax-5 encodes the transcription factor BSAP and is expressed in B lymphocytes, the developing CNS, and adult testis. 1992 , 6, 1589-607		432

491	In vivo footprint analysis of the HLA-DRA gene promoter: cell-specific interaction at the octamer site and up-regulation of X box binding by interferon gamma. 1992 , 89, 7601-5	60
490	Mechanism of transcriptional antirepression by GAL4-VP16. 1992 , 6, 2270-81	87
489	Heparin inhibits collagenase gene expression mediated by phorbol ester-responsive element in primate arterial smooth muscle cells. 1992 , 70, 1062-9	54
488	Trans-repressor activity of nuclear glycosaminoglycans on Fos and Jun/AP-1 oncoprotein-mediated transcription. 1992 , 116, 31-42	113
487	Casein kinase II inhibits the DNA-binding activity of Max homodimers but not Myc/Max heterodimers. 1992 , 6, 166-76	188
486	The erythroid-specific protein cGATA-1 mediates distal enhancer activity through a specialized beta-globin TATA box. 1992 , 6, 521-32	93
485	Switch recombination breakpoints are strictly correlated with DNA recognition motifs for immunoglobulin S gamma 3 DNA-binding proteins. 1992 , 176, 339-49	48
484	Characterization of apolipoprotein B mRNA editing from rabbit intestine. 1992 , 12, 172-9	36
483	Site-specific cross-linking of mammalian U5 snRNP to the 5' splice site before the first step of pre-mRNA splicing. 1992 , 6, 2542-53	239
482	Transcriptional analysis of the mts1 gene with specific reference to 5' flanking sequences. 1992 , 89, 9146-50	28
481	The dyad palindromic glutathione transferase P enhancer binds multiple factors including AP1. <i>Nucleic Acids Research</i> , 1992 , 20, 5153-8	20.1 44
480	Recognition of the surface of a homeo domain protein. 1992 , 6, 2047-57	106
479	Evidence for a HeLa cell splicing activity that is necessary for activation of a regulated adenovirus 3' splice site. <i>Nucleic Acids Research</i> , 1992 , 20, 3955-61	20.1 9
478	The expression of the amyloid precursor protein (APP) is regulated by two GC-elements in the promoter. <i>Nucleic Acids Research</i> , 1992 , 20, 63-8	20.1 86
477	SR proteins: a conserved family of pre-mRNA splicing factors. 1992 , 6, 837-47	629
476	Thymidine kinase transcription is regulated at G1/S phase by a complex that contains retinoblastoma-like protein and a cdc2 kinase. 1992 , 89, 3256-60	94
475	Structure of the small nuclear RNP particle U1: identification of the two structural protuberances with RNP-antigens A and 70K. 1992 , 116, 839-49	46
474	The AP-1 site at -150 bp, but not the NF-kappa B site, is likely to represent the major target of protein kinase C in the interleukin 2 promoter. 1992 , 175, 853-62	68

473	Molecular cloning and characterization of PEA3, a new member of the Ets oncogene family that is differentially expressed in mouse embryonic cells. 1992 , 6, 481-96		288
472	Purification and characterization of NF-R1 that regulates the expression of the human multidrug resistance (MDR1) gene. <i>Nucleic Acids Research</i> , 1992 , 20, 5811-7	20.1	31
471	Optimal tRNA((Ser)Sec) gene activity requires an upstream SPH motif. <i>Nucleic Acids Research</i> , 1992 , 20, 203-9	20.1	57
470	In vitro reconstitution of active influenza virus ribonucleoprotein complexes using viral proteins purified from infected cells. 1992 , 73 (Pt 7), 1855-9		23
469	Sequence-specific DNA-binding proteins are components of a nuclear matrix-attachment site. 1992 , 89, 4178-82		114
468	A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. 1992 , 12, 5447-54		2001
467	Subunit of an alpha-interferon-responsive transcription factor is related to interferon regulatory factor and Myb families of DNA-binding proteins. 1992 , 12, 3315-24		371
466	Association of U6 snRNA with the 5'-splice site region of pre-mRNA in the spliceosome. 1992 , 6, 244-54		120
465	Evolutionary conserved multiprotein complexes interact with the 3' untranslated region of histone transcripts. <i>Nucleic Acids Research</i> , 1992 , 20, 1023-30	20.1	12
464	Specific binding of chromosomal protein HMG1 to DNA damaged by the anticancer drug cisplatin. 1992 , 256, 234-7		489
463	Holo-TFIID supports transcriptional stimulation by diverse activators and from a TATA-less promoter. 1992 , 6, 1964-74		352
462	Reduction of a conserved Cys is essential for Myb DNA-binding. <i>Nucleic Acids Research</i> , 1992 , 20, 2279-86	20.1	113
461	Inhibition of Moloney Murine Leukaemia Virus Transcription by a Phospholipase-C Inhibitor Affecting Trans-Acting Factors. 1992 , 3, 283-291		
460	Ser/Thr-specific protein phosphatases are required for both catalytic steps of pre-mRNA splicing. <i>Nucleic Acids Research</i> , 1992 , 20, 5263-9	20.1	182
459	A cap binding protein that may mediate nuclear export of RNA polymerase II-transcribed RNAs. 1992 , 118, 1287-95		137
458	Induction of NF-kappa B-like activity by platelet-derived growth factor in mouse fibroblasts. 1992 , 3, 1131-9		35
457	Regulation of glycine tRNA gene expression in the posterior silk glands of the silkworm <i>Bombyx mori</i> . 1992 , 89, 1070-4		23
456	The mechanism of somatic inhibition of <i>Drosophila</i> P-element pre-mRNA splicing: multiprotein complexes at an exon pseudo-5' splice site control U1 snRNP binding. 1992 , 6, 1386-401		129

455	Differential ASF/SF2 activity in extracts from normal WI38 and transformed WI38VA13 cells. <i>Nucleic Acids Research</i> , 1992 , 20, 5197-204	20.1	12
454	SPT3 interacts with TFIID to allow normal transcription in <i>Saccharomyces cerevisiae</i> . 1992 , 6, 1319-31		194
453	Characterization of an enhancer upstream from the muscle-type promoter of a gene encoding 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase. <i>Nucleic Acids Research</i> , 1992 , 20, 3575-83	20.1	20
452	Extension of the DNA binding consensus of the chicken c-Myb and v-Myb proteins. <i>Nucleic Acids Research</i> , 1992 , 20, 3043-9	20.1	81
451	A rapid method for the isolation of DNA-binding proteins from purified nuclei of tissues and cells in culture. <i>Nucleic Acids Research</i> , 1992 , 20, 3555-9	20.1	48
450	A downstream sequence of the rpl32 promoter competes with the glucocorticoid responsive element for a protein factor. 1992 , 70, 787-91		5
449	Double-stranded RNA and interferon-alpha induce transcription through different molecular mechanisms. 1992 , 12, 445-8		16
448	Interferon-alpha activates binding of nuclear factors to a sequence element in the c-fos proto-oncogene 5'-flanking region. 1992 , 12, 355-61		8
447	Characterization and uses of monoclonal antibody derived against DNA binding domain of the ets family of genes. 1992 , 11, 277-94		11
446	Interferon-gamma potentiates the antiviral activity and the expression of interferon-stimulated genes induced by interferon-alpha in U937 cells. 1992 , 12, 87-94		42
445	Direct effect of basic fibroblast growth factor on gene transcription in a cell-free system. 1992 , 89, 5216-20		88
444	cis and trans elements differ among mouse strains with high and low extrahepatic complement factor B gene expression. 1992 , 175, 471-9		16
443	HIV-1 Tat acts as a processivity factor in vitro in conjunction with cellular elongation factors. 1992 , 6, 655-66		176
442	The Drosophila RNA-binding protein RBP1 is localized to transcriptionally active sites of chromosomes and shows a functional similarity to human splicing factor ASF/SF2. 1992 , 6, 2569-79		77
441	Identification of a negative regulatory element involved in tissue-specific expression of mouse renin genes. 1992 , 89, 885-9		37
440	Independent deposition of heterogeneous nuclear ribonucleoproteins and small nuclear ribonucleoprotein particles at sites of transcription. 1992 , 89, 8409-13		46
439	Endogenous retroviral sequences are required for tissue-specific expression of a human salivary amylase gene. 1992 , 6, 1457-65		203
438	Functional role of the ultraviolet light responsive element (URE; TGACAACA) in the transcription and replication of polyoma DNA. <i>Nucleic Acids Research</i> , 1992 , 20, 4305-10	20.1	18

437	Analysis of spliceosome assembly and the structure of a regulated intron in <i>Drosophila</i> in vitro splicing extracts. <i>Nucleic Acids Research</i> , 1992 , 20, 5719-27	20.1	18
436	An ATP-independent U2 small nuclear ribonucleoprotein particle/precursor mRNA complex requires both splice sites and the polypyrimidine tract. 1992 , 89, 5482-6		35
435	Identification of a peroxisome proliferator-responsive element upstream of the gene encoding rat peroxisomal enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase. 1992 , 89, 7541-5		217
434	Myocardial activation of the human cardiac alpha-actin promoter by helix-loop-helix proteins. 1992 , 89, 4047-51		96
433	Antisense probes targeted to an internal domain in U2 snRNP specifically inhibit the second step of pre-mRNA splicing. <i>Nucleic Acids Research</i> , 1992 , 20, 4457-64	20.1	30
432	Activation of human papillomavirus type 18 E6-E7 oncogene expression by transcription factor Sp1. <i>Nucleic Acids Research</i> , 1992 , 20, 6701-6	20.1	47
431	Heat shock gene regulation by nascent polypeptides and denatured proteins: hsp70 as a potential autoregulatory factor. 1992 , 117, 1151-9		299
430	A multiple cytokine- and second messenger-responsive element in the enhancer of the human interleukin-6 gene: similarities with c-fos gene regulation. 1989 , 9, 5537-5547		54
429	A B-cell-specific nuclear protein that binds to DNA sites 5' to immunoglobulin S alpha tandem repeats is regulated during differentiation. 1989 , 9, 5594-5601		19
428	The C-terminal domain of the largest subunit of RNA polymerase II and transcription initiation. 1989 , 9, 5750-5753		6
427	Functional identification of regulatory elements within the promoter region of platelet-derived growth factor 2. 1989 , 9, 396-405		12
426	NF-kappa B binds within a region required for B-cell-specific expression of major histocompatibility complex class II gene E alpha d. 1989 , 9, 844-846		23
425	Regulation of 2',5'-oligoadenylate synthetase gene expression by interferons and platelet-derived growth factor. 1989 , 9, 1060-1068		5
424	Cell-specific activity of the modulator region in the human cytomegalovirus major immediate-early gene. 1989 , 9, 1342-1345		28
423	Interleukin 1 and cyclic AMP induce kappa immunoglobulin light-chain expression via activation of an NF-kappa B-like DNA-binding protein. 1989 , 9, 959-964		81
422	Role of poly(A) polymerase in the cleavage and polyadenylation of mRNA precursor. 1989 , 9, 1435-1444		20
421	Sea urchin early and late H4 histone genes bind a specific transcription factor in a stable preinitiation complex. 1989 , 9, 1476-1487		6
420	Forskolin inducibility and tissue-specific expression of the fibronectin promoter. 1989 , 9, 1498-1506		34

4 ¹⁹	Purification and characterization of multiple nuclear factors that bind to the TAX-inducible enhancer within the human T-cell leukemia virus type 1 long terminal repeat. 1989 , 9, 1733-1745	47
4 ¹⁸	Interaction of nuclear proteins with a positive cis-acting element of rat embryonic myosin heavy-chain promoter: identification of a new transcriptional factor. 1989 , 9, 1839-1849	13
4 ¹⁷	Factor interactions with the simian virus 40 early pre-mRNA influence branch site selection and alternative splicing. 1989 , 9, 2007-2017	8
4 ¹⁶	Cross-binding of factors to functionally different promoter elements in c-fos and skeletal actin genes. 1989 , 9, 2191-2201	83
4 ¹⁵	Remarkable intron and exon sequence conservation in human and mouse homeobox Hox 1.3 genes. 1989 , 9, 2273-2278	8
4 ¹⁴	In vitro activation and nuclear translocation of NF-kappa B catalyzed by cyclic AMP-dependent protein kinase and protein kinase C. 1989 , 9, 2424-2430	171
4 ¹³	Major histocompatibility complex class I genes in murine fibrosarcoma IC9 are down regulated at the level of the chromatin structure. 1989 , 9, 3136-3142	5
4 ¹²	A lipopolysaccharide-induced DNA-binding protein for a class II gene in B cells is distinct from NF-kappa B. 1989 , 9, 3184-3192	15
4 ¹¹	Alteration of a single nucleotide allows efficient binding of H2TF1/KBF1 to the immunoglobulin kappa enhancer B motif. 1989 , 9, 3548-3552	15
4 ¹⁰	Identification of HeLa cell nuclear factors that bind to and activate the early promoter of human polyomavirus BK in vitro. 1989 , 9, 3821-3828	11
4 ⁰⁹	Unusual levels of heat shock element-binding activity in embryonal carcinoma cells. 1989 , 9, 3888-3896	17
4 ⁰⁸	Negative transcriptional regulatory element that functions in embryonal carcinoma cells. 1989 , 9, 4032-4037	8
4 ⁰⁷	The differentiation-specific factor CDP/Cut represses transcription and replication of human papillomaviruses through a conserved silencing element. 2000 , 74, 401-10	31
4 ⁰⁶	Non-radioactive detection of m3G capped RNAs by immunoblotting. <i>Nucleic Acids Research</i> , 1994 , 22, 252-3	20.1 1
4 ⁰⁵	The carboxyl terminus of vertebrate poly(A) polymerase interacts with U2AF 65 to couple 3'-end processing and splicing. 2000 , 14, 403-13	114
4 ⁰⁴	Pre-mRNA splicing alters mRNP composition: evidence for stable association of proteins at exon-exon junctions. 2000 , 14, 1098-108	247
4 ⁰³	Sequence-specific functions of the early palindrome domain within the SV40 core origin of replication. <i>Nucleic Acids Research</i> , 1989 , 17, 9279-89	20.1 1
4 ⁰²	Experimental evidence for RNA trans-splicing in mammalian cells. 1995 , 14, 3226-35	22

401	The human splicing factors ASF/SF2 and SC35 possess distinct, functionally significant RNA binding specificities. 1995 , 14, 3540-51	169
400	In vitro reconstitution of mammalian U2 and U5 snRNPs active in splicing: Sm proteins are functionally interchangeable and are essential for the formation of functional U2 and U5 snRNPs. 1995 , 14, 4010-21	49
399	The cellular factor TRP-185 regulates RNA polymerase II binding to HIV-1 TAR RNA. 1995 , 14, 5995-6009	18
398	The HMG box-containing nucleolar transcription factor UBF interacts with a specific subunit of RNA polymerase I. 1994 , 13, 190-9	32
397	The length and the secondary structure of the D-stem of human selenocysteine tRNA are the major identity determinants for serine phosphorylation. 1994 , 13, 241-8	14
396	A common factor regulates both Th1- and Th2-specific cytokine gene expression. 1994 , 13, 625-33	38
395	IDX-1: a new homeodomain transcription factor expressed in rat pancreatic islets and duodenum that transactivates the somatostatin gene. 1994 , 13, 1145-56	117
394	The E6/E7 promoter of extrachromosomal HPV16 DNA in cervical cancers escapes from cellular repression by mutation of target sequences for YY1. 1994 , 13, 1460-6	51
393	Differential transactivation potential of Oct1 and Oct2 is determined by additional B cell-specific activities. 1994 , 13, 1654-63	16
392	The site of 3' end formation of histone messenger RNA is a fixed distance from the downstream element recognized by the U7 snRNP. 1994 , 13, 2432-40	40
391	Characterization and cloning of the human splicing factor 9G8: a novel 35 kDa factor of the serine/arginine protein family. 1994 , 13, 2639-49	63
390	A cell-specific enhancer far upstream of the mouse tyrosinase gene confers high level and copy number-related expression in transgenic mice. 1994 , 13, 3083-93	29
389	Targeted disruption of the c-fos gene demonstrates c-fos-dependent and -independent pathways for gene expression stimulated by growth factors or oncogenes. 1994 , 13, 3094-103	69
388	TATA-binding protein-associated factor(s) in TFIID function through the initiator to direct basal transcription from a TATA-less class II promoter. 1994 , 13, 3115-26	60
387	The exchange of the discriminator base A73 for G is alone sufficient to convert human tRNA(Leu) into a serine-acceptor in vitro. 1994 , 13, 3166-9	23
386	The thyroid transcription factor-1 gene is a candidate target for regulation by Hox proteins. 1994 , 13, 3339-47	20
385	Two distinct mechanisms contribute to the constitutive activation of RelB in lymphoid cells. 1994 , 13, 4060-9	30
384	The Spi-1/PU.1 and Spi-B ets family transcription factors and the recombination signal binding protein RBP-J kappa interact with an Epstein-Barr virus nuclear antigen 2 responsive cis-element. 1994 , 13, 5624-32	71

383	MAZ-dependent termination between closely spaced human complement genes. 1994 , 13, 5656-67	51
382	Response to cAMP levels of the Epstein-Barr virus EBNA2-inducible LMP1 oncogene and EBNA2 inhibition of a PP1-like activity. 1994 , 13, 6041-51	11
381	Assembly and disassembly of spliceosomes along a specific pre-messenger RNP fiber. 1994 , 13, 6052-61	32
380	Hemoglobin switching in man and chicken is mediated by a heteromeric complex between the ubiquitous transcription factor CP2 and a developmentally specific protein. 1995 , 14, 97-105	38
379	NF-E2 and GATA binding motifs are required for the formation of DNase I hypersensitive site 4 of the human beta-globin locus control region. 1995 , 14, 106-16	55
378	The bovine papillomavirus 1 E2 protein contains two activation domains: one that interacts with TBP and another that functions after TBP binding. 1995 , 14, 329-40	38
377	Purification of the sequence-specific transcription factor CTCBF, involved in the control of human collagen IV genes: subunits with homology to Ku antigen. 1995 , 14, 791-800	23
376	GATA1 and YY1 are developmental repressors of the human epsilon-globin gene. 1995 , 14, 801-9	46
375	Interleukin-3 signals through multiple isoforms of Stat5. 1995 , 14, 1402-11	64
374	Electron microscopy of U1 small nuclear ribonucleoprotein particles: shape of the particle and position of the 5' RNA terminus. 1989 , 8, 277-86	26
373	In vitro synthesis of vertebrate U1 snRNA. 1989 , 8, 287-92	29
372	Non-enzymatic excision of pre-tRNA introns?. 1989 , 8, 293-300	9
371	Human immunodeficiency virus type 1 LTR TATA and TAR region sequences required for transcriptional regulation. 1989 , 8, 765-78	146
370	Transcription in the reverse orientation at either terminus of the adenovirus type 5 genome. 1989 , 8, 863-8	7
369	Double-stranded RNA activates binding of NF-kappa B to an inducible element in the human beta-interferon promoter. 1989 , 8, 1129-38	134
368	Two distinct cellular phosphoproteins bind to the c-fos serum response element. 1989 , 8, 1785-92	78
367	Interactions of alpha- and gamma-interferon in the transcriptional regulation of the gene encoding a guanylate-binding protein. 1989 , 8, 2009-14	78
366	A thyroid-specific nuclear protein essential for tissue-specific expression of the thyroglobulin promoter. 1989 , 8, 2537-42	86

365	A family of octamer-specific proteins present during mouse embryogenesis: evidence for germline-specific expression of an Oct factor. 1989 , 8, 2543-50	234
364	Octamer binding proteins confer transcriptional activity in early mouse embryogenesis. 1989 , 8, 2551-7	104
363	Repression of c-fos transcription is mediated through p67SRF bound to the SRE. 1989 , 8, 2567-74	86
362	Transcriptional enhancement by upstream activators is brought about by different molecular mechanisms for class I and II RNA polymerase genes. 1989 , 8, 3011-7	4
361	Evidence from complementation assays in vitro that U5 snRNP is required for both steps of mRNA splicing. 1989 , 8, 3105-12	16
360	U3, U8 and U13 comprise a new class of mammalian snRNPs localized in the cell nucleolus. 1989 , 8, 3113-9	214
359	Suppression of MHC class I gene expression by N-myc through enhancer inactivation. 1989 , 8, 3351-5	43
358	Identification of the RNA binding segment of human U1 A protein and definition of its binding site on U1 snRNA. 1989 , 8, 4163-70	153
357	Mapping U2 snRNP--pre-mRNA interactions using biotinylated oligonucleotides made of 2'-OMe RNA. 1989 , 8, 4171-8	35
356	An abundant U6 snRNP found in germ cells and embryos of <i>Xenopus laevis</i> . 1989 , 8, 4179-87	50
355	The SV40 TC-II(kappa B) enhanson binds ubiquitous and cell type specifically inducible nuclear proteins from lymphoid and non-lymphoid cell lines. 1989 , 8, 4215-27	28
354	The octamer-binding proteins form multi-protein--DNA complexes with the HSV alpha TIF regulatory protein. 1989 , 8, 4229-38	123
353	TPA can overcome the requirement for Ela and together act synergistically in stimulating expression of the adenovirus E111 promoter. 1989 , 8, 4239-50	11
352	NF-kappa B activation of the cytomegalovirus enhancer is mediated by a viral transactivator and by T cell stimulation. 1989 , 8, 4251-8	151
351	DNA replication origin and transcriptional enhancer in c-myc gene share the c-myc protein binding sequences. 1989 , 8, 4273-9	39
350	The Epstein-Barr virus nuclear antigen 2 interacts with an EBNA2 responsive cis-element of the terminal protein 1 gene promoter. 1993 , 12, 167-75	86
349	The NF-kappa B precursor p105 and the proto-oncogene product Bcl-3 are I kappa B molecules and control nuclear translocation of NF-kappa B. 1993 , 12, 213-22	67
348	Interferon-beta promoters contain a DNA element that acts as a position-independent silencer on the NF-kappa B site. 1993 , 12, 451-9	22

347	Distinct TFIID complexes mediate the effect of different transcriptional activators. 1993 , 12, 489-99	96
346	DNA methylation in the Alu sequences of diploid and haploid primary human cells. 1993 , 12, 1141-51	78
345	Insect immunity. Two 17 bp repeats nesting a kappa B-related sequence confer inducibility to the dipterin gene and bind a polypeptide in bacteria-challenged <i>Drosophila</i> . 1993 , 12, 1561-8	38
344	H ₂ O ₂ and antioxidants have opposite effects on activation of NF-kappa B and AP-1 in intact cells: AP-1 as secondary antioxidant-responsive factor. 1993 , 12, 2005-15	240
343	A novel transcription factor, OB2-1, is required for overexpression of the proto-oncogene c-erbB-2 in mammary tumour lines. 1993 , 12, 2369-75	31
342	Unique TATA-binding protein-containing complexes and cofactors involved in transcription by RNA polymerases II and III. 1993 , 12, 2749-62	90
341	Cytokine-inducible expression in endothelial cells of an I kappa B alpha-like gene is regulated by NF kappa B. 1993 , 12, 2773-9	97
340	Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF. 1993 , 12, 3095-104	121
339	Identity determinants of human tRNA(Ser): sequence elements necessary for serylation and maturation of a tRNA with a long extra arm. 1993 , 12, 3333-8	35
338	Pathways for selection of 5' splice sites by U1 snRNPs and SF2/ASF. 1993 , 12, 3607-17	108
337	NF-kappa B activates the HIV promoter in neurons. 1993 , 12, 4261-7	20
336	Repression of adenovirus E1A enhancer activity by a novel zinc finger-containing DNA-binding protein related to the GLI-Kruppel protein. 1993 , 12, 4985-92	11
335	Promoter analysis of the gene encoding the I kappa B-alpha/MAD3 inhibitor of NF-kappa B: positive regulation by members of the rel/NF-kappa B family. 1993 , 12, 5043-9	114
334	LEF-1 contains an activation domain that stimulates transcription only in a specific context of factor-binding sites. 1993 , 12, 4667-76	56
333	Synthesis of reinitiated transcripts by mammalian RNA polymerase II is controlled by elongation factor SII. 1993 , 12, 4677-84	8
332	Functional analysis of pre-mRNA splicing factor SF2/ASF structural domains. 1993 , 12, 4715-26	118
331	Identification and properties of the RNA-dependent RNA polymerase of hepatitis C virus. 1996 , 15, 12-22	223
330	Editing of the GLuR-B ion channel RNA in vitro by recombinant double-stranded RNA adenosine deaminase. 1996 , 15, 34-45	34

329	Cross-linking CD40 on B cells preferentially induces stress-activated protein kinases rather than mitogen-activated protein kinases. 1996 , 15, 92-101	42
328	Interferon-alpha-induced phosphorylation and activation of cytosolic phospholipase A2 is required for the formation of interferon-stimulated gene factor three. 1996 , 15, 1566-71	10
327	The p38/RK mitogen-activated protein kinase pathway regulates interleukin-6 synthesis response to tumor necrosis factor. 1996 , 15, 1914-23	119
326	Tissue-specific factors additively increase the probability of the all-or-none formation of a hypersensitive site. 1996 , 15, 2496-507	67
325	pX, the HBV-encoded coactivator, interacts with components of the transcription machinery and stimulates transcription in a TAF-independent manner. 1996 , 15, 3413-20	34
324	The HeLa 200 kDa U5 snRNP-specific protein and its homologue in <i>Saccharomyces cerevisiae</i> are members of the DEXH-box protein family of putative RNA helicases. 1996 , 15, 4001-15	63
323	Purification and biochemical heterogeneity of the mammalian SWI-SNF complex. 1996 , 15, 5370-82	341
322	An IL-2 response element in the human IL-2 receptor alpha chain promoter is a composite element that binds Stat5, Elf-1, HMG-I(Y) and a GATA family protein. 1996 , 15, 5627-35	36
321	hPop1: an autoantigenic protein subunit shared by the human RNase P and RNase MRP ribonucleoproteins. 1996 , 15, 5936-48	48
320	Replication of poliovirus in <i>Xenopus</i> oocytes requires two human factors. 1996 , 15, 5988-98	23
319	Novel downstream elements upregulate transcription initiated from an Epstein-Barr virus latent promoter. 1991 , 10, 143-51	11
318	Mammalian nuclei contain foci which are highly enriched in components of the pre-mRNA splicing machinery. 1991 , 10, 195-206	105
317	Biochemical characterization of U2 snRNP auxiliary factor: an essential pre-mRNA splicing factor with a novel intranuclear distribution. 1991 , 10, 207-14	119
316	The interaction of ICP4 with cell/infected-cell factors and its state of phosphorylation modulate differential recognition of leader sequences in herpes simplex virus DNA. 1991 , 10, 397-406	17
315	A novel splicing factor is an integral component of 200S large nuclear ribonucleoprotein (InRNP) particles. 1991 , 10, 425-32	19
314	Cytoplasmic activation of GAF, an IFN-gamma-regulated DNA-binding factor. 1991 , 10, 927-32	132
313	N-myc disrupts protein kinase C-mediated signal transduction in neuroblastoma. 1991 , 10, 1119-25	18
312	Pax-3, a novel murine DNA binding protein expressed during early neurogenesis. 1991 , 10, 1135-47	211

311	The 5'HS2 of the globin locus control region enhances transcription through the interaction of a multimeric complex binding at two functionally distinct NF-E2 binding sites. 1991 , 10, 1391-8	82
310	Three protein factors (SF1, SF3 and U2AF) function in pre-splicing complex formation in addition to snRNPs. 1991 , 10, 1503-9	47
309	TFIID is required for in vitro transcription of the human U6 gene by RNA polymerase III. 1991 , 10, 1853-62	70
308	Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. 1991 , 10, 2247-58	804
307	Affinity purification of transcription factor IIA from HeLa cell nuclear extracts. 1991 , 10, 2305-10	17
306	An intact Box C sequence in the U3 snRNA is required for binding of fibrillarin, the protein common to the major family of nucleolar snRNPs. 1991 , 10, 2645-51	86
305	Poliovirus proteinase 3C converts an active form of transcription factor IIIc to an inactive form: a mechanism for inhibition of host cell polymerase III transcription by poliovirus. 1991 , 10, 2941-7	81
304	In vitro deadenylation of mammalian mRNA by a HeLa cell 3' exonuclease. 1991 , 10, 3067-71	40
303	A novel lineage-specific nuclear factor regulates mb-1 gene transcription at the early stages of B cell differentiation. 1991 , 10, 3409-17	58
302	Changes in NF-kappa B and ISGF3 DNA binding activities are responsible for differences in MHC and beta-2-microglobulin gene expression in Ad5- versus Ad12-transformed cells. 1991 , 10, 4169-75	18
301	Transcriptional termination between the closely linked human complement genes C2 and factor B: common termination factor for C2 and c-myc?. 1991 , 10, 4197-207	48
300	Interferon-induced binding of nuclear factors to promoter elements of the 2-5A synthetase gene. 1988 , 7, 751-9	72
299	The role of the polypyrimidine stretch at the SV40 early pre-mRNA 3' splice site in alternative splicing. 1988 , 7, 809-17	54
298	Interaction of a protein from rat liver nuclei with cruciform DNA. 1988 , 7, 843-9	39
297	Two different liver-specific factors stimulate in vitro transcription from the human alpha 1-antitrypsin promoter. 1988 , 7, 2075-87	123
296	Purification of the human immunodeficiency virus type 1 enhancer and TAR binding proteins EBP-1 and UBP-1. 1988 , 7, 2117-30	86
295	The -117 mutation in Greek HPFH affects the binding of three nuclear factors to the CCAAT region of the gamma-globin gene. 1988 , 7, 3099-107	61
294	Nuclear factor NF-kappa B can interact functionally with its cognate binding site to provide lymphoid-specific promoter function. 1988 , 7, 3109-13	38

293	Purification of a NF1-like DNA-binding protein from rat liver and cloning of the corresponding cDNA. 1988 , 7, 3115-23	103
292	Evidence for a nuclear factor(s), IRF-1, mediating induction and silencing properties to human IFN-beta gene regulatory elements. 1988 , 7, 3397-405	122
291	Heat shock but not other stress inducers leads to the disruption of a sub-set of snRNPs and inhibition of in vitro splicing in HeLa cells. 1988 , 7, 3509-18	80
290	RNA binding specificity of hnRNP proteins: a subset bind to the 3' end of introns. 1988 , 7, 3519-29	130
289	A zinc-responsive factor interacts with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I gene. 1988 , 7, 3763-70	94
288	Cells resistant to interferon are defective in activation of a promoter-binding factor. 1988 , 7, 3779-83	29
287	Proteins bound at adjacent DNA elements act synergistically to regulate human proenkephalin cAMP inducible transcription. 1988 , 7, 3793-805	91
286	Wheat germ splicing endonuclease is highly specific for plant pre-tRNAs. 1988 , 7, 3823-8	21
285	The human beta-globin promoter; nuclear protein factors and erythroid specific induction of transcription. 1988 , 7, 4203-12	95
284	Complementation between two cell lines lacking kappa enhancer activity: implications for the developmental control of immunoglobulin transcription. 1988 , 7, 4213-20	18
283	Identification of a novel lymphoid specific octamer binding protein (OTF-2B) by proteolytic clipping bandshift assay (PCBA). 1988 , 7, 4221-9	126
282	Direct combinatorial interaction between a herpes simplex virus regulatory protein and a cellular octamer-binding factor mediates specific induction of virus immediate-early gene expression. 1988 , 7, 4231-8	88
281	The promoter of Alzheimer's disease amyloid A4 precursor gene. 1988 , 7, 2807-13	60
280	A cell-specific activator in the Xenopus A2 vitellogenin gene: promoter elements functioning with rat liver nuclear extracts. 1988 , 7, 2495-501	20
279	Antisense RNA inhibits splicing of pre-mRNA in vitro. 1988 , 7, 2523-32	30
278	Physical identification of branched intron side-products of splicing in Trypanosoma brucei. 1988 , 7, 2539-45	16
277	Two distinct factors interact with the promoter regions of several liver-specific genes. 1988 , 7, 1711-9	87
276	Tissue specific trans-acting factor interaction with proximal rat prolactin gene promoter sequences. 1988 , 7, 1721-33	5

275	Interaction of a nodule specific, trans-acting factor with distinct DNA elements in the soybean leghaemoglobin lbc(3) 5' upstream region. 1988 , 7, 1265-71	74
274	Enhancer-like interferon responsive sequences of the human and murine (2'-5') oligoadenylate synthetase gene promoters. 1988 , 7, 1411-9	68
273	Processing at immunoglobulin polyadenylation sites in lymphoid cell extracts. 1988 , 7, 1421-9	5
272	Trypanosome trans-splicing utilizes 2'-5' branches and a corresponding debranching activity. 1988 , 7, 1431-7	18
271	The mouse ribosomal gene terminator consists of three functionally separable sequence elements. 1988 , 7, 1497-502	24
270	The specific DNA binding activity of the dioxin receptor is modulated by the 90 kd heat shock protein. 1990 , 9, 69-76	56
269	Tissue-specific nuclear factors mediate expression of the CD3 delta gene during T cell development. 1990 , 9, 109-15	16
268	Identification of sequences responsible for positive and negative regulation by E1A in the promoter of H-2Kbm1 class I MHC gene. 1990 , 9, 127-35	22
267	In vitro splicing of mutually exclusive exons from the chicken beta-tropomyosin gene: role of the branch point location and very long pyrimidine stretch. 1990 , 9, 241-9	39
266	Discrete domains of human U6 snRNA required for the assembly of U4/U6 snRNP and splicing complexes. 1990 , 9, 251-5	22
265	Amino-terminal domain of NF1 binds to DNA as a dimer and activates adenovirus DNA replication. 1990 , 9, 559-66	64
264	Expression of a human 7S K RNA gene in vivo requires a novel pol III upstream element. 1990 , 9, 711-8	21
263	Transcription factor E4TF1 contains two subunits with different functions. 1990 , 9, 841-7	26
262	Extinction of Ig genes expression in myeloma x fibroblast somatic cell hybrids is accompanied by repression of the oct-2 gene encoding a B-cell specific transcription factor. 1990 , 9, 849-55	24
261	Protein recognition sites in polyomavirus enhancer: formation of a novel site for NF-1 factor in an enhancer mutant and characterization of a site in the enhancer D domain. 1990 , 9, 947-55	14
260	Tax1 induction of the HTLV-I 21 bp enhancer requires cooperation between two cellular DNA-binding proteins. 1990 , 9, 957-64	41
259	Synergistic interaction between interferon-alpha and interferon-gamma through induced synthesis of one subunit of the transcription factor ISGF3. 1990 , 9, 1105-11	93
258	The presence of a DNA binding complex correlates with E beta class II MHC gene expression. 1990 , 9, 1543-9	11

257	A mitigator sequence in the downstream region of the major late promoter of adenovirus type 12 DNA. 1990 , 9, 1615-23	12
256	A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family. 1990 , 9, 1897-906	475
255	The beta-globin dominant control region: hypersensitive site 2. 1990 , 9, 2159-67	123
254	Detailed analysis of the site 3 region of the human beta-globin dominant control region. 1990 , 9, 2169-77	131
253	A role for the adenovirus inducible E2F transcription factor in a proliferation dependent signal transduction pathway. 1990 , 9, 2179-84	106
252	A growth-dependent transcription initiation factor (TIF-IA) interacting with RNA polymerase I regulates mouse ribosomal RNA synthesis. 1990 , 9, 2857-63	64
251	A novel upstream element compensates for an ineffectual octamer motif in an immunoglobulin V kappa promoter. 1990 , 9, 3109-17	21
250	A 3' transcriptional enhancer regulates tissue-specific expression of the human CD2 gene. 1990 , 9, 3129-36	25
249	Splicing factor SF4 is dispensable for the assembly of a functional splicing complex and participates in the subsequent steps of the splicing reaction. 1990 , 9, 4119-26	9
248	Lymphoid specific gene expression of the adenovirus early region 3 promoter is mediated by NF-kappa B binding motifs. 1990 , 9, 4435-42	23
247	The SIF binding element confers sis/PDGF inducibility onto the c-fos promoter. 1990 , 9, 4477-84	169
246	A human and a plant intron-containing tRNA ^{Tyr} gene are both transcribed in a HeLa cell extract but spliced along different pathways. 1987 , 6, 35-41	38
245	Formation of mRNA 3' termini: stability and dissociation of a complex involving the AAUAAA sequence. 1987 , 6, 177-86	46
244	Xenopus cytoskeletal actin and human c-fos gene promoters share a conserved protein-binding site. 1987 , 6, 667-73	96
243	RNA 3' processing regulates histone mRNA levels in a mammalian cell cycle mutant. A processing factor becomes limiting in G1-arrested cells. 1987 , 6, 1721-6	74
242	Two spliceosomes can form simultaneously and independently on synthetic double-intron messenger RNA precursors. 1987 , 6, 1747-55	12
241	Promoter interaction of the E1A-inducible factor E2F and its potential role in the formation of a multi-component complex. 1987 , 6, 2061-8	72
240	Linear oligomeric potato spindle tuber viroid (PSTV) RNAs are accurately processed in vitro to the monomeric circular viroid proper when incubated with a nuclear extract from healthy potato cells. 1987 , 6, 2173-83	17

239	Identification and purification of a polypeptide that binds to the c-fos serum response element. 1987 , 6, 2711-7	192
238	Cis- and trans-acting elements responsible for the cell-specific expression of the human alpha 1-antitrypsin gene. 1987 , 6, 2759-66	83
237	A cell-free plant extract for accurate pre-tRNA processing, splicing and modification. 1987 , 6, 2811-8	39
236	One cell-specific and three ubiquitous nuclear proteins bind in vitro to overlapping motifs in the domain B1 of the SV40 enhancer. 1987 , 6, 3005-13	62
235	In vitro binding of cell-specific and ubiquitous nuclear proteins to the octamer motif of the SV40 enhancer and related motifs present in other promoters and enhancers. 1987 , 6, 3015-25	76
234	Functional characterization of X. laevis U5 snRNA genes. 1987 , 6, 3071-8	36
233	A novel promoter in the mouse rDNA spacer is active in vivo and in vitro. 1987 , 6, 3487-92	36
232	Interactions of cellular proteins involved in the transcriptional regulation of the human immunodeficiency virus. 1987 , 6, 3761-70	132
231	Cleavage and polyadenylation of messenger RNA precursors in vitro occurs within large and specific 3' processing complexes. 1987 , 6, 4159-68	53
230	Analysis of mRNA 3' end formation by modification interference: the only modifications which prevent processing lie in AAUAAA and the poly(A) site. 1987 , 6, 4177-84	39
229	Specific pre-cleavage and post-cleavage complexes involved in the formation of SV40 late mRNA 3' termini in vitro. 1987 , 6, 4185-92	38
228	Cell type-specificity elements of the immunoglobulin heavy chain gene enhancer. 1987 , 6, 1323-30	154
227	A cellular transcription factor E4F1 interacts with an E1a-inducible enhancer and mediates constitutive enhancer function in vitro. 1987 , 6, 1345-53	106
226	The two intervening sequences of human beta- and gamma-globin pre-mRNAs are excised in a preferred temporal order in vitro. 1985 , 4, 1991-6	15
225	Cell type-specific transcriptional enhancement in vitro requires the presence of trans-acting factors. 1985 , 4, 3005-13	77
224	Specific switching on of silent egg protein genes in vitro by an S-100 fraction in isolated nuclei from male Xenopus. 1985 , 4, 3253-8	2
223	Specific interaction between a transcription factor and the upstream element of the adenovirus-2 major late promoter. 1985 , 4, 3563-70	89
222	Purification of a protein required for the splicing of pre-mRNA and its separation from the lariat debranching enzyme. 1985 , 4, 3571-81	34

221	Functional domains of the <i>Xenopus laevis</i> 5S gene promoter. 1985 , 4, 3751-6	23
220	The characterization of the promoter of the gene encoding the p50 subunit of NF-kappa B indicates that it participates in its own regulation. 1992 , 11, 195-203	104
219	A novel complex between the p65 subunit of NF-kappa B and c-Rel binds to a DNA element involved in the phorbol ester induction of the human urokinase gene. 1992 , 11, 205-13	56
218	The upstream region of the human homeobox gene HOX3D is a target for regulation by retinoic acid and HOX homeoproteins. 1992 , 11, 265-77	32
217	Repression of genes by DNA methylation depends on CpG density and promoter strength: evidence for involvement of a methyl-CpG binding protein. 1992 , 11, 327-33	154
216	Reconstituted mammalian U4/U6 snRNP complements splicing: a mutational analysis. 1992 , 11, 345-59	33
215	Identification and characterization of Ref-1, a nuclear protein that facilitates AP-1 DNA-binding activity. 1992 , 11, 653-65	218
214	Dissection of the interferon gamma-MHC class II signal transduction pathway reveals that type I and type II interferon systems share common signalling component(s). 1992 , 11, 1351-63	22
213	RXR alpha, a promiscuous partner of retinoic acid and thyroid hormone receptors. 1992 , 11, 1409-18	144
212	H-2RIIBP (RXR beta) heterodimerization provides a mechanism for combinatorial diversity in the regulation of retinoic acid and thyroid hormone responsive genes. 1992 , 11, 1419-35	147
211	Identification of NF-jun, a novel inducible transcription factor that regulates c-jun gene transcription. 1992 , 11, 1479-86	11
210	Sequence and structure requirements for the biosynthesis of pseudouridine (psi 35) in plant pre-tRNA(Tyr). 1992 , 11, 1907-12	12
209	The nucleolar transcription factor mUBF is phosphorylated by casein kinase II in the C-terminal hyperacidic tail which is essential for transactivation. 1992 , 11, 2211-8	75
208	Transcription factor AP-2 activates gene expression of HTLV-I. 1992 , 11, 2573-81	12
207	Identification of a stage selector element in the human gamma-globin gene promoter that fosters preferential interaction with the 5' HS2 enhancer when in competition with the beta-promoter. 1992 , 11, 2961-9	48
206	Cell-specific transcription of the peripherin gene in neuronal cell lines involves a cis-acting element surrounding the TATA box. 1992 , 11, 2971-80	14
205	Hepatitis B virus transactivator MHBst: activation of NF-kappa B, selective inhibition by antioxidants and integral membrane localization. 1992 , 11, 2991-3001	43
204	An autoregulatory element of the murine Hox-4.2 gene. 1992 , 11, 3673-80	35

203	Hold back of RNA polymerase II at the transcription start site mediates down-regulation of c-myc in vivo. 1992 , 11, 3307-14	88
202	Phosphorylation of CREB affects its binding to high and low affinity sites: implications for cAMP induced gene transcription. 1992 , 11, 3337-46	94
201	TAR-independent transactivation by Tat in cells derived from the CNS: a novel mechanism of HIV-1 gene regulation. 1992 , 11, 3395-403	42
200	Erythroid-specific activity of the glycophorin B promoter requires GATA-1 mediated displacement of a repressor. 1992 , 11, 4095-102	28
199	Activation of HIV-1 pre-mRNA 3' processing in vitro requires both an upstream element and TAR. 1992 , 11, 4419-28	47
198	Identification of a negative regulatory domain in the human papillomavirus type 18 promoter: interaction with the transcriptional repressor YY1. 1992 , 11, 4607-17	66
197	Hormone-regulated v-rel estrogen receptor fusion protein: reversible induction of cell transformation and cellular gene expression. 1992 , 11, 4641-52	26
196	Is actin a transcription initiation factor for RNA polymerase B?. 1984 , 3, 2363-71	80
195	Stimulation of in vitro transcription from the SV40 early promoter by the enhancer involves a specific trans-acting factor. 1984 , 3, 3129-33	121
194	Human genes and pseudogenes for the 7SL RNA component of signal recognition particle. 1984 , 3, 3303-10	69
193	In vitro transcription by Xenopus oocytes RNA polymerase III requires a DNA topoisomerase II activity. 1986 , 5, 151-5	7
192	Generation of histone mRNA 3' ends by endonucleolytic cleavage of the pre-mRNA in a snRNP-dependent in vitro reaction. 1986 , 5, 1319-26	103
191	DNA topoisomerase II cleaves at specific sites in the 5' flanking region of c-fos proto-oncogenes in vitro. 1986 , 5, 2257-65	15
190	Accurate in vitro splicing of two pre-mRNA plant introns in a HeLa cell nuclear extract. 1986 , 5, 2749-58	52
189	Proteins that bind to RNA polymerase II are required for accurate initiation of transcription at the adenovirus 2 major late promoter. 1986 , 5, 2923-30	31
188	UV-A-induced decrease in nuclear factor-kappaB activity in human keratinocytes. 1999 , 338 (Pt 3), 607-13	7
187	Roles of an Ets motif and a novel CACGAC direct repeat in transcription of the murine dihydrolipoamide dehydrogenase (Dld) gene. 1999 , 338 (Pt 3), 667-75	
186	Positive and negative regulatory elements in the upstream region of the rat Cu/Zn-superoxide dismutase gene. 1999 , 339 (Pt 2), 335-41	6

185	Efficient transcription of the human angiotensin II type 2 receptor gene requires intronic sequence elements. 1999 , 340 (Pt 1), 17-24	5
184	Glucocorticoid inhibition of human SP-A1 promoter activity in NCI-H441 cells. 1999 , 340 (Pt 1), 69-76	4
183	Structural and functional characterization of the mouse p8 gene: promotion of transcription by the CAAT-enhancer binding protein alpha (C/EBPalpha) and C/EBPbeta trans-acting factors involves a C/EBP cis-acting element and other regions of the promoter. 1999 , 343 Pt 2, 377-83	16
182	Mitogen-activated protein kinase mediates erythropoietin-induced phosphorylation of the TAL1/SCL transcription factor in murine proerythroblasts. 1999 , 343 Pt 3, 615-20	6
181	Functional analysis of the promoter region of the human phosphotyrosine phosphatase activator gene: Yin Yang 1 is essential for core promoter activity. 1999 , 344 Pt 3, 755-63	2
180	Transcription from the P2 promoter of the growth hormone receptor gene involves members of the Sp transcription factor family. 1999 , 344 Pt 3, 867-72	3
179	Mitochondrial 3-hydroxy-3-methylglutaryl-CoA synthase promoter contains a CREB binding site that regulates cAMP action in Caco-2 cells. 2000 , 345 Pt 2, 201-6	4
178	Export and transport of tRNA are coupled to a multi-protein complex. 2000 , 346 Pt 1, 107-15	10
177	Regulation of the gene promoter for extracellular signal-regulated protein kinase 2 by transcription factors NF-Y and Sp3. 2000 , 347 Pt 1, 155-61	4
176	The human vasoactive intestinal peptide/pituitary adenylate cyclase-activating peptide receptor 1 (VPAC1) promoter: characterization and role in receptor expression during enterocytic differentiation of the colon cancer cell line Caco-2Cl.20. 2000 , 347 Pt 3, 623-32	
175	Identification of an involucrin promoter transcriptional response element with activity restricted to keratinocytes. 2000 , 348 Pt 1, 45-53	5
174	cAMP attenuates interleukin-1-stimulated macrophage colony-stimulating factor (M-CSF) expression. 2000 , 350 Pt 1, 115-22	6
173	Insulin effects on sterol regulatory-element-binding protein-1c (SREBP-1c) transcriptional activity in rat hepatocytes. 2000 , 350 Pt 2, 389-93	65
172	Transcription of genes encoding pregnancy-specific glycoproteins is regulated by negative promoter-selective elements. 2000 , 350 Pt 2, 511-9	1
171	Nuclear factor kappaB-dependent mechanisms coordinate the synergistic effect of PMA and cytokines on the induction of superoxide dismutase 2. 2001 , 353, 147-156	39
170	Differentiation-dependent mechanisms of transcriptional regulation of the catalytic subunit of phosphorylase kinase. 2002 , 362, 199-211	
169	The 20kD protein of human [U4/U6.U5] tri-snRNPs is a novel cyclophilin that forms a complex with the U4/U6-specific 60kD and 90kD proteins. 1998 , 4, 127-41	53
168	Transcriptional and post-transcriptional mechanisms involved in the differential expression of LPS-induced IL-1 and TNF mRNA. 1991 , 73, 460-5	51

167	Novel region within the V kappa gene promoter is responsible for tissue and stage-specific expression of immunoglobulin genes in human lymphoid neoplasms. 1989 , 66, 328-34	1
166	MRL/lpr mice produce anti-Ro 52,000 MW antibodies: detection, analysis of specificity and site of production. 1994 , 83, 9-15	35
165	Sequential recognition of the pre-mRNA branch point by U2AF65 and a novel spliceosome-associated 28-kDa protein. 1995 , 1, 407-17	35
164	Autoantibody production by patients infected with Leishmania. 1989 , 76, 190-7	48
163	The detection of anti-Sm-RNP activity in sera of patients with monoclonal gammopathies. 1989 , 75, 349-53	16
162	C-myc gene binding factors in peripheral blood mononuclear cells from patients with systemic lupus erythematosus (SLE). 1989 , 77, 157-62	1
161	RelB regulation of chemokine expression modulates local inflammation. 1997 , 151, 375-87	124
160	NF-kappaB activation during IgG immune complex-induced lung injury: requirements for TNF-alpha and IL-1beta but not complement. 1998 , 152, 1327-36	64
159	Cytoplasmic sequestration of p53 in cytomegalovirus-infected human endothelial cells. 1996 , 149, 1531-9	57
158	CAG trinucleotide RNA repeats interact with RNA-binding proteins. 1996 , 59, 561-9	32
157	Pentoxifylline and other protein kinase C inhibitors down-regulate HIV-LTR NF-kappa B induced gene expression. 1994 , 1, 31-43	11
156	Interleukin 1 induces HIV-1 expression in chronically infected U1 cells: blockade by interleukin 1 receptor antagonist and tumor necrosis factor binding protein type 1. 1995 , 1, 667-77	20
155	Engagement of tumor necrosis factor mRNA by an endotoxin-inducible cytoplasmic protein. 1996 , 2, 479-88	11
154	Classification of breast cancer cells on the basis of a functional assay for estrogen receptor. 1998 , 4, 454-67	2
153	Loss of NF-kappaB activity during cerebral ischemia and TNF cytotoxicity. 1999 , 5, 372-81	19
152	beta-lapachone induces cell cycle arrest and apoptosis in human colon cancer cells. 1999 , 5, 711-20	15
151	Phorbol 12-myristate 13-acetate (PMA) responsive sequence in Galphaq promoter during megakaryocytic differentiation. Regulation by EGR-1 and MAP kinase pathway. 2008 , 100, 821-8	2
150	Purification of the major histocompatibility complex class I transcription factor H2TF1. The full-length product of the nfkb2 gene. 1993 , 268, 18882-90	12

149	Culture of K562 human myeloid leukemia cells in presence of fibronectin expresses and secretes MMP-9 in serum-free culture medium. 2010 , 3, 288-302	5
148	Functional evaluation of TNFAIP3 (A20) in rheumatoid arthritis. 2010 , 28, 708-14	41
147	Clusterin silencing sensitizes pancreatic cancer MIA-PaCa-2 cells to gmcitabine via regulation of NF-kB/Bcl-2 signaling. 2015 , 8, 12476-86	8
146	Heat shock protein 90 inhibition by 17-Dimethylaminoethylamino-17-demethoxygeldanamycin protects blood-brain barrier integrity in cerebral ischemic stroke. 2015 , 7, 1826-37	19
145	Preferential distribution of active RNA polymerase II molecules in the nuclear periphery. 1991 , 1, 61-70	12
144	Sp1 activation of RNA polymerase II transcription complexes involves a heat-labile DNA-binding component. 1991 , 1, 137-48	8
143	Multiple activities of the human splicing factor ASF. 1992 , 2, 19-29	19
142	Identification of cis- and trans-acting factors regulating the expression of rat salivary-specific RP4 gene. 1992 , 2, 365-77	7
141	Tissue-specific expression of the skeletal alpha-actin gene involves sequences that can function independently of MyoD and Id. 1992 , 2, 241-57	8
140	The human skeletal alpha-actin gene is regulated by a muscle-specific enhancer that binds three nuclear factors. 1992 , 2, 111-26	33
139	The SV40 early transcriptional regulatory element is unable to direct gene expression in pituitary GH-3 cells. 1992 , 2, 175-89	3
138	Comparative utilization of transcription factor GABP by the promoters of ribosomal protein genes rpl30 and rpl32. 1993 , 3, 279-88	27
137	Distinct negative regulation of the human embryonic globin genes zeta and epsilon. 1993 , 3, 61-75	10
136	Analysis of the promoter element of the serum amyloid A gene and its interaction with constitutive and inducible nuclear factors from rabbit liver. 1993 , 3, 151-62	11
135	Molecular organization of the rat glia-derived nexin/protease nexin-1 promoter. 1993 , 3, 163-74	7
134	Identification of a stable RNA encoded by the H-strand of the mouse mitochondrial D-loop region and a conserved sequence motif immediately upstream of its polyadenylation site. 1995 , 4, 125-41	10
133	E1BF/Ku interacts physically and functionally with the core promoter binding factor CPBF and promotes the basal transcription of rat and human ribosomal RNA genes. 1995 , 4, 111-24	13
132	Transcriptional activity of domain A of the rat glucagon G3 element conferred by an islet-specific nuclear protein that also binds to similar pancreatic islet cell-specific enhancer sequences (PISCES). 1995 , 4, 205-16	9

131	A novel transcriptional initiator activity of the GABP factor binding ets sequence repeat from the murine cytochrome c oxidase Vb gene. 1995 , 5, 93-111	23
130	Nuclear factor YY1 activates the mammalian F0F1 ATP synthase alpha-subunit gene. 1996 , 5, 181-91	12
129	Gene transcription through activation of G-protein-coupled chemoattractant receptors. 1996 , 5, 205-15	5
128	C/EBP proteins contain nuclear localization signals imbedded in their basic regions. 1997 , 6, 371-85	40
127	TATA box and Sp1 sites mediate the activation of c-myc promoter P1 by immunoglobulin kappa enhancers. 1996 , 6, 113-27	17
126	Analysis of the negative transcriptional regulatory element in the angiotensin-converting enzyme gene. 1996 , 6, 73-85	3
125	Specific binding sites for a pol III transcriptional repressor and pol II transcription factor YY1 within the internucleosomal spacer region in primate Alu repetitive elements. 1996 , 6, 151-68	24
124	Induction of basic helix-loop-helix protein-containing complexes during erythroid differentiation. 1998 , 7, 25-38	10
123	RNA polymerase II transcription complex assembly in nuclear extracts. 1998 , 7, 191-204	4
122	The myogenic regulatory circuit that controls cardiac/slow twitch troponin C gene transcription in skeletal muscle involves E-box, MEF-2, and MEF-3 motifs. 1999 , 8, 247-61	8
121	Dual tandem promoter elements containing CCAC-like motifs from the tetrodotoxin-resistant voltage-sensitive Na ⁺ channel (rSkM2) gene can independently drive muscle-specific transcription in L6 cells. 1999 , 8, 85-103	2
120	Two internal sequence elements modulate transcription from the external human 7S K RNA gene promoter in vivo. 1999 , 8, 105-14	2
119	FOXP1 Interacts with MyoD to Repress its Transcription and Myoblast Conversion. 2021 , 2, 9-26	
118	Using quantitative reconstitution to investigate multi-component condensates. 2021 ,	2
117	UBR7 acts as a histone chaperone for post-nucleosomal histone H3. 2021 , e108307	2
116	5' Half of specific tRNAs feeds back to promote corresponding tRNA gene transcription in vertebrate embryos. 2021 , 7, eabh0494	1
115	Methods to separate nuclear soluble fractions reflecting localizations in living cells.. 2021 , 24, 103503	2
114	Runx2 Regulates Chromatin Accessibility to Direct Skeletal Cell Programs.	2

113	A Model for DHX15 Mediated Disassembly of A-Complex Spliceosomes.. 2022 ,	0
112	SMARCA5 interacts with NUP98-NSD1 oncofusion protein and sustains hematopoietic cells transformation.. 2022 , 41, 34	2
111	Structural basis of branch site recognition by the human spliceosome. 2022 , 375, 50-57	3
110	Epigenetic anti-cancer treatment with a stabilized carbocyclic Decitabine analogue.	
109	MYC assembles and stimulates topoisomerases 1 and 2 in a "topoisome". 2021 ,	3
108	Lucidin 3-methyl ether from <i>Rubia philippinensis</i> suppresses the proliferation of multiple myeloma cells through the promotion of β -catenin degradation.. 2022 , 99, 153971	
107	Comprehensive comparison between azacytidine and decitabine treatment in an acute myeloid leukemia cell line.	
106	CRISPR and biochemical screens identify MAZ as a cofactor in CTCF-mediated insulation at Hox clusters.. 2022 , 54, 202-212	2
105	The Selective Serotonin 2A Receptor Antagonist Sarpogrelate Prevents Cardiac Hypertrophy and Systolic Dysfunction via Inhibition of the ERK1/2-GATA4 Signaling Pathway.. 2021 , 14,	1
104	An A/G-rich motif in the rat fibroblast growth factor-2 gene confers enhancer activity on a heterologous promoter in neonatal rat cardiac myocytes. 1998 , 188, 169-76	2
103	Regulation of tyrosine hydroxylase gene transcription by the cAMP-signaling pathway: involvement of multiple transcription factors. 2000 , 212, 51-60	17
102	cAMP increases the expression of human angiotensinogen gene through a combination of cyclic AMP responsive element binding protein and a liver specific transcription factor. 2000 , 212, 81-90	4
101	Exon Skipping Through Chimeric Antisense snRNAs to Correct Retinitis Pigmentosa GTPase-Regulator () Splice Defect.. 2022 ,	1
100	LncRNA RUS shapes the gene expression program towards neurogenesis.	
99	Epigenetic anti-cancer treatment with a stabilized carbocyclic Decitabine analogue.. 2022 ,	
98	CMT α cap-adjacent 2'-O-ribose mRNA methyltransferases are required for reward learning and mRNA localization to synapses.. 2022 , 13, 1209	1
97	Principles and pitfalls of high-throughput analysis of microRNA-binding thermodynamics and kinetics by RNA Bind-n-Seq.. 2022 , 2, 100185	0
96	The PAF1 complex promotes 3' processing of pervasive transcripts.. 2022 , 38, 110519	0

- 95 Altered BAF occupancy and transcription factor dynamics in PBAF-deficient melanoma.. **2022**, 39, 110637 ○
- 94 Long noncoding RNAs transcribed downstream of the human β globin locus regulate β globin gene expression. **2021**, ○
- 93 Splicing efficiency of minor introns in a mouse model of SMA predominantly depends on their branchpoint sequence and can involve the contribution of major spliceosome components. **2021**, ○
- 92 Hormonal Regulation of the MHC Class I Gene in Thyroid Cells: Role of the Promoter "Tissue-Specific" Region.. **2021**, 12, 749609 ○
- 91 Galectin-3-U1 snRNP Complexes Initiate Splicing Activity in U1-Depleted Nuclear Extracts.. **2022**, 2442, 713-726
- 90 Regulation of the cell cycle under anoxia stress in tail muscle and hepatopancreas of the freshwater crayfish, *Orconectes virilis*.. **2022**, 111215 ○
- 89 Novel role of UHRF1 in the epigenetic repression of the latent HIV-1.. **2022**, 79, 103985 ○
- 88 Image_1.TIF. **2018**,
- 87 Image_2.TIF. **2018**,
- 86 Image_3.TIF. **2018**,
- 85 Table_1.docx. **2020**,
- 84 Inositol polyphosphate multikinase physically binds to the SWI/SNF complex and modulates BRG1 occupancy in mouse embryonic stem cells.. **2022**, 11, ○
- 83 IKAROS and MENIN coordinate therapeutically actionable leukemogenic gene expression in MLL-r acute myeloid leukemia.. **2022**, ○
- 82 The Cytoskeleton Differentially Localizes the Early Growth Response Gene-1 Protein in Cancer and Benign Cells of the Prostate. **2004**, 2, 115-128 14
- 81 HMG-I/Y Is a c-Jun/Activator Protein-1 Target Gene and Is Necessary for c-Jun-Induced Anchorage-Independent Growth in Rat1a Cells. **2004**, 2, 305-314 26
- 80 Mediator-RNA Polymerase II Interactions Critical for Transcriptional Activation Are Mediated by the N-terminal Half of MED14 and C-terminal Domain of RPB1.
- 79 Screening of organoids derived from patients with breast cancer implicates the repressor NCOR2 in cytotoxic stress response and antitumor immunity. 1
- 78 RNF219 regulates CCR4-NOT function in mRNA translation and deadenylation. **2022**, 12, ○

77	Establishment and Fractionation of Metastatic Axillary Lymph Node Cell Suspension for Determination of Protein Expression Levels of Nuclear cFOS and Cytosolic TGF β 1 from Breast Cancer Patients. 2022 , 24,	
76	Identifying Specific Protein Interactors of Nucleosomes Carrying Methylated Histones Using Quantitative Mass Spectrometry. 2022 , 327-403	
75	In Vitro Histone Demethylase Assays. 2022 , 63-88	
74	LncRNA RUS shapes the gene expression program towards neurogenesis. 2022 , 5, e202201504	0
73	Tandem immuno-purification of affinity-tagged proteins from mouse testis extracts for MS analysis. 2022 , 3, 101452	0
72	ITRAQ-based quantitative proteomic analysis reveals that VPS35 promotes the expression of MCM2-7 genes in HeLa cells. 2022 , 12,	0
71	Intrinsic ROS Drive Hair Follicle Cycle Progression by Modulating DNA Damage and Repair and Subsequently Hair Follicle Apoptosis and Macrophage Polarization. 2022 , 2022, 1-35	0
70	Structural analysis of the basal state of the Artemis:DNA-PKcs complex. <i>Nucleic Acids Research</i> ,	20.1 0
69	Novel Prognostic Value of Nuclear Epidermal Growth Factor Receptor in Breast Cancer. 2005 , 65, 338-348	66
68	Curcumin sensitizes prostate cancer cells to tumor necrosis factor-related apoptosis-inducing ligand/Apo2L by inhibiting nuclear factor- κ B through suppression of I κ B phosphorylation. 2004 , 3, 803-812	42
67	Epigenetic and transcriptome responsiveness to ER modulation by tissue selective estrogen complexes in breast epithelial and breast cancer cells. 2022 , 17, e0271725	
66	HSF1 phosphorylation establishes an active chromatin state via the TRRAP β IP60 complex and promotes tumorigenesis. 2022 , 13,	0
65	Estrogen restores cellular immunity in injured male mice via suppression of interleukin-6 production. 2001 , 70, 887-895	8
64	Adenoviral gene delivery can inactivate Kupffer cells: role of oxidants in NF- κ B activation and cytokine production. 2001 , 69, 622-630	10
63	Compare analysis of the PRNP gene indels polymorphism and expression among Yellow cattle, Zhongdian Yak and their hybrid.	0
62	Comprehensive comparison between azacytidine and decitabine treatment in an acute myeloid leukemia cell line. 2022 , 14,	0
61	Runx2 regulates chromatin accessibility to direct the osteoblast program at neonatal stages. 2022 , 40, 111315	1
60	A Light-Inducible Bidirectional Promoter Initiates Expression of Both Genes SHH2 and CFM3 in <i>Brassica napus</i> L..	0

59	The inVivo Interaction Landscape of Histones H3.1 and H3.3. 2022 , 21, 100411	0
58	Quantification of Farnesylated Progerin in Hutchinson-Gilford Progeria Patient Cells by Mass Spectrometry. 2022 , 23, 11733	0
57	Mechanisms of mineralocorticoid receptor-associated hypertension in diabetes mellitus: the role of O-GlcNAc modification.	0
56	Development of Macrocyclic PRMT5Adaptor Protein Interaction Inhibitors.	0
55	The POLR3G Subunit of Human RNA Polymerase III Regulates Tumorigenesis and Metastasis in Triple-Negative Breast Cancer. 2022 , 14, 5732	0
54	=?UTF-8?B?VGhlIGVmZmVjdHMgb2YgcmViYW1pcGlkZSwgc3VjcmFsZmF0ZSwgYW5kIHJpZmF4aW1pb2FpbnN0IGU=	0
53	A Two-Step Mechanism for Recruitment of Pip by PU.1. 1998 , 160, 241-252	17
52	Regulation of IL-6 Synthesis in Human Peripheral Blood Mononuclear Cells by C3a and C3adesArg. 1999 , 162, 453-459	21
51	Nuclear Targeted Suppression of NF- κ B Activity by the Novel Quinone Derivative E3330. 1998 , 160, 810-819	14
50	Role of the Rho GTPase in Bradykinin-Stimulated Nuclear Factor- κ B Activation and IL-1 β Gene Expression in Cultured Human Epithelial Cells. 1998 , 160, 3038-3045	13
49	PU.1/Spi-1 Is Essential for the B Cell-Specific Activity of the Mouse CD72 Promoter. 1998 , 160, 2287-2296	5
48	Involvement of the IL-2 Receptor β Chain (β) in the Control by IL-4 of Human Monocyte and Macrophage Proinflammatory Mediator Production. 1998 , 160, 4048-4056	10
47	Extracellular HIV-1 Tat Protein Induces the Rapid Ser133 Phosphorylation and Activation of CREB Transcription Factor in Both Jurkat Lymphoblastoid T Cells and Primary Peripheral Blood Mononuclear Cells. 1998 , 160, 3891-3898	16
46	IFN- γ Induction of the Human Monocyte Chemoattractant Protein (hMCP)-1 Gene in Astrocytoma Cells: Functional Interaction Between an IFN- γ Activated Site and a GC-Rich Element. 1998 , 160, 3908-3916	18
45	Regulation of Transcription of the TATA-less Human Complement Component C4 Gene. 1998 , 160, 4353-4360	5
44	Expression of Bright at Two Distinct Stages of B Lymphocyte Development. 1998 , 160, 4747-4754	6
43	CpG Motifs in Bacterial DNA Activate Leukocytes Through the pH-Dependent Generation of Reactive Oxygen Species. 1998 , 160, 4755-4761	65
42	Involvement of CD14 and Complement Receptors CR3 and CR4 in Nuclear Factor- κ B Activation and TNF Production Induced by Lipopolysaccharide and Group B Streptococcal Cell Walls. 1998 , 160, 4535-4542	23

41	Receptors for the Anaphylatoxin C5a (CD88) on Human Mesangial Cells. 1998 , 160, 5646-5652	7
40	The Regulation of Murine H-2Dd Expression by Activation Transcription Factor 1 and cAMP Response Element Binding Protein. 1998 , 160, 5907-5914	2
39	Angiotensin II Participates in Mononuclear Cell Recruitment in Experimental Immune Complex Nephritis Through Nuclear Factor- κ B Activation and Monocyte Chemoattractant Protein-1 Synthesis. 1998 , 161, 430-439	31
38	Regulatory Elements in the Promoter of a Murine TCRD V Gene Segment. 1998 , 161, 791-804	2
37	Generation of Switch Hybrid DNA Between Ig Heavy Chain- γ and Downstream Switch Regions in B Lymphocytes. 1998 , 161, 1354-1362	1
36	The Invariant Chain Gene Intronic Enhancer Shows Homology to Class II Promoter Elements. 1998 , 161, 1844-1852	1
35	Role of Nuclear Factor- κ B and Mitogen-Activated Protein Kinase Signaling Pathways in IL-1 β -Mediated Induction of PDGF Receptor Expression in Rat Pulmonary Myofibroblasts. 1998 , 161, 3464-3468	3
34	Mef2 Proteins, Required for Muscle Differentiation, Bind an Essential Site in the Ig γ Enhancer. 1998 , 161, 4795-4802	4
33	Fc γ R1-Mediated Induction of TNF- α Gene Expression in the RBL-2H3 Mast Cell Line: Regulation by a Novel NF- κ B-Like Nuclear Binding Complex. 1998 , 161, 4768-4776	6
32	p48/STAT-1 β -Containing Complexes Play a Predominant Role in Induction of IFN- γ -Inducible Protein, 10 kDa (IP-10) by IFN- γ Alone or in Synergy with TNF- α 1998 , 161, 4736-4744	32
31	Regulation of Mouse CD72 Gene Expression During B Lymphocyte Development. 1998 , 161, 4760-4767	6
30	Human Cytomegalovirus Inhibits IFN- γ -Stimulated Antiviral and Immunoregulatory Responses by Blocking Multiple Levels of IFN- γ Signal Transduction. 1999 , 162, 6107-6113	21
29	CD40 Ligand-CD40 Interaction Induces Chemokines in Cervical Carcinoma Cells in Synergism with IFN- γ 1999 , 162, 4140-4147	4
28	Membrane-Anchored Forms of Lipopolysaccharide (LPS)-Binding Protein Do Not Mediate Cellular Responses to LPS Independently of CD14. 1999 , 162, 5483-5489	4
27	Osteoclast Differentiation Factor Acts as a Multifunctional Regulator in Murine Osteoclast Differentiation and Function. 1999 , 163, 434-442	52
26	Constitutive Nuclear Translocation of NF- κ B in B Cells in the Absence of I κ B Degradation. 1999 , 163, 269-277	12
25	Complex Regulation of Ly-6E Gene Transcription in T Cells by IFNs. 1999 , 163, 811-819	4
24	Bruton's Tyrosine Kinase Deficiency in Macrophages Inhibits Nitric Oxide Generation Leading to Enhancement of IL-12 Induction. 1999 , 163, 1786-1792	7

23	Roles of IFN Consensus Sequence Binding Protein and PU.1 in Regulating IL-18 Gene Expression. 1999 , 163, 2000-2007	16
22	An Activation-Responsive Element in Single C Motif-1/Lymphotactin Promoter Is a Site of Constitutive and Inducible DNA-Protein Interactions Involving Nuclear Factor of Activated T Cell. 1999 , 163, 3295-3303	3
21	TNF- α -Induced Growth Suppression of CD34+ Myeloid Leukemic Cell Lines Signals Through TNF Receptor Type I and Is Associated with NF- κ B Activation. 1999 , 163, 3106-3115	3
20	The Signal Transduction Pathway of CD23 (Fc β RIIb) Targets I κ B Kinase. 1999 , 163, 3851-3857	1
19	Characterization of RNA Binding Proteins Associated with CD40 Ligand (CD154) mRNA Turnover in Human T Lymphocytes. 1999 , 163, 4199-4206	4
18	The Epithelial Cell Response to Rotavirus Infection. 1999 , 163, 4442-4452	16
17	Functional Synergism of STAT6 with Either NF- κ B or PU.1 to Mediate IL-4-Induced Activation of IgE Germline Gene Transcription. 1999 , 163, 4383-4391	21
16	Recruitment of CREB-Binding Protein by PU.1, IFN-Regulatory Factor-1, and the IFN Consensus Sequence-Binding Protein Is Necessary for IFN- γ -Induced p67phox and gp91phox Expression. 1999 , 163, 6095-6105	31
15	Analysis of the Defect in IFN- γ -Induction of MHC Class II Genes in G1B Cells: Identification of a Novel and Functionally Critical Leucine-Rich Motif (62-LYLYLQL-68) in the Regulatory Factor X 5 Transcription Factor. 1999 , 163, 6622-6630	4
14	NF- κ B/Rel Transcription Factors: c-Rel Promotes Airway Hyperresponsiveness and Allergic Pulmonary Inflammation. 1999 , 163, 6827-6833	28
13	Transcriptional and Translational Regulation of Inflammatory Mediator Production by Endogenous TGF- β in Macrophages That Have Ingested Apoptotic Cells. 1999 , 163, 6164-6172	59
12	SPARC is a decoy counterpart for c-Fos and is associated with osteoblastic differentiation of bone marrow stromal cells by inhibiting adipogenesis. 2023 , 27,	0
11	A Novel Chalcone Derivative Regulates the Expression and Phosphorylation of ERK1/2 by Inhibiting Fli-1 Promoter Activity for Preventing the Malignant Progression of Erythroleukemia. 2023 , 13, 84	0
10	Characterization of the TBR1 interactome: variants associated with neurodevelopmental disorders disrupt novel protein interactions.	0
9	Transcriptional Regulation of Math1 by Aryl Hydrocarbon Receptor: Effect on Math1+ Progenitor Cells in Mouse Small Intestine. 2023 , 43, 43-63	0
8	Exposure to brefeldin A induces unusual expression of hybrid- and complex-type free N-glycans in HepG2 cells. 2023 , 1867, 130331	0
7	Human DNA polymerase δ promotes RNA-templated error-free repair of DNA double-strand breaks. 2023 , 299, 102991	0
6	Structural basis of paralog-specific KDM2A/B nucleosome recognition.	0

- 5 Regulation of 3' splice site selection after step 1 of splicing by spliceosomal C* proteins. **2023**, 9, o
- 4 Phosphorylation of pICln by the autophagy activating kinase ULK1 regulates snRNP biogenesis and splice activity of the cell. **2023**, 21, 2100-2109 o
- 3 Functionally distinct promoter classes initiate transcription via different mechanisms reflected in focused versus dispersed initiation patterns. o
- 2 Functional analysis of a conserved site mutation in the DNA processing enzyme PNKP leading to Ataxia with Oculomotor Apraxia type 4 (AOA4) in humans. **2023**, 104714 o
- 1 Novel Chromatin Insulating Activities Uncovered upon Eliminating Known Insulators *in vivo*. o