

Mercury concentrations in canned and fresh fish and its port moresby residents

Science of the Total Environment

26, 157-162

DOI: [10.1016/0048-9697\(83\)90108-0](https://doi.org/10.1016/0048-9697(83)90108-0)

Citation Report

#	ARTICLE	IF	CITATIONS
1	South Pacific regional environment programme. Environmental Policy and Law, 1982, 9, 63-63.	0.2	7
2	Interindividual variations of blood total mercury levels according to sex, age and area of residence. Science of the Total Environment, 1985, 44, 101-109.	8.0	9
3	Mercury in scalp hair of healthy Singapore residents. Science of the Total Environment, 1988, 72, 113-122.	8.0	27
4	Use of hair analysis for evaluating mercury intoxication of the human body: A review. Journal of Applied Toxicology, 1992, 12, 79-84.	2.8	65
5	Concentrations of heavy metals in maternal and umbilical cord blood. BioMetals, 1993, 6, 61-6.	4.1	45
6	Mercury in human hair and relation to fish consumption in Bangladesh. Science of the Total Environment, 1996, 186, 181-188.	8.0	59
7	Factors influencing the total mercury and methyl mercury in the hair of the fishermen of Kuwait. Environmental Pollution, 2000, 109, 239-250.	7.5	88
8	A Review of Mercury in Seafood. Journal of Aquatic Food Product Technology, 2005, 14, 71-100.	1.4	33
10	Mercury Content of Hair in Different Populations Relative to Fish Consumption. Reviews of Environmental Contamination and Toxicology, 2007, 189, 107-130.	1.3	15
11	Total mercury in canned tuna sold in Canada in 2006. Food Additives and Contaminants: Part B Surveillance, 2014, 7, 110-114.	2.8	12
12	Mercury in canned fish from local markets in the Czech Republic. Food Additives and Contaminants: Part B Surveillance, 2017, 10, 149-154.	2.8	11