

CITATION REPORT

List of articles citing

Improved spectral resolution in cosy ^1H NMR spectra of proteins via double quantum filtering

DOI: 10.1016/0006-291x(83)91225-1
Biochemical and Biophysical Research
Communications, 1983, 117, 479-85.

Source: <https://exaly.com/paper-pdf/16388264/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
2376	Chemical Synthesis Demonstrates That Dynamic Oglycosylation Regulates the Folding and Functional Conformation of a Pivotal EGF12 Domain of the Human NOTCH1 Receptor.		
2375	NMR Relaxation Dispersion Reveals Macrocycle Breathing Dynamics in a Cyclodextrin-based Rotaxane.		
2374	The Configuration of the WWW Motif of a Short Trp-Rich Peptide Is Critical for Targeting Bacterial Membranes, Disrupting Preformed Biofilms, and Killing Methicillin-Resistant <i>Staphylococcus aureus</i> .		
2373	Pattern recognition in two-dimensional NMR spectra. 1984 , 60, 161-163		11
2372	Longitudinal two-spin order in 2D exchange spectroscopy (NOESY). 1984 , 59, 542-550		8
2371	Application of π -decoupled 2D correlation spectra to the study of proteins. 1984 , 59, 250-261		15
2370	Sequential assignment of the ^1H and ^{31}P resonances of the double stranded deoxynucleotide d(ATGCAT) ₂ by 2D-NMR correlation spectroscopy. <i>Biochemical and Biophysical Research Communications</i> , 1984 , 124, 774-83	3.4	51
2369	The immunodominant site of a synthetic immunogen has a conformational preference in water for a type-II reverse turn. 1985 , 318, 480-3		221
2368	The selection of coherence-transfer pathways by inhomogeneous z pulses. 1985 , 64, 470-478		5
2367	The appearance of forbidden cross peaks in two-dimensional nuclear magnetic resonance spectra due to multiexponential T2 relaxation. 1985 , 65, 531-534		12
2366	Design of pulse sequences sensitive to relative signs of J coupling constants. Corrections to the SLAP experiment. 1985 , 63, 219-224		4
2365	Effect of strong coupling on multiple-quantum filtering. 1985 , 61, 540-544		
2364	Correlation of aliphatic proton chemical shifts in proteins using two-dimensional double-quantum correlated spectroscopy. 1985 , 62, 543-550		3
2363	Comparison and evaluation of methods for two-dimensional NMR spectra with absorption-mode lineshapes. 1985 , 63, 454-472		35
2362	The application of a simple analog phase-shifting device in multiple-quantum NMR. 1985 , 63, 392-396		1
2361	Multiple-quantum excitation and spin topology filtration in high-resolution NMR. 1985 , 83, 3297-3310		63
2360	Systematic application of high-resolution, phase-sensitive two-dimensional ^1H -NMR techniques for the identification of the amino-acid-proton spin systems in proteins. Rabbit metallothionein-2. <i>FEBS Journal</i> , 1985 , 151, 257-73		248

2359	Simplification of Spectra for the Determination of Coupling Constants from Homonuclear Correlated 2D NMR Spectra. 1985 , 24, 690-692	11
2358	Sequence-specific assignment of the backbone 1H-and 31P-NMR lines in a short DNA duplex with homo- and heteronuclear correlated spectroscopy. 1985 , 24, 2371-80	37
2357	Differences and sums of traces within, COSY spectra (DISCO) for the extraction of coupling constants: Decoupling after the measurement. 1985 , 23, 844-852	73
2356	Spektrvereinfachung zur Ermittlung von Kopplungskonstanten aus homonuclear-korrelierten 2D-NMR-Spektren. 1985 , 97, 689-690	7
2355	Secondary structure of the alpha-amylase polypeptide inhibitor tendamistat from Streptomyces tendae determined in solution by 1H nuclear magnetic resonance. 1985 , 183, 503-7	146
2354	Identification of glycine spin systems in 1H NMR spectra of proteins using multiple quantum coherences. 1985 , 186, 35-40	24
2353	1H nuclear magnetic resonance assignments for d-(GCATTAATGC) ₂ using experimental refinements of established procedures. 1986 , 190, 439-53	88
2352	Two-dimensional nuclear magnetic resonance spectroscopy. 1986 , 232, 960-7	105
2351	Complete sequence-specific 1H nuclear magnetic resonance assignments for the alpha-amylase polypeptide inhibitor tendamistat from Streptomyces tendae. 1986 , 192, 869-90	44
2350	Residue-specific assignments of resonances in the 1H nuclear magnetic resonance spectrum of ribosomal protein E-L30 by systematic application of two-dimensional Fourier transform nuclear magnetic resonance methods. 1986 , 192, 389-417	20
2349	Nuclear magnetic resonance study of the globular domain of chicken histone H5: resonance assignment and secondary structure. 1986 , 83, 7628-32	29
2348	Elucidation of glycolipid structure by proton nuclear magnetic resonance spectroscopy. 1986 , 42, 27-48	47
2347	Conformational study of bacterial lipopeptides: refinement of the structure of iturin A in solution by two-dimensional 1H-NMR and energy calculations. 1986 , 25, 153-70	43
2346	Modern NMR techniques for structure elucidation. 1986 , 24, 371-403	150
2345	1H NMR studies of heme pocket conformation in zinc-substituted leghemoglobin, a diamagnetic analog of deoxyleghemoglobin. 1986 , 28, 303-9	9
2344	Simulation of two-dimensional NMR experiments using numerical density matrix calculations. 1986 , 70, 270-279	6
2343	Complete protein fingerprints by double-quantum spectroscopy. 1986 , 66, 359-363	8
2342	Analysis of 1H NMR spectra of proteins using multiple-quantum coherence. 1986 , 66, 372-378	15

2341	COSY representation of two-dimensional homonuclear double-quantum spectra. 1986 , 66, 153-156	1
2340	Two-dimensional correlation of directly and remotely connected transitions by z-filtered COSY. 1986 , 69, 559-566	14
2339	Analysis of cross peak fine structure in multiple-quantum filtered COSY spectra. 1986 , 68, 67-84	7
2338	Transformation of homonuclear two-dimensional NMR techniques into one-dimensional techniques using Gaussian pulses. 1986 , 70, 106-133	69
2337	¹ H-2D-nuclear magnetic resonance applied to the primary structure determination of a novel octasaccharide glycolipid isolated from the spermatozoa of bivalves. 1986 , 155, 45-56	11
2336	Sequence-specific ¹ H-NMR assignments in rabbit-liver metallothionein-2. <i>FEBS Journal</i> , 1986 , 157, 275-89	51
2335	¹ H-NMR assignment and secondary structure of a herpes simplex virus glycoprotein D-1 antigenic domain. <i>FEBS Journal</i> , 1986 , 158, 527-36	44
2334	Correlation of connected transitions by two-dimensional NMR spectroscopy. 1986 , 85, 6837-6852	261
2333	A Two-Dimensional ¹ H NMR Approach for Peptide Backbone Assignments: Gramicidin S. 1987 , 34, 309-315	0
2332	Conformational analysis of the deoxyribofuranose ring in DNA by means of sums of proton-proton coupling constants: a graphical method. 1987 , 4, 621-49	277
2331	Oligosaccharide structure by two-dimensional proton nuclear magnetic resonance spectroscopy. 1987 , 138, 38-59	50
2330	¹ H nuclear magnetic resonance studies of an integral membrane protein: subunit c of the F1F0 ATP synthase. 1987 , 193, 759-74	31
2329	Nuclear magnetic resonance study of the solution structure of alpha 1-purothionin. Sequential resonance assignment, secondary structure and low resolution tertiary structure. 1987 , 193, 571-8	42
2328	Assignment of resonances in the ¹ H nuclear magnetic resonance spectrum of the carbon monoxide complex of sperm whale myoglobin by phase-sensitive two-dimensional techniques. 1987 , 194, 313-27	80
2327	Macrocycles containing tin. Through space cooperative binding and high size selectivity in the complexation of chloride ion by Lewis acidic macrobicyclic hosts. 1987 , 109, 7878-7879	53
2326	NMR of a synthetic peptide spanning the triphosphate binding site of adenosine 5'-triphosphate in actin. 1987 , 26, 1471-8	9
2325	Three-dimensional structure of phoratoxin in solution: combined use of nuclear magnetic resonance, distance geometry, and restrained molecular dynamics. 1987 , 26, 1732-1745	76
2324	Sequence specific assignment of the proton nuclear magnetic resonance spectrum of barley serine proteinase inhibitor 2. 1987 , 52, 327-354	26

2323	Nuclear magnetic resonance studies of ionomers: 2. Proton transfer in polyurethane-poly(styrene-co-styrene sulphonic acid) mixtures in solution. 1987 , 28, 885-888	16
2322	Two-dimensional n.m.r. spectra of 6- O - D -glucopyranosylcyclomalto-hexaose; 1 H and 13 C resonance assignments and primary structure determination. 1987 , 166, 156-161	11
2321	Nuclear magnetic resonance studies of ionomers. 6. Polyurethane-polyphosphonate blends. 1987 , 27, 1504-1511	2
2320	Optimized detection of small spin-spin couplings in two-dimensional chemical shift correlation experiments?COSY and DEPT. 1987 , 25, 125-128	7
2319	Identification of constituent sugar residues in oligosaccharides by two-dimensional 1H NMR phase-sensitive correlated spectroscopy. 1987 , 25, 338-346	17
2318	Assignments of the 1H- and 13C-n.m.r. spectra of tobramycin at low and high pH. 1987 , 170, 1-17	26
2317	z-Filtered double-quantum NMR spectra and automated analysis by pattern recognition. 1987 , 73, 493-511	5
2316	Application of multiple-quantum filters in zero-quantum spectroscopy. 1987 , 73, 534-538	
2315	Relaxation-induced violations of coherence transfer selection rules in nuclear magnetic resonance. 1987 , 75, 297-334	36
2314	Practical aspects of the E.COSY technique. Measurement of scalar spin-spin coupling constants in peptides. 1987 , 75, 474-492	50
2313	Strong coupling effects in z-filtered two-dimensional NMR correlation spectra. 1987 , 72, 475-492	4
2312	P.E.COSY, a simple alternative to E.COSY. 1987 , 72, 191-196	35
2311	A method for pure absorption 2D spectroscopy using complex fourier transformation in both dimensions. 1987 , 72, 279-297	3
2310	Simulated two-dimensional nmr cross-peak fine structures for 1H spin systems in polypeptides and polydeoxynucleotides. 1987 , 74, 316-336	12
2309	Stereospecific assignments of side-chain protons and characterization of torsion angles in Eglin c. <i>FEBS Journal</i> , 1987 , 164, 625-35	161
2308	Monitoring the purification by high-performance liquid chromatography of cardiotoxins from <i>Naja mossambica mossambica</i> using phase-sensitive two-dimensional nuclear magnetic resonance. <i>FEBS Journal</i> , 1987 , 168, 603-7	10
2307	Sequence-specific 1H-NMR assignments and determination of the secondary structure in aqueous solution of the cardiotoxins CTXIIa and CTXIIb from <i>Naja mossambica mossambica</i> . <i>FEBS Journal</i> , 1987 , 168, 609-20	25
2306	1H-NMR sequential assignments and cation-binding studies of spinach plastocyanin. <i>FEBS Journal</i> , 1987 , 170, 279-92	33

2305	Two-dimensional ¹ H-NMR studies of the solution structure of plasminogen kringle 4. <i>FEBS Journal</i> , 1988 , 170, 539-48	13
2304	The secondary structure of the toxin ATX Ia from <i>Anemonia sulcata</i> in aqueous solution determined on the basis of complete sequence-specific ¹ H-NMR assignments. <i>FEBS Journal</i> , 1988 , 171, 177-92	24
2303	The structure of melittin. A ¹ H-NMR study in methanol. <i>FEBS Journal</i> , 1988 , 173, 139-46	209
2302	cDNA cloning and functional expression in yeast <i>Saccharomyces cerevisiae</i> of beta-naphthoflavone-induced rabbit liver P-450 LM4 and LM6. <i>FEBS Journal</i> , 1988 , 177, 285-93	68
2301	Secondary structure determination for alpha-neurotoxin from <i>Dendroaspis polylepsis polylepsis</i> based on sequence-specific ¹ H-nuclear-magnetic-resonance assignments. <i>FEBS Journal</i> , 1988 , 177, 295-305	23
2300	NMR studies of the conformations of leghemoglobins from soybean and lupin. <i>FEBS Journal</i> , 1988 , 178, 419-35	14
2299	Zweidimensionale NMR-Spektroskopie, Grundlagen und Übersicht über die Experimente. 1988 , 100, 507-554	159
2298	Suppression of direct peaks in two-dimensional heteronuclear NOE spectra. Sensitivity-enhanced detection of weak indirect and relay peaks. 1988 , 26, 181-184	3
2297	Assignment of proton and carbon NMR signals of didemnin A in solution. 1988 , 26, 501-506	7
2296	Two-dimensional NMR spectra of sesquiterpenes. HMQC study of some sesquiterpene lactones with pseudoguaianolide, 3,4-seco-pseudoguaianolide and guaianolide skeletons. 1988 , 26, 725-728	4
2295	Improvement of 2D NMR spectra by matching symmetry-related spectral features. 1988 , 26, 848-851	19
2294	Two-Dimensional NMR Spectroscopy: Background and Overview of the Experiments [New Analytical Methods (36)]. 1988 , 27, 490-536	363
2293	Complete ¹ H-NMR spectral assignments for globotriaosyl-Z- and isoglobotriaosyl-E-ceramide. 1988 , 49, 31-7	9
2292	Lavandulifolioside: A New Phenylpropanoid Glycoside from <i>Stachys lavandulifolia</i> . 1988 , 71, 1483-1490	45
2291	Beiträge zur Chemie der Pyrrolpigmente, 75. Mitt. Phytochrom-Modell-Studien: Stereochemische Untersuchungen an einem zwischen den pyrrolischen Ringen A und C peptidartig verbrückten 2,3-Dihydrobilatrien-abc. 1988 , 119, 113-126	4
2290	Conformation of a hexahydrocannabinol derivative via 1- and 2-D noe experiments. evaluation of ¹ JCH coupling constants by semiselective 2-D J-spectroscopy and their correlation with the ring geometry. 1988 , 44, 2329-2338	3
2289	Diversity of the gramicidin a spatial structure: two-dimensional ¹ H nmr study in solution. 1988 , 44, 925-940	81
2288	Conformational study of super-active analogues of somatostatin with reduced ring size by XXXH nmr. 1988 , 44, 941-973	14

2287	Structure of P401 (mast cell degranulating peptide) in solution. 1988 , 31, 113-9	22
2286	Computer-aided sequential assignment of protein ¹ H NMR spectra. 1988 , 80, 119-127	7
2285	Application of homonuclear double-acquisition two-dimensional shift correlation spectroscopy. 1988 , 76, 316-320	1
2284	Effect of pulse angle imperfection and resonance offset in the phase-modulated rotating-frame spectroscopic imaging experiment. 1988 , 77, 203-222	
2283	Steady-state dqf-cosy spectra using a variable relaxation delay. 1988 , 78, 314-320	3
2282	Two-dimensional NMR studies of Kazal proteinase inhibitors. 1. Sequence-specific assignments and secondary structure of turkey ovomucoid third domain. 1988 , 27, 2519-29	34
2281	Structure elucidation of the core regions from Citrobacter O4 and O36 lipopolysaccharides by chemical and enzymatic methods, gas chromatography/mass spectrometry, and NMR spectroscopy at 500 MHz. 1988 , 27, 4153-61	32
2280	Structural differences between oxidized and reduced thioredoxin monitored by two-dimensional ¹ H NMR spectroscopy. 1988 , 228, 254-8	35
2279	Two-dimensional ¹ H-NMR study of bacterioopsin-(3485)-polypeptide conformation. 1988 , 231, 81-88	32
2278	A 500 MHz study of peptide T in a DMSO solution. 1988 , 231, 159-63	26
2277	Folding of immunogenic peptide fragments of proteins in water solution. I. Sequence requirements for the formation of a reverse turn. 1988 , 201, 161-200	634
2276	¹ H n.m.r. study of. 1988 , 10, 277-281	3
2275	Principles of Multiple-Quantum Spectroscopy. 1988 , 259-341	
2274	Folding of immunogenic peptide fragments of proteins in water solution. II. The nascent helix. 1988 , 201, 201-17	449
2273	Characterisation of four lactose monophosphates by application of ³¹ P-, ¹³ C-, and ¹ H-n.m.r. spectroscopy. 1988 , 174, 23-36	25
2272	Complete assignment of the ¹ H nuclear magnetic resonance spectrum of French bean plastocyanin. Application of an integrated approach to spin system identification in proteins. 1988 , 202, 603-22	78
2271	Secondary structure of acylphosphatase from rabbit skeletal muscle. A nuclear magnetic resonance study. 1988 , 199, 233-7	14
2270	Determination of the complete three-dimensional structure of the alpha-amylase inhibitor tendamistat in aqueous solution by nuclear magnetic resonance and distance geometry. 1988 , 204, 675-724	208

2269	Three-dimensional structure of rabbit liver [Cd7]metallothionein-2a in aqueous solution determined by nuclear magnetic resonance. 1988 , 201, 637-57		275
2268	Complete sequence-specific 1H nuclear magnetic resonance assignments for mouse epidermal growth factor. 1988 , 103, 554-71		25
2267	Secondary structure determination for the Antennapedia homeodomain by nuclear magnetic resonance and evidence for a helix-turn-helix motif.. <i>EMBO Journal</i> , 1988 , 7, 4305-4309	13	107
2266	Proton nuclear magnetic resonance study of the solution conformation of C3a-Arg69, a 69-residue N-terminal fragment of the third component of complement. 1989 , 105, 330-7		
2265	Homonuclear 2D Shift Correlated Spectroscopy. 1989 , 269-358		2
2264	2D Multiple-Quantum Spectroscopy. 1989 , 447-491		
2263	Spin topology filtration in N.M.R.. 1989 , 66, 161-197		16
2262	Non-contiguous regions of Z-DNA in a DNA dodecamer. 1989 , 17, 7923-44		2
2261	1H-nuclear magnetic resonance studies of the neuropeptide head activator. 1989 , 997, 144-53		10
2260	Conformational analysis of the sialyl alpha(2----3/6)N-acetyllactosamine structural element occurring in glycoproteins, by two-dimensional NOE 1H-NMR spectroscopy in combination with energy calculations by hard-sphere exo-anomeric and molecular mechanics force-field with hydrogen-bonding potential. <i>FEBS Journal</i> , 1989 , 178, 727-39		91
2259	A high-resolution 1H-NMR study of human transforming growth factor alpha. Structure and pH-dependent conformational interconversion. <i>FEBS Journal</i> , 1989 , 179, 629-37		27
2258	Conformation of two somatostatin analogues in aqueous solution. Study by NMR methods and circular dichroism. <i>FEBS Journal</i> , 1989 , 185, 371-81		13
2257	O-specific polysaccharides of <i>Hafnia alvei</i> lipopolysaccharides isolated from two serologically related strains: ATCC 13337 and 1187. A serological and structural study using chemical methods, gas chromatography/mass spectrometry and NMR spectroscopy at 500 MHz. <i>FEBS Journal</i> , 1989 , 186, 611-20		22
2256	Conformational Analysis of Didemnins. A multidisciplinary approach by means of X-Ray, NMR, molecular-dynamics, and molecular-mechanics techniques. 1989 , 72, 530-555		40
2255	Two-dimensional correlated spectroscopy (COSY) of intact frog muscle: spectral pattern characterization and lactate quantitation. 1989 , 11, 316-30		28
2254	Isolation and structural studies of phosphate-containing oligosaccharides from alkaline and acid hydrolysates of <i>Streptococcus pneumoniae</i> type 6B capsular polysaccharide. 1989 , 187, 267-86		12
2253	1H- and 13C-n.m.r. assignments for structural elements of xylose-containing N-linked oligosaccharides, using 1D-and 2D-n.m.r. experiments. 1989 , 186, 39-49		10
2252	Bioactive conformation of linear peptides in solution: an elusive goal?. 1989 , 28, 91-107		40

2251	Ala analogues of the cyclolinopeptide A. 1989 , 28, 371-83	12
2250	Conformational analysis of peptide T and of its C-pentapeptide fragment. 1989 , 28, 479-86	24
2249	Comparison of the conformations of cyclolinopeptide A in the solid state and in solution. 1989 , 28, 513-23	21
2248	A sesquiterpene alcohol from <i>Pallenis spinosa</i> . 1989 , 28, 849-850	16
2247	Triple-strand formation in the homopurine:homopyrimidine DNA oligonucleotides d(G-A) ₄ and d(T-C) ₄ . 1989 , 339, 637-40	247
2246	Identification and description of alpha-helical regions in horse muscle acylphosphatase by ¹ H nuclear magnetic resonance spectroscopy. 1989 , 205, 229-39	22
2245	Polypeptide chain fold of human transforming growth factor alpha analogous to those of mouse and human epidermal growth factors as studied by two-dimensional ¹ H NMR. 1989 , 28, 953-8	46
2244	Three-dimensional solution structure of a DNA duplex containing the BclI restriction sequence: two-dimensional NMR studies, distance geometry calculations, and refinement by back-calculation of the NOESY spectrum. 1989 , 28, 6996-7010	65
2243	Assignment of the proton NMR spectrum of reduced and oxidized thioredoxin: sequence-specific assignments, secondary structure, and global fold. 1989 , 28, 7074-87	95
2242	A proton nuclear magnetic resonance assignment and secondary structure determination of recombinant human thioredoxin. 1989 , 28, 7088-97	26
2241	Solution conformation of conotoxin GI determined by ¹ H nuclear magnetic resonance spectroscopy and distance geometry calculations. 1989 , 28, 4853-60	60
2240	Identification of 3,N ⁴ -propanodeoxycytidine 5'-monophosphate formed by the reaction of acrolein with deoxycytidine 5'-monophosphate. 1989 , 2, 267-71	25
2239	NMR studies of triple-strand formation from the homopurine-homopyrimidine deoxyribonucleotides d(GA) ₄ and d(TC) ₄ . 1989 , 28, 7859-70	168
2238	Sequential ¹ H NMR assignment and secondary structure determination of salmon calcitonin in solution. 1989 , 28, 7996-8002	39
2237	Spin-labeling proton NMR study on aromatic amino acid residues in the guanine nucleotide binding site of human c-Ha-ras(1-171) protein. 1989 , 28, 9550-6	12
2236	Preparative in vitro generation of lacto-series type 1 chain glycolipids catalyzed by beta 1----3-galactosyltransferase from human colonic adenocarcinoma Colo 205 cells. 1989 , 274, 14-25	22
2235	NMR studies of lantibiotics Assignment of the ¹ H-NMR spectrum of nisin and identification of interresidual contacts. 1989 , 252, 22-28	52
2234	Complete assignment of the ¹ H NMR spectrum of a synthetic zinc finger from Xfin. Sequential resonance assignments and secondary structure. 1989 , 254, 159-64	21

2233	Conformational properties of deltorphin: new features of the delta-opioid receptor. 1989 , 247, 283-8	33
2232	¹ H-NMR study of endothelin, sequence-specific assignment of the spectrum and a solution structure. 1989 , 257, 145-8	54
2231	Solution conformation of endothelin determined by nuclear magnetic resonance and distance geometry. 1989 , 257, 149-54	63
2230	Isolation and structure analysis of components from venom of the spider <i>Argiope lobata</i> . 1989 , 27, 541-9	59
2229	Homonuclear correlated spectroscopy (COSY): The basics of two-dimensional NMR. 1989 , 1, 59-72	8
2228	A ¹ H NMR determination of the solution conformation of a synthetic peptide analogue of calcium-binding site III of rabbit skeletal troponin C. 1989 , 28, 8839-47	37
2227	Determination of three-dimensional structures of proteins and nucleic acids in solution by nuclear magnetic resonance spectroscopy. 1989 , 24, 479-564	282
2226	Multiple-quantum nuclear magnetic resonance. 1989 , 176, 114-34	21
2225	Structural characterization of protein folding intermediates by proton magnetic resonance and hydrogen exchange. 1989 , 176, 446-73	88
2224	Conformational studies of d-(AAAAATTTTT) ₂ using constraints from nuclear overhauser effects and from quantitative analysis of the cross-peak fine structures in two-dimensional ¹ H nuclear magnetic resonance spectra. 1989 , 28, 1462-71	74
2223	N-terminal half of a mitochondrial presequence peptide takes a helical conformation when bound to dodecylphosphocholine micelles: a proton nuclear magnetic resonance study. 1989 , 106, 396-400	70
2222	Unstable Hoogsteen base pairs adjacent to echinomycin binding sites within a DNA duplex. 1989 , 86, 3006-10	74
2221	Two-dimensional nuclear magnetic resonance spectroscopy of proteins: an overview. 1989 , 176, 12-64	38
2220	Applications of modern-high-field NMR spectroscopy in medicinal chemistry. 1989 , 26, 355-436	6
2219	The structure of vacidin A, an aromatic heptaene macrolide antibiotic. I. Complete assignment of the ¹ H NMR spectrum and geometry of the polyene chromophore. 1989 , 42, 1631-8	24
2218	Proton nuclear magnetic resonance assignments and secondary structure determination of the ColE1 rop (rom) protein. 1990 , 29, 7402-7	21
2217	A two-dimensional ¹ H NMR study on <i>Megasphaera elsdenii</i> flavodoxin in the reduced state. Sequential assignments. <i>FEBS Journal</i> , 1990 , 187, 521-41	18
2216	NMR identification of a partial helical conformation for bombesin in solution. <i>FEBS Journal</i> , 1990 , 187, 645-50	33

2215	Secondary and tertiary structure characteristics of Megasphaera elsdenii flavodoxin in the reduced state as determined by two-dimensional ¹ H NMR. <i>FEBS Journal</i> , 1990 , 189, 589-600	12
2214	Conformational studies on the N-linked carbohydrate chain of bromelain. <i>FEBS Journal</i> , 1990 , 190, 113-22	51
2213	Proton NMR studies of apo-neocarzinostatin from Streptomyces carzinostaticus. Sequence-specific assignment and secondary structure. <i>FEBS Journal</i> , 1990 , 190, 263-71	22
2212	Sequence-specific ¹ H-NMR assignment and conformation of proteolytic fragment 163-231 of bacterioopsin. <i>FEBS Journal</i> , 1990 , 192, 321-7	33
2211	Conformation in solution of porcine brain natriuretic peptide determined by combined use of nuclear magnetic resonance and distance geometry. <i>FEBS Journal</i> , 1990 , 193, 127-34	11
2210	Nuclear-magnetic-resonance studies of human epidermal growth factor. <i>FEBS Journal</i> , 1990 , 193, 807-15	13
2209	Tertiary structure of two-electron reduced Megasphaera elsdenii flavodoxin and some implications, as determined by two-dimensional ¹ H-NMR and restrained molecular dynamics. <i>FEBS Journal</i> , 1990 , 194, 185-98	33
2208	A two-dimensional ¹ H-NMR study on Megasphaera elsdenii flavodoxin in the oxidized state and some comparisons with the two-electron-reduced state. <i>FEBS Journal</i> , 1990 , 194, 199-216	18
2207	Opioid conformations: The remarkable effect of C-14 substituents on ring c conformations of 6β and 6α epimers of 6-substituted epoxymorphinan opioid ligands. 1990 , 27, 385-389	6
2206	Solution phase conformation studies of the prekallikrein binding domain of high molecular weight kininogen. 1990 , 9, 647-59	15
2205	Cis-trans isomerization of an angiotensin I-converting enzyme inhibitor. An enzyme kinetic and nuclear magnetic resonance study. 1990 , 1041, 22-30	12
2204	Proton nuclear magnetic resonance study of human interleukin 6: chemical modifications and partial spectral assignments for the aromatic residues. 1990 , 1041, 243-9	16
2203	Structural characterization of an arabinose-containing heptadecasaccharide enzymically isolated from sycamore extracellular xyloglucan. 1990 , 197, 139-158	39
2202	NMR investigations of protein structure. 1990 , 22, 101-139	109
2201	Conformational studies of cyclic peptide structures in solution from ¹ H-Nmr data by distance geometry calculation and restrained energy minimization. 1990 , 29, 1387-400	9
2200	Determination of the conformation of Lewis blood group oligosaccharides by simulation of two-dimensional nuclear Overhauser data. 1990 , 30, 1123-38	66
2199	Entflechtung Berlagernder Multipletts in zweidimensionalen NMR-Korrelationspektren durch selektive Inversion der Spins der Kopplungspartner. 1990 , 102, 576-579	5
2198	Solution Structure of [Me-L-Leu7]Didemnin B Determined by NMR Spectroscopy and Refined by MD Calculation. 1990 , 73, 25-47	23

2197	Identification by n.m.r. spectroscopy of oligosaccharides obtained by treatment of the hairy regions of apple pectin with rhamnogalacturonase. 1990 , 206, 131-44	104
2196	Structural analysis of xyloglucan oligosaccharides by ¹ H-n.m.r. spectroscopy and fast-atom-bombardment mass spectrometry. 1990 , 200, 9-31	196
2195	Structure determination of polypeptides and proteins by two-dimensional nuclear magnetic resonance spectroscopy. 1990 , 164, 180-192	3
2194	The time dependence of coherence transfer in homonuclear isotropic mixing experiments. 1990 , 87, 110-131	5
2193	Rapid-pulsing artifacts in double-quantum-filtered COSY. 1990 , 88, 177-185	53
2192	3D TOCSY-TOCSY processing using linear prediction, as a potential technique for automated assignment. 1990 , 89, 184-190	1
2191	Cluster analysis and multiplet pattern recognition in two-dimensional NMR spectra. 1990 , 89, 543-552	1
2190	Oligosaccharide conformations: Application of NMR and energy calculations. 1990 , 22, 55-81	90
2189	Conformational requirement of signal sequences functioning in yeast: circular dichroism and ¹ H nuclear magnetic resonance studies of synthetic peptides. 1990 , 29, 8998-9006	36
2188	Complex formation of peptide antibiotic Ro09-0198 with lysophosphatidylethanolamine: ¹ H NMR analyses in dimethyl sulfoxide solution. 1990 , 29, 113-8	59
2187	Solution structures of human transforming growth factor alpha derived from ¹ H NMR data. 1990 , 29, 7805-13	64
2186	Complete structure of the polysaccharide from <i>Streptococcus sanguis</i> J22. 1990 , 29, 234-48	61
2185	Sequential ¹ H NMR assignments and secondary structure of the B domain of staphylococcal protein A: structural changes between the free B domain in solution and the Fc-bound B domain in crystal. 1990 , 29, 8787-93	68
2184	¹ H NMR studies of the glucocorticoid receptor DNA-binding domain: sequential assignments and identification of secondary structure elements. 1990 , 29, 9015-23	56
2183	Conformational preferences of synthetic peptides derived from the immunodominant site of the circumsporozoite protein of <i>Plasmodium falciparum</i> by ¹ H NMR. 1990 , 29, 7828-37	79
2182	Multinuclear magnetic resonance studies of the 2Fe.2S* ferredoxin from <i>Anabaena</i> species strain PCC 7120. 1. Sequence-specific hydrogen-1 resonance assignments and secondary structure in solution of the oxidized form. 1990 , 29, 3993-4004	57
2181	Assignment of the ¹ H and ¹⁵ N NMR spectra of <i>Rhodobacter capsulatus</i> ferrocyclochrome c2. 1990 , 29, 2278-90	30
2180	Structural properties of apocytochrome b5: presence of a stable native core. 1990 , 29, 1984-9	32

2179	Mechanism of adenylate kinase. Structural and functional demonstration of arginine-138 as a key catalytic residue that cannot be replaced by lysine. 1990 , 29, 6385-92		37
2178	Structural Studies of Biomolecules at High Field. 1990 , 91-141		
2177	Studies of synthetic helical peptides using circular dichroism and nuclear magnetic resonance. 1990 , 215, 607-22		157
2176	Secondary structure of a leucine zipper determined by nuclear magnetic resonance spectroscopy. 1990 , 29, 2891-4		145
2175	Determination of the three-dimensional structure of the Antennapedia homeodomain from <i>Drosophila</i> in solution by 1H nuclear magnetic resonance spectroscopy. 1990 , 214, 183-97		116
2174	Determining local conformational variations in DNA. Nuclear magnetic resonance structures of the DNA duplexes d(CGCCTAATCG) and d(CGTCACGCGC) generated using back-calculation of the nuclear Overhauser effect spectra, a distance geometry algorithm and constrained molecular dynamics. 1990 , 214, 711-21		66
2173	Amide proton exchange in human metallothionein-2 measured by nuclear magnetic resonance spectroscopy. 1990 , 214, 781-6		20
2172	Complete sequence-specific 1H NMR assignments for human insulin. 1990 , 29, 2906-13		76
2171	Structural elements pertinent to the interaction of cyclosporin A with its specific receptor protein, cyclophilin. 1990 , 40, 131-40		20
2170	Solution conformation of leiurotoxin I (scyllatoxin) by 1H nuclear magnetic resonance. Resonance assignment and secondary structure. 1990 , 260, 249-53		45
2169	Structure of the DNA binding wing of the gene-V encoded single- stranded DNA binding protein of the filamentous bacteriophage M13. 1990 , 261, 1-4		28
2168	1H nuclear magnetic resonance study of the solution conformation of an antibacterial protein, sapecin. 1990 , 269, 413-20		84
2167	New features of the delta opioid receptor: conformational properties of deltorphin I analogues. <i>Biochemical and Biophysical Research Communications</i> , 1990 , 169, 617-22	3-4	40
2166	A 1H NMR study of the solution conformation of the neuropeptide galanin. <i>Biochemical and Biophysical Research Communications</i> , 1990 , 166, 1102-9	3-4	43
2165	A 1H-NMR study of the solution conformation of Icaria chemotactic peptide and its [Lys7] analog: effects on the physiological activity of a substitution of proline to lysine at position 7. <i>Biochemical and Biophysical Research Communications</i> , 1990 , 168, 596-603	3-4	1
2164	Three-dimensional structure of human [¹¹³ Cd7]metallothionein-2 in solution determined by nuclear magnetic resonance spectroscopy. 1990 , 214, 765-79		144
2163	The conformation of porcine-brain natriuretic peptide by two-dimensional NMR spectroscopy. <i>FEBS Journal</i> , 1991 , 201, 183-90		8
2162	Solution conformation of endothelin-3 by (1)H NMR and distance geometry calculations. 1991 , 18, 491-6		27

2161	Structure of the smooth muscle myosin light-chain kinase calmodulin-binding domain peptide bound to calmodulin. 1991 , 30, 10078-84	56
2160	Design of a helix-bundle cross-link: NMR and UV-visible spectroscopic analyses and molecular modeling of ring-oxidized retinals. 1991 , 30, 2976-88	1
2159	Determination of the three-dimensional solution structure of barnase using nuclear magnetic resonance spectroscopy. 1991 , 30, 8697-701	108
2158	¹ H and ¹⁵ N resonance assignments of oxidized flavodoxin from <i>Anacystis nidulans</i> with 3D NMR. 1991 , 30, 7718-30	32
2157	Complete structure of the cell surface polysaccharide of <i>Streptococcus oralis</i> ATCC 10557: a receptor for lectin-mediated interbacterial adherence. 1991 , 30, 6528-40	39
2156	Solution structure of the basic region from the transcriptional activator GCN4. 1991 , 30, 1310-7	80
2155	Evidence for two interconverting protein isomers in the methotrexate complex of dihydrofolate reductase from <i>Escherichia coli</i> . 1991 , 30, 2184-91	48
2154	¹ H assignments and secondary structure determination of the soybean trypsin/chymotrypsin Bowman-Birk inhibitor. 1991 , 30, 3356-64	48
2153	The DNA sequence at echinomycin binding sites determines the structural changes induced by drug binding: NMR studies of echinomycin binding to [d(ACGTACGT)] ₂ and [d(TCGATCGA)] ₂ . 1991 , 30, 2483-94	60
2152	Structure-function relationships in human epidermal growth factor studied by site-directed mutagenesis and ¹ H NMR. 1991 , 30, 8891-8	49
2151	Complete structure of the cell surface polysaccharide of <i>Streptococcus oralis</i> C104: a 600-MHz NMR study. 1991 , 30, 8568-77	49
2150	Structure of the propeptide of prothrombin containing the gamma-carboxylation recognition site determined by two-dimensional NMR spectroscopy. 1991 , 30, 9835-41	36
2149	C-terminal structure and mobility of rabbit skeletal muscle light meromyosin as studied by one- and two-dimensional ¹ H NMR spectroscopy and X-ray small-angle scattering. 1991 , 30, 8083-91	12
2148	Sequential NMR resonance assignment and structure determination of the Kunitz-type inhibitor domain of the Alzheimer's beta-amyloid precursor protein. 1991 , 30, 10467-78	15
2147	ADR1a, a zinc finger peptide, exists in two folded conformations. 1991 , 30, 3365-71	20
2146	The secondary structure in solution of acyl-coenzyme A binding protein from bovine liver using ¹ H nuclear magnetic resonance spectroscopy. 1991 , 30, 10654-63	28
2145	Structural determination of oligosaccharides derived from lipooligosaccharide of <i>Neisseria gonorrhoeae</i> F62 by chemical, enzymatic, and two-dimensional NMR methods. 1991 , 30, 10566-75	83
2144	¹ H, ¹⁵ N, and ¹³ C NMR signal assignments of IIIgIc, a signal-transducing protein of <i>Escherichia coli</i> , using three-dimensional triple-resonance techniques. 1991 , 30, 10043-57	67

2143	Tertiary structure of conotoxin GlIIA in aqueous solution. 1991 , 30, 6908-16	87
2142	Dihydrofolate reductase: sequential resonance assignments using 2D and 3D NMR and secondary structure determination in solution. 1991 , 30, 6330-41	30
2141	Polypeptide backbone resonance assignments and secondary structure of <i>Bacillus subtilis</i> enzyme IIIglc determined by two-dimensional and three-dimensional heteronuclear NMR spectroscopy. 1991 , 30, 6896-907	52
2140	Domain interactions in human plasminogen studied by proton NMR. 1991 , 278, 17-22	13
2139	Zinc is required for folding and binding of a single zinc finger to DNA. 1991 , 279, 289-94	20
2138	Syringopeptins, new phytotoxic lipodepsipeptides of <i>Pseudomonas syringae</i> pv. <i>syringae</i> . 1991 , 291, 109-12	111
2137	(14-38, 30-51) double-disulphide intermediate in folding of bovine pancreatic trypsin inhibitor: a two-dimensional ¹ H nuclear magnetic resonance study. 1991 , 222, 353-71	75
2136	Two-dimensional ¹ H nuclear magnetic resonance study of the (5-55) single-disulphide folding intermediate of bovine pancreatic trypsin inhibitor. 1991 , 222, 373-90	82
2135	Interaction of troponin I and troponin C. Use of the two-dimensional nuclear magnetic resonance transferred nuclear Overhauser effect to determine the structure of the inhibitory troponin I peptide when bound to skeletal troponin C. 1991 , 222, 405-21	71
2134	Sequence-specific ¹ H n.m.r. assignments and determination of the three-dimensional structure of reduced <i>Escherichia coli</i> glutaredoxin. 1991 , 221, 1311-24	89
2133	Three-dimensional structural analysis of the group B polysaccharide of <i>Neisseria meningitidis</i> 6275 by two-dimensional NMR: the polysaccharide is suggested to exist in helical conformations in solution. 1991 , 30, 851-7	71
2132	Accurate measurements of coupling constants from two-dimensional nuclear magnetic resonance spectra of proteins and determination of phi-angles. 1991 , 217, 731-6	150
2131	Comparative NMR studies on cardiac troponin C and a mutant incapable of binding calcium at site II. 1991 , 30, 10236-45	30
2130	Purification and structure determination of the bacterial mating inhibitor for <i>Chlamydomonas reinhardtii</i> and <i>Alexandrium catenella</i> .. 1991 , 57, 307-314	6
2129	Anantin—a peptide antagonist of the atrial natriuretic factor (ANF). II. Determination of the primary sequence by NMR on the basis of proton assignments. 1991 , 44, 172-80	60
2128	Identification of valine/leucine/isoleucine and threonine/alanine/glycine proton-spin systems of <i>Escherichia coli</i> adenylate kinase by selective deuteration and selective protonation. <i>Biochemical Journal</i> , 1991 , 273(Pt 2), 311-6	3.8 2
2127	Solvent signal suppression in NMR. 1991 , 23, 135-209	176
2126	Cyanidin 3-[6-(p-coumaroyl)-2-(xylosyl)-glucoside]-5-glucoside and other anthocyanins from fruits of <i>Sambucus canadensis</i> . 1991 , 30, 4137-41	42

2125	Applications of COSY to paramagnetic heme-containing systems. 1991 , 95, 244-252	7
2124	Optimization of two-dimensional NMR by matched accumulation. 1991 , 95, 1-9	4
2123	Solution structure of the major binding protein for the immunosuppressant FK506. 1991 , 351, 248-50	138
2122	Synthesis and characterization of cyclic peptides with hydrolytic activity. 1991 , 32, 5337-5340	7
2121	Three-dimensional structure of natural charybdotoxin in aqueous solution by 1H-NMR. Charybdotoxin possesses a structural motif found in other scorpion toxins. <i>FEBS Journal</i> , 1991 , 196, 19-28	151
2120	NMR and circular dichroic studies on the solution conformation of a synthetic peptide derived from the calmodulin-binding domain of Bordetella pertussis adenylate cyclase. <i>FEBS Journal</i> , 1991 , 196, 67-72	11
2119	A spectroscopic analysis of the Pro35----Ala mutant of Rhodobacter capsulatus cytochrome c2. The strictly conserved Pro35 is not structurally essential. <i>FEBS Journal</i> , 1991 , 196, 653-61	15
2118	Proton nuclear magnetic resonance investigation of adrenodoxin. Assignment of aromatic resonances and evidence for a conformational similarity with ferredoxin from Spirulina platensis. <i>FEBS Journal</i> , 1991 , 197, 747-57	23
2117	Nuclear magnetic resonance studies of recombinant Escherichia coli glutaredoxin. Sequence-specific assignments and secondary structure determination of the oxidized form. <i>FEBS Journal</i> , 1991 , 200, 369-77	16
2116	Sequence-specific 1H-NMR assignments and secondary structure of the lipoyl domain of the Bacillus stearothermophilus pyruvate dehydrogenase multienzyme complex. <i>FEBS Journal</i> , 1991 , 201, 203-9	30
2115	Assignment of the 600-MHz 1H-NMR spectrum of amicyanin from Thiobacillus versutus by two-dimensional NMR methods provides information on secondary structure. <i>FEBS Journal</i> , 1991 , 201, 695-702	13
2114	Isolation and characterization of a new variant of surfactin, the [Val7]surfactin. <i>FEBS Journal</i> , 1991 , 202, 101-6	79
2113	1H NMR studies of echistatin in solution. Sequential resonance assignments and secondary structure. <i>FEBS Journal</i> , 1991 , 202, 315-21	33
2112	Nuclear magnetic resonance studies of the snake toxin echistatin. 1H resonance assignments and secondary structure. <i>FEBS Journal</i> , 1991 , 202, 323-8	40
2111	The secondary structure of echistatin from 1H-NMR, circular-dichroism and Raman spectroscopy. <i>FEBS Journal</i> , 1991 , 202, 329-38	37
2110	Sequence-specific 1H-NMR assignment and secondary structure of the Tyr41----His mutant of the single-stranded DNA binding protein, gene V protein, encoded by the filamentous bacteriophage M13. <i>FEBS Journal</i> , 1991 , 202, 349-60	31
2109	Characterization of wild-type and mutant M13 gene V proteins by means of 1H-NMR. <i>FEBS Journal</i> , 1991 , 200, 139-48	22
2108	Assignment of structures to oligosaccharides produced by enzymic degradation of a beta-D-glucan from barley by 1H- and 13C-n.m.r. spectroscopy. 1991 , 211, 219-33	43

2107	Structural analysis of a second acidic exopolysaccharide of <i>Rhizobium meliloti</i> that can function in alfalfa root nodule invasion. 1991 , 210, 339-47	23
2106	Coenzyme F430 from Methanogenic Bacteria: Complete Assignment of Configuration Based on an X-Ray Analysis of 12,13-Diepi-F430 Pentamethyl Ester and on NMR Spectroscopy. 1991 , 74, 697-716	94
2105	Solution structure of salmon calcitonin. 1991 , 31, 233-41	63
2104	New insights on mu/delta selectivity of opioid peptides: conformational analysis of deltorphin analogues. 1991 , 31, 751-60	43
2103	Ion binding of cyclolinopeptide A: an NMR and CD conformational study. 1991 , 31, 761-7	27
2102	The solution structure of the lantibiotic gallidermin. 1991 , 31, 803-11	69
2101	Zum R�ssel des Chlorophyllabbaus: Die Konstitution eines secoporphinoiden Kataboliten. 1991 , 103, 1354-1357	53
2100	Detection of long-range couplings in oligomers and evaluation of coupling constants in polymers. Application of the 1D COSY technique. 1991 , 29, 22-28	2
2099	Structural elucidation of germine esters isolated from <i>Veratrum nigrum</i> L. Total assignment of ¹ H and ¹³ C nuclear magnetic resonances by two-dimensional NMR techniques. 1991 , 29, 100-112	9
2098	Structure of two stereoisomers of Pt(trans-MeO ₂ CCH ₂ CHCO ₂ Me) ₂ [P(cyclohexyl) ₃]: Complete assignment of the 11 non-equivalent tricyclohexylphosphine protons using DQF-COSY and ¹³ C, ¹ H two-dimensional methods. 1991 , 29, 113-119	4
2097	¹ H and ¹³ C NMR assignments of the antitumor macrolide bryostatin 1. 1991 , 29, 366-374	7
2096	Assignment of proton NMR resonances and conformational analysis of [Lys ⁸]-vasopressin homologues. 1991 , 29, 580-586	13
2095	Solution structure of phalloidin obtained by NMR spectroscopy in [D ₆]DMSO and molecular dynamics calculation in vacuo and in water. 1991 , 1991, 179-184	11
2094	Controlled reduction of acarbose: conformational analysis of acarbose and the resulting saturated products. 1991 , 221, 1-16	21
2093	Removal of diagonal peaks in two-dimensional NMR spectra by means of digital filtering. 1991 , 95, 178-183	2
2092	Enhancement of global symmetries in two-dimensional NMR spectra. 1991 , 91, 155-164	
2091	Computer-assisted assignment of 2D ¹ H NMR spectra of proteins: basic algorithms and application to phoratoxin B. 1991 , 1, 23-47	39
2090	Two-dimensional NMR study of the conformation of (34-65)bacterioopsin polypeptide in SDS micelles. 1991 , 1, 313-22	36

2089	Proton NMR studies on the chromophore binding structure in neocarzinostatin complex. 1991 , 32, 3175-3178	20
2088	Structure of petanin, an acylated anthocyanin isolated from solanum tuberosum, using homo- and hetero-nuclear two-dimensional nuclear magnetic resonance techniques. 1991 , 2, 230-236	35
2087	Photoreaction of Phenacyl Aryl Ether Type Lignols. 1991 , 11, 263-289	13
2086	Characterization of a posttranslational fucosylation in the growth factor domain of urinary plasminogen activator. 1991 , 88, 3992-6	59
2085	Solution conformation of endothelin determined by means of 1H-NMR spectroscopy and distance geometry calculations. 1991 , 4, 509-18	67
2084	Active conformation of an insect neuropeptide family. 1991 , 88, 4518-22	76
2083	Nuclear magnetic resonance studies and molecular dynamics simulations of the solution conformation of a 'designed', alpha-helical peptide. 1992 , 5, 333-41	3
2082	Bivalent-metal binding to CheY protein. Effect on protein conformation. <i>Biochemical Journal</i> , 1992 , 287 (Pt 2), 521-31	3.8 21
2081	Purification and determination of the structure of capsular polysaccharide of <i>Vibrio vulnificus</i> M06-24. 1992 , 174, 2620-30	80
2080	Solution structure of a DNA-binding unit of Myb: a helix-turn-helix-related motif with conserved tryptophans forming a hydrophobic core. 1992 , 89, 6428-32	202
2079	NMR studies of amyloid beta-peptides: proton assignments, secondary structure, and mechanism of an alpha-helix----beta-sheet conversion for a homologous, 28-residue, N-terminal fragment. 1992 , 31, 5621-31	190
2078	1H NMR-based determination of the secondary structure of porcine pancreatic spasmolytic polypeptide: one of a new family of "trefoil" motif containing cell growth factors. 1992 , 31, 1998-2004	46
2077	NMR characterization of a diamagnetic model of unliganded alpha chains from human hemoglobin. 1992 , 74, 845-51	2
2076	Analysis of side-chain organization on a refined model of charybdotoxin: structural and functional implications. 1992 , 31, 7756-64	154
2075	Determination of the three-dimensional structure of iberiotoxin in solution by 1H nuclear magnetic resonance spectroscopy. 1992 , 31, 8151-9	99
2074	1H NMR assignment and secondary structure of the cell adhesion type III module of fibronectin. 1992 , 31, 2068-73	91
2073	Structural comparison of apomyoglobin and metaquomyoglobin: pH titration of histidines by NMR spectroscopy. 1992 , 31, 6481-91	83
2072	Characterization of a 12-mer duplex d(GGCGGAGTTAGG).d(CCTAACTCCGCC) containing a highly reactive (+)-CC-1065 sequence by 1H and 31P NMR, hydroxyl-radical footprinting, and NOESY restrained molecular dynamics calculations. 1992 , 5, 167-82	29

2071	1H and 15N NMR characterization of free and bound states of an amphiphilic peptide interacting with calmodulin. 1992 , 31, 229-36		18
2070	Structural characterization of the cell surface lipooligosaccharides from a nontypable strain of Haemophilus influenzae. 1992 , 31, 4515-26		130
2069	Characterization of the secondary structure of calmodulin in complex with a calmodulin-binding domain peptide. 1992 , 31, 1443-51		42
2068	Conformation of a pentacosapeptide representing the RNA-binding N-terminus of cowpea chlorotic mottle virus coat protein in the presence of oligophosphates: a two-dimensional proton nuclear magnetic resonance and distance geometry study. 1992 , 31, 9177-82		17
2067	Alpha-cobratoxin: proton NMR assignments and solution structure. 1992 , 31, 4867-75		41
2066	Two-dimensional NMR, circular dichroism, and fluorescence studies of PP-50, a synthetic ATP-binding peptide from the beta-subunit of mitochondrial ATP synthase. 1992 , 31, 7915-21		14
2065	1H NMR studies of a biosynthetic lacto-ganglio hybrid glycosphingolipid: confirmation of structure, interpretation of "anomalous" chemical shifts, and evidence for interresidue amide-amide hydrogen bonding. 1992 , 31, 1069-80		18
2064	Assignment of the 1H NMR spectrum and secondary structure elucidation of the single-stranded DNA binding protein encoded by the filamentous bacteriophage IKe. 1992 , 31, 1254-62		8
2063	Immunogenic peptides corresponding to the dominant antigenic region alanine-597 to cysteine-619 in the transmembrane protein of simian immunodeficiency virus have a propensity to fold in aqueous solution. 1992 , 31, 1458-63		13
2062	Structural analysis of lipooligosaccharide produced by Neisseria gonorrhoeae, strain MS11mk (variant A): a precursor for a gonococcal lipooligosaccharide associated with virulence. 1992 , 31, 12760-8		42
2061	The secondary structure of the colicin E3 immunity protein as studied by 1H-1H and 1H-15N two-dimensional NMR spectroscopy. 1992 , 31, 5578-86		9
2060	Two-dimensional NMR studies of d(GGTTAATGCGGT).d(ACCGCATTAACC) complexed with the minor groove binding drug SN-6999. 1992 , 31, 4406-13		30
2059	Structural consequences of the natural substitution, E9K, on reactive-site-hydrolyzed squash (Cucurbita maxima) trypsin inhibitor (CMTI), as studied by two-dimensional NMR. 1992 , 31, 4965-9		7
2058	A receptor-binding peptide from human interleukin-6: isolation and a proton nuclear magnetic resonance study. <i>Biochemical and Biophysical Research Communications</i> , 1992 , 189, 211-20	3-4	13
2057	A 1H-NMR study of the solution conformation of cyclo(GRGDSPA): conformational effects on the physiological activity. <i>Biochemical and Biophysical Research Communications</i> , 1992 , 182, 966-73	3-4	12
2056	Design and synthesis of an alpha-helical peptide containing periodic proline residues. 1992 , 298, 233-6		6
2055	Solution structure of FK506 bound to FKBP-12. 1992 , 302, 89-96		34
2054	A study of D52S hen lysozyme-GlcNAc oligosaccharide complexes by NMR spectroscopy and electrospray mass spectrometry. 1992 , 296, 153-7		27

2053	Structural determinants of Cys2His2 zinc fingers. 1992 , 296, 11-5	18
2052	Beta-turns induced in bradykinin by (S)-alpha-methylproline. 1992 , 297, 216-20	27
2051	¹ H NMR study on the conformation of bacitracin A in aqueous solution. 1992 , 305, 105-9	14
2050	The fine structure of two DNA dodecamers containing the cAMP responsive element sequence and its inverse. Nuclear magnetic resonance and molecular simulation studies. 1992 , 227, 852-75	48
2049	Nuclear magnetic resonance solution structure of hirudin(1-51) and comparison with corresponding three-dimensional structures determined using the complete 65-residue hirudin polypeptide chain. 1992 , 228, 1193-205	50
2048	Solution structure of the fibrin binding finger domain of tissue-type plasminogen activator determined by ¹ H nuclear magnetic resonance. 1992 , 225, 821-33	47
2047	Nuclear magnetic resonance solution structure of the alpha-neurotoxin from the black mamba (<i>Dendroaspis polylepis polylepis</i>). 1992 , 227, 1118-35	33
2046	Three-dimensional solution structure of an insulin dimer. A study of the B9(Asp) mutant of human insulin using nuclear magnetic resonance, distance geometry and restrained molecular dynamics. 1992 , 227, 1146-63	48
2045	Folding of peptide fragments comprising the complete sequence of proteins. Models for initiation of protein folding. I. Myohemerythrin. 1992 , 226, 795-817	351
2044	Folding of peptide fragments comprising the complete sequence of proteins. Models for initiation of protein folding. II. Plastocyanin. 1992 , 226, 819-35	215
2043	¹ H nuclear magnetic resonance studies of the interaction of urea with hen lysozyme. Origins of the conformational change induced in hen lysozyme by N-acetylglucosamine oligosaccharides. 1992 , 227, 9-14	25
2042	Comparison of the solution conformations of human [Zn7]-metallothionein-2 and [Cd7]-metallothionein-2 using nuclear magnetic resonance spectroscopy. 1992 , 225, 433-43	91
2041	Determination of the nuclear magnetic resonance solution structure of the DNA-binding domain (residues 1 to 69) of the 434 repressor and comparison with the X-ray crystal structure. 1992 , 223, 743-67	62
2040	The internalization signal in the cytoplasmic tail of lysosomal acid phosphatase consists of the hexapeptide PGRHV.. <i>EMBO Journal</i> , 1992 , 11, 4391-4399	13 52
2039	Determination of the structure of the nucleocapsid protein NCp7 from the human immunodeficiency virus type 1 by ¹ H NMR.. <i>EMBO Journal</i> , 1992 , 11, 3059-3065	13 198
2038	A novel difucosylated neutral glycosphingolipid from the eggs of the sea urchin, <i>Hemicentrotus pulcherrimus</i> : II. Structural determination by two-dimensional NMR. 1992 , 112, 286-9	7
2037	The application of two-dimensional nuclear magnetic resonance methodologies to the structure solution of the new natural product tenerol acetate from <i>Laurencia tenera</i> . 1992 , 3, 263-269	10
2036	The three-dimensional structure of the first EGF-like module of human factor IX: comparison with EGF and TGF-alpha. 1992 , 1, 81-90	65

2035	Comparison of solution structures of mutant bovine pancreatic trypsin inhibitor proteins using two-dimensional nuclear magnetic resonance. 1992 , 1, 91-106	21
2034	Characterization of taxol in methylene chloride by nmr spectroscopy. 1992 , 33, 1169-1172	45
2033	Structure elucidation of a plant metabolite of 4-desoxynivalenol. 1992 , 3, 953-960	65
2032	Hydrogenation of vermistatin. 1992 , 123, 251-256	5
2031	NMR studies of a tetrasaccharide from hyaluronic acid. 1992 , 237, 271-81	25
2030	Analysis of conformationally restricted models for the (1---6)-branch of asparagine-linked oligosaccharides by n.m.r.-spectroscopy and HSEA calculation. 1992 , 228, 1-20	27
2029	A nuclear magnetic resonance spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of Chlamydia lipopolysaccharides. 1992 , 229, 213-24	61
2028	Multiple-quantum NMR methods. 1992 , 24, 295-375	50
2027	Diterpenoids from Isodon adenoloma. 1992 , 31, 4237-4240	24
2026	NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. 1992 , 31, 3307-30	1173
2025	Suppression of artifacts in phase-sensitive COSY. 1992 , 100, 469-483	1
2024	Structural and functional characterization of the mutant Escherichia coli glutaredoxin (C14---S) and its mixed disulfide with glutathione. 1992 , 31, 9288-93	199
2023	Interaction of troponin I and troponin C: use of the two-dimensional transferred nuclear Overhauser effect to determine the structure of a Gly-110 inhibitory troponin I peptide analog when bound to cardiac troponin C. 1992 , 1160, 35-54	23
2022	Toxin III of the scorpion Androctonus australis Hector: proton nuclear magnetic resonance assignments and secondary structure. 1992 , 2, 57-70	16
2021	Solution conformation of human big endothelin-1. 1992 , 2, 407-20	12
2020	¹ H resonance assignments, secondary structure and general topology of single-chain monellin in solution as determined by ¹ H 2D-NMR. 1992 , 2, 557-72	15
2019	Phase cycling in double-quantum-filtered COSY. 1992 , 96, 551-562	1
2018	Characterization of seven xyloglucan oligosaccharides containing from seventeen to twenty glycosyl residues. 1992 , 227, 45-71	44

2017	Relationship between ¹ H and ¹³ C NMR chemical shifts and the secondary and tertiary structure of a zinc finger peptide. 1992 , 2, 307-22	37
2016	Positive theta-angles in proteins by nuclear magnetic resonance spectroscopy. 1992 , 2, 227-33	85
2015	Sequence-specific resonance assignment and secondary structure of (1-71) bacterioopsin. 1992 , 2, 161-71	19
2014	Extended type-1 chain glycosphingolipid antigens. Isolation and characterization of trifucosyl-Leb antigen (III4V4VI2Fuc3Lc6). <i>FEBS Journal</i> , 1992 , 203, 577-86	23
2013	Sequence specific ¹ H-NMR assignments and secondary structure of a carboxy-terminal functional fragment of apolipoprotein CII. <i>FEBS Journal</i> , 1992 , 205, 223-31	11
2012	Proton NMR studies of bovine serum albumin. Assignment of spin systems. <i>FEBS Journal</i> , 1992 , 205, 631-43	26
2011	Sequential assignment of the proton NMR spectrum of isolated alpha(CO) chains from human adult hemoglobin. <i>FEBS Journal</i> , 1992 , 205, 661-70	11
2010	The three-dimensional structure of guanine-specific ribonuclease F1 in solution determined by NMR spectroscopy and distance geometry. <i>FEBS Journal</i> , 1992 , 208, 41-51	5
2009	¹ H-NMR assignments and local environments of aromatic residues in bovine, human and guinea pig variants of alpha-lactalbumin. <i>FEBS Journal</i> , 1992 , 210, 699-709	43
2008	Elucidation of the structure and conformation of a methylated tetrasaccharide-alditol acetate by n.m.r. spectroscopy. 1992 , 226, 15-27	2
2007	¹ H-N.m.r. spectroscopy of O-carboxymethyl derivatives of d-glucose. 1992 , 228, 433-437	10
2006	Complete assignment of the ¹ H, ¹³ C and ¹⁵ N spectra of bradykinin derivatives. 1992 , 30, 683-685	2
2005	Application of HMQC to the measurement of J(H, H) homonuclear coupling constants. 1992 , 30, 711-715	6
2004	Proton and carbon chemical shift assignment and dynamic investigation of the macrolide antibiotic tylosin. 1992 , 30, 954-957	6
2003	Spectral assignment of 2'-O-β-D-glucopyranosylsaikosaponin b2 by two-dimensional NMR techniques. 1992 , 30, 1097-1103	10
2002	¹³ C and ¹ H NMR study of the metabolic degradation of 4-pentenoate in different dog nephron segments. 1992 , 28, 137-44	1
2001	Structural analysis of tamarind seed xyloglucan oligosaccharides using beta-galactosidase digestion and spectroscopic methods. 1993 , 248, 285-301	108
2000	Synthesis and NMR spectroscopic investigation of oligosaccharides containing Kdo and L-glycero-D-manno-heptopyranosyl residues. 1993 , 243, 273-91	8

1999	Synthesis, conformation, and activity of HCO-Met-delta Z Leu-Phe-OMe, an active analogue of chemotactic N-formyltripeptides. 1993 , 33, 437-51	31
1998	Bioactive peptides: conformational study of a cystinyl cycloheptapeptide in its free and calcium complexed forms. 1993 , 33, 1083-91	15
1997	The solution conformation of Ac-Pen-Arg-Gly-Asp-Cys-OH, a potent fibrinogen receptor antagonist. 1993 , 33, 1287-97	13
1996	Warum pentose-und nicht hexose-nucleinsäuren? Teil III. Oligo(2',3'-dideoxy-β-D-glucopyranosyl) nucleotide (βomo-DNS) Paarungseigenschaften. 1993 , 76, 259-352	109
1995	The Dynamic Behaviour and NMR Solution structures of complexes of the type (Bisphosphine)(cycloocta-1,5-diene)iridium(I) and the X-ray crystal structure of (cycloocta-1,5-diene)(P-norphos)iridium(I) hexafluorophosphate. 1993 , 76, 788-803	16
1994	The Effect of Electron-Donating and Electron-Withdrawing Substituents on 1H-and 13C-NMR chemical shifts of novel 7?-aryl-substituted 7?-apo-β-carotenes. 1993 , 76, 1928-1938	6
1993	Synthesis and NMR-Spectroscopic Structure Determination of Novel 7,7?-Diphenyl-7,7?-diapocarotenoids. 1993 , 76, 1939-1948	12
1992	Assignment of the 1H and 13C NMR spectra of the C21 steroids 12-β-acetyltenacigenin A and tenacigenin A by two-dimensional NMR techniques and computer modeling. 1993 , 31, 215-221	14
1991	NMR chemical shift assignments for 7?-substituted aromatic-7?-apo-β-carotenes. 1993 , 31, 516-517	1
1990	Determination of the structure of alkoxy silane polymers in solution by natural abundance two dimensional 29Si-29Si INEPT DQF COSY NMR. 1993 , 31, 623-631	26
1989	Solution structures of [IrH2(1,5-cyclooctadiene) (bisphosphine)](CF3SO3) complexes. Homo- and heteronuclear long-range couplings from hydride and phosphorus spins to cyclooctadiene protons. 1993 , 31, 677-684	9
1988	Conformational analysis of four β-methylphenylalanine stereoisomers in a bioactive peptide by z-filtered relay NMR spectroscopy. 1993 , 31, 1072-1076	16
1987	Sequential assignments by 1H 2D NMR of oxidized ferredoxins from Clostridium pasteurianum and Clostridium acidurici. 1993 , 31, S27-S33	13
1986	Saikosaponin derivatives from Bupleurum wenchuanense. 1993 , 33, 1197-1205	21
1985	600 MHz 1H-NMR spectroscopy of human cerebrospinal fluid: effects of sample manipulation and assignment of resonances. 1993 , 11, 651-64	74
1984	3-O-Acetylbeiwutine from Aconitum liaotungeuse. 1993 , 32, 767-770	4
1983	1H nuclear magnetic resonance determination of the membrane-bound conformation of senktide, a highly selective neurokinin B agonist. 1993 , 3, 443-61	17
1982	Sequence-specific 1H and 15N resonance assignments and secondary structure of GDP-bound human c-Ha-Ras protein in solution. 1993 , 3, 165-84	24

1981	1H NMR studies of the mercuric ion binding protein MerP: sequential assignment, secondary structure and global fold of oxidized MerP. 1993 , 3, 613-26	25
1980	Oxytocin solution structure changes upon protonation of the N-terminus in dimethyl sulfoxide. 1993 , 3, 653-73	17
1979	Extensive distance geometry calculations with different NOE calibrations: new criteria for structure selection applied to Sandostatin and BPTI. 1993 , 3, 307-24	44
1978	Measurement of proton relaxation rates in proteins. 1993 , 3, 335-348	32
1977	The solution structure of the human retinoic acid receptor-beta DNA-binding domain. 1993 , 3, 1-17	49
1976	Secondary structural features of modules M2 and M3 of barnase in solution by NMR experiment and distance geometry calculation. 1993 , 16, 341-56	20
1975	Solution structure of tertiapin determined using nuclear magnetic resonance and distance geometry. 1993 , 17, 124-37	27
1974	Use of proline mutants to help solve the NMR solution structure of type III antifreeze protein. 1993 , 2, 1411-28	48
1973	Solution structure of the calcium channel antagonist omega-conotoxin GVIA. 1993 , 2, 1591-603	57
1972	A peptide analog of the calmodulin-binding domain of myosin light chain kinase adopts an alpha-helical structure in aqueous trifluoroethanol. 1993 , 2, 1931-7	50
1971	Action of rat liver Gal beta 1-4GlcNAc alpha(2-6)-sialyltransferase on Man beta 1-4GlcNAc beta-OMe, GalNAc beta 1-4GlcNAc beta-OMe, Glc beta 1-4GlcNAc beta-OMe and GlcNAc beta 1-4GlcNAc beta-OMe as synthetic substrates. 1993 , 10, 82-90	19
1970	A structural study of the asparagine-linked oligosaccharide moiety of duck ovomucoid. 1993 , 10, 425-34	13
1969	Recombinant human IL-6 expressed in E. coli undergoes selective N-terminal degradation: evidence that the protein consists of a stable core and a nonessential flexible N-terminal. 1993 , 12, 489-97	22
1968	Polyoxypregnanes from <i>Marsdenia tenacissima</i> . 1993 , 34, 1615-1620	80
1967	Sequence-specific 1H-NMR assignment and secondary structure of black mamba dendrotoxin I, a highly selective blocker of voltage-gated potassium channels. <i>FEBS Journal</i> , 1993 , 211, 813-20	28
1966	Effects of subsite alterations on substrate-binding mode in the active site of hen egg-white lysozyme. <i>FEBS Journal</i> , 1993 , 212, 151-6	21
1965	Primary structure of N-linked carbohydrate chains of a human chimeric plasminogen activator K2tu-PA expressed in Chinese hamster ovary cells. <i>FEBS Journal</i> , 1993 , 212, 639-56	31
1964	The NMR solution structure of a Kunitz-type proteinase inhibitor from the sea anemone <i>Stichodactyla helianthus</i> . <i>FEBS Journal</i> , 1993 , 212, 675-84	50

1963	Homonuclear and heteronuclear NMR studies of oxidized <i>Desulfovibrio vulgaris</i> flavodoxin. Sequential assignments and identification of secondary structure elements. <i>FEBS Journal</i> , 1993 , 213, 167-84	35
1962	¹ H-NMR spectroscopy of beta B2-crystallin from bovine eye lens. Conformation of the N- and C-terminal extensions. <i>FEBS Journal</i> , 1993 , 213, 313-20	27
1961	Determination of the nuclear-magnetic-resonance solution structure of cardiotoxin CTX IIb from <i>Naja mossambica mossambica</i> . <i>FEBS Journal</i> , 1993 , 213, 891-900	26
1960	Sequential assignment of proton resonances in the NMR spectrum of Zn-substituted alpha chains from human hemoglobin. Ligand-induced tertiary changes in the heme pocket. <i>FEBS Journal</i> , 1993 , 214, 383-93	5
1959	Structure of a novel glycopeptidolipid antigen containing a O-methylated serine isolated from <i>Mycobacterium xenopi</i> . Complete ¹ H-NMR and ¹³ C-NMR assignment. <i>FEBS Journal</i> , 1993 , 214, 395-403	8
1958	The solution structure of the leucine zipper motif of the Jun oncoprotein homodimer. <i>FEBS Journal</i> , 1993 , 214, 415-24	14
1957	¹ H-NMR analysis of turkey egg-white lysozyme and comparison with hen egg-white lysozyme. <i>FEBS Journal</i> , 1993 , 215, 255-66	14
1956	The solution structure of the histidine-containing protein (HPr) from <i>Staphylococcus aureus</i> as determined by two-dimensional ¹ H-NMR spectroscopy. <i>FEBS Journal</i> , 1993 , 216, 205-14	39
1955	A modified uridine in the first position of the anticodon of a minor species of arginine tRNA, the argU gene product, from <i>Escherichia coli</i> . <i>FEBS Journal</i> , 1993 , 216, 369-75	33
1954	Solution conformation of an antibacterial peptide, sarcotoxin IA, as determined by ¹ H-NMR. <i>FEBS Journal</i> , 1993 , 217, 639-44	43
1953	The structure of the carbohydrate backbone of the core-lipid-A region of the lipopolysaccharide from <i>Vibrio cholerae</i> strain H11 (non-O1). <i>FEBS Journal</i> , 1993 , 218, 543-54	13
1952	¹ H-NMR studies and secondary structure of the RGD-containing snake toxin, albolabrin. <i>FEBS Journal</i> , 1993 , 218, 853-60	11
1951	Structure of the glycosylated adhesion domain of human T lymphocyte glycoprotein CD2. 1993 , 1, 69-81	64
1950	Refined solution structure of bovine pancreatic Ribonuclease A by ¹ H NMR methods. Sidechain dynamics. 1993 , 4, 385-415	14
1949	The structure of new cis and trans 3-(3-phenyl-3,4,5,6,7-tetrahydro-2,1-benzisoxazole-7-yl)-spiro-2-(3-phenylaziridine). 1993 , 30, 97-104	1
1948	Structure of the novel steroidal antibiotic squalamine determined by two-dimensional NMR spectroscopy. 1993 , 58, 370-8	63
1947	Replacement of His23 by Cys in a zinc finger of HIV-1 NCp7 led to a change in ¹ H NMR-derived 3D structure and to a loss of biological activity. 1993 , 331, 43-8	16
1946	¹ H assignment and secondary structure determination of human melanoma growth stimulating activity (MGSA) by NMR spectroscopy. 1993 , 330, 302-6	16

1945	A peptide corresponding to the N-terminal 13 residues of T4 lysozyme forms an alpha-helix. 1993 , 315, 323-8	15
1944	Identification of the adsorbing site of lysozyme onto the hydroxyapatite surface using hydrogen exchange and ¹ H NMR. 1993 , 317, 128-30	20
1943	Overexpression, purification, and mechanistic study of UDP-N-acetylenolpyruvylglucosamine reductase. 1993 , 32, 2024-30	90
1942	Peptide models of protein folding initiation sites. 3. The G-H helical hairpin of myoglobin. 1993 , 32, 6356-64	101
1941	Peptide models of protein folding initiation sites. 1. Secondary structure formation by peptides corresponding to the G- and H-helices of myoglobin. 1993 , 32, 6337-47	198
1940	Peptide models of protein folding initiation sites. 2. The G-H turn region of myoglobin acts as a helix stop signal. 1993 , 32, 6348-55	86
1939	Structure of ten free N-glycans in ripening tomato fruit. Arabinose is a constituent of a plant N-glycan. 1993 , 102, 445-58	102
1938	Membrane-bound conformation of a signal peptide: a transferred nuclear Overhauser effect analysis. 1993 , 32, 13991-9	74
1937	Characterization of an independent structural unit in apocytochrome b5. 1993 , 32, 199-207	40
1936	Capping interactions in isolated alpha helices: position-dependent substitution effects and structure of a serine-capped peptide helix. 1993 , 32, 421-5	125
1935	¹ H resonance assignments and secondary structure of the 13.6 kDa glycosylated adhesion domain of human CD2. 1993 , 32, 10995-1006	42
1934	Transforming growth factor beta 1: NMR signal assignments of the recombinant protein expressed and isotopically enriched using Chinese hamster ovary cells. 1993 , 32, 1152-63	49
1933	Collapse of parallel folding channels in dihydrofolate reductase from Escherichia coli by site-directed mutagenesis. 1993 , 32, 13566-74	20
1932	Solution conformation of cobrotoxin: a nuclear magnetic resonance and hybrid distance geometry-dynamical simulated annealing study. 1993 , 32, 2131-6	51
1931	Determination of the solution structure of neuropeptide K by high-resolution nuclear magnetic resonance spectroscopy. 1993 , 32, 7406-12	21
1930	Solution structure of the parallel-stranded hairpin d(T8C4A8) as determined by two-dimensional NMR. 1993 , 32, 646-56	27
1929	Interaction of phospholipase A2 with thioether amide containing phospholipid analogues. 1993 , 32, 5009-16	31
1928	Conformational analysis of [D-Ala9]alpha-factor and [L-Ala9]alpha-factor in solution and in the presence of lipid. 1993 , 32, 908-17	25

1927	Mobility at the TpA cleavage site in the T3A3-containing AhaIII and PmeI restriction sequences. 1993 , 32, 8022-35		40
1926	Two-dimensional NMR and restrained molecular dynamics studies of the hairpin d(T8C4A8): detection of an extraloop cytosine. 1993 , 32, 637-45		13
1925	Isolation, identification and synthesis of an endogenous arachidonic amide that inhibits calcium channel antagonist 1,4-dihydropyridine binding. 1993 , 48, 429-37		64
1924	NMR studies of the human retinoic acid receptor-beta DNA-binding domain. Metal coordination and three-dimensional structure. 1993 , 684, 49-62		2
1923	NMR studies of the dynamics of the multidomain protein urokinase-type plasminogen activator. 1993 , 32, 298-309		27
1922	Solution structure of omega-conotoxin GVIA using 2-D NMR spectroscopy and relaxation matrix analysis. 1993 , 32, 7396-405		123
1921	Involvement of various amino- and carboxyl-terminal residues in the active site of the histidine-containing protein HPr of the phosphoenolpyruvate-dependent phosphotransferase system of <i>Staphylococcus carnosus</i> : site-directed mutagenesis with the ptsH gene, biochemical characterization and NMR studies of the mutant proteins. 1993 , 6, 417-23		15
1920	Confirmation of the Structure of Kessane by NMR Spectroscopy. 1993 , 5, 169-178		11
1919	Cyanopeptolins, new depsipeptides from the cyanobacterium <i>Microcystis</i> sp. PCC 7806. 1993 , 46, 1550-6		143
1918	Characterization of structural and folding properties of streptokinase by n.m.r. spectroscopy. <i>Biochemical Journal</i> , 1993 , 290 (Pt 2), 313-9	3.8	19
1917	Solution structure of the epidermal growth factor-like domain of heregulin-alpha, a ligand for p180erbB-4.. <i>EMBO Journal</i> , 1994 , 13, 3517-3523	13	35
1916	Solution structure of human insulin-like growth factor II; recognition sites for receptors and binding proteins.. <i>EMBO Journal</i> , 1994 , 13, 5590-5597	13	36
1915	Formation of sheared G:A base pairs in an RNA duplex modelled after ribozymes, as revealed by NMR. 1994 , 22, 2752-9		39
1914	The cell wall polysaccharide of <i>Streptococcus gordonii</i> 38: structure and immunochemical comparison with the receptor polysaccharides of <i>Streptococcus oralis</i> 34 and <i>Streptococcus mitis</i> J22. 1994 , 4, 183-92		21
1913	Conformation of the cytoplasmic domain of phospholamban by NMR and CD. 1994 , 11, 263-9		16
1912	Structures in Solution of Toxins from Taiwan Cobra Venom, <i>Naja naja atra</i> , Derived from NMR Spectra. 1994 , 13, 291-315		17
1911	Nuclear magnetic resonance study and secondary structure determination of the antibiotic peptide, aibellin. 1994 , 58, 2188-92		2
1910	Determination of structure and conformation in solution of syringotoxin, a lipodepsipeptide from <i>Pseudomonas syringae</i> pv. <i>syringae</i> by 2D NMR and molecular dynamics. 1994 , 5, 43-50		18

1909	¹ H and ¹⁵ N resonance assignments and secondary structure of the carbon monoxide complex of sperm whale myoglobin. 1994 , 4, 491-504	48
1908	Three-dimensional ¹ H NMR structure of the nucleocapsid protein NCp10 of Moloney murine leukemia virus. 1994 , 4, 153-70	33
1907	Filtering methods for selection of singlet and doublet signals in NMR spectra of DNA oligomers. 1994 , 4, 181-91	1
1906	¹ H and ¹⁵ N resonance assignments and secondary structure of cellular retinoic acid-binding protein with and without bound ligand. 1994 , 4, 741-60	38
1905	An evaluation of least-squares fits to COSY spectra as a means of estimating proton-proton coupling constants. II. Applications to polypeptides. 1994 , 4, 827-44	5
1904	Application of 1D and 2D NMR techniques to the structure elucidation of the O-polysaccharide from <i>Proteus mirabilis</i> O:57. 1994 , 4, 615-30	34
1903	Molecular dynamics and NMR studies of single-stranded PNAs. 1994 , 35, 5105-5108	38
1902	¹³ C NMR Spectra of pentacyclic triterpenoids— compilation and some salient features. 1994 , 37, 1517-1575	1060
1901	Peptidomimetic antagonists designed to inhibit the binding of CD4 to HIV GP120. 1994 , 2, 1007-13	12
1900	Role of the Cys 2-Cys 10 disulfide bond for the structure, stability, and folding kinetics of ribonuclease T1. 1994 , 3, 227-39	25
1899	The native state of apomyoglobin described by proton NMR spectroscopy: interaction with the paramagnetic probe HyTEMPO and the fluorescent dye ANS. 1994 , 3, 267-81	61
1898	An ¹ H NMR determination of the three-dimensional structures of mirror-image forms of a Leu-5 variant of the trypsin inhibitor from <i>Ecballium elaterium</i> (EETI-II). 1994 , 3, 291-302	28
1897	Determination of the structure of the DNA binding domain of gamma delta resolvase in solution. 1994 , 3, 1286-95	14
1896	The NMR solution structure of the pheromone Er-2 from the ciliated protozoan <i>Euplotes raikovi</i> . 1994 , 3, 1515-26	38
1895	The NMR solution structure of the pheromone Er-1 from the ciliated protozoan <i>Euplotes raikovi</i> . 1994 , 3, 1527-36	38
1894	Structure-function relationship in the globular type III antifreeze protein: identification of a cluster of surface residues required for binding to ice. 1994 , 3, 1760-9	104
1893	Solution structure of the DNA-binding domain of the heat shock transcription factor determined by multidimensional heteronuclear magnetic resonance spectroscopy. 1994 , 3, 1806-21	66
1892	Formation of a native-like subdomain in a partially folded intermediate of bovine pancreatic trypsin inhibitor. 1994 , 3, 1822-32	116

1891	Quantification of the calcium-induced secondary structural changes in the regulatory domain of troponin-C. 1994 , 3, 1961-74	174
1890	Solution structure of a core peptide derived from scyllatoxin. 1994 , 18, 205-15	13
1889	A model for the structure of a homodimeric prohormone: the precursor to the locust neuropeptide AKH I. 1994 , 20, 356-66	3
1888	Structural analysis of novel rhamnose-branched oligosaccharides from the glycoposphosphingolipids of <i>Leptomonas samueli</i> . 1994 , 11, 23-33	10
1887	NMR spectroscopy of organophosphorus compounds II: Stereochemistry of [1-(2H-azirin-2-yl)alkyl]phosphonates. 1994 , 125, 1101-1111	1
1886	¹ H NMR study of the diastereomeric forms of the protease inhibitor antipain. 1994 , 32, 220-224	2
1885	Will the real pinene please stand up. 1994 , 32, 316-319	4
1884	Structure identification of feruloylated oligosaccharides from sugar-beet pulp by NMR spectroscopy. 1994 , 263, 243-56	186
1883	NMR analysis of succinoglycans from different microbial sources: partial assignment of their ¹ H and ¹³ C NMR spectra and location of the succinate and the acetate groups. 1994 , 265, 167-79	31
1882	NMR assignments for the aldopentoses. 1994 , 258, 27-33	21
1881	NMR spectral analysis of a water-insoluble (1→3)-beta-D-glucan isolated from <i>Saccharomyces cerevisiae</i> . 1994 , 258, 307-11	70
1880	Identification of a novel sugar, 4-amino-4,6-dideoxy-2-O-methylmannose in the lipopolysaccharide of <i>Vibrio cholerae</i> O1 serotype Ogawa. 1994 , 256, 113-28	67
1879	¹ H NMR spectroscopic characterization of perchloric acid extracts from breast carcinomas and non-involved breast tissue. 1994 , 7, 181-94	102
1878	Synthesis and NMR Spectroscopic Investigation of a Macrocyclic Diphosphine Ligand and its nickel(II) and palladium(II) complexes. 1994 , 77, 409-418	15
1877	Synthesis of Tri-, Penta-, and Heptapeptides Containing and (R)-2-alkyl-2-amino-3-(methylamino)-propionic acid residue in the central position. 1994 , 77, 2035-2050	23
1876	ENC, a motor of progress in NMR. 1994 , 6, 201-229	2
1875	Conformation of the oligosaccharide receptor for E-selectin. 1994 , 34, 21-9	44
1874	Conformational analysis of CCK-B agonists using ¹ H-NMR and restrained molecular dynamics: comparison of biologically active Boc-Trp-(N-Me) Nle-Asp-Phe-NH ₂ and inactive Boc-Trp-(N-Me)Phe-Asp-Phe-NH ₂ . 1994 , 34, 155-69	21

1873	Solution conformation of a pectic acid fragment by 1H-NMR and molecular dynamics. 1994 , 34, 457-62	14
1872	NMR investigation of cyclo7,10[C7,X9,C10,Nle12] analogues of the alpha-factor from <i>Saccharomyces cerevisiae</i> . 1994 , 34, 709-20	9
1871	Conformation and molecular dynamics calculations on uteroglobin fragment 18-47. 1994 , 34, 773-82	12
1870	Design, synthesis, and conformation of a model peptide of endothelin with cystine-stabilized alpha-helix motif. 1994 , 34, 963-7	7
1869	Solution three-dimensional structure of surfactin: a cyclic lipopeptide studied by 1H-NMR, distance geometry, and molecular dynamics. 1994 , 34, 975-86	94
1868	Proteinstrukturaufklärung mit drei- und vierdimensionaler NMR-Spektroskopie. 1994 , 106, 284-300	6
1867	Conformational analysis of DDAVP and DDAVP-OH in methanol. 1994 , 328, 145-151	1
1866	Purification and secondary structural analysis of tissue inhibitor of metalloproteinases-1. 1994 , 1208, 94-100	3
1865	Resonance assignment strategies for the analysis of NMR spectra of proteins. 1994 , 2, 61-93	17
1864	Synthesis, molecular modelling, and NMR structure determination of four cyclic peptide antagonists of endothelin. 1994 , 2, 279-96	5
1863	Biosynthesis of the pseudobactin chromophore from tyrosine. 1994 , 50, 9865-9872	12
1862	Solution structure and DNA-binding properties of a thermostable protein from the archaeon <i>Sulfolobus solfataricus</i> . 1994 , 1, 808-19	143
1861	Structure of the N-terminal SH3 domain of GRB2 complexed with a peptide from the guanine nucleotide releasing factor Sos. 1994 , 1, 891-7	88
1860	NMR structure of the N-terminal SH3 domain of GRB2 and its complex with a proline-rich peptide from Sos. 1994 , 1, 898-907	94
1859	Solution structure of a peptide fragment of human alpha-lactalbumin in trifluoroethanol: a model for local structure in the molten globule. 1994 , 2, 703-12	27
1858	Solution structure and ligand-binding site of the carboxy-terminal SH3 domain of GRB2. 1994 , 2, 1029-40	71
1857	The immunologically reactive O-linked polysaccharide chains derived from circulating cathodic antigen isolated from the human blood fluke <i>Schistosoma mansoni</i> have Lewis x as repeating unit. <i>FEBS Journal</i> , 1994 , 225, 467-82	97
1856	NMR assignments and secondary structure of the retinoid X receptor alpha DNA-binding domain. Evidence for the novel C-terminal helix. <i>FEBS Journal</i> , 1994 , 224, 639-50	13

1855	Sequence-specific assignment of the ¹ H and ¹⁵ N nuclear magnetic resonance spectra of the reduced recombinant high-potential iron-sulfur protein I from <i>Ectothiorhodospira halophila</i> . <i>FEBS Journal</i> , 1994 , 225, 703-14	22
1854	¹ H-NMR assignments and secondary structure of dendroaspin, an RGD-containing glycoprotein IIb-IIIa (alpha IIb-beta 3) antagonist with a neurotoxin fold. <i>FEBS Journal</i> , 1994 , 226, 861-8	11
1853	Equilibrium unfolding studies of horse muscle acylphosphatase. <i>FEBS Journal</i> , 1994 , 225, 811-7	18
1852	Trifluoroethanol stabilizes a helix-turn-helix motif in equine infectious-anemia-virus trans-activator protein. <i>FEBS Journal</i> , 1994 , 225, 855-61	20
1851	Isolation and structural analysis of oligosaccharide phosphates containing the complete carbohydrate chain of the lipopolysaccharide from <i>Vibrio cholerae</i> strain H11 (non-O1). <i>FEBS Journal</i> , 1994 , 225, 1029-39	52
1850	Cloning, expression, and spectroscopic studies of the Jun leucine zipper domain. <i>FEBS Journal</i> , 1994 , 219, 877-86	12
1849	NMR and circular dichroic studies of the solution structure of conformationally constrained antigenic peptides. <i>FEBS Journal</i> , 1994 , 220, 415-25	5
1848	The nuclear-magnetic-resonance solution structure of the mutant alpha-amylase inhibitor [R19L] Tendamistat and comparison with wild-type Tendamistat. <i>FEBS Journal</i> , 1994 , 220, 763-70	13
1847	Sequential ¹ H and ¹⁵ N nuclear magnetic resonance assignments and secondary structure of the N-terminal lipoyl domain of the dihydrolipoyl transacetylase component of the pyruvate dehydrogenase complex from <i>Azotobacter vinelandii</i> . <i>FEBS Journal</i> , 1994 , 221, 87-100	14
1846	Conformational analysis of elcatonin in solution. <i>FEBS Journal</i> , 1994 , 222, 659-66	11
1845	Solution structure of the variant-3 neurotoxin from <i>Centruroides sculpturatus</i> Ewing. <i>FEBS Journal</i> , 1994 , 219, 89-95	29
1844	Solution structure of synthetic peptides corresponding to the C-terminal helix of interleukin-6. <i>FEBS Journal</i> , 1994 , 219, 97-107	9
1843	Structure determination of extracellular fragments of amyloid proteins involved in Alzheimer's disease and Dutch-type hereditary cerebral haemorrhage with amyloidosis. <i>FEBS Journal</i> , 1994 , 219, 237-51	46
1842	¹ H resonance assignments and secondary structure of the carbon monoxide complex of soybean leghemoglobin determined by homonuclear two-dimensional and three-dimensional NMR spectroscopy. <i>FEBS Journal</i> , 1994 , 219, 611-26	8
1841	[Ala4]surfactin, a novel isoform from <i>Bacillus subtilis</i> studied by mass and NMR spectroscopies. <i>FEBS Journal</i> , 1994 , 224, 89-96	67
1840	Phospholipase A2 engineering. Structural and functional roles of the highly conserved active site residue aspartate-49. 1994 , 33, 14714-22	65
1839	Kaliotoxin (1-37) shows structural differences with related potassium channel blockers. 1994 , 33, 14256-63	51
1838	Structures of wild-type and mutant signal sequences of <i>Escherichia coli</i> ribose binding protein. 1994 , 66, 1604-11	13

1837	¹³ C NMR and fluorescence analysis of tryptophan dynamics in wild-type and two single-Trp variants of <i>Escherichia coli</i> thioredoxin. 1994 , 66, 2111-26	30
1836	A thermodynamic scale for the beta-sheet forming tendencies of the amino acids. 1994 , 33, 5510-7	375
1835	Structural analysis of the lipid A derived from the lipopolysaccharide of <i>Brucella abortus</i> . 1994 , 1, 137-148	25
1834	NMR solution structure of the recombinant tick anticoagulant protein (rTAP), a factor Xa inhibitor from the tick <i>Ornithodoros moubata</i> . 1994 , 352, 251-7	40
1833	The 3D structure of a cyclosporin analogue in water is nearly identical to the cyclophilin-bound cyclosporin conformation. 1994 , 340, 255-9	66
1832	Characterization by ¹ H NMR of a C32S,C35S double mutant of <i>Escherichia coli</i> thioredoxin confirms its resemblance to the reduced wild-type protein. 1994 , 339, 11-7	18
1831	NMR structures of ferredoxin chloroplastic transit peptide from <i>Chlamydomonas reinhardtii</i> promoted by trifluoroethanol in aqueous solution. 1994 , 343, 261-6	36
1830	Proton NMR studies of the structural and dynamical effect of chemical modification of a single aromatic side-chain in a snake cardiotoxin. Relation to the structure of the putative binding site and the cytolytic activity of the toxin. 1994 , 243, 719-35	19
1829	Stabilization of a type VI turn in a family of linear peptides in water solution. 1994 , 243, 736-53	143
1828	Conformational behaviour of the active and inactive forms of the nucleocapsid NCp7 of HIV-1 studied by ¹ H NMR. 1994 , 235, 287-301	133
1827	NMR and dynamical simulated annealing studies on the solution conformation of urotensin II. 1994 , 1199, 115-22	10
1826	Hairpin folding of subunit c of F ₁ F ₀ ATP synthase: ¹ H distance measurements to nitroxide-derivatized aspartyl-61. 1994 , 33, 665-74	80
1825	A determination of the solution conformation of the nonmammalian tachykinin eledoisin by NMR and CD spectroscopy. 1994 , 33, 6802-11	22
1824	Solution structure of RP 71955, a new 21 amino acid tricyclic peptide active against HIV-1 virus. 1994 , 33, 42-50	93
1823	The solution structure of a monocyclic analogue of endothelin [1,15 Aba]-ET-1, determined by ¹ H NMR spectroscopy. 1994 , 37, 656-64	16
1822	Solution structure of bound trimethoprim in its complex with <i>Lactobacillus casei</i> dihydrofolate reductase. 1994 , 33, 12416-26	36
1821	¹ H and ¹⁵ N nuclear magnetic resonance assignments, secondary structure in solution, and solvent exchange properties of azurin from <i>Alcaligenes denitrificans</i> . 1994 , 33, 3560-71	19
1820	Control of Calix[6]arene Conformations by Self-Inclusion of 1,3,5-Tri-O-alkyl Substituents: Synthesis and NMR Studies. 1994 , 116, 5814-5822	99

1819	Solution structure of the active domain of tissue inhibitor of metalloproteinases-2. A new member of the OB fold protein family. 1994 , 33, 11745-59		97
1818	Proton nuclear magnetic resonance and distance geometry/simulated annealing studies on the variant-1 neurotoxin from the New World scorpion <i>Centruroides sculpturatus</i> Ewing. 1994 , 33, 2468-75		26
1817	High-resolution structure of an engineered biologically potent insulin monomer, B16 Tyr-->His, as determined by nuclear magnetic resonance spectroscopy. 1994 , 33, 7998-8006		72
1816	¹ H NMR structure and biological studies of the His23-->Cys mutant nucleocapsid protein of HIV-1 indicate that the conformation of the first zinc finger is critical for virus infectivity. 1994 , 33, 11707-16		97
1815	Structure of the signal sequences for two mitochondrial matrix proteins that are not proteolytically processed upon import. 1994 , 33, 8610-7		73
1814	Solution structure of horse heart ferrocycytochrome c determined by high-resolution NMR and restrained simulated annealing. 1994 , 33, 6408-17		75
1813	Structural analysis of peptide fragment 71-93 of transthyretin by NMR spectroscopy and electron microscopy: insight into amyloid fibril formation. 1994 , 33, 33-41		26
1812	Sequential proton resonance assignments and metal cluster topology of lobster metallothionein-1. 1994 , 33, 8858-65		34
1811	Lipo-oligosaccharidic antigen from <i>Mycobacterium gastris</i> . Complete structure of a novel C4-branched 3,6-dideoxy-alpha-xylo-hexopyranose. 1994 , 33, 1930-7		19
1810	The NMR structure of the pulmonary surfactant-associated polypeptide SP-C in an apolar solvent contains a valyl-rich alpha-helix. 1994 , 33, 6015-23		184
1809	One-disulfide intermediates of apamin exhibit native-like structure. 1994 , 33, 5253-61		16
1808	¹ H and ¹⁵ N NMR assignments of PsaE, a photosystem I subunit from the cyanobacterium <i>Synechococcus</i> sp. strain PCC 7002. 1994 , 33, 6043-51		20
1807	O-glycosidically linked N-acetylglucosamine-bound oligosaccharides from glycoproteins of <i>Trypanosoma cruzi</i> . <i>Biochemical Journal</i> , 1994 , 301 (Pt 1), 151-9	3.8	102
1806	Elongation of helix III of the NK-2 homeodomain upon binding to DNA: a secondary structure study by NMR. 1994 , 33, 15053-60		72
1805	Solution structure of a trefoil-motif-containing cell growth factor, porcine spasmodic protein. 1994 , 91, 2206-10		45
1804	Multiple steroid-binding orientations: alteration of regiospecificity of dehydroepiandrosterone 2- and 7-hydroxylase activities of cytochrome P-450 2a-5 by mutation of residue 209. <i>Biochemical Journal</i> , 1995 , 306 (Pt 1), 29-33	3.8	16
1803	Solution structure of DNA polymerases and DNA polymerase-substrate complexes. 1995 , 262, 147-71		1
1802	Solution structures of horse ferro- and ferricytochrome c using 2D and 3D ¹ H NMR and restrained simulated annealing. 1995 , 6, 511-519		

1801	Solution structure of the cardiostimulant polypeptide anthopleurin-B and comparison with anthopleurin-A. 1995 , 3, 791-803	54
1800	The solution structure of Abl SH3, and its relationship to SH2 in the SH(32) construct. 1995 , 3, 1075-86	41
1799	Structural comparison of the free and DNA-bound forms of the purine repressor DNA-binding domain. 1995 , 3, 1217-24	40
1798	Overproduction, Purification and Characterization of the Cellulose-Binding Domain of the <i>Erwinia Chrysanthemi</i> Secreted Endoglucanase EGZ. <i>FEBS Journal</i> , 1995 , 231, 142-148	13
1797	An NMR Study of the Interaction of Cardiotoxin β from <i>Naja nigricollis</i> with Perdeuterated Dodecylphosphocholine Micelles. <i>FEBS Journal</i> , 1995 , 230, 213-220	37
1796	¹ H, ¹³ C, ¹⁵ N-NMR Resonance Assignments of Oxidized Thioredoxin h from the Eukaryotic Green Alga <i>Chlamydomonas reinhardtii</i> Using New Methods based on Two-Dimensional Triple-Resonance NMR Spectroscopy and Computer-Assisted Backbone Assignment. <i>FEBS Journal</i> , 1995 , 229, 473-485	1
1795	Elucidation of the Primary Structure of the Lantibiotic Epilancin K7 from <i>Staphylococcus epidermidis</i> K7. Cloning and Characterisation of the Epilancin-K7-Encoding Gene and NMR Analysis of Mature Epilancin K7. <i>FEBS Journal</i> , 1995 , 230, 587-600	2
1794	¹ H and ¹⁵ N Assignment of NMR Spectrum, Secondary Structure and Global Folding of the Immunophilin-Like Domain of the 59-kDa FK506-Binding Protein. <i>FEBS Journal</i> , 1995 , 231, 761-772	4
1793	Proton NMR Study of the Trimannosyl Unit in a Pentaantennary N-Linked Decasaccharide Structure. Complete Assignment of the Proton Resonances and Conformational Characterization. <i>FEBS Journal</i> , 1995 , 228, 822-829	
1792	Sequential ¹ H and ¹⁵ N nuclear magnetic resonance assignments and secondary structure of the lipoyl domain of the 2-oxoglutarate dehydrogenase complex from <i>Azotobacter vinelandii</i> . Evidence for high structural similarity with the lipoyl domain of the pyruvate dehydrogenase complex. <i>FEBS Journal</i> , 1995 , 234, 148-59	6
1791	Primary Structure, Sequence-Specific ¹ H-NMR Assignments and Secondary Structure in Solution of Bromelain Inhibitor VI from Pineapple Stem. <i>FEBS Journal</i> , 1995 , 232, 335-343	14
1790	NMR studies of the major coat protein of bacteriophage M13. Structural information of gVIIIp in dodecylphosphocholine micelles. <i>FEBS Journal</i> , 1995 , 232, 490-500	24
1789	¹ H and ¹⁵ N assignments and secondary structure of the starch-binding domain of glucoamylase from <i>Aspergillus niger</i> . <i>FEBS Journal</i> , 1995 , 233, 568-78	13
1788	Comparative Study on the Conformation of Phalloidin, Viroisin, and Related Derivatives in Aqueous Solution. <i>FEBS Journal</i> , 1995 , 232, 726-736	
1787	NMR-based structural studies of the pNR-2/pS2 single domain trefoil peptide. Similarities to porcine spasmolytic peptide and evidence for a monomeric structure. <i>FEBS Journal</i> , 1995 , 233, 847-55	14
1786	¹ H and ¹³ C NMR assignments and structural aspects of a ferrocyclochrome c-551 from the purple phototrophic bacterium <i>Ectothiorhodospira halophila</i> . <i>FEBS Journal</i> , 1995 , 227, 249-60	7
1785	NMR spectroscopy of steroidal sapogenins and steroidal saponins: An update. 1995 , 33, 923-953	95
1784	Solution structures of the DP IV (CD26) inhibitor Val-boroPro determined by NMR spectroscopy. 1995 , 33, 959-970	2

1783	Helix formation in model peptides based on nucleolin TPAKK motifs. 1995 , 35, 93-102	9
1782	Conformation of cyclobradysin by NMR and distance geometry calculations. 1995 , 36, 461-72	11
1781	Conformation and interactions of bombolitin I analogues with SDS micelles and phospholipid vesicles: CD, fluorescence, two-dimensional NMR and computer simulations. 1995 , 36, 473-84	17
1780	Conformation of parathyroid hormone antagonists by CD, NMR, and molecular dynamics simulations. 1995 , 36, 485-95	14
1779	Eleven newly characterized xyloglucan oligoglycosyl alditols: the specific effects of sidechain structure and location on ¹ H NMR chemical shifts. 1995 , 267, 79-104	26
1778	Full assignment of the NMR spectrum of the capsular polysaccharide from <i>Streptococcus pneumoniae</i> serotype 10A. 1995 , 269, 175-81	18
1777	NMR studies of acetan and the related bacterial polysaccharide, CR1/4, produced by a mutant strain of <i>Acetobacter xylinum</i> . 1995 , 269, 319-31	25
1776	Structure of the O-deacetylated glucuronoxylomannan from <i>Cryptococcus neoformans</i> serotype C as determined by 2D ¹ H NMR spectroscopy. 1995 , 276, 365-86	13
1775	Structural analysis of the carbohydrate moiety of arabinogalactan-proteins from stigmas and styles of <i>Nicotiana glauca</i> . 1995 , 277, 67-85	102
1774	A conformational model for cyclic beta-(1-->2)-linked glucans based on NMR analysis of the beta-glucans produced by <i>Xanthomonas campestris</i> . 1995 , 278, 205-25	44
1773	RES-701-1, comparative study of the synthetic and the microbial-origin compounds. 1995 , 5, 1595-1600	11
1772	Structure-function studies of mEGF: probing the type I beta-turn between residues 25 and 26. 1995 , 14, 753-62	1
1771	Lipooligosaccharidic antigens from <i>Mycobacterium kansasii</i> and <i>Mycobacterium gastri</i> . 1995 , 12, 298-308	13
1770	Structure of Leu5-enkephalin bound to a model membrane as determined by high-resolution NMR. 1995 , 2, 59-70	17
1769	Three-dimensional structure of gurmarin, a sweet taste-suppressing polypeptide. 1995 , 5, 297-305	36
1768	Backbone assignment, secondary structure and protein A binding of an isolated, human antibody VH domain. 1995 , 6, 141-52	23
1767	A variable target intensity-restrained global optimization (VARTIGO) procedure for determining three-dimensional structures of polypeptides from NOESY data: application to gramicidin-S. 1995 , 5, 37-48	50
1766	Structural studies of a synthetic peptide derived from the carboxyl-terminal domain of RNA polymerase II. 1995 , 21, 149-60	43

1765	Prosapogenins from <i>Dodonaea attenuata</i> . 1995 , 6, 157-163	2
1764	Structural analysis of the N- and C-termini in a peptide with consensus sequence. 1995 , 4, 1446-56	39
1763	Solution structure of the potassium channel inhibitor agitoxin 2: caliper for probing channel geometry. 1995 , 4, 1478-89	112
1762	¹ H, ¹⁵ N, and ¹³ C backbone chemical shift assignments, secondary structure, and magnesium-binding characteristics of the <i>Bacillus subtilis</i> response regulator, Spo0F, determined by heteronuclear high-resolution NMR. 1995 , 4, 1801-14	38
1761	Comparison of the free and DNA-complexed forms of the DNA-binding domain from c-Myb. 1995 , 2, 309-20	136
1760	Structure of the carboxy-terminal domain of bovine rhodopsin. 1995 , 2, 832-4	57
1759	Solution structure of a mammalian PCB-binding protein in complex with a PCB. 1995 , 2, 983-9	35
1758	Solution structure of endothelin B receptor selective antagonist RES-701-1 determined by ¹ H NMR spectroscopy. 1995 , 3, 1273-80	52
1757	Heteronuclear NMR studies of cobalt corrinoids 17. characterization of neopentylcobinamide and neopentyl-13-epicobinamide by ¹ H and ¹³ C NMR spectroscopy: Inferred corrin ring conformations from chemical shift differentials. 1995 , 14, 2961-2977	3
1756	Conformation studies of aerosol-OT in reverse micelles using phase sensitive double quantum COSY NMR spectroscopy. 1995 , 96, 165-169	11
1755	Investigation of the impurities found in methamphetamine synthesised from pseudoephedrine by reduction with hydriodic acid and red phosphorus. 1995 , 76, 97-114	40
1754	Diterpenoids from <i>Isodon eriocalyx</i> var. <i>laxiflora</i> . 1995 , 38, 1451-5	26
1753	Cytotoxic ent-kaurene diterpenoids from <i>Isodon gesneroides</i> . 1995 , 40, 1461-7	11
1752	Chemical characterisation of glycosylinositolphospholipids of <i>Herpetomonas samuelpessoai</i> . 1995 , 69, 81-92	20
1751	NMR studies of the solution properties of recombinant murine interleukin-6. 1995 , 1249, 189-203	7
1750	CD and NMR determination of the solution structure of a peptide corresponding to T4 lysozyme residues 38-51. 1995 , 1250, 163-70	13
1749	Conformation of polymyxin B analogs in DMSO from NMR spectra and molecular modeling. 1995 , 1252, 312-20	9
1748	Conformation of A82846B, a glycopeptide antibiotic, complexed with its cell wall fragment: an asymmetric homodimer determined using NMR spectroscopy. 1995 , 34, 9632-44	36

1747	Solution structure of cysteine-rich domain of protein kinase C alpha. 1995 , 117, 566-74		55
1746	Structural analysis of the mannan region of lipoarabinomannan from Mycobacterium bovis BCG. Heterogeneity in phosphorylation state. <i>Journal of Biological Chemistry</i> , 1995 , 270, 15012-21	5.4	26
1745	Development of a receptor peptide antagonist to human gamma-interferon and characterization of its ligand-bound conformation using transferred nuclear Overhauser effect spectroscopy. <i>Journal of Biological Chemistry</i> , 1995 , 270, 9241-9	5.4	6
1744	The structure of a 19-residue fragment from the C-loop of the fourth epidermal growth factor-like domain of thrombomodulin. <i>Journal of Biological Chemistry</i> , 1995 , 270, 23366-72	5.4	21
1743	Solution structure of the acetylated and noncleavable mitochondrial targeting signal of rat chaperonin 10. <i>Journal of Biological Chemistry</i> , 1995 , 270, 1323-31	5.4	60
1742	Structural characterization of the major glycosylphosphatidylinositol membrane-anchored glycoprotein from epimastigote forms of Trypanosoma cruzi Y-strain. <i>Journal of Biological Chemistry</i> , 1995 , 270, 7241-50	5.4	117
1741	Enhanced transglycosylation activity of Arthrobacter protophormiae endo-beta-N-acetylglucosaminidase in media containing organic solvents. <i>Journal of Biological Chemistry</i> , 1995 , 270, 17723-9	5.4	103
1740	NMR investigation of echinomycin binding to d(ACGTAAACGT) ₂ : Hoogsteen versus Watson-Crick A.T base pairing between echinomycin binding sites. 1995 , 118, 989-95		7
1739	Evidence for DAPI intercalation in CG sites of DNA oligomer [d(CGACGTCC)] ₂ : a ¹ H NMR study. 1995 , 23, 1333-40		44
1738	Calmodulin binding of a peptide derived from the regulatory domain of Bordetella pertussis adenylate cyclase. <i>Journal of Biological Chemistry</i> , 1995 , 270, 7088-96	5.4	20
1737	Conformation analysis of non-pepsin-type acid proteinase A from the fungus Aspergillus niger by NMR. 1995 , 362, 611-5		
1736	NMR behavior of the aromatic protons of bovine neurophysin-I and its peptide complexes: implications for solution structure and for function. 1995 , 34, 2137-47		12
1735	Membrane and Solution Conformations of Monogalactosyldiacylglycerol Using NMR/Molecular Modeling Methods. 1995 , 117, 5031-5040		35
1734	NMR and restrained molecular dynamics study of the three-dimensional solution structure of toxin FS2, a specific blocker of the L-type calcium channel, isolated from black mamba venom. 1995 , 34, 5923-37		60
1733	Three-dimensional structure in solution of the polypeptide cardiac stimulant anothopleurin-A. 1995 , 34, 3782-94		44
1732	Gene cloning, expression, and characterization of the Sac7 proteins from the hyperthermophile Sulfolobus acidocaldarius. 1995 , 34, 10063-77		72
1731	Study of human calcitonin fibrillation by proton nuclear magnetic resonance spectroscopy. 1995 , 34, 12138-43		39
1730	Elucidation of the primary and three-dimensional structure of the uterotonic polypeptide kalata B1. 1995 , 34, 4147-58		362

1729	Structure of a compact peptide from staphylococcal nuclease determined by circular dichroism and NMR spectroscopy. 1995 , 34, 5795-800	12
1728	[Lys(-2)-Arg(-1)]endothelin-1 solution structure by two-dimensional ¹ H-NMR: possible involvement of electrostatic interactions in native disulfide bridge formation and in biological activity decrease. 1995 , 34, 4546-61	23
1727	¹ H NMR studies of sarafotoxin SRTb, a nonselective endothelin receptor agonist, and IRL 1620, an ETB receptor-specific agonist. 1995 , 34, 2026-33	25
1726	Solution structure of the cytoplasmic domain of phospholamban: phosphorylation leads to a local perturbation in secondary structure. 1995 , 34, 7603-13	76
1725	Structure and stability of monomeric lambda repressor: NMR evidence for two-state folding. 1995 , 34, 3884-92	87
1724	NMR solution structure of the RNA-binding peptide from human immunodeficiency virus (type 1) Rev. 1995 , 34, 8242-9	25
1723	Solution structure of the DNA-binding protein Sac7d from the hyperthermophile Sulfolobus acidocaldarius. 1995 , 34, 13289-304	75
1722	Conformation of micellar phospholipid bound to the active site of phospholipase A2. 1995 , 34, 4943-51	22
1721	Reactive-site hydrolyzed Cucurbita maxima trypsin inhibitor-V: function, thermodynamic stability, and NMR solution structure. 1995 , 34, 12087-94	15
1720	Conformation of Two Peptides Corresponding to Human Apolipoprotein C-I Residues 7-24 and 35-53 in the Presence of Sodium Dodecyl Sulfate by CD and NMR Spectroscopy. 1995 , 34, 7401-7408	64
1719	Structure effects of double D-amino acid replacements: a nuclear magnetic resonance and circular dichroism study using amphipathic model helices. 1995 , 34, 12954-62	69
1718	Characterization of the calmodulin binding domain of SIV transmembrane glycoprotein by NMR and CD spectroscopy. 1995 , 34, 10690-6	39
1717	Topology of the pore-region of a K ⁺ channel revealed by the NMR-derived structures of scorpion toxins. 1995 , 15, 1169-81	251
1716	Conformation of the propeptide domain of factor IX. 1995 , 1245, 227-31	3
1715	Structural characterization by nuclear magnetic resonance spectroscopy of a genetically engineered high-affinity calmodulin-binding peptide derived from Bordetella pertussis adenylate cyclase. 1995 , 320, 224-35	5
1714	Solution structure determination by NMR spectroscopy of a synthetic peptide corresponding to a putative amphipathic alpha-helix of spiralin: resonance assignment, distance geometry and simulated annealing. 1995 , 1235, 169-77	5
1713	The physical properties of local interactions of tyrosine residues in peptides and unfolded proteins. 1995 , 245, 251-60	61
1712	The three-dimensional solution structure of human stefin A. 1995 , 246, 331-43	100

1711	Solution structure of omega-conotoxin MVIIIC, a high affinity ligand of P-type calcium channels, using 1H NMR spectroscopy and complete relaxation matrix analysis. 1995 , 248, 106-24	70
1710	Three-dimensional solution structure of the calcium channel antagonist omega-agatoxin IVA: consensus molecular folding of calcium channel blockers. 1995 , 250, 659-71	71
1709	Complete 1H nuclear magnetic resonance assignments and structural characterization of a fusion protein of the alpha-amylase inhibitor tendamistat with the activation domain of the human immunodeficiency virus type 1 Tat protein. 1995 , 250, 672-88	11
1708	The three-dimensional solution structure of the NK-2 homeodomain from <i>Drosophila</i> . 1995 , 251, 297-307	59
1707	Determination of the folding topology of the SL1 RNA from <i>Caenorhabditis elegans</i> by multidimensional heteronuclear NMR. 1995 , 252, 314-27	18
1706	Three-dimensional solution structure of bombyxin-II an insulin-like peptide of the silkworm <i>Bombyx mori</i> : structural comparison with insulin and relaxin. 1995 , 253, 749-58	47
1705	Identification of the receptor-recognition surface of bombyxin-II, an insulin-like peptide of the silkworm <i>Bombyx mori</i> : critical importance of the B-chain central part. 1995 , 253, 759-70	14
1704	Search for nucleation sites in smaller fragments of chymotrypsin inhibitor 2. 1995 , 254, 289-304	69
1703	Refined solution structure and dynamics of the DNA-binding domain of the heat shock factor from <i>Kluyveromyces lactis</i> . 1995 , 254, 704-19	19
1702	The three-dimensional solution structure of a constrained peptidomimetic in water and in chloroform. Observation of solvent induced hydrophobic cluster. 1995 , 359, 113-8	5
1701	Solution structure of the DNA binding domain of a nucleoid-associated protein, H-NS, from <i>Escherichia coli</i> . 1995 , 360, 125-31	102
1700	Ligand-induced perturbations in <i>Urtica dioica</i> agglutinin. 1995 , 361, 157-61	16
1699	Pulmonary surfactant-associated polypeptide SP-C in lipid micelles: CD studies of intact SP-C and NMR secondary structure determination of depalmitoyl-SP-C(1-17). 1995 , 362, 261-5	52
1698	1H and 13C NMR assignment and secondary structure of <i>Chlorobium limicola</i> f. <i>thiosulfatophilum</i> ferrocytochrome c555. 1995 , 365, 172-8	5
1697	Complement assembly of two fragments of the streptococcal protein G B1 domain in aqueous solution. 1995 , 366, 99-103	55
1696	Structures of revertant signal sequences of <i>Escherichia coli</i> ribose binding protein. 1995 , 69, 2703-9	9
1695	Solution structure of a green mamba toxin that activates muscarinic acetylcholine receptors, as studied by nuclear magnetic resonance and molecular modeling. 1995 , 34, 1248-60	47
1694	Determination of the three-dimensional solution structure of noxiustoxin: analysis of structural differences with related short-chain scorpion toxins. 1995 , 34, 16563-73	77

1693	NMR studies of DNA duplexes singly cross-linked by different synthetic linkers. 1995 , 23, 4827-35	31
1692	Solution structure of bovine neutrophil beta-defensin-12: the peptide fold of the beta-defensins is identical to that of the classical defensins. 1995 , 34, 13663-71	144
1691	Palmitoylation-induced conformational changes of specific side chains in the gramicidin transmembrane channel. 1995 , 34, 9299-306	33
1690	Decapeptide agonists of human C5a: the relationship between conformation and neutrophil response. 1995 , 38, 3669-75	27
1689	Structural characterization of a minimal functional transactivation domain from the human glucocorticoid receptor. 1995 , 92, 1699-703	146
1688	Structure refinement of the glucocorticoid receptor-DNA binding domain from NMR data by relaxation matrix calculations. 1995 , 247, 689-700	13
1687	Solution structure of omega-conotoxin MVIIA using 2D NMR spectroscopy. 1995 , 370, 163-9	59
1686	Proton and carbon-13 NMR studies of steroids and triterpenes. 1996 , 405, 281-97	1
1685	Conformation of sulfoquinovosyldiacylglycerol bound to a magnetically oriented membrane system. 1996 , 71, 2573-82	12
1684	The conformation of substance P in lipid environments. 1996 , 70, 1716-27	55
1683	Three-dimensional structure analysis of mu-agatoxins: further evidence for common motifs among neurotoxins with diverse ion channel specificities. 1996 , 35, 2836-44	74
1682	Conformational preferences of a chimeric peptide HIV-1 immunogen from the C4-V3 domains of gp120 envelope protein of HIV-1 CAN0A based on solution NMR: comparison to a related immunogenic peptide from HIV-1 RF. 1996 , 35, 5158-65	32
1681	Structural determinants of the catalytic reactivity of the buried cysteine of Escherichia coli thioredoxin. 1996 , 35, 14876-81	33
1680	Site-specific d(GpG) intrastrand cross-links formed by dinuclear platinum complexes. Bending and NMR studies. 1996 , 35, 16705-13	82
1679	Novel marine di-Z-carotenoids: cucumariaxanthins A, B, and C from the sea cucumber Cucumaria japonica. 1996 , 59, 30-4	23
1678	Solution structure of LSIII, a long neurotoxin from the venom of Laticauda semifasciata. 1996 , 35, 418-26	19
1677	Characterization of two transposon mutants from Haemophilus influenzae type b with altered lipooligosaccharide biosynthesis. 1996 , 35, 5937-47	40
1676	Structural and thermodynamic characterization of the interaction of the SH3 domain from Fyn with the proline-rich binding site on the p85 subunit of PI3-kinase. 1996 , 35, 15646-53	92

1675	Stabilization of a Cis Amide Bond in a Host-Guest Complex. 1996 , 118, 10220-10227	61
1674	Complexes formed between calmodulin and the antagonists J-8 and TFP in solution. 1996 , 35, 10287-99	38
1673	Solution structure of horse heart ferricytochrome c and detection of redox-related structural changes by high-resolution ¹ H NMR. 1996 , 35, 12275-86	123
1672	Lactam bridge stabilization of alpha-helices: the role of hydrophobicity in controlling dimeric versus monomeric alpha-helices. 1996 , 35, 10041-50	53
1671	Stereostructure of (R)-Chloropeptin I, a Novel Inhibitor of gp120-CD4 Binding, via High-Temperature Molecular Dynamics, Monte Carlo Conformational Searching, and NMR Spectroscopy. 1996 , 118, 13087-13088	41
1670	Conformation and Dynamics of Membrane-Bound Digalactosyldiacylglycerol. 1996 , 118, 3345-3353	22
1669	Three-dimensional solution structure of the N-terminal domain of DNA polymerase beta and mapping of the ssDNA interaction interface. 1996 , 35, 6188-200	52
1668	Photochemistry of Dihydrido(hydrotris(3,5-dimethylpyrazolyl)borato)(Z-cyclooctene)iridium. Synthetic Intermediates and Mechanism of the Photochemical Formation of Hydridophenyl(hydrotris(3,5-dimethylpyrazolyl)borato)(trimethyl phosphite)iridium. 1996 , 35, 1602-1608	21
1667	Conformations of human apolipoprotein E(263-286) and E(267-289) in aqueous solutions of sodium dodecyl sulfate by CD and ¹ H NMR. 1996 , 35, 10358-66	50
1666	Solution and membrane bound structure of a peptide derived from the protein kinase C substrate domain of neuromodulin. 1996 , 35, 11104-12	21
1665	Solution structure of reduced plastocyanin from the blue-green alga <i>Anabaena variabilis</i> . 1996 , 35, 7021-31	73
1664	Metabolism of Pentyl 2-Chloro-4-fluoro-5-(3,4,5,6- tetrahydrophthalimido)phenoxyacetate (Flumiclorac Pentyl, S-23031) in Rats. 1. Identification of Metabolites in Feces and Urine of Rats. 1996 , 44, 1910-1914	7
1663	Three-dimensional solution structure of mu-conotoxin GIIIB, a specific blocker of skeletal muscle sodium channels. 1996 , 35, 8824-35	93
1662	Binding of phenylphosphocholine-carrier conjugates to the combining site of antibodies maintains a conformation of the hapten. 1996 , 35, 2958-67	6
1661	Transfer of a beta-hairpin from the functional site of snake curaremimetic toxins to the alpha/beta scaffold of scorpion toxins: three-dimensional solution structure of the chimeric protein. 1996 , 35, 8535-43	25
1660	NMR characterization of side chain flexibility and backbone structure in the type I antifreeze protein at near freezing temperatures. 1996 , 35, 16698-704	50
1659	Structure of the membrane binding domain of CTP:phosphocholine cytidyltransferase. 1996 , 35, 11975-84	90
1658	Nuclear magnetic resonance solution structure of the growth factor receptor-bound protein 2 Src homology 2 domain. 1996 , 35, 11852-64	24

1657	Absolute Configuration of Phorbaxozoles A and B from the Marine Sponge Phorbasp. 1. Macrolide and Hemiketal Rings. 1996 , 118, 9422-9423		134
1656	Characterization of human calcitonin fibrillation in aqueous urea solution by ¹ H NMR spectroscopy. 1996 , 35, 12671-6		10
1655	A model of the iron responsive element RNA hairpin loop structure determined from NMR and thermodynamic data. 1996 , 35, 13586-96		62
1654	Solution structure of bromelain inhibitor IV from pineapple stem: structural similarity with Bowman-Birk trypsin/chymotrypsin inhibitor from soybean. 1996 , 35, 5379-84		34
1653	The conformational properties of the highly selective cannabinoid receptor ligand CP-55,940. <i>Journal of Biological Chemistry</i> , 1996 , 271, 10640-7	5-4	48
1652	Evidence for isomerization in myotoxin a from the prairie rattlesnake (<i>Crotalus viridis viridis</i>). 1996 , 34, 417-34		16
1651	Spectroscopic study of an HIV-1 capsid protein major homology region peptide analog. 1996 , 378, 43-7		5
1650	Secondary structure of an armadillo single repeat from the APC protein. 1996 , 383, 31-6		12
1649	Investigation of oxidation state-dependent conformational changes in <i>Desulfovibrio vulgaris</i> Hildenborough cytochrome c553 by two-dimensional H-NMR spectra. 1996 , 389, 203-9		9
1648	NMR structures of a mitochondrial transit peptide from the green alga <i>Chlamydomonas reinhardtii</i> . 1996 , 391, 203-8		15
1647	Conserved secondary structure in the actinorhodin polyketide synthase acyl carrier protein from <i>Streptomyces coelicolor</i> A3(2) and the fatty acid synthase acyl carrier protein from <i>Escherichia coli</i> . 1996 , 391, 302-6		16
1646	Structural comparison of a 15 residue peptide from the V3 loop of HIV-1IIIb and an O-glycosylated analogue. 1996 , 393, 280-6		31
1645	Structure of a methionine-rich segment of <i>Escherichia coli</i> Ffh protein. 1996 , 395, 160-4		2
1644	Structural characterization of a monomeric chemokine: monocyte chemoattractant protein-3. 1996 , 395, 277-82		57
1643	Structure of the cysteine-rich intestinal protein, CRIP. 1996 , 257, 153-74		54
1642	Comparison of the (30-51, 14-38) two-disulphide folding intermediates of the homologous proteins dendrotoxin K and bovine pancreatic trypsin inhibitor by two-dimensional ¹ H nuclear magnetic resonance. 1996 , 257, 188-98		9
1641	Solution structure of the superactive monomeric des-[Phe(B25)] human insulin mutant: elucidation of the structural basis for the monomerization of des-[Phe(B25)] insulin and the dimerization of native insulin. 1996 , 257, 684-99		44
1640	Solution structure of PMP-C: a new fold in the group of small serine proteinase inhibitors. 1996 , 258, 158-71		48

1639	H NMR study of the solution structure of Ac-AMP2, a sugar binding antimicrobial protein isolated from <i>Amaranthus caudatus</i> . 1996 , 258, 322-33			73
1638	Structural studies of synthetic peptides dissected from the voltage-gated sodium channel. 1996 , 258, 672-87			32
1637	Insight into a random coil conformation and an isolated helix: structural and dynamical characterisation of the C-helix peptide from hen lysozyme. 1996 , 261, 443-53			43
1636	A network of heterogeneous hydrogen bonds in GNRA tetraloops. 1996 , 264, 968-80			331
1635	Structural role of a buried salt bridge in the 434 repressor DNA-binding domain. 1996 , 264, 1002-12			42
1634	Conformation of human serum apolipoprotein A-I(166-185) in the presence of sodium dodecyl sulfate or dodecylphosphocholine by 1H-NMR and CD. Evidence for specific peptide-SDS interactions. 1996 , 1301, 174-84			51
1633	The solution structure of a psoralen cross-linked DNA duplex by NMR and relaxation matrix refinement. <i>Biochemical and Biophysical Research Communications</i> , 1996 , 219, 191-7	3-4		26
1632	Conformational study of two linear hexapeptides by two-dimensional NMR and computer-simulated modeling: implication for peptide cyclization in solution. <i>Biochemical and Biophysical Research Communications</i> , 1996 , 219, 572-9	3-4		2
1631	Structural analysis of a biologically active echistatin analogue des(46-49)-[Ala8,37]-echistatin gamma with three disulfide bonds by 2D-NMR and computer graphics. <i>Biochemical and Biophysical Research Communications</i> , 1996 , 220, 246-54	3-4		7
1630	Structural characterization of a new galactofuranose-containing glycolipid antigen of <i>Paracoccidioides brasiliensis</i> . <i>Biochemical and Biophysical Research Communications</i> , 1996 , 222, 639-45	3-4		29
1629	Important 2D NMR Experiments. 1996 , 213-343			3
1628	¹ H NMR evidence that Glu-38 interacts with the N-terminal functional domain in interleukin-8. 1996 , 399, 43-6			14
1627	Three-dimensional structures of the amyloid beta peptide (25-35) in membrane-mimicking environment. 1996 , 35, 16094-104			80
1626	¹ H-Detected, Gradient-Enhanced ¹⁵ N and ¹³ C NMR Experiments for the Measurement of Small Heteronuclear Coupling Constants and Isotopic Shifts. 1996 , 118, 5096-5102			19
1625	High-resolution solution structure of the EGF-like domain of heregulin-alpha. 1996 , 35, 3402-17			61
1624	Trimeric assembly and three-dimensional structure model of the FACIT collagen COL1-NC1 junction from CD and NMR analysis. 1996 , 35, 9647-60			32
1623	Three-dimensional solution structure of <i>Saccharomyces cerevisiae</i> reduced iso-1-cytochrome c. 1996 , 35, 13788-96			85
1622	FUNDAMENTALS OF NUCLEAR MAGNETIC RESONANCE. 1996 , 1-115			5

1621	THE SOLUTION CHARACTERIZATION OF POLYMERS. 1996 , 155-241		5
1620	Aciculitins AII: Cytotoxic and Antifungal Cyclic Peptides from the Lithistid Sponge <i>Aciculites orientalis</i> . 1996 , 118, 4314-4321		67
1619	Stereodynamics of 9,11-Diphenyl-10-azatetracyclo[6.3.0.0.(4,11)0.(5,9)]undecanes. Highly Restricted Nitrogen Inversion and Isolated Phenyl Rotation. X-ray Crystallographic, Dynamic NMR, and Molecular Mechanics Studies. 1996 , 61, 4319-4327		8
1618	Solution structure of an engineered insulin monomer at neutral pH. 1996 , 35, 8836-45		120
1617	A systematic NMR approach for the determination of the molecular structure of steroidal saponins. 1996 , 405, 299-315		10
1616	Chymotrypsin inhibitory conformation induced by amino acid side chain-side chain intramolecular CH/ π interaction. 1996 , 2479-2485		17
1615	Synthesis of 3-(2,2,2-trifluoroethylidene)-lactams. First examples of 1,3-dipolar cycloaddition with diazomethane and N-methyl- β -phenylnitron. 1996 , 1853-1858		6
1614	Absolute Stereochemistry of Precorrin-3x and Its Relevance to the Dichotomy of Ring Contraction Mechanism in Vitamin B12 Biosynthesis. 1996 , 118, 1657-1662		16
1613	Structure of the core oligosaccharide in the serotype O8 lipopolysaccharide from <i>Klebsiella pneumoniae</i> . 1996 , 178, 1731-41		26
1612	Chymotrypsin inhibition induced by side chain-side chain intramolecular CH/ π interaction in D-Thr-L-Phe benzylamide. 1996 , 119, 870-7		9
1611	A synthetic peptide study on the molten globule of alpha-lactalbumin. 1996 , 119, 947-52		19
1610	Determination of the three-dimensional structure of hordothionin-alpha by nuclear magnetic resonance. <i>Biochemical Journal</i> , 1996 , 313 (Pt 3), 885-92	3.8	17
1609	Base decomposition of erythromycin A methoxime. 1996 , 49, 318-20		1
1608	The solution structure of the Raf-1 cysteine-rich domain: a novel ras and phospholipid binding site. 1996 , 93, 8312-7		170
1607	Metal binding properties and secondary structure of the zinc-binding domain of Nup475. 1996 , 93, 13754-9		98
1606	Solution Structure of the Human CD4 (403-419) Receptor Peptide. 1996 , 3, 435-441		11
1605	Conformation of <i>Vespa basalis</i> mastoparan-B in trifluoroethanol-containing aqueous solution. 1996 , 1292, 1-8		22
1604	Determination of three-dimensional solution structure of waglerin I, a toxin from <i>Trimeresurus wagleri</i> , using 2D-NMR and molecular dynamics simulation. 1996 , 1292, 145-55		7

1603	Metabolite profiling by one- and two-dimensional NMR analysis of complex mixtures. 1996 , 28, 161-219	400
1602	NMR structural studies of DNA decamer duplex containing the Dewar photoproduct of thymidylyl(3'-->5')thymidine. Conformational changes of the oligonucleotide duplex by photoconversion of a (6-4) adduct to its Dewar valence isomer. <i>FEBS Journal</i> , 1996 , 235, 359-65	17
1601	Three-dimensional structure of the lantibiotic nisin in the presence of membrane-mimetic micelles of dodecylphosphocholine and of sodium dodecylsulphate. <i>FEBS Journal</i> , 1996 , 235, 382-93	74
1600	Structural studies of the equine infectious anemia virus trans-activator protein. <i>FEBS Journal</i> , 1996 , 240, 45-52	3
1599	Structure of the B-Myb DNA-binding domain in solution and evidence for multiple conformations in the region of repeat-2 involved in DNA binding: implications for sequence-specific DNA binding by Myb proteins. <i>FEBS Journal</i> , 1996 , 235, 721-35	24
1598	Solution structure of the DNA-binding domain of the tomato heat-stress transcription factor HSF24. <i>FEBS Journal</i> , 1996 , 236, 911-21	49
1597	Analysis of the 1H-NMR chemical shifts of Cu(I)-, Cu(II)- and Cd-substituted pea plastocyanin. Metal-dependent differences in the hydrogen-bond network around the copper site. <i>FEBS Journal</i> , 1996 , 242, 132-47	42
1596	Secondary structural elements as a basis for antibody recognition in the immunodominant region of human immunodeficiency viruses 1 and 2. <i>FEBS Journal</i> , 1996 , 237, 188-204	18
1595	Methoxyamine-induced mutagenesis of nucleic acids. A proton NMR study of oligonucleotides containing N4-methoxycytosine paired with adenine or guanine. <i>FEBS Journal</i> , 1996 , 242, 271-9	22
1594	Sequence-specific resonance assignments of the 1H-NMR spectra and structural characterization in solution of the HIV-1 transframe protein p6. <i>FEBS Journal</i> , 1996 , 237, 383-92	28
1593	Conformational study of [Met5]enkephalin-Arg-Phe in the presence of phosphatidylserine vesicles. <i>FEBS Journal</i> , 1996 , 240, 540-9	20
1592	Assignment of 13C resonances and analysis of relaxation properties and internal dynamics of pike parvalbumin by 13C-NMR at natural abundance. <i>FEBS Journal</i> , 1996 , 237, 561-74	10
1591	Synthesis and solution structure of the antimicrobial peptide protegrin-1. <i>FEBS Journal</i> , 1996 , 237, 575-83	145
1590	Structure and multiple conformations of the kunitz-type domain from human type VI collagen alpha3(VI) chain in solution. 1996 , 4, 195-209	19
1589	Solution structure and peptide binding of the SH3 domain from human Fyn. 1996 , 4, 705-14	86
1588	Refined solution structure of type III antifreeze protein: hydrophobic groups may be involved in the energetics of the protein-ice interaction. 1996 , 4, 1325-37	164
1587	Solution conformation of an immunogenic peptide derived from the principal neutralizing determinant of the HIV-2 envelope glycoprotein gp125. 1996 , 1, 157-65	8
1586	Native-like secondary structure in a peptide from the alpha-domain of hen lysozyme. 1996 , 1, 473-84	14

1585	The native state of apomyoglobin described by proton NMR spectroscopy: the A-B-G-H interface of wild-type sperm whale apomyoglobin. 1996 , 25, 267-85	26
1584	Threonine6-bradykinin: Conformational study of a flexible peptide in dimethyl sulfoxide by NMR and ensemble calculations. 1996 , 40, 561-569	13
1583	Conformational characterization of a peptide mimetic of the third cytoplasmic loop of the G-protein coupled parathyroid hormone/parathyroid hormone related protein receptor. 1996 , 40, 653-66	34
1582	Conformational analysis of cyclic hexapeptides designed as constrained ligands for the SH2 domain of the p85 subunit of phosphatidylinositol-3-OH kinase. 1996 , 38, 191-208	2
1581	NMR Assignment of a 2?-Hydroxyl Proton from the UUCG Tetraloop Through Long-Range Correlations with 13C. 1996 , 34, S11-S17	15
1580	Determination of Homo- and Heteronuclear Coupling Constants in Uniformly 13C,15N-Labeled DNA Oligonucleotides. 1996 , 34, S177-S186	23
1579	Structure of the saponin adjuvant QS-21 and its base-catalyzed isomerization product by 1H and natural abundance 13C NMR spectroscopy. 1996 , 280, 1-14	93
1578	Structure of the O-deacetylated glucuronoxylomannan from <i>Cryptococcus neoformans</i> Cap70 as determined by 2D NMR spectroscopy. 1996 , 283, 95-110	26
1577	NMR analysis of galactoglucan from <i>Pseudomonas marginalis</i> : assignment of the 1H and 13C NMR spectra and location of succinate groups. 1996 , 283, 195-205	12
1576	Structural characterisation of xyloglucan secreted by suspension-cultured cells of <i>Nicotiana glumabaginifolia</i> . 1996 , 293, 147-72	73
1575	Comparative conformational analysis of CCK-B agonist Boc-Trp-Phg-Asp-(1-Nal)-NH ₂ and CCK1-B antagonist Boc-Trp-Phg-Asp-(1-Nal)-N(Me) ₂ using 1HNMR spectroscopy and restrained molecular dynamics. 1996 , 329, 197-204	2
1574	Nitrogen-15 NMR chemical shifts in oligopeptides coordinated to cobalt(III). 1996 , 62, 117-26	5
1573	MS-271, a novel inhibitor of calmodulin-activated myosin light chain kinase from <i>Streptomyces</i> sp.-II. Solution structure of MS-271: characteristic features of the "lasso" structure. 1996 , 4, 121-9	41
1572	Full 1H NMR assignment and detailed O-acetylation patterns of capsular polysaccharides from <i>Neisseria meningitidis</i> used in vaccine production. 1996 , 296, 83-96	94
1571	The structures of arabinoxyloglucans produced by solanaceous plants. 1996 , 285, 99-128	99
1570	Complete 1H NMR assignments of synthetic glycopeptides from the carbohydrate-protein linkage region of serglycins. 1996 , 13, 599-607	3
1569	The secondary structure of a pyrimidine-guanine sequence-specific ribonuclease possessing cytotoxic activity from the oocytes of <i>Rana catesbeiana</i> . 1996 , 8, 331-44	6
1568	Determination of the solution structure of the SH3 domain of human p56 Lck tyrosine kinase. 1996 , 8, 105-22	20

1567	Elucidation of the origin of multiple conformations of the human alpha 3-chain type VI collagen C-terminal Kunitz domain: the reorientation of the Trp21 ring. 1996 , 8, 391-403	8
1566	Spiroannulation reactions of 3-trifluoroacetyl- and other 3-acyl-1-methyl-lactams. 1996 , 338, 538-543	0
1565	2D 1H and 3D 1H-15N NMR of zinc-rubredoxins: contributions of the beta-sheet to thermostability. 1996 , 5, 883-94	30
1564	The NMR solution structure of the pheromone Er-11 from the ciliated protozoan <i>Euplotes raikovi</i> . 1996 , 5, 1512-22	21
1563	Three-dimensional structure of the complexes of ribonuclease A with 2',5'-CpA and 3',5'-d(CpA) in aqueous solution, as obtained by NMR and restrained molecular dynamics. 1996 , 5, 1633-47	38
1562	1H NMR assignments of apo calyculin and comparative structural analysis with calbindin D9k and S100 beta. 1996 , 5, 2162-74	32
1561	NMR studies of structure, hydrogen exchange, and main-chain dynamics in a disrupted-core mutant of thioredoxin. 1996 , 5, 2552-65	21
1560	Conformational analysis of neuropeptide Y segments by CD, NMR spectroscopy and restrained molecular dynamics. 1996 , 2, 176-93	5
1559	Characteristic Anthocyanin Pattern from Onions and other <i>Allium</i> spp. 1996 , 61, 703-706	98
1558	Stabilization of proteins by glycosylation examined by NMR analysis of a fucosylated proteinase inhibitor. 1996 , 3, 45-53	104
1557	The N-terminal domain of TFIIB from <i>Pyrococcus furiosus</i> forms a zinc ribbon. 1996 , 3, 122-4	117
1556	The three-dimensional structure of the first domain of neural cell adhesion molecule. 1996 , 3, 581-5	50
1555	Solution structure of a cyclic RGD peptide that inhibits platelet aggregation. 1996 , 14, 1-11	18
1554	Glutaredoxin-3 from <i>Escherichia coli</i> . Amino acid sequence, 1H AND 15N NMR assignments, and structural analysis. <i>Journal of Biological Chemistry</i> , 1996 , 271, 6736-45	5-4 56
1553	The recombinant dehydrin-like desiccation stress protein from the resurrection plant <i>Craterostigma plantagineum</i> displays no defined three-dimensional structure in its native state. 1996 , 377, 555-61	47
1552	Changing the structural context of a functional beta-hairpin. Synthesis and characterization of a chimera containing the curaremimetic loop of a snake toxin in the scorpion alpha/beta scaffold. <i>Journal of Biological Chemistry</i> , 1996 , 271, 11979-87	5-4 60
1551	Simultaneous and different binding mechanisms of 4',6-diamidino-2-phenylindole to DNA hexamer (d(CGATCG)) ₂ . A 1H NMR study. <i>Journal of Biological Chemistry</i> , 1996 , 271, 27608-14	5-4 28
1550	Kaempferol 3-O-(2-O-β-D-Glucopyranosyl-6-O-α-L-Rhamnopyranosyl-β-D-Glucopyranoside) from the African Plant <i>Erythrina abyssinica</i> . 1996 , 34, 370-373	7

1549	Two-Dimensional NMR Spectroscopy: A Graphical, Top-Down Description. 1996 , 1-84		
1548	Structural studies of a peptide activator of human lecithin-cholesterol acyltransferase. <i>Journal of Biological Chemistry</i> , 1996 , 271, 3039-45	5.4	16
1547	The Role of Positive Charges and Structural Segments in the Presequence of Rat Liver Aldehyde Dehydrogenase in Import into Mitochondria. <i>Journal of Biological Chemistry</i> , 1996 , 271, 21041-21048	5.4	41
1546	Replacement of Trp28 in Escherichia coli thioredoxin by site-directed mutagenesis affects thermodynamic stability but not function. <i>Journal of Biological Chemistry</i> , 1996 , 271, 3091-6	5.4	24
1545	Sequential ¹ H and ¹⁵ N NMR resonance assignment and secondary structure of ferrocytochrome c2 from Rhodobacter sphaeroides. 1996 , 119, 1131-42		1
1544	A case history. NMR studies of the structure of a small protein, omega-conotoxin MVIIA. <i>Methods in Molecular Biology</i> , 1997 , 60, 337-62	1.4	5
1543	Leishmania adleri, a lizard parasite, expresses structurally similar glycoinositolphospholipids to mammalian Leishmania. 1997 , 7, 687-95		19
1542	Effect of geometric isomerism in dinuclear platinum antitumour complexes on the rate of formation and structure of intrastrand adducts with oligonucleotides. 1997 , 25, 1265-71		21
1541	Improvement of the refolding yield and solubility of hen egg-white lysozyme by altering the Met residue attached to its N-terminus to Ser. 1997 , 10, 1333-8		30
1540	Structural features of the DNA hairpin d(ATCCTA-GTTA-TAGGAT): formation of a G-A base pair in the loop. 1997 , 25, 1537-47		48
1539	Structural studies on tRNA acceptor stem microhelices: exchange of the discriminator base A73 for G in human tRNA ^{Leu} switches the acceptor specificity from leucine to serine possibly by decreasing the stability of the terminal G1-C72 base pair. 1997 , 25, 4551-6		18
1538	NMR spectroscopic studies of the DNA-binding domain of the monomer-binding nuclear orphan receptor, human estrogen related receptor-2. The carboxyl-terminal extension to the zinc-finger region is unstructured in the free form of the protein. <i>Journal of Biological Chemistry</i> , 1997 , 272, 18038-43	5.4	25
1537	Specificity of the ubiquitin isopeptidase in the PA700 regulatory complex of 26 S proteasomes. <i>Journal of Biological Chemistry</i> , 1997 , 272, 28438-46	5.4	86
1536	Protein NMR Techniques. 1997 ,		2
1535	Structural and functional roles of tyrosine 78 of yeast guanylate kinase. <i>Journal of Biological Chemistry</i> , 1997 , 272, 19343-50	5.4	6
1534	Structure of the C-terminal fragment 300-320 of the rat angiotensin II AT1A receptor and its relevance with respect to G-protein coupling. <i>Journal of Biological Chemistry</i> , 1997 , 272, 9734-41	5.4	58
1533	On the convergent evolution of animal toxins. Conservation of a diad of functional residues in potassium channel-blocking toxins with unrelated structures. <i>Journal of Biological Chemistry</i> , 1997 , 272, 4302-9	5.4	283
1532	Circular dichroism and ¹ H nuclear magnetic resonance studies on the solution and membrane structures of GAP-43 calmodulin-binding domain. <i>Journal of Biological Chemistry</i> , 1997 , 272, 7639-45	5.4	28

1531	An alpha-helical signal in the cytosolic domain of the interleukin 2 receptor beta chain mediates sorting towards degradation after endocytosis. 1997 , 136, 583-95		46
1530	Proton NMR study of triantennary complex type N-linked glycan chains: assignment of proton chemical shifts of the beta-Man residue in a basic unit of the triantennary glycan chain having a GlcNAc beta 1->6 Man alpha 1->6 Man beta->sequence. 1997 , 7, 31-6		5
1529	Metal-ion-binding properties of synthetic conantokin-G. <i>Biochemical Journal</i> , 1997 , 328 (Pt 3), 777-83	3.8	21
1528	Chapter 6 Nuclear magnetic resonance spectroscopy (NMR): Principles and applications. 1997 , 179-237		
1527	Determination of the solution structures of conantokin-G and conantokin-T by CD and NMR spectroscopy. <i>Journal of Biological Chemistry</i> , 1997 , 272, 2291-9	5.4	68
1526	Involvement of basic amphiphilic helical domain in the reversible membrane interaction of amphitropic proteins: Structural studies by mass spectrometry, circular dichroism, and nuclear magnetic resonance. 1997 , 555-564		
1525	¹ H-NMR evidence for two buried ASN side-chains in the c-Myc-Max heterodimeric helical coiled-coil. 1997 , 617-624		2
1524	Human nucleotide excision repair protein XPA: ¹ H NMR and CD solution studies of a synthetic peptide fragment corresponding to the zinc-binding domain (101-141). 1997 , 14, 677-90		3
1523	Novel saponins hainaneosides A and B isolated from <i>Marsdenia hainanensis</i> . 1997 , 60, 134-8		11
1522	Role of gamma-carboxyglutamic acid in the calcium-induced structural transition of conantokin G, a conotoxin from the marine snail <i>Conus geographus</i> . 1997 , 36, 15677-84		52
1521	MIXED-LIGAND COMPLEXES OF COBALT(III) WITH DITHIOCARBAMATES AND A CYCLIC TETRADENTATE SECONDARY AMINE. 1997 , 41, 19-25		25
1520	Glycosylation affects both the three-dimensional structure and antibody binding properties of the HIV-1IIIB GP120 peptide RP135. 1997 , 36, 10846-56		74
1519	Structure of the carboxy-terminal fragment of the apo-biotin carboxyl carrier subunit of <i>Escherichia coli</i> acetyl-CoA carboxylase. 1997 , 36, 15089-100		68
1518	4,5-Cyclopropanocholestan-3βol Substrates for Cholesterol Oxidase and Their ¹ H NMR Assignments. 1997 , 62, 5893-5897		2
1517	Solution structure of a polypeptide from the N terminus of the HIV protein Nef. 1997 , 36, 5970-80		44
1516	Mitosene-DNA adducts. Characterization of two major DNA monoadducts formed by 1,10-bis(acetoxy)-7-methoxymitosene upon reductive activation. 1997 , 36, 9211-20		20
1515	Effects of buried charged groups on cysteine thiol ionization and reactivity in <i>Escherichia coli</i> thioredoxin: structural and functional characterization of mutants of Asp 26 and Lys 57. 1997 , 36, 2622-36		176
1514	Ab-Initio Calculations on ArginineDisulfide Complexes Modeling the One-Electron Reduction of Lysozyme. Comparison to an Experimental Reinvestigation. 1997 , 101, 7809-7817		30

1513	Solution structure of the actinorhodin polyketide synthase acyl carrier protein from <i>Streptomyces coelicolor</i> A3(2). 1997 , 36, 6000-8	129
1512	Solution structure for Pandinus toxin K-alpha (PiTX-K alpha), a selective blocker of A-type potassium channels. 1997 , 36, 2763-71	18
1511	The helix-hinge-helix structural motif in human apolipoprotein A-I determined by NMR spectroscopy. 1997 , 36, 13657-66	49
1510	Characterization of the unbound 2[Fe4S4]-ferredoxin-like photosystem I subunit PsaC from the <i>Cyanobacterium synechococcus elongatus</i> . 1997 , 36, 13629-37	17
1509	Conformational analysis of LYS(11-36), a peptide derived from the beta-sheet region of T4 lysozyme, in TFE and SDS. 1997 , 36, 11525-33	43
1508	Absence of observable biotin-protein interactions in the 1.3S subunit of transcarboxylase: an NMR study. 1997 , 36, 14676-82	15
1507	Solution structure of the cellulose-binding domain of the endoglucanase Z secreted by <i>Erwinia chrysanthemi</i> . 1997 , 36, 16074-86	92
1506	Three-dimensional structure of a gamma-carboxyglutamic acid-containing conotoxin, conantokin G, from the marine snail <i>Conus geographus</i> : the metal-free conformer. 1997 , 36, 6906-14	41
1505	Conformational studies of mono- and bicyclic parathyroid hormone-related protein-derived agonists. 1997 , 36, 10372-83	32
1504	Structure of hexadienoyl-CoA bound to enoyl-CoA hydratase determined by transferred nuclear Overhauser effect measurements: mechanistic predictions based on the X-ray structure of 4-(chlorobenzoyl)-CoA dehalogenase. 1997 , 36, 2211-20	36
1503	Mono- and bicyclic analogs of parathyroid hormone-related protein. 2. Conformational analysis of antagonists by CD, NMR, and distance geometry calculations. 1997 , 36, 3300-7	29
1502	Contribution of increased length and intact capping sequences to the conformational preference for helix in a 31-residue peptide from the C terminus of myohemerythrin. 1997 , 36, 5234-44	41
1501	Solution structure and backbone dynamics of the human alpha3-chain type VI collagen C-terminal Kunitz domain,. 1997 , 36, 10439-50	15
1500	High-resolution NMR structure and backbone dynamics of the <i>Bacillus subtilis</i> response regulator, Spo0F: implications for phosphorylation and molecular recognition. 1997 , 36, 10015-25	62
1499	Mechanism of direct coupling between binding and induced structural change in regulatory calcium binding proteins. 1997 , 36, 4386-92	116
1498	NMR structure determination of a novel conotoxin, [Pro 7,13] alpha A-conotoxin PIVA. 1997 , 36, 1669-77	28
1497	Solution structure of the paramagnetic complex of the N-terminal domain of calmodulin with two Ce ³⁺ ions by ¹ H NMR. 1997 , 36, 11605-18	82
1496	Collagen-based structures containing the peptoid residue N-isobutylglycine (Nleu): conformational analysis of Gly-Nleu-Pro sequences by ¹ H-NMR and molecular modeling. 1997 , 36, 8725-32	31

1495	Is the structure of the N-domain of phosphoglycerate kinase affected by isolation from the intact molecule?. 1997 , 36, 333-40	21
1494	Conformational studies of RS-66271, an analog of parathyroid hormone-related protein with pronounced bone anabolic activity. 1997 , 40, 3025-31	22
1493	Three-dimensional solution structure of alpha-conotoxin MII, an alpha3beta2 neuronal nicotinic acetylcholine receptor-targeted ligand. 1997 , 36, 15693-700	55
1492	Novel Cyclic Biphenyl Ether Peptide β -Strand Mimetics and HIV-Protease Inhibitors. 1997 , 119, 441-442	53
1491	Pregnane glycosides. 1997 , 71, 169-325	22
1490	Solution structures of the R6 human insulin hexamer,. 1997 , 36, 9409-22	116
1489	Conformation of a protein kinase C substrate NG(28-43), and its analog in aqueous and sodium dodecyl sulfate micelle solutions. 1997 , 72, 554-66	10
1488	Three-dimensional structure of leucocin A in trifluoroethanol and dodecylphosphocholine micelles: spatial location of residues critical for biological activity in type IIa bacteriocins from lactic acid bacteria. 1997 , 36, 15062-72	148
1487	Comparison of the hemolytic activity and solution structures of two snake venom cardiotoxin analogues which only differ in their N-terminal amino acid. 1997 , 36, 14635-41	28
1486	NMR solution structure of a two-disulfide derivative of charybdotoxin: structural evidence for conservation of scorpion toxin alpha/beta motif and its hydrophobic side chain packing. 1997 , 36, 3760-6	25
1485	Design, Synthesis, and Evaluation of a Depsipeptide Mimic of Tendamistat. 1997 , 62, 93-102	36
1484	A diminished role for hydrogen bonds in antifreeze protein binding to ice. 1997 , 36, 14652-60	188
1483	Oxysporidinone: a novel, antifungal N-methyl-4-hydroxy-2-pyridone from <i>Fusarium oxysporum</i> . 1997 , 60, 33-5	45
1482	Structure of the chromatin binding (chromo) domain from mouse modifier protein 1. <i>EMBO Journal</i> , 1997 , 16, 2473-81	13 140
1481	A Trisubstituted 1,4-Diazepine-3-one-Based Dipeptidomimetic: Conformational Characterization by NMR and Computer Simulation. 1997 , 119, 2430-2436	11
1480	Multiconformational NMR analysis of sandostatin (octreotide): equilibrium between beta-sheet and partially helical structures. 1997 , 36, 1233-41	88
1479	Design of a potent combined pseudopeptide endothelin-A/endothelin-B receptor antagonist, Ac-DBhg16-Leu-Asp-Ile-[NMe]Ile-Trp21 (PD 156252): examination of its pharmacokinetic and spectral properties. 1997 , 40, 2228-40	89
1478	Threonine6-bradykinin: structural characterization in the presence of micelles by nuclear magnetic resonance and distance geometry. 1997 , 40, 92-8	37

1477	Accounting for Conformational Variability in NMR Structure of Cyclopeptides: Ensemble Averaging of Interproton Distance and Coupling Constant Restraints. 1997 , 119, 5239-5248	37
1476	Structure-based design of a constrained peptide mimic of the HIV-1 V3 loop neutralization site. 1997 , 266, 31-9	72
1475	NMR solution structure of the pathogenesis-related protein P14a. 1997 , 266, 576-93	123
1474	Interaction of the receptor binding domains of <i>Pseudomonas aeruginosa</i> pili strains PAK, PAO, KB7 and P1 to a cross-reactive antibody and receptor analog: implications for synthetic vaccine design. 1997 , 267, 382-402	34
1473	Solution structure of the conserved 16 S-like ribosomal RNA UGAA tetraloop. 1997 , 268, 348-58	51
1472	Structure of neurotoxin B-IV from the marine worm <i>Cerebratulus lacteus</i> : a helical hairpin cross-linked by disulphide bonding. 1997 , 268, 886-902	28
1471	Structural characterisation and comparison of the native and A-states of equine lysozyme. 1997 , 268, 903-21	70
1470	Solution structure of ferredoxin from the thermophilic cyanobacterium <i>Synechococcus elongatus</i> and its thermostability. 1997 , 268, 922-33	43
1469	Enhanced protein flexibility caused by a destabilizing amino acid replacement in BPTI. 1997 , 269, 154-64	54
1468	NMR structural studies of human cystatin C dimers and monomers. 1997 , 271, 266-77	97
1467	The detailed structure of tandem G.A mismatched base-pair motifs in RNA duplexes is context dependent. 1997 , 271, 147-58	51
1466	A characteristic arrangement of aromatic amino acid residues in the solution structure of the amino-terminal RNA-binding domain of <i>Drosophila</i> sex-lethal. 1997 , 272, 82-94	23
1465	NMR studies of a viral protein that mimics the regulators of complement activation. 1997 , 272, 253-65	119
1464	Structure and dynamics of the iron responsive element RNA: implications for binding of the RNA by iron regulatory binding proteins. 1997 , 274, 72-83	169
1463	Solution structure of human p8MTCP1, a cysteine-rich protein encoded by the MTCP1 oncogene, reveals a new alpha-helical assembly motif. 1997 , 274, 801-15	23
1462	Role of modified glutamic acid in the helical structure of conantokin-T. 1997 , 407, 243-8	16
1461	A hairpin-loop conformation in tandem repeat sequence of the ice nucleation protein revealed by NMR spectroscopy. 1997 , 409, 227-31	14
1460	The NMR solution structure of the NMDA receptor antagonist, conantokin-T, in the absence of divalent metal ions. 1997 , 411, 19-26	26

1459	A cytoplasmic peptide of the neurotrophin receptor p75NTR: induction of apoptosis and NMR determined helical conformation. 1997 , 415, 145-54	9
1458	Solution structure of robustoxin, the lethal neurotoxin from the funnel-web spider <i>Atrax robustus</i> . 1997 , 419, 191-6	60
1457	A cyclic disulfide peptide reproduces in solution the main structural features of a native antigenic site of foot-and-mouth disease virus. 1997 , 20, 209-19	11
1456	Glycosylated flavones as selective inhibitors of topoisomerase IV. 1997 , 41, 992-8	95
1455	Purification, characterization, and structural analysis of a plant low-temperature-induced protein. 1997 , 113, 367-76	27
1454	Structure of the repeating oligosaccharide from the lipopolysaccharide of the nitrogen-fixing bacterium <i>Acetobacter diazotrophicus</i> strain PAL 5. 1997 , 298, 311-8	7
1453	The structure of a novel insecticidal neurotoxin, omega-atracotoxin-HV1, from the venom of an Australian funnel web spider. 1997 , 4, 559-66	149
1452	Interlocking structural motifs mediate molecular discrimination by a theophylline-binding RNA. 1997 , 4, 644-9	236
1451	Photodegradation of irinotecan (CPT-11) in aqueous solutions: identification of fluorescent products and influence of solution composition. 1997 , 86, 1410-6	21
1450	A conserved infection pathway for filamentous bacteriophages is suggested by the structure of the membrane penetration domain of the minor coat protein g3p from phage fd. 1997 , 5, 265-75	71
1449	Solution structure of a type 2 module from fibronectin: implications for the structure and function of the gelatin-binding domain. 1997 , 5, 359-70	90
1448	Solution structure of the granular starch binding domain of <i>Aspergillus niger</i> glucoamylase bound to beta-cyclodextrin. 1997 , 5, 647-61	162
1447	The SH2 domain from the tyrosine kinase Fyn in complex with a phosphotyrosyl peptide reveals insights into domain stability and binding specificity. 1997 , 5, 1313-23	36
1446	The structure of versutoxin (delta-atracotoxin-Hv1) provides insights into the binding of site 3 neurotoxins to the voltage-gated sodium channel. 1997 , 5, 1525-35	108
1445	Solution structure and proposed binding mechanism of a novel potassium channel toxin kappa-conotoxin PVIIA. 1997 , 5, 1585-97	82
1444	Solution structure of Lqh-8/6, a toxin-like peptide from a scorpion venom--structural heterogeneity induced by proline cis/trans isomerization. <i>FEBS Journal</i> , 1997 , 246, 218-27	34
1443	Constitution and solution conformation of the antibiotic mersacidin determined by NMR and molecular dynamics. <i>FEBS Journal</i> , 1997 , 244, 501-12	39
1442	The solution structure and activity of caerin 1.1, an antimicrobial peptide from the Australian green tree frog, <i>Litoria splendida</i> . <i>FEBS Journal</i> , 1997 , 247, 545-57	112

1441	Three-dimensional solution structure of human angiogenin determined by ¹ H, ¹⁵ N-NMR spectroscopy—characterization of histidine protonation states and pKa values. <i>FEBS Journal</i> , 1997 , 250, 712-26		35
1440	Comparative study of chicken and human parathyroid hormone-(1-34)-peptides in solution with SDS. <i>FEBS Journal</i> , 1997 , 249, 878-85		10
1439	The FLG motif in the N-terminal region of glucoprotein 41 of human immunodeficiency virus type 1 adopts a type-I beta turn in aqueous solution and serves as the initiation site for helix formation. <i>FEBS Journal</i> , 1997 , 247, 896-905		17
1438	Efficient photoaffinity labeling of the rat V1a vasopressin receptor using a linear azidopeptidic antagonist. <i>FEBS Journal</i> , 1997 , 247, 906-13		10
1437	Spectroscopic studies of the C-terminal secretion signal of the <i>Serratia marcescens</i> haem acquisition protein (HasA) in various membrane-mimetic environments. <i>FEBS Journal</i> , 1997 , 243, 400-7		19
1436	Solution structure of reduced microsomal rat cytochrome b5. <i>FEBS Journal</i> , 1997 , 249, 270-9		25
1435	Secondary structure and tertiary fold of the human immunodeficiency virus protein U (Vpu) cytoplasmic domain in solution. <i>FEBS Journal</i> , 1997 , 245, 581-8		70
1434	A conformational study of the human and rat encephalitogenic myelin oligodendrocyte glycoprotein peptides 35-55. <i>FEBS Journal</i> , 1997 , 246, 59-70		20
1433	Solution structure of alpha-conotoxin MI determined by ¹ H-NMR spectroscopy and molecular dynamics simulation with the explicit solvent water. 1997 , 1343, 327-34		28
1432	Solution structure of midkine, a new heparin-binding growth factor. <i>EMBO Journal</i> , 1997 , 16, 6936-46	13	122
1431	Solution structure of a GAAA tetraloop receptor RNA. <i>EMBO Journal</i> , 1997 , 16, 7490-9	13	96
1430	Proton nuclear magnetic resonance studies on huwentoxin-I from the venom of the spider <i>Selenocosmia huwena</i> : 2. Three-dimensional structure in solution. 1997 , 16, 565-74		51
1429	A new method to determine DNA sugar conformation from the joint use of 2D and 3D NMR data. 1997 , 10, 187-192		1
1428	The effects of guanidine hydrochloride on the 'random coil' conformations and NMR chemical shifts of the peptide series GGXGG. 1997 , 10, 221-30		89
1427	Solution structures of staphylococcal nuclease from multidimensional, multinuclear NMR: nuclease-H124L and its ternary complex with Ca ²⁺ and thymidine-3',5'-bisphosphate. 1997 , 10, 143-64		26
1426	Two-dimensional ¹ H NMR experiments show that the 23-residue magainin antibiotic peptide is an alpha-helix in dodecylphosphocholine micelles, sodium dodecylsulfate micelles, and trifluoroethanol/water solution. 1997 , 9, 127-35		207
1425	NMR structural characterization of oligo-N-substituted glycine lead compounds from a combinatorial library. 1997 , 3, 1-15		25
1424	Complete ¹ H, ¹³ C, and ¹⁵ N NMR resonance assignments and secondary structure of human glutaredoxin in the fully reduced form. 1997 , 6, 383-90		10

1423	The interaction of beta-amyloid protein fragment (12-28) with lipid environments. 1997 , 6, 666-75	44	
1422	Multiple conformations of a human interleukin-3 variant. 1997 , 6, 1777-82	29	
1421	Conformational studies of the N-terminal lipid-associating domain of human apolipoprotein C-I by CD and ¹ H NMR spectroscopy. 1997 , 6, 1858-68	22	
1420	Solution structure of alpha t alpha, a helical hairpin peptide of de novo design. 1997 , 6, 1869-77	45	
1419	A novel RGD containing dodecapeptidomimetic which exhibits selective binding to the $\alpha_5\beta_1$ receptor. 1997 , 7, 997-1002	13	
1418	The conformational study of β -D-GlcA-(1,4)-l-Rha in solution by NMR and molecular dynamics simulations. 1997 , 224, 81-94	3	
1417	NMR study and comparison of the antigenic properties of a peptide recognized by two HIV-1 neutralizing antibodies. 1997 , 10, 256-61	6	
1416	Conformational analysis of the dipeptide taste ligand L-aspartyl-D-2-aminobutyric acid-(S)-alpha-ethylbenzylamide and its analogues by NMR spectroscopy, computer simulations and X-ray diffraction studies. 1997 , 3, 231-41	10	
1415	Structural comparison in solution of a native and retro peptide derived from the third helix of Staphylococcus aureus protein A, domain B: retro peptides, a useful tool for the discrimination of helix stabilization factors dependent on the peptide chain orientation. 1997 , 3, 299-313	11	
1414	Conformation of four peptides corresponding to the β helical segments of human GM128SF. 1997 , 3, 336-346	4	
1413	Pulsed field gradient spectroscopy (PFGS): application to the structure elucidation of (+)-(10S)-10-bromo- β -chamigrene. 1997 , 8, 167-172	7	
1412	Polymyxin B nonapeptide: Conformations in water and in the lipopolysaccharide-bound state determined by two-dimensional NMR and molecular dynamics. 1997 , 41, 251-265	54	
1411	Structural comparison between retro-inverso and parent peptides: molecular basis for the biological activity of a retro-inverso analogue of the immunodominant fragment of VP1 coat protein from foot-and-mouth disease virus. 1997 , 41, 569-90	25	
1410	Hydrophobic forces are responsible for the folding of a highly potent natriuretic peptide analogue at a membrane mimetic surface: an NMR study. 1997 , 42, 37-48	12	
1409	Structure, dynamics, and topological orientation of the polyether, ionophore antibiotic monensin, in a micellar environment. 1997 , 42, 759-769	9	
1408	Measurement of the degree of coupled isotopic enrichment of different positions in an antibiotic peptide by NMR. <i>Journal of Magnetic Resonance</i> , 1997 , 125, 120-31	3	
1407	Artifact-Free Pure Absorption PFG-Enhanced DQF-COSY Spectra Including a Gradient Pulse in the Evolution Period. <i>Journal of Magnetic Resonance</i> , 1997 , 125, 348-354	3	39
1406	Improvements for measuring ¹ H- ¹ H coupling constants in DNA via new stripe-COSY and superstripe-COSY pulse sequences combined with a novel strategy of selective deuteration. <i>Journal of Magnetic Resonance</i> , 1997 , 129, 212-8	3	10

1405	Determination of the absolute configuration of monosaccharides by ¹ H NMR spectroscopy of their per-O-(S)-2-methylbutyrate derivatives. 1997 , 300, 199-206	17	
1404	Isolation and structure elucidation of a highly haemolytic saponin from the Merck saponin extract using high-field gradient-enhanced NMR techniques. 1997 , 302, 67-78	26	
1403	Structural characterisation of galactoglucomannan secreted by suspension-cultured cells of <i>Nicotiana plumbaginifolia</i> . 1997 , 303, 79-92	51	
1402	Structural characterization of novel L-galactose-containing oligosaccharide subunits of jojoba seed xyloglucans. 1997 , 304, 11-20	48	
1401	Structural studies on a sulfated polysaccharide from an <i>Arthrobacter</i> sp. by NMR spectroscopy and methylation analysis. 1997 , 305, 253-60	5	
1400	Structure of the capsular polysaccharide of <i>Clostridium perfringens</i> Hobbs 10 determined by NMR spectroscopy. 1997 , 305, 65-72	18	
1399	NMR observation of two states of triple helix in the thermal transition of (Pro-Pro-Gly) ₁₀ . 1997 , 281, 86-91	6	
1398	Structural characterization of the galactoxylomannan of <i>Cryptococcus neoformans</i> Cap67. 1998 , 306, 315-30	109	
1397	Chemical and spectroscopic characterisation of the phosphatidylinositol manno-oligosaccharides from <i>Mycobacterium bovis</i> AN5 and WAg201 and <i>Mycobacterium smegmatis</i> mc2 155. 1998 , 308, 397-408	25	
1396	Local helix content and RNA-binding activity of the N-terminal leucine-repeat region of hepatitis delta antigen. 1998 , 12, 183-8	7	
1395	Solution structure of the recombinant human oncoprotein p13MTCP1. 1998 , 11, 337-54	10	
1394	Reduction of ortho-aminobenzoyl-proline fluorescence and formation of pyrrolobenzodiazepine-5,11-dione. 1998 , 5, 19-28	1	
1393	Solution structure of cytochrome c6 from the thermophilic cyanobacterium <i>Synechococcus elongatus</i> . <i>EMBO Journal</i> , 1998 , 17, 27-36	13	41
1392	Structure of the IGF-binding domain of the insulin-like growth factor-binding protein-5 (IGFBP-5): implications for IGF and IGF-I receptor interactions. <i>EMBO Journal</i> , 1998 , 17, 6558-72	13	131
1391	A helix-turn-helix structure unit in human centromere protein B (CENP-B). <i>EMBO Journal</i> , 1998 , 17, 827-37	33	
1390	Structure of a human DNA repair protein UBA domain that interacts with HIV-1 Vpr. 1998 , 5, 1042-7	114	
1389	Structural studies on phospholamban and implications for regulation of the Ca(2+)-ATPase. 1998 , 853, 63-78	9	
1388	Binding of peptides in solution by the <i>Escherichia coli</i> chaperone PapD as revealed using an inhibition ELISA and NMR spectroscopy. 1998 , 6, 2085-101	12	

1387	Isoquinolinium N-arylimides and electrophilic ethylenes: Structures and NMR spectra of cycloadducts. 1998 , 54, 3735-3744	9
1386	Two-dimensional filter-diagonalization: spectral inversion of 2D NMR time-correlation signals including degeneracies. 1998 , 291, 465-470	16
1385	Homology predicted structure and comparison with the secondary structure from NMR data for plastocyanin for the cyanobacterium <i>Synechocystis</i> sp. PCC 6803. 1998 , 275-276, 73-89	5
1384	Secondary structure of an antibacterial peptide Abp3 studied by two-dimensional proton-NMR. 1998 , 1384, 299-305	5
1383	Proton resonance assignments and ligand exchange kinetics in high-spin and mixed-spin myoglobin complexes using two-dimensional exchange spectroscopy. 1998 , 1388, 66-76	2
1382	Solution structure of a fragment of the dimerization domain of DP-1 determined by ¹ H-nuclear magnetic resonance and distance geometry. 1998 , 1429, 18-28	
1381	Solution structures of the melanocyte-stimulating hormones by two-dimensional NMR spectroscopy and dynamical simulated-annealing calculations. <i>FEBS Journal</i> , 1998 , 257, 31-40	21
1380	Solution structures of human immunodeficiency virus type 1 (HIV-1) and moloney murine leukemia virus (MoMLV) capsid protein major-homology-region peptide analogs by NMR spectroscopy. <i>FEBS Journal</i> , 1998 , 257, 69-77	9
1379	NMR characterization of a single-cysteine mutant of <i>Escherichia coli</i> thioredoxin and a covalent thioredoxin-peptide complex. <i>FEBS Journal</i> , 1998 , 257, 299-308	14
1378	Conformation analysis of eel calcitonin--comparison with the conformation of elcatonin. <i>FEBS Journal</i> , 1998 , 257, 331-6	9
1377	The solution NMR structure of a blue-green algae hepatotoxin, microcystin-RR--a comparison with the structure of microcystin-LR. <i>FEBS Journal</i> , 1998 , 258, 301-12	11
1376	The solution structure of parsley [2Fe-2S]ferredoxin. <i>FEBS Journal</i> , 1998 , 258, 465-77	22
1375	Solution structure of an artificial Fe ₈ S ₈ ferredoxin: the D ¹³ C variant of <i>Bacillus schlegelii</i> Fe ₇ S ₈ ferredoxin. <i>FEBS Journal</i> , 1998 , 258, 502-14	9
1374	Neutral core oligosaccharides of bovine submaxillary mucin--use of lead tetraacetate in the cold for establishing branch positions. <i>FEBS Journal</i> , 1998 , 258, 603-22	33
1373	Solution structure of the DNA-binding domain of human telomeric protein, hTRF1. 1998 , 6, 1057-65	47
1372	Spectral editing technique for the in vitro and in vivo detection of taurine. <i>Journal of Magnetic Resonance</i> , 1998 , 133, 70-8	3 27
1371	Quantum mechanical calculations and experimental measurement of N-terminal charge effects on ¹ H _N and ¹ H _{Cα chemical shifts in peptides. 1998, 38, 573-581}	4
1370	Conformational analysis of cyclic analogues of the <i>Saccharomyces cerevisiae</i> alpha-factor pheromone. 1998 , 45, 21-34	10

- 1369 How to Predict Conformations Accessible to a Molecule in Solution: Validation of a Force Field-Based Prediction of NOE Distances by Comparison with the Experimental Data for the Series of Compounds $\text{CH}_3\text{C}[\text{CH}_2\text{P}(\text{Bzl})\text{R}]_3\text{Mo}(\text{CO})_3$ (R = Ph, m-Xyl). **1998**, 1998, 1641-1653 5
- 1368 SIAM, ein neues NMR-Experiment für die Bestimmung homonuclearer Kopplungskonstanten. **1998**, 110, 817-821
- 1367 Synthesis, Spectroscopic Characterisation, and Metal Ion Interaction of a New α -Helical Peptide. **1998**, 4, 1791-1798 9
- 1366 ^1H NMR structure of an antifungal gamma-thionin protein Slalpha1: similarity to scorpion toxins. **1998**, 32, 334-49 32
- 1365 Structure and dynamic properties of the single disulfide-deficient alpha-amylase inhibitor [C45A/C73A]tendamistat: an NMR study. **1998**, 33, 285-94 9
- 1364 Reduction of ortho-aminobenzoyl-proline fluorescence and formation of pyrrolobenzodiazepine-5,11-dione. **1998**, 5, 19-28 2
- 1363 Crown ether styryl dyes. **1998**, 47, 97-106 5
- 1362 Structure of the inactivating gate from the Shaker voltage gated K^+ channel analyzed by NMR spectroscopy. **1998**, 27, 99-104 21
- 1361 NMR structures of the C-terminal end of human complement serine protease C1s. **1998**, 54, 171-8 1
- 1360 Protonation of alkene-rhodium(I) complexes leads to alkyl-rhodium(III) - An NMR study. **1998**, 558, 123-130 8
- 1359 Flavonoids from red onion (*Allium cepa*). **1998**, 47, 281-285 140
- 1358 A cycloartane triterpene 3 β ,16 β -diglucoside from *Astragalus trigonus* and its non natural 6-hydroxy epimer. **1998**, 49, 845-852 13
- 1357 Cycloartane triterpene glycosides from *Astragalus sieberi*. **1998**, 48, 1403-1409 17
- 1356 Structural study on silkworm eclosion hormone fragment (1-34) in solution by proton nuclear magnetic resonance spectroscopy. **1998**, 6, 465-72 4
- 1355 Glycosyl-transferase catalyzed assemblage of sialyl-Lewis(x)-saccharopeptides. **1998**, 6, 1673-82 27
- 1354 Interhelical contacts are required for the helix bundle fold of apolipoprotein III and its ability to interact with lipoproteins. **1998**, 7, 336-41 9
- 1353 Solution structure of Compstatin, a potent complement inhibitor. **1998**, 7, 619-27 75
- 1352 Solution structure and dynamics of a designed monomeric variant of the lambda Cro repressor. **1998**, 7, 983-93 9

1351	A conformational equilibrium in a protein fragment caused by two consecutive capping boxes: 1H-, 13C-NMR, and mutational analysis. 1998 , 7, 1506-15	12
1350	Role of the 6-20 disulfide bridge in the structure and activity of epidermal growth factor. 1998 , 7, 1738-49	33
1349	Pulmonary surfactant-associated polypeptide C in a mixed organic solvent transforms from a monomeric alpha-helical state into insoluble beta-sheet aggregates. 1998 , 7, 2533-40	76
1348	Probing the Scope of the Asymmetric Dihydroxylation of Polymer-Bound Olefins. Monitoring by HRMAS NMR Allows for Reaction Control and On-Bead Measurement of Enantiomeric Excess. 1998 , 120, 8994-9000	39
1347	Solution structure and backbone dynamics of Mason-Pfizer monkey virus (MPMV) nucleocapsid protein. 1998 , 7, 2265-80	29
1346	Solution and solid state conformation of the human EGF receptor transmembrane region. 1998 , 1371, 241-53	11
1345	Adsorption of human lysozyme onto hydroxyapatite. Identification of its adsorbing site using site-directed mutagenesis. 1998 , 422, 175-8	36
1344	Three-dimensional structure of binase in solution. 1998 , 431, 250-4	11
1343	Accessibility of selenomethionine proteins by total chemical synthesis: structural studies of human herpesvirus-8 MIP-II. 1998 , 441, 77-82	11
1342	NMR and CD conformational studies of bradykinin and its agonists and antagonists: application to receptor binding. 1998 , 76, 257-266	49
1341	Solution structures of the Ca ²⁺ -free and Ca ²⁺ -bound C2A domain of synaptotagmin I: does Ca ²⁺ induce a conformational change?. 1998 , 37, 16106-15	201
1340	Stereospecific 1H and 13C NMR Assignments of Crotonyl CoA and Hexadienoyl CoA: Conformational Analysis and Comparison with Protein-CoA Complexes. 1998 , 120, 9988-9994	9
1339	Solution structure of the epidermal growth factor (EGF)-like module of human complement protease C1r, an atypical member of the EGF family. 1998 , 37, 1204-14	46
1338	Structural characterization and optimization of antibody-selected phage library mimotopes of an antigen associated with autoimmune recurrent thrombosis. 1998 , 37, 16069-81	18
1337	Site-specific phosphorylation of Lys-Ser-Pro repeat peptides from neurofilament H by cyclin-dependent kinase 5: structural basis for substrate recognition. 1998 , 37, 4759-66	25
1336	Analysis of the Structure and Stability of a Backbone-Modified Oligonucleotide: Implications for Avoiding Product Inhibition in Catalytic Template-Directed Synthesis. 1998 , 120, 3019-3031	44
1335	Novel branched nod factor structure results from alpha-(1->3) fucosyl transferase activity: the major lipo-chitin oligosaccharides from Mesorhizobium loti strain NZP2213 bear an alpha-(1->3) fucosyl substituent on a nonterminal backbone residue. 1998 , 37, 9024-32	50
1334	The Rubredoxin from Clostridium pasteurianum: Mutation of the Iron Cysteinyll Ligands to Serine. Crystal and Molecular Structures of Oxidized and Dithionite-Treated Forms of the Cys42Ser Mutant. 1998 , 120, 4135-4150	70

1333	Solution structure of the oxidized Fe7S8 ferredoxin from the thermophilic bacterium <i>Bacillus schlegelii</i> by 1H NMR spectroscopy. 1998 , 37, 9812-26	45
1332	Solution Structure of β D, a Nativelike de Novo Designed Protein. 1998 , 120, 1138-1145	132
1331	NMR study suggests a major role for Arg111 in maintaining the structure and dynamical properties of type II human cellular retinoic acid binding protein. 1998 , 37, 13021-32	12
1330	Structural studies of <i>Impatiens balsamina</i> antimicrobial protein (Ib-AMP1). 1998 , 37, 983-90	51
1329	NMR study of the conformation and localization of porcine galanin in SDS micelles. Comparison with an inactive analog and a galanin receptor antagonist. 1998 , 37, 9169-78	28
1328	Structural characterization and topology of the second potential membrane anchor region in the thromboxane A2 synthase amino-terminal domain. 1998 , 37, 822-30	19
1327	Three-dimensional solution structure of alpha-conotoxin MII by NMR spectroscopy: effects of solution environment on helicity. 1998 , 37, 15621-30	55
1326	Structure determination of the Ras-binding domain of the Ral-specific guanine nucleotide exchange factor Rlf. 1998 , 37, 13453-62	34
1325	Dihydroxylation of the Triene Subunit of Rapamycin. 1998 , 63, 10069-10073	4
1324	Human low-molecular-weight salivary mucin expresses the sialyl lewisx determinant and has L-selectin ligand activity. 1998 , 37, 4916-27	56
1323	HIV-1 Reverse Transcriptase Can Discriminate between Two Conformationally Locked Carbocyclic AZT Triphosphate Analogues. 1998 , 120, 2780-2789	138
1322	Helix-stabilizing nonpolar interactions between tyrosine and leucine in aqueous and TFE solutions: 2D-1H NMR and CD studies in alanine-lysine peptides. 1998 , 37, 17318-30	34
1321	Measurement of Deoxyribose 3JHH Scalar Couplings Reveals Protein Binding-Induced Changes in the Sugar Puckers of the DNA. 1998 , 120, 821-822	16
1320	Thermal Denaturation of β Lactoglobulin. A 1H NMR Study. 1998 , 46, 1805-1813	62
1319	Solution structure of the SH3 domain from Bruton's tyrosine kinase. 1998 , 37, 2912-24	68
1318	Fluorescence and NMR Studies of Human Seminal Plasma Prostatic Inhibin: Association of Lifetimes with Sterically Constrained Tryptophans. 1998 , 102, 5520-5528	1
1317	Determination of the magnetic axes of cobalt(II) and nickel(II) azurins from 1H NMR data: influence of the metal and axial ligands on the origin of magnetic anisotropy in blue copper proteins. 1998 , 37, 8659-73	51
1316	Conformational studies on a diastereoisomeric pair of tricyclic nonclassical cannabinoids by NMR spectroscopy and computer molecular modeling. 1998 , 41, 167-74	17

1315	Structure elucidation of sphingolipids from the mycopathogen <i>Paracoccidioides brasiliensis</i> : an immunodominant beta-galactofuranose residue is carried by a novel glycosylinositol phosphorylceramide antigen. 1998 , 37, 8764-75	80
1314	Self-Assembling Cyclic β -Peptide Nanotubes as Artificial Transmembrane Ion Channels. 1998 , 120, 651-656	270
1313	Solution structure of murine macrophage inflammatory protein-2. 1998 , 37, 8303-13	34
1312	The solution structure of type II antifreeze protein reveals a new member of the lectin family. 1998 , 37, 4712-21	108
1311	Formation of native disulfide bonds in endothelin-1. Structural evidence for the involvement of a highly specific salt bridge between the prosequence and the endothelin-1 sequence. 1998 , 37, 5220-30	7
1310	NMR solution structure of type II human cellular retinoic acid binding protein: implications for ligand binding. 1998 , 37, 12727-36	34
1309	High-resolution solution NMR structure of the minimal active domain of the human immunodeficiency virus type-2 nucleocapsid protein. 1998 , 37, 17704-13	19
1308	Aggregation behaviour and Zn ²⁺ binding properties of secretin. 1998 , 37, 16967-74	6
1307	Three-dimensional solution structure of conotoxin psi-PIIIe, an acetylcholine gated ion channel antagonist. 1998 , 37, 1215-20	18
1306	Structure-function relationships in side chain lactam cross-linked peptide models of a conserved N-terminal domain of apolipoprotein E. 1998 , 37, 13222-9	7
1305	¹ H nuclear magnetic resonance study on equilibrium between two four-stranded solution conformations of short d(CnT). 1998 , 37, 12979-86	19
1304	Interaction of a novel GDP exchange inhibitor with the Ras protein. 1998 , 37, 15631-7	34
1303	NMR structure and comparison of the archaeal histone HFoB from the mesophile <i>Methanobacterium formicicum</i> with HMfB from the hyperthermophile <i>Methanothermus fervidus</i> . 1998 , 37, 10573-80	24
1302	Solution structure of calmodulin-W-7 complex: the basis of diversity in molecular recognition. 1998 , 276, 165-76	137
1301	Solution structure of the glycosylated second type 2 module of fibronectin. 1998 , 276, 177-87	50
1300	Dynamical behavior of the HIV-1 nucleocapsid protein. 1998 , 279, 633-49	129
1299	Determination of the three-dimensional solution structure of <i>Raphanus sativus</i> antifungal protein 1 by ¹ H NMR. 1998 , 279, 257-70	137
1298	Structure determination of the small ubiquitin-related modifier SUMO-1. 1998 , 280, 275-86	321

1297	Solution structure of the M13 major coat protein in detergent micelles: a basis for a model of phage assembly involving specific residues. 1998 , 282, 401-19	74
1296	The role of structure in antibody cross-reactivity between peptides and folded proteins. 1998 , 281, 183-201	81
1295	Side-chains in native and random coil protein conformations. Analysis of NMR coupling constants and chi1 torsion angle preferences. 1998 , 280, 867-77	35
1294	Insights into the mechanism of heterodimerization from the 1H-NMR solution structure of the c-Myc-Max heterodimeric leucine zipper. 1998 , 281, 165-81	88
1293	Alpha-helix mimicry of a beta-turn. 1998 , 281, 235-40	17
1292	Solution structure of <i>Desulfovibrio vulgaris</i> (Hildenborough) ferrocyclochrome c3: structural basis for functional cooperativity. 1998 , 281, 719-39	54
1291	A structural homologue of colipase in black mamba venom revealed by NMR floating disulphide bridge analysis. 1998 , 283, 205-19	62
1290	The solution structure of a cytotoxic ribonuclease from the oocytes of <i>Rana catesbeiana</i> (bullfrog). 1998 , 283, 231-44	26
1289	Structure determination and analysis of helix parameters in the DNA decamer d(CATGGCCATG) ₂ comparison of results from NMR and crystallography. 1998 , 284, 1453-63	35
1288	Wine analysis by 1D and 2D NMR spectroscopy. 1998 , 26, 97-101	29
1287	Intramolecular Diels-Alder Reactions of 3-(Tetrahydropyridinyl)indoles: Stereoselective Synthesis of Novel Pentacyclic Ring Systems. 1998 , 63, 1974-1980	10
1286	Three-dimensional structure of kappa-conotoxin PVIIA, a novel potassium channel-blocking toxin from cone snails. 1998 , 37, 5407-16	78
1285	Cylindrical Sheet Peptide Assemblies. 1998 , 120, 8949-8962	163
1284	Oxidative Addition of Dihydrogen to [Ir(bisphosphine)(1,5-cyclooctadiene)]BF ₄ Complexes: Kinetic and Thermodynamic Selectivity. 1998 , 120, 10115-10125	29
1283	Three-dimensional solution structure of lactoferricin B, an antimicrobial peptide derived from bovine lactoferrin. 1998 , 37, 4288-98	208
1282	Self-association of disulfide-dimerized melittin analogues. 1998 , 37, 5699-708	17
1281	Vancomycin: conformational consequences of the sugar substituent. 1998 , 41, 2090-9	30
1280	Model of the complex formed between the H2 domain of the TATA box binding protein and the L(281-301) fragment of the human transcription factor TFIIA. 1998 , 11, 729-38	6

1279	Mapping of a protein-RNA kissing hairpin interface: Rom and Tar-Tar*. 1998 , 26, 4688-95		31
1278	A membrane setting for the sorting motifs present in the adenovirus E3-13.7 protein which down-regulates the epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 1998 , 273, 17343-50	5.4	16
1277	Solution structure and internal motion of a bioactive peptide derived from nerve growth factor. <i>Journal of Biological Chemistry</i> , 1998 , 273, 23652-8	5.4	28
1276	Solution structure of a syndecan-4 cytoplasmic domain and its interaction with phosphatidylinositol 4,5-bisphosphate. <i>Journal of Biological Chemistry</i> , 1998 , 273, 13022-9	5.4	78
1275	Identification of a novel heptoglycan of alpha1-->2-linked D-glycero-D-manno-heptopyranose. Chemical and antigenic structure of lipopolysaccharides from <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i> rough strain R20 (O1:K20-). <i>Journal of Biological Chemistry</i> , 1998 , 273, 7006-17	5.4	69
1274	An NMR study of ligand binding by maltodextrin binding protein. 1998 , 76, 189-197		7
1273	Delineation of conformational preferences in human salivary statherin by ¹ H, ³¹ P NMR and CD studies: sequential assignment and structure-function correlations. 1998 , 16, 91-107		44
1272	Addressing the tertiary structure of human parathyroid hormone-(1-34). <i>Journal of Biological Chemistry</i> , 1998 , 273, 10420-7	5.4	90
1271	Conformational changes in conantokin-G induced upon binding of calcium and magnesium as revealed by NMR structural analysis. <i>Journal of Biological Chemistry</i> , 1998 , 273, 16248-58	5.4	35
1270	Identification of a novel core type in <i>Salmonella</i> lipopolysaccharide. Complete structural analysis of the core region of the lipopolysaccharide from <i>Salmonella enterica</i> sv. <i>Arizonae</i> O62. <i>Journal of Biological Chemistry</i> , 1998 , 273, 3817-29	5.4	42
1269	Sequence-specific ¹ H NMR resonance assignments and secondary structure of human apolipoprotein C-I in the presence of sodium dodecyl sulfate. 1998 , 76, 267-275		12
1268	Structural and energetic factors of the increased thermal stability in a genetically engineered <i>Escherichia coli</i> adenylate kinase. <i>Journal of Biological Chemistry</i> , 1998 , 273, 19102-7	5.4	21
1267	Characterization of a novel branched tetrasaccharide of 3-deoxy-D-manno-oct-2-ulopyranosonic acid. The structure of the carbohydrate backbone of the lipopolysaccharide from <i>Acinetobacter baumannii</i> strain nctc 10303 (atcc 17904). <i>Journal of Biological Chemistry</i> , 1998 , 273, 28122-31	5.4	29
1266	Novel 3'-Spiro Nucleoside Analogues of TSAO-T. Part II. a Comparative Study Based on NMR Conformational Analysis in Solution and Theoretical Calculations. 1998 , 9, 49-56		18
1265	NMR determination of the major solution conformation of a peptoid pentamer with chiral side chains. 1998 , 95, 4309-14		268
1264	Interaction of Pyrylium Dye with Self-Complementary DNA Oligomer as Studied by H NMR Spectroscopy 1998 , 17, 603-611		1
1263	Structure and topography of the membrane-binding C2 domain of factor VIII in the presence of dodecylphosphocholine micelles. <i>Biochemical Journal</i> , 1998 , 332 (Pt 2), 549-55	3.8	19
1262	Germacrene C synthase from <i>Lycopersicon esculentum</i> cv. VFNT cherry tomato: cDNA isolation, characterization, and bacterial expression of the multiple product sesquiterpene cyclase. 1998 , 95, 2216-21		106

1261	Assignment of the helical structure in neuropeptide Y by HPLC studies of methionine replacement analogues and ¹ H-NMR spectroscopy. 1998 , 39, 207-219		3
1260	Single atom modification (O→S) of tRNA confers ribosome binding. 1999 , 5, 188-94		106
1259	Solution structure and dynamics of a serpin reactive site loop using interleukin 1beta as a presentation scaffold. 1999 , 12, 189-202		3
1258	Assignment of the L30-mRNA complex using selective isotopic labeling and RNA mutants. 1999 , 27, 4059-70		13
1257	A conotoxin from <i>Conus textile</i> with unusual posttranslational modifications reduces presynaptic Ca ²⁺ influx. 1999 , 96, 5758-63		114
1256	Direct kinetic evidence for folding via a highly compact, misfolded state. <i>Journal of Biological Chemistry</i> , 1999 , 274, 26828-37	5-4	16
1255	Direct NMR observation of the thioredoxin-mediated reduction of the chloroplast NADP-malate dehydrogenase provides a structural basis for the relief of autoinhibition. <i>Journal of Biological Chemistry</i> , 1999 , 274, 34539-42	5-4	33
1254	NMR solution structure of domain 1 of human annexin I shows an autonomous folding unit. <i>Journal of Biological Chemistry</i> , 1999 , 274, 2971-7	5-4	12
1253	NMR structures of a nonapeptide from DNA binding domain of human polymerase-alpha determined by iterative complete-relaxation-matrix approach. 1999 , 16, 1075-85		5
1252	Solution structure of an insect growth factor, growth-blocking peptide. <i>Journal of Biological Chemistry</i> , 1999 , 274, 1887-90	5-4	33
1251	NMR methods utilized in combinatorial chemistry research. 1999 , 35, 153-200		49
1250	Conformational studies on the specific cleavage site of Type I collagen (alpha-1) fragment (157-192) by cathepsins K and L by proton NMR spectroscopy. 1999 , 7, 375-9		21
1249	C-terminal capping motifs in model helical peptides. 1999 , 7, 143-51		14
1248	A 30-residue fragment of the carp granulin-1 protein folds into a stack of two beta-hairpins similar to that found in the native protein. 1999 , 53, 590-7		18
1247	The effect of conformation of the acyloxyalkoxy-based cyclic prodrugs of opioid peptides on their membrane permeability. 1999 , 53, 403-13		28
1246	Structural recognition of an ICAM-1 peptide by its receptor on the surface of T cells: conformational studies of cyclo (1, 12)-Pen-Pro-Arg-Gly-Gly-Ser-Val-Leu-Val-Thr-Gly-Cys-OH. 1999 , 53, 422-31		25
1245	Effect of conformation on the conversion of cyclo-(1,7)-Gly-Arg-Gly-Asp-Ser-Pro-Asp-Gly-OH to its cyclic imide degradation product. 1999 , 54, 43-53		6
1244	The solution structure of uperin 3.6, an antibiotic peptide from the granular dorsal glands of the Australian toadlet, <i>Uperoleia mjobergii</i> . 1999 , 54, 137-45		29

1243	NMR analysis of human salivary mucin (MUC7) derived O-linked model glycopeptides: comparison of structural features and carbohydrate-peptide interactions. 1999 , 54, 290-310	41
1242	Characterization of monomeric and dimeric B domain of Staphylococcal protein A. 1999 , 54, 344-52	4
1241	NMR structure of active and inactive forms of the sterol-dependent antifungal antibiotic bacillomycin L. <i>FEBS Journal</i> , 1999 , 264, 200-10	36
1240	Multinuclear magnetic resonance studies of Escherichia coli adenylate kinase in free and bound forms. Resonance assignment, secondary structure and ligand binding. <i>FEBS Journal</i> , 1999 , 264, 765-74	12
1239	High-resolution solution structure of gurmarin, a sweet-taste-suppressing plant polypeptide. <i>FEBS Journal</i> , 1999 , 264, 525-33	26
1238	A coil-helix instead of a helix-coil motif can be induced in a chloroplast transit peptide from Chlamydomonas reinhardtii. <i>FEBS Journal</i> , 1999 , 265, 171-80	32
1237	Host defence peptides from the skin glands of the Australian blue mountains tree-frog Litoria citropa. Solution structure of the antibacterial peptide citropin 1.1. <i>FEBS Journal</i> , 1999 , 265, 627-37	65
1236	Type I beta-turn conformation is important for biological activity of the melanocyte-stimulating hormone analogues. <i>FEBS Journal</i> , 1999 , 265, 430-40	28
1235	Solution structure of a new hypothalamic neuropeptide, human hypocretin-2/orexin-B. <i>FEBS Journal</i> , 1999 , 266, 831-9	54
1234	The chimeric peptide [Lys(-2)-Arg(-1)]-sarafotoxin-S6b, composed of the endothelin pro-sequence and sarafotoxin, retains the salt-bridge staple between Arg(-1) and Asp8 previously observed in [Lys(-2)-Arg(-1)]-endothelin. Implications of this salt-bridge in the contractile activity and the oxidative folding reaction. <i>FEBS Journal</i> , 1999 , 266, 977-85	5
1233	Determination of solution conformations of PrP106-126, a neurotoxic fragment of prion protein, by 1H NMR and restrained molecular dynamics. <i>FEBS Journal</i> , 1999 , 266, 1192-201	30
1232	A synthetic peptide adhesion epitope as a novel antimicrobial agent. 1999 , 17, 42-7	89
1231	A novel loop-loop recognition motif in the yeast ribosomal protein L30 autoregulatory RNA complex. 1999 , 6, 1139-47	99
1230	The structure of the specific capsular polysaccharide of Rhodococcus equi serotype 4. 1999 , 320, 209-22	14
1229	An NMR solution study of the mega-oligosaccharide, rhamnogalacturonan II. 1999 , 14, 253-71	25
1228	Refined solution structure and backbone dynamics of 15N-labeled C12A-p8MTCP1 studied by NMR relaxation. 1999 , 15, 271-88	32
1227	NMR structure of the chimeric hybrid duplex r(gcaguggc).r(gcca)d(CTGC) comprising the tRNA-DNA junction formed during initiation of HIV-1 reverse transcription. 1999 , 13, 343-55	36
1226	Folding propensities of synthetic peptide fragments covering the entire sequence of phage 434 Cro protein. 1999 , 8, 1675-88	12

1225	Solution structures of the first and second RNA-binding domains of human U2 small nuclear ribonucleoprotein particle auxiliary factor (U2AF(65)). <i>EMBO Journal</i> , 1999 , 18, 4523-34	13	37
1224	Conformational studies of a bicyclic, lactam-constrained parathyroid hormone-related protein-derived agonist. 1999 , 5, 330-7		6
1223	Solution structure of the histidine-containing phosphocarrier protein from <i>Staphylococcus carnosus</i> . 1999 , 17, 465-480		12
1222	Solution structure of alpha-conotoxin Iml determined by two-dimensional NMR spectroscopy. 1999 , 1431, 384-94		8
1221	NMR study of the binding of all-trans-retinoic acid to type II human cellular retinoic acid binding protein. 1999 , 1433, 240-52		5
1220	Solution structure of a pair of modules from the gelatin-binding domain of fibronectin. 1999 , 7, 1451-60		19
1219	Structure of a putative ancestral protein encoded by a single sequence repeat from a multidomain proteinase inhibitor gene from <i>Nicotiana glauca</i> . 1999 , 7, 793-802		17
1218	Solution structure of a conformationally constrained Arg-Gly-Asp-like motif inserted into the alpha/beta scaffold of leurotoxin I. <i>FEBS Journal</i> , 1999 , 260, 810-7		5
1217	The structure of the carbohydrate backbone of the core-lipid A region of the lipopolysaccharide from a clinical isolate of <i>Yersinia enterocolitica</i> O:9. <i>FEBS Journal</i> , 1999 , 261, 19-24		19
1216	¹ H-NMR structural studies of a cystine-linked peptide containing residues 71-93 of transthyretin and effects of a Ser84 substitution implicated in familial amyloidotic polyneuropathy. <i>FEBS Journal</i> , 1999 , 262, 586-94		8
1215	Suppression of diagonal peaks in TROSY-type ¹ H NMR NOESY spectra of ¹⁵ N-labeled proteins. <i>Journal of Magnetic Resonance</i> , 1999 , 140, 499-503	3	23
1214	Cycloadditions of adamantanethione S-methylide to CC multiple bonds. 1999 , 36, 959-968		15
1213	Complete assignment of the ¹ H, ¹³ C and ¹⁹ F NMR spectra of fluoro-substituted 3H-naphthopyrans. 1999 , 37, 159-162		13
1212	Three-dimensional solution structures of two DNA dodecamers through full relaxation matrix analysis. 1999 , 37, 564-572		1
1211	Peptides and proteins in neurodegenerative disease: helix propensity of a polypeptide containing helix 1 of the mouse prion protein studied by NMR and CD spectroscopy. 1999 , 51, 145-52		30
1210	An nmr conformational analysis of a synthetic peptide Cn2(1-15)NH ₂ -S-S-acetyl-Cn2(52-66)NH ₂ from the New World <i>Centruroides noxius</i> 2 (Cn2) scorpion toxin: comparison of the structure with those of the <i>Centruroides</i> scorpion toxins. 1999 , 49, 277-86		1
1209	A light-harvesting antenna protein retains its folded conformation in the absence of protein-lipid and protein-pigment interactions. 1999 , 49, 361-72		17
1208	Conformational analysis of potent sweet taste ligands by NMR and computer simulations. 1999 , 49, 525-539		2

1207	Solid phase synthesis and NMR conformational studies on cyclic decapeptide template molecule. 1999 , 49, 565-74	6
1206	Conformational studies of parathyroid hormone (PTH)/PTH-related protein (PTHrP) point-mutated hybrids. 1999 , 50, 525-35	14
1205	Design, synthesis, and conformational studies of the hGM-CSF derived peptide (13-27)-Gly-(75-87). 1999 , 50, 545-54	
1204	Conformational and functional variability supported by the BPTI fold: Solution structure of the Ca ²⁺ channel blocker calcicludine. 1999 , 34, 520-532	37
1203	Solution structure and RNA-binding activity of the N-terminal leucine-repeat region of hepatitis delta antigen. 1999 , 37, 121-9	27
1202	The three-dimensional solution structure of Aesculus hippocastanum antimicrobial protein 1 determined by 1H nuclear magnetic resonance. 1999 , 37, 388-403	52
1201	Conformational analyses of cyclic hexapeptide analogs of somatostatin containing arylalkyl peptoid and naphthylalanine residues. 1999 , 5, 161-75	11
1200	Conformations and pharmacophores of cyclic RGD containing peptides which selectively bind integrin alpha(v)beta3. 1999 , 5, 491-506	18
1199	Photomodulierung der Konformation cyclischer Peptide mit Azobenzol-Bausteinen im Peptidrückgrat. 1999 , 111, 2941-2943	31
1198	Covalent Capture and Stabilization of Cylindrical β -Sheet Peptide Assemblies. 1999 , 5, 782-792	61
1197	Photomodulation of the Conformation of Cyclic Peptides with Azobenzene Moieties in the Peptide Backbone. 1999 , 38, 2771-2774	127
1196	Saponins in alfalfa (<i>Medicago sativa</i> L.) root and their structural elucidation. 1999 , 47, 3185-92	64
1195	Conformational studies by NMR of the antimicrobial peptide, drosocin, and its non-glycosylated derivative: effects of glycosylation on solution conformation. 1999 , 38, 705-14	65
1194	Retention of the Cis Proline Conformation in Tripeptide Fragments of Bovine Pancreatic Ribonuclease A Containing a Non-natural Proline Analogue, 5,5-Dimethylproline. 1999 , 121, 11558-11566	92
1193	Solid phase organic synthesis of piperazinone containing enkephalin mimetics: a readily derivatized, traceless scaffold. 1999 , 1, 383-7	25
1192	NMR conformational analysis on cyclic decapeptide template molecule. 1999 , 77, 1394-1404	2
1191	Structure, modelling, and molecular dynamics studies of the inhibition of protein tyrosine phosphatase 1B by sulfotyrosine peptides. 1999 , 77, 469-486	8
1190	Comparison of the structures of beta amyloid peptide (25-35) and substance P in trifluoroethanol/water solution. 1999 , 17, 381-91	24

1189	NMR spatial structure of alpha-conotoxin Iml reveals a common scaffold in snail and snake toxins recognizing neuronal nicotinic acetylcholine receptors. 1999 , 444, 275-80	52
1188	Minimal conformation of the alpha-conotoxin Iml for the alpha7 neuronal nicotinic acetylcholine receptor recognition: correlated CD, NMR and binding studies. 1999 , 454, 293-8	38
1187	Solution structure of neuromedin B by (1)H nuclear magnetic resonance spectroscopy. 1999 , 460, 263-9	17
1186	Conformational analyses by 1H NMR and computer simulations of cyclic hexapeptides related to somatostatin containing acidic and basic peptoid residues. 1999 , 53, 146-60	4
1185	1H and (13)C NMR and Molecular Dynamics Study of Chiral Recognition of Camphor Enantiomers by alpha-Cyclodextrin. 1999 , 64, 1503-1507	56
1184	Nuclear magnetic resonance and restrained molecular dynamics studies of the interaction of an epidermal growth factor-derived peptide with protein tyrosine phosphatase 1B. 1999 , 38, 5256-71	15
1183	Comparison of the solution conformations of a cell-adhesive peptide LBE and its reverse sequence EBL. 1999 , 17, 429-44	9
1182	Direct effects of phosphorylation on the preferred backbone conformation of peptides: a nuclear magnetic resonance study. 1999 , 76, 76-87	68
1181	The effect of O-fucosylation on the first EGF-like domain from human blood coagulation factor VII. 1999 , 38, 7097-110	45
1180	Structure of the antimicrobial peptide tritripticin bound to micelles: a distinct membrane-bound peptide fold. 1999 , 38, 16749-55	137
1179	NMR and molecular mechanics study of pyrethrins I and II. 1999 , 47, 3402-10	16
1178	The solution structure of oxidized Escherichia coli cytochrome b562. 1999 , 38, 8657-70	74
1177	Vulcanization of Butadiene Rubber by Means of Cyclic Disulfides. 2. A 2D Solid State HRMAS NMR Study on Cross-Link Structures in BR Vulcanizates. 1999 , 32, 7509-7520	9
1176	NMR solution structure of alpha-conotoxin Iml and comparison to other conotoxins specific for neuronal nicotinic acetylcholine receptors. 1999 , 38, 3874-82	47
1175	Kinetics and Mechanism of the Stereochemical Isomerization of an AreneRuthenium Complex of the Atropisomeric Ligand 1,1'-Biphenyl-2,2'-diamine. 1999 , 18, 5112-5119	22
1174	Structure of steroidal saponins from underground parts of Allium nutans L. 1999 , 47, 3193-6	21
1173	Biochemical characterization and nuclear magnetic resonance structure of novel alpha-conotoxins isolated from the venom of Conus consors. 1999 , 38, 6317-26	51
1172	Phenylethanoid glycosides from Globularia trichosantha. 1999 , 62, 1165-8	27

1171	NMR solution conformation of an antitoxic analogue of alpha-conotoxin GI: identification of a common nicotinic acetylcholine receptor alpha 1-subunit binding surface for small ligands and alpha-conotoxins. 1999 , 38, 11895-904	16
1170	Solution structure of a five-adenine bulge loop within a DNA duplex. 1999 , 38, 12860-8	38
1169	Solution structures of apo and holo biotinyl domains from acetyl coenzyme A carboxylase of <i>Escherichia coli</i> determined by triple-resonance nuclear magnetic resonance spectroscopy. 1999 , 38, 5045-53	67
1168	Structural and conformational requirements for high-affinity binding to the SH2 domain of Grb2(1). 1999 , 42, 971-80	79
1167	Solution structure of the N-terminal F1 module pair from human fibronectin. 1999 , 38, 8304-12	35
1166	Low-temperature-induced structural changes in the Apo regulatory domain of skeletal muscle troponin C. 1999 , 38, 5693-700	23
1165	Probing the topological arrangement of the N- and C-terminal residues of bradykinin for agonist activity at the B1 receptor. 1999 , 42, 3369-77	18
1164	Modeling Studies of the Interactions between the Insulin Receptor Kinase Domain and Protein Tyrosine Phosphatase 1B. 1999 , 121, 3579-3589	7
1163	Solution structure of a sweet protein single-chain monellin determined by nuclear magnetic resonance and dynamical simulated annealing calculations. 1999 , 38, 2340-6	32
1162	NMR structure of the complex between the zinc finger protein NCp10 of Moloney murine leukemia virus and the single-stranded pentanucleotide d(ACGCC): comparison with HIV-NCp7 complexes. 1999 , 38, 12984-94	24
1161	Structure analysis of a fusogenic peptide sequence from the sea urchin fertilization protein bindin. 1999 , 38, 2560-9	53
1160	Vulcanization of Butadiene Rubber by Means of Cyclic Disulfides. 1. A 2D NMR Study on the Cross-Link Structure of a BR Model Compound Vulcanizate. 1999 , 32, 7504-7508	9
1159	Vulcanization of Butadiene Rubber by Means of Cyclic Disulfides. 3. A 2D Solid State HRMAS NMR Study on Accelerated Sulfur Vulcanizates of BR Rubber. 1999 , 32, 7521-7529	10
1158	Surface Recognition and Helix Stabilization of a Tetraaspartate Peptide by Shape and Electrostatic Complementarity of an Artificial Receptor. 1999 , 121, 11813-11820	39
1157	Reactions of beta-carotene with cigarette smoke oxidants. Identification of carotenoid oxidation products and evaluation of the prooxidant/antioxidant effect. 1999 , 12, 535-43	72
1156	Conformation of human apolipoprotein C-I in a lipid-mimetic environment determined by CD and NMR spectroscopy. 1999 , 38, 14475-84	42
1155	Solution structure by NMR of circulin A: a macrocyclic knotted peptide having anti-HIV activity. 1999 , 285, 333-45	102
1154	Stabilization of the anticodon stem-loop of tRNA ^{Lys,3} by an A+C base-pair and by pseudouridine. 1999 , 285, 115-31	145

1153	NMR structures and activity of a novel alpha-like toxin from the scorpion <i>Leiurus quinquestriatus hebraeus</i> . 1999 , 285, 1749-63	67
1152	Conformational analysis of a set of peptides corresponding to the entire primary sequence of the N-terminal domain of the ribosomal protein L9: evidence for stable native-like secondary structure in the unfolded state. 1999 , 287, 395-407	32
1151	Structure, backbone dynamics and interactions with RNA of the C-terminal RNA-binding domain of a mouse neural RNA-binding protein, Musashi1. 1999 , 287, 315-30	43
1150	Structure and interactions with RNA of the N-terminal UUAG-specific RNA-binding domain of hnRNP D0. 1999 , 287, 221-37	42
1149	Solution structure of the transactivation domain of ATF-2 comprising a zinc finger-like subdomain and a flexible subdomain. 1999 , 287, 593-607	49
1148	Structure of the anchor-domain of myristoylated and non-myristoylated HIV-1 Nef protein. 1999 , 289, 123-38	99
1147	Effects of varying the local propensity to form secondary structure on the stability and folding kinetics of a rapid folding mixed alpha/beta protein: characterization of a truncation mutant of the N-terminal domain of the ribosomal protein L9. 1999 , 289, 167-74	25
1146	Structure-activity relationships of omega-conotoxins MVIIA, MVIIIC and 14 loop splice hybrids at N and P/Q-type calcium channels. 1999 , 289, 1405-21	73
1145	Molecular and cellular analysis of Grb2 SH3 domain mutants: interaction with Sos and dynamin. 1999 , 290, 717-30	41
1144	Structure, dynamics and hydration of the nogalamycin-d(ATGCAT) ₂ Complex determined by NMR and molecular dynamics simulations in solution. 1999 , 290, 699-716	41
1143	Solution structure of the DNA-binding domain of NtrC with three alanine substitutions. 1999 , 292, 1095-110	71
1142	MiAMP1, a novel protein from <i>Macadamia integrifolia</i> adopts a Greek key beta-barrel fold unique amongst plant antimicrobial proteins. 1999 , 293, 629-38	53
1141	BBRC Journals Cited 500 Times or More during the Period 1960-1995. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 266, 621-623	3-4
1140	Solution structure of a defensin-like peptide from platypus venom. <i>Biochemical Journal</i> , 1999 , 341, 785-794	52
1139	Solution structure of a defensin-like peptide from platypus venom. <i>Biochemical Journal</i> , 1999 , 341, 785	3.8 24
1138	A Ca ²⁺ binding cyclic peptide derived from the alpha-subunit of LFA-1: inhibitor of ICAM-1/LFA-1-mediated T-cell adhesion. 1999 , 53, 18-29	17
1137	Solution Structure of a Designed Four-Helix Bundle Maquette Scaffold. 1999 , 121, 4941-4951	75
1136	Nuclear magnetic resonance spectroscopy of peptide ion channel ligands: cloning and expression as aid to evaluation of structural and dynamic properties. 1999 , 294, 92-117	3

1135	Chapter 5 Correlations through the chemical bond I: Homonuclear shift correlation. 1999 , 19, 147-220		
1134	Determination of the solution structure of the N-domain plus linker of Antarctic eel pout antifreeze protein RD3. 1999 , 126, 387-94		10
1133	Novel Schizosaccharomyces pombe N-linked GalMan9GlcNAc isomers: role of the Golgi GMA12 galactosyltransferase in core glycan galactosylation. 1999 , 9, 497-505		22
1132	Kodaistatins, novel inhibitors of glucose-6-phosphate translocase T1 from Aspergillus terreus thom DSM 11247. Isolation and structural elucidation. 2000 , 53, 677-86		18
1131	Defensin-like peptide-2 from platypus venom: member of a class of peptides with a distinct structural fold. <i>Biochemical Journal</i> , 2000 , 348, 649	3.8	22
1130	Human T-lymphotrophic virus type I nucleocapsid protein NCp15: structural study and stability of the N-terminal zinc-finger. <i>Biochemical Journal</i> , 2000 , 352, 293	3.8	1
1129	Defensin-like peptide-2 from platypus venom: member of a class of peptides with a distinct structural fold. <i>Biochemical Journal</i> , 2000 , 348, 649-656	3.8	61
1128	Conformational analyses of sandostatin analogs containing stereochemical changes in positions 6 or 8. 2000 , 53, 506-22		7
1127	Structural basis for the biological activity of dendrotoxin-I, a potent potassium channel blocker. 2000 , 54, 44-57		20
1126	Structural characterization of a methionine-rich, emulsifying protein from sunflower seed. 2000 , 38, 341-349		24
1125	Structural and functional differences of two toxins from the scorpion Pandinus imperator. 2000 , 38, 441-9		6
1124	N-1 and N-2 positional effects in the propagation of 310-type fold in the helical model peptide Boc-(D)GluPro[Ala]Lys[Ala]Leu[Ala]OMe. 2000 , 38, 257-264		
1123	Hydrogen exchange study of canine milk lysozyme: stabilization mechanism of the molten globule. 2000 , 40, 579-89		29
1122	The physical and NMR characterizations of allyl- and crotylcelluloses. 2000 , 38, 1889-1902		18
1121	. 2000 , 2000, 597-615		5
1120	Solid State and Solution Conformation of 6-[4-[N-tert-ButoxycarbonylN-(N?-ethyl)propanamide]imidazolyl]-6-deoxycyclomaltoheptaose: Evidence of Self-Inclusion of the Boc Group within the β -Cyclodextrin Cavity. 2000 , 2000, 1065-1076		18
1119	Chemical synthesis, characterization and activity of RK-1, a novel alpha-defensin-related peptide. 2000 , 6, 19-25		14
1118	A novel acylated flavonol glycoside isolated from Brunfelsia grandiflora ssp. grandiflora. Structure elucidation by gradient accelerated NMR spectroscopy at 14T. 2000 , 11, 29-33		9

1117	Conformational studies of parathyroid hormone (PTH)/PTH-related protein (PTHrp) chimeric peptides. 2000 , 54, 429-47		17
1116	Photomodulation of conformational states. I. Mono- and bicyclic peptides with (4-amino)phenylazobenzoic acid as backbone constituent. 2000 , 54, 489-500		66
1115	Photomodulation of conformational states. II. Mono- and bicyclic peptides with (4-aminomethyl)phenylazobenzoic acid as backbone constituent. 2000 , 54, 501-14		55
1114	¶Phenylsulfonyl-N-arylacetamides (¶phenylsulfonylacetanilides): ¹ H, ¹³ C and ¹⁵ N NMR spectral characterization. 2000 , 38, 384-385		1
1113	Three-dimensional electrophoretic NMR correlation spectroscopy. <i>Journal of Magnetic Resonance</i> , 2000 , 147, 361-5	3	18
1112	The structure of the membrane-binding 38 C-terminal residues from bovine PP3 determined by liquid- and solid-state NMR spectroscopy. <i>FEBS Journal</i> , 2000 , 267, 188-99		11
1111	The structure of the linkage between the O-specific polysaccharide and the core region of the lipopolysaccharide from <i>Salmonella enterica</i> serovar Typhimurium revisited. <i>FEBS Journal</i> , 2000 , 267, 2014-27		38
1110	Structural requirements for thymosin beta4 in its contact with actin. An NMR-analysis of thymosin beta4 mutants in solution and correlation with their biological activity. <i>FEBS Journal</i> , 2000 , 267, 3530-8		26
1109	Conotoxin TVIIA, a novel peptide from the venom of <i>Conus tulipa</i> 2. Three-dimensional solution structure. <i>FEBS Journal</i> , 2000 , 267, 4649-57		12
1108	The antibiotic and anticancer active aurein peptides from the Australian Bell Frogs <i>Litoria aurea</i> and <i>Litoria raniformis</i> the solution structure of aurein 1.2. <i>FEBS Journal</i> , 2000 , 267, 5330-41		197
1107	Four steroidal alkaloids from the leaves of <i>Buxus sempervirens</i> . 2000 , 54, 951-7		19
1106	¹ H-NMR determination of the solution structure and absolute configuration of FR134043, a novel inhibitor of human leukocyte elastase. 2000 , 55, 120-8		3
1105	Three-dimensional solution structure of a disulfide bond isomer of the human insulin-like growth factor-I. 2000 , 56, 218-30		19
1104	The conformational and biological analysis of a cyclic anti-obesity peptide from the C-terminal domain of human growth hormone. 2000 , 56, 388-97		9
1103	Solution structure of a HNA-RNA hybrid. 2000 , 7, 719-31		56
1102	A novel artificial loop scaffold for the noncovalent constraint of peptides. 2000 , 7, 515-27		23
1101	Solution configuration of stereogenic hydroxide bridges in the di-¶hydroxo-tetrakis(S-arginine)dicobalt(III) complex ion. 2000 , 298, 103-106		2
1100	Preparation and characterisation of gels based on sucrose modified with glycidyl methacrylate. <i>Carbohydrate Polymers</i> , 2000 , 41, 15-24	10.3	53

1099	NMR studies on metal complexes of DNA oligomers. 2000 , 79, 353-8	10
1098	Another piece of the ribosome: solution structure of S16 and its location in the 30S subunit. 2000 , 8, 875-82	14
1097	Solution structures of two CCHC zinc fingers from the FOG family protein U-shaped that mediate protein-protein interactions. 2000 , 8, 1157-66	36
1096	Functional Correlation in Amino Acid Residue Mutations of Yeast Iso-2-Cytochrome c that is Consistent with the Prediction of the Concomitantly Variable Codon Theory in Cytochrome c Evolution. 2000 , 38, 177-196	3
1095	Backbone dynamics and solution structure refinement of the ¹⁵ N-labeled human oncogenic protein p13MTCP1: comparison with X-ray data. 2000 , 17, 215-30	20
1094	Random coil chemical shifts in acidic 8 M urea: implementation of random coil shift data in NMRView. 2000 , 18, 43-8	258
1093	Glycosylations versus conformational preferences of cancer associated mucin core. 2000 , 17, 835-48	17
1092	The acid-induced folded state of Sac7d is the native state. 2000 , 9, 1878-88	20
1091	The structure and dynamics in solution of Cu(I) pseudoazurin from <i>Paracoccus pantotrophus</i> . 2000 , 9, 846-58	20
1090	Synthesis and NMR solution structure of an alpha-helical hairpin stapled with two disulfide bridges. 2000 , 9, 942-55	29
1089	NMR solution structure of <i>Apis mellifera</i> chymotrypsin/cathepsin G inhibitor-1 (AMCI-1): structural similarity with <i>Ascaris</i> protease inhibitors. 2000 , 9, 976-84	38
1088	NMR structure of a concatemer of the first and second ligand-binding modules of the human low-density lipoprotein receptor. 2000 , 9, 1282-93	38
1087	The active site and substrates binding mode of malonyl-CoA synthetase determined by transferred nuclear Overhauser effect spectroscopy, site-directed mutagenesis, and comparative modeling studies. 2000 , 9, 1294-303	14
1086	Recombinant decorsin: dynamics of the RGD recognition site. 2000 , 9, 1428-38	15
1085	Structural studies of a baboon (<i>Papio</i> sp.) plasma protein inhibitor of cholesteryl ester transferase. 2000 , 9, 1548-58	5
1084	Structure of a (Cys3His) zinc ribbon, a ubiquitous motif in archaeal and eucaryal transcription. 2000 , 9, 1743-52	53
1083	Redox-related conformational changes in <i>Rhodobacter capsulatus</i> cytochrome c2. 2000 , 9, 1828-37	9
1082	Structure of the central core domain of TFIIIEbeta with a novel double-stranded DNA-binding surface. <i>EMBO Journal</i> , 2000 , 19, 1346-56	13 51

1081	The binding of a glycoprotein 120 V3 loop peptide to HIV-1 neutralizing antibodies. Structural implications. <i>Journal of Biological Chemistry</i> , 2000 , 275, 36645-52	5-4	19
1080	A structural requirement for activation of skeletal ryanodine receptors by peptides of the dihydropyridine receptor II-III loop. <i>Journal of Biological Chemistry</i> , 2000 , 275, 11631-7	5-4	42
1079	Nuclear magnetic resonance solution conformation of alpha-conotoxin AulB, an alpha(3)beta(4) subtype-selective neuronal nicotinic acetylcholine receptor antagonist. <i>Journal of Biological Chemistry</i> , 2000 , 275, 8680-5	5-4	27
1078	The accumulation of Man(6)GlcNAc(2)-PP-dolichol in the <i>Saccharomyces cerevisiae</i> Deltaalg9 mutant reveals a regulatory role for the Alg3p alpha1,3-Man middle-arm addition in downstream oligosaccharide-lipid and glycoprotein glycan processing. <i>Journal of Biological Chemistry</i> , 2000 , 275, 4267-77	5-4	22
1077	Selected mutations in a mesophilic cytochrome c confer the stability of a thermophilic counterpart. <i>Journal of Biological Chemistry</i> , 2000 , 275, 37824-8	5-4	46
1076	Preparation of oligomeric beta-glycosides from cellulose and hemicellulosic polysaccharides via the glycosyl transferase activity of a <i>trichoderma reesei</i> cellulase. 2000 , 10, 193-201		15
1075	Molecular organization of the alkali-insoluble fraction of <i>Aspergillus fumigatus</i> cell wall. <i>Journal of Biological Chemistry</i> , 2000 , 275, 27594-607	5-4	274
1074	Synthesis and properties of O-beta-D-ribofuranosyl-(1''-2'')-adenosine-5''-O-phosphate and its derivatives. 2000 , 19, 1847-59		12
1073	Kinetic Element Effect for Atropisomerization of an Organometallic Complex of the Misdirected Ligand 1,1-Biisoquinoline 2000 , 19, 547-552		29
1072	Anaerobic chlorophyll isocyclic ring formation in <i>Rhodobacter capsulatus</i> requires a cobalamin cofactor. 2000 , 97, 6908-13		97
1071	Acyclic permutants of naturally occurring cyclic proteins. Characterization of cystine knot and beta-sheet formation in the macrocyclic polypeptide kalata B1. <i>Journal of Biological Chemistry</i> , 2000 , 275, 19068-75	5-4	88
1070	Solution structure of an N-capping peptide from the N-terminal leucine-repeat region of hepatitis delta antigen. 2000 , 377, 219-27		1
1069	NMR solution structure of butantoxin. 2000 , 379, 18-27		28
1068	NMR studies on novel antitumor drug candidates, deoxoartemisinin and carboxypropyldeoxoartemisinin. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 274, 359-63	4	29
1067	Solution structure of a designed amphipathic antimicrobial synthetic peptide, PGAa. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 276, 1278-85	3-4	2
1066	Structural comparison of the PhoB and OmpR DNA-binding/transactivation domains and the arrangement of PhoB molecules on the phosphate box. 2000 , 295, 1225-36		83
1065	NMR structure of activated CheY. 2000 , 297, 543-51		119
1064	Solution structure of hanatoxin1, a gating modifier of voltage-dependent K(+) channels: common surface features of gating modifier toxins. 2000 , 297, 771-80		133

1063	The solution structure of Rhodobacter sphaeroides LH1beta reveals two helical domains separated by a more flexible region: structural consequences for the LH1 complex. 2000 , 298, 83-94	49
1062	Structural basis for the network of functional cooperativities in cytochrome c(3) from Desulfovibrio gigas: solution structures of the oxidised and reduced states. 2000 , 298, 61-82	66
1061	Molten globule structure of equine beta-lactoglobulin probed by hydrogen exchange. 2000 , 299, 757-70	51
1060	Determination of a high precision structure of a novel protein, Linum usitatissimum trypsin inhibitor (LUTI), using computer-aided assignment of NOESY cross-peaks. 2000 , 302, 1179-92	24
1059	NMR structure of oxidized glutaredoxin 3 from Escherichia coli. 2000 , 303, 423-32	39
1058	Solution structure of the module X2 1 of unknown function of the cellulosomal scaffolding protein CipC of Clostridium cellulolyticum. 2000 , 304, 201-17	40
1057	Structure-activity relationships in a peptidic alpha7 nicotinic acetylcholine receptor antagonist. 2000 , 304, 911-26	22
1056	Salt effects on ionization equilibria of histidines in myoglobin. 2000 , 79, 1637-54	89
1055	Solution conformations of a trimannoside from nuclear magnetic resonance and molecular dynamics simulations. 2000 , 79, 3313-29	40
1054	An advantage for use of isotope labeling and NMR chemical shifts to analyze the structure of four homologous IgG-binding domains of staphylococcal protein A. 2000 , 42, 35-47	11
1053	Structure of a biologically active fragment of human serum apolipoprotein C-II in the presence of sodium dodecyl sulfate and dodecylphosphocholine. 2000 , 1486, 253-64	14
1052	Solution structure of the NADP(H)-binding component (dIII) of proton-translocating transhydrogenase from Rhodospirillum rubrum. 2000 , 1459, 248-57	26
1051	NMR structure of antibiotics plipastatins A and B from Bacillus subtilis inhibitors of phospholipase A(2). 2000 , 485, 76-80	60
1050	NMR structure of the channel-former zervamicin IIB in isotropic solvents. 2000 , 466, 333-6	42
1049	Structural analysis of multifunctional peptide motifs in human bifunctional tRNA synthetase: identification of RNA-binding residues and functional implications for tandem repeats. 2000 , 39, 15775-82	51
1048	Location of cyanine-3 on double-stranded DNA: importance for fluorescence resonance energy transfer studies. 2000 , 39, 6317-24	214
1047	Three-dimensional structure of RK-1: a novel alpha-defensin peptide. 2000 , 39, 15757-64	31
1046	Synthesis and Characterization of a Chiral, Aza-15-Crown-5-Functionalized Ferrocenyldiphosphine Ligand for Asymmetric Catalysis. 2000 , 19, 994-1002	29

1045	Two novel Diels-Alder adducts from Hippocratea celastroides roots and their insecticidal activity. 2000 , 78, 248-254	14
1044	Solution structure of O-glycosylated C-terminal leucine zipper domain of human salivary mucin (MUC7). 2000 , 18, 145-54	3
1043	Structure and stability effects of the mutation of glycine 34 to serine in Rhodobacter capsulatus cytochrome c(2). 2000 , 39, 4053-61	6
1042	Solution structure and activity of the synthetic four-disulfide bond Mediterranean mussel defensin (MGD-1). 2000 , 39, 14436-47	93
1041	Ring opening of benzo[a]pyrene in the germ-free rat is a novel pathway for formation of potentially genotoxic metabolites. 2000 , 39, 15585-91	16
1040	Solution structures of the N-terminal domain of yeast calmodulin: Ca ²⁺ -dependent conformational change and its functional implication. 2000 , 39, 13660-8	24
1039	Three-dimensional solution structure of omega-conotoxin TxVII, an L-type calcium channel blocker. 2000 , 39, 14761-7	30
1038	Self-assembly of polar functionalities using noncovalent platforms. 2000 , 2, 4121-4	18
1037	Insensitivity of perturbed carboxyl pK(a) values in the ovomucoid third domain to charge replacement at a neighboring residue. 2000 , 39, 8067-72	39
1036	Structure of the Bovine Antimicrobial Peptide Indolicidin Bound to Dodecylphosphocholine and Sodium Dodecyl Sulfate Micelles. 2000 , 39, 15765-15774	247
1035	Nuclear magnetic resonance studies of the N-terminal fragment of adenosine diphosphate ribosylation factor 1 in micelles and bicelles: influence of N-myristoylation. 2000 , 39, 3804-16	33
1034	Determination of biophysical parameters of polypeptide retro-inverso isomers and their analogues by capillary electrophoresis. 2000 , 72, 1964-72	29
1033	Recognition properties of V3-specific antibodies to V3 loop peptides derived from HIV-1 gp120 presented in multiple conformations. 2000 , 39, 10866-76	6
1032	A rational approach to the design and synthesis of a new bradykinin B(1) receptor antagonist. 2000 , 43, 2387-94	21
1031	Solution structure by NMR and molecular dynamics of a duplex containing a guanine opposite a N-(2-deoxy-beta-D-erythro-pentofuranosyl)formamide lesion. 2000 , 39, 5614-21	11
1030	Solution Conformations of Helix-Forming α -Amino Acid Homooligomers. 2000 , 122, 2711-2718	111
1029	Cyclohexene Nucleic Acids (CeNA): Serum Stable Oligonucleotides that Activate RNase H and Increase Duplex Stability with Complementary RNA. 2000 , 122, 8595-8602	117
1028	NMR Study of the Interaction of Monomeric and Polymeric Chiral Surfactants with (R)- and (S)-1,1'-Binaphthyl-2,2'-diyl Hydrogen Phosphate. 2000 , 16, 3022-3029	25

1027	Role of phosphorylation in the conformation of tau peptides implicated in Alzheimer's disease. 2000 , 39, 9039-46	68
1026	Opiate aromatic pharmacophore structure-activity relationships in CTAP analogues determined by topographical bias, two-dimensional NMR, and biological activity assays. 2000 , 43, 569-80	21
1025	Identification of initiation sites for T4 lysozyme folding using CD and NMR spectroscopy of peptide fragments. 2000 , 39, 5911-20	17
1024	Three-dimensional solution structure of oryzacystatin-I, a cysteine proteinase inhibitor of the rice, <i>Oryza sativa</i> L. japonica. 2000 , 39, 14753-60	97
1023	Solution structure of monomeric peptide YY supports the functional significance of the PP-fold. 2000 , 39, 9935-42	39
1022	Dual structural requirements for multilineage hematopoietic-suppressive activity of chemokine-derived peptides. 2000 , 39, 9612-22	5
1021	Synthesis and physical properties of protein core mimetics. 2001 , 66, 8328-35	11
1020	Solution structure of the satiety factor, CART, reveals new functionality of a well-known fold. 2001 , 40, 9082-8	28
1019	Limonoids from Nigerian <i>Harrisonia abyssinica</i> and their stimulatory activity against <i>Striga hermonthica</i> seeds. 2001 , 64, 1434-8	20
1018	Solution structure of the dimeric cytoplasmic domain of syndecan-4. 2001 , 40, 8471-8	42
1017	NMR structure of a 2',5' RNA favors A type duplex with compact C2'endo nucleotide repeat. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 283, 537-43	3-4 20
1016	Structure and dynamics of micelle-bound neuropeptide Y: comparison with unligated NPY and implications for receptor selection. 2001 , 305, 307-29	122
1015	The solution structure of C1-T1, a two-domain proteinase inhibitor derived from a circular precursor protein from <i>Nicotiana glauca</i> . 2001 , 306, 69-79	18
1014	Solution structure of an A-tract DNA bend. 2001 , 306, 1081-98	133
1013	Structure of the C-terminal RNA-binding domain of hnRNP D0 (AUF1), its interactions with RNA and DNA, and change in backbone dynamics upon complex formation with DNA. 2001 , 311, 973-88	35
1012	NMR structure and backbone dynamics of a concatemer of epidermal growth factor homology modules of the human low-density lipoprotein receptor. 2001 , 311, 341-56	29
1011	Solution structures by 1H NMR of the novel cyclic trypsin inhibitor SFTI-1 from sunflower seeds and an acyclic permutant. 2001 , 311, 579-91	195
1010	NMR structure of the hRap1 Myb motif reveals a canonical three-helix bundle lacking the positive surface charge typical of Myb DNA-binding domains. 2001 , 312, 167-75	42

1009	An intramolecular quadruplex of (GGA)(4) triplet repeat DNA with a G:G:G:G tetrad and a G(:A):G(:A):G(:A):G heptad, and its dimeric interaction. 2001 , 313, 255-69	127
1008	NMR structure of the <i>Euplotes raikovi</i> pheromone Er-23 and identification of its five disulfide bonds. 2001 , 313, 923-31	32
1007	Three-dimensional structure of RTD-1, a cyclic antimicrobial defensin from Rhesus macaque leukocytes. 2001 , 40, 4211-21	145
1006	Parallel sheet secondary structure in gamma-peptides. 2001 , 123, 11077-8	86
1005	Dynamics of membrane penetration of the fluorescent 7-nitrobenz-2-oxa-1,3-diazol-4-yl (NBD) group attached to an acyl chain of phosphatidylcholine. 2001 , 80, 822-31	101
1004	Structural determinants for activation or inhibition of ryanodine receptors by basic residues in the dihydropyridine receptor II-III loop. 2001 , 80, 2715-26	28
1003	Solution NMR structure of a model class A (apolipoprotein) amphipathic alpha helical peptide. 2001 , 22, 567-73	4
1002	Solution conformation of an RNA-DNA hybrid duplex containing a pyrimidine RNA strand and a purine DNA strand. 2001 , 28, 273-84	26
1001	Structure of an RNA hairpin from HRV-14. 2001 , 40, 8055-64	16
1000	Re-face stereospecificity of NADP dependent methylenetetrahydromethanopterin dehydrogenase from <i>Methylobacterium extorquens</i> AM1 as determined by NMR spectroscopy. 2001 , 494, 95-8	5
999	Structural analysis of emerin, an inner nuclear membrane protein mutated in X-linked Emery-Dreifuss muscular dystrophy. 2001 , 501, 171-6	58
998	Three-dimensional solution structure and stability of thioredoxin m from spinach. 2001 , 40, 15246-56	12
997	Dynamic characterization of the water binding loop in the P-type cardiotoxin: implication for the role of the bound water molecule. 2001 , 40, 12782-94	18
996	Aib-rich peptides containing lactam-bridged side chains as models of the 3(10)-helix. 2001 , 123, 2743-51	31
995	Design, synthesis and pharmacological characterization of a potent radioiodinated and photoactivatable peptidic oxytocin antagonist. 2001 , 44, 3022-30	7
994	NMR solution studies of hamster galectin-3 and electron microscopic visualization of surface-adsorbed complexes: evidence for interactions between the N- and C-terminal domains. 2001 , 40, 4859-66	51
993	Solution structure of BSTI: a new trypsin inhibitor from skin secretions of <i>Bombina bombina</i> . 2001 , 40, 4601-9	19
992	Solution structure of the carboxyl-terminal cysteine-rich domain of the VHv1.1 polydnviral gene product: comparison with other cystine knot structural folds. 2001 , 40, 14404-12	15

991	Symmetrical stabilization of bound Ca ²⁺ ions in a cooperative pair of EF-hands through hydrogen bonding of coordinating water molecules in calbindin D(9k). 2001 , 40, 9887-95		8
990	Solution structure of an immunoactive peptide fragment of Staphylococcal protein-A. 2001 , 18, 773-81		4
989	Solution structure of 2-(pyrido[1,2-e]purin-4-yl)amino-ethanol intercalated in the DNA duplex d(CGATCG) ₂ . 2001 , 40, 8717-26		26
988	Solution structure of (+)-discodermolide. 2001 , 3, 695-8		51
987	Solution structure and backbone dynamics of the human DNA ligase IIIalpha BRCT domain. 2001 , 40, 13158-66		39
986	NMR structure of bucandin, a neurotoxin from the venom of the Malayan krait (<i>Bungarus candidus</i>). <i>Biochemical Journal</i> , 2001 , 360, 539-548	3.8	28
985	Polyproline II helix is a key structural motif of the elastic PEVK segment of titin. 2001 , 40, 3427-38		128
984	High-resolution solution structure of human intestinal trefoil factor and functional insights from detailed structural comparisons with the other members of the trefoil family of mammalian cell motility factors. 2001 , 40, 9552-9		23
983	NMR structure of bucandin, a neurotoxin from the venom of the Malayan krait (<i>Bungarus candidus</i>). <i>Biochemical Journal</i> , 2001 , 360, 539-48	3.8	14
982	The chemistry of oligosaccharide ligands of selectins: significance for the development of new immunomodulatory medicines. 2001 , 57, 207-435		24
981	Solution structure, dimerization, and dynamics of a lipophilic alpha/3(10)-helical, C alpha-methylated peptide. Implications for folding of membrane proteins. 2001 , 123, 6678-86		31
980	Conformational analysis of the nonapeptide leuprorelin using NMR and molecular modeling. 2001 , 8, 77-87		1
979	Solution structure of human immunodeficiency virus type 1 Vpr(13-33) peptide in micelles. <i>FEBS Journal</i> , 2001 , 268, 389-95		19
978	Impact of C5-cytosine methylation on the solution structure of d(GAAAACGTTTTC) ₂ . An NMR and molecular modelling investigation. <i>FEBS Journal</i> , 1999 , 265, 1032-42		19
977	Isolation and structural characterization of an R-form lipopolysaccharide from <i>Yersinia enterocolitica</i> serotype O:8. <i>FEBS Journal</i> , 2001 , 268, 554-64		41
976	Three-dimensional structure of the histidine-containing phosphocarrier protein (HPr) from <i>Enterococcus faecalis</i> in solution. <i>FEBS Journal</i> , 2001 , 268, 635-44		18
975	Solution structure of cytochrome b5 mutant (E44/48/56A/D60A) and its interaction with cytochrome c. <i>FEBS Journal</i> , 2001 , 268, 1620-1630		11
974	Solution structure of microcin J25, the single macrocyclic antimicrobial peptide from <i>Escherichia coli</i> . <i>FEBS Journal</i> , 2001 , 268, 2124-33		33

973	Chemosensory protein from the moth <i>Mamestra brassicae</i> . Expression and secondary structure from ¹ H and ¹⁵ N NMR. <i>FEBS Journal</i> , 2001 , 268, 4731-9	27
972	Synthesis of linear-B saccharopeptides via enzymatic galactosylation of non-natural glucosamide acceptors. 2001 , 9, 1857-69	5
971	A novel mechanism of antibiotic resistance: study of the complex state of peptides with bacterial <i>Staphylococcus aureus</i> ribosomes. 2001 , 4, 745-750	
970	NMR and CD conformational studies of the C-terminal 16-peptides of <i>Pseudomonas aeruginosa</i> c551 and <i>Hydrogenobacter thermophilus</i> c552 cytochromes. 2001 , 57, 39-47	12
969	Solution structures of the cytoplasmic linkers between segments S4 and S5 (S4-S5) in domains III and IV of human brain sodium channels in SDS micelles. 2001 , 58, 193-203	8
968	Solution structures of the N-terminal domain of histone H4. 2001 , 58, 389-98	6
967	Long-range substituent and temperature effect on prototropic tautomerism in 2-(acylmethyl)quinolines. 2001 , 14, 201-204	11
966	Structure of rat parvalbumin with deleted AB domain: implications for the evolution of EF hand calcium-binding proteins and possible physiological relevance. 2001 , 45, 117-28	13
965	Solution structure of the cytoplasmic linker between domain III-S6 and domain IV-S1 (III-IV linker) of the rat brain sodium channel in SDS micelles. 2001 , 59, 380-93	12
964	Stabilization of an alpha-helical conformation in an isolated hexapeptide inhibitor of calmodulin. 2001 , 59, 467-76	17
963	Synthesis and Coordination Chemistry of Chelate Ligands Containing Cyclopentadienyl, Indenyl and Fluorenyl Donors Diastereoselectivity and NMR Structure Analysis. 2001 , 2001, 949-971	11
962	The 2-naphthylmethyl (NAP) group in carbohydrate synthesis: first total synthesis of the GlyCAM-1 oligosaccharide structures. 2001 , 7, 356-67	35
961	Structural consequences of metal complexation of cyclo[Pro-Phe-Phe-Ala-Xaa] ₂ decapeptides. 2001 , 7, 1176-83	9
960	Solid state and solution conformation of [Ala7]-phalloidin: a synthetic phallotoxin analogue. 2001 , 7, 1479-85	23
959	Phalloidin synthetic analogues: structural requirements in the interaction with F-actin. 2001 , 7, 4665-73	18
958	Alpha-homo-DNA and RNA form a parallel oriented non-A, non-B-type double helical structure. 2001 , 7, 5183-94	16
957	Purification and characterisation of a galactoglucomannan from kiwifruit (<i>Actinidia deliciosa</i>). 2001 , 331, 291-306	81
956	Alfalfa (<i>Medicago sativa</i> L.) flavonoids. 1. Apigenin and luteolin glycosides from aerial parts. 2001 , 49, 753-8	99

955	Dipolar filtered ¹ H- ¹³ C heteronuclear correlation spectroscopy for resonance assignment of proteins. 2001 , 20, 263-74		16
954	Structure and Conformational Heterogeneity of a Weak Toxin from the Cobra <i>Naja kaouthia</i> Venom. 2001 , 27, 72-83		4
953	Structure of the <i>Bacillus subtilis</i> peptide antibiotic subtilisin A determined by ¹ H-NMR and matrix assisted laser desorption/ionization time-of-flight mass spectrometry. 2001 , 20, 501-6		52
952	Inhibition of <i>Helicobacter pylori</i> adherence by a peptide derived from neuraminyl lactose binding adhesin. 2001 , 228, 83-9		7
951	Variability in automated assignment of NOESY spectra and three-dimensional structure determination: a test case on three small disulfide-bonded proteins. 2001 , 19, 49-62		24
950	A novel family of RNA tetraloop structure forms the recognition site for <i>Saccharomyces cerevisiae</i> RNase III. <i>EMBO Journal</i> , 2001 , 20, 7240-9	13	71
949	The extracellular human melanoma inhibitory activity (MIA) protein adopts an SH3 domain-like fold. <i>EMBO Journal</i> , 2001 , 20, 340-9	13	62
948	The hairpin structure of the (6)F1(1)F2(2)F2 fragment from human fibronectin enhances gelatin binding. <i>EMBO Journal</i> , 2001 , 20, 1519-29	13	66
947	Isolation and NMR characterization of rosacelose, a novel sulfated polysaccharide from the sponge <i>Mixylla rosacea</i> . 2001 , 334, 39-47		21
946	Solution structures of the inactivation gate particle peptides of rat brain type-IIA and human heart sodium channels in SDS micelles. 2001 , 57, 203-14		7
945	The Val-210-Ile pathogenic Creutzfeldt-Jakob disease mutation increases both the helical and aggregation propensities of a sequence corresponding to helix-3 of PrP(C). 2001 , 1544, 242-54		10
944	The hydrophobin EAS is largely unstructured in solution and functions by forming amyloid-like structures. 2001 , 9, 83-91		126
943	Structural characterization of the LEM motif common to three human inner nuclear membrane proteins. 2001 , 9, 503-11		97
942	Conformational variation of the central CG site in d(ATGACGTCAT) ₂ and d(GAAAACGTTTTTC) ₂ . An NMR, molecular modelling and 3D-homology investigation. <i>FEBS Journal</i> , 1999 , 261, 722-33		7
941	Role of the substrate conformation and of the S1 protein in the cleavage efficiency of the T4 endoribonuclease RegB. <i>Journal of Biological Chemistry</i> , 2001 , 276, 13264-72	5-4	21
940	The yeast ALG11 gene specifies addition of the terminal alpha 1,2-Man to the Man5GlcNAc2-PP-dolichol N-glycosylation intermediate formed on the cytosolic side of the endoplasmic reticulum. <i>Journal of Biological Chemistry</i> , 2001 , 276, 21828-40	5-4	53
939	Structure and activity of the insect cytokine growth-blocking peptide. Essential regions for mitogenic and hemocyte-stimulating activities are separate. <i>Journal of Biological Chemistry</i> , 2001 , 276, 31813-8	5-4	36
938	NMR structure of the "ball-and-chain" domain of KCNB2, the beta 2-subunit of large conductance Ca ²⁺ - and voltage-activated potassium channels. <i>Journal of Biological Chemistry</i> , 2001 , 276, 42116-21	5-4	55

937	Biosynthesis of 1,4,5,6-Tetrahydro-2-Methyl-4-Pyrimidinecarboxylic Acid (Ectoine) from <i>Marinococcus halophilus</i> into Tobacco. 2001 , 2, 15-42		2
936	Structural properties of carnation mottle virus p7 movement protein and its RNA-binding domain. <i>Journal of Biological Chemistry</i> , 2001 , 276, 18122-9	5-4	27
935	Solution structure of the N-terminal domain of the human TFIIH MAT1 subunit: new insights into the RING finger family. <i>Journal of Biological Chemistry</i> , 2001 , 276, 7457-64	5-4	39
934	A novel endo-beta-galactosidase from <i>Clostridium perfringens</i> that liberates the disaccharide GlcNAc α 1 \rightarrow Gal from glycans specifically expressed in the gastric gland mucous cell-type mucin. <i>Journal of Biological Chemistry</i> , 2001 , 276, 28226-32	5-4	19
933	Structure and mechanism of action of an indolicidin peptide derivative with improved activity against gram-positive bacteria. <i>Journal of Biological Chemistry</i> , 2001 , 276, 24015-22	5-4	162
932	Structure and properties of the exopolysaccharide produced by <i>Streptococcus macedonicus</i> Sc136. 2001 , 11, 131-9		56
931	Solution structure of a Kunitz-type chymotrypsin inhibitor isolated from the elapid snake <i>Bungarus fasciatus</i> . <i>Journal of Biological Chemistry</i> , 2001 , 276, 45079-87	5-4	52
930	An evolutionarily conserved motif in the TAB1 C-terminal region is necessary for interaction with and activation of TAK1 MAPKKK. <i>Journal of Biological Chemistry</i> , 2001 , 276, 24396-400	5-4	53
929	RNase H mediated cleavage of RNA by cyclohexene nucleic acid (CeNA). 2001 , 29, 4941-7		46
928	Multiple four-stranded conformations of human telomere sequence d(CCCTAA) in solution. 2001 , 29, 831-40		27
927	A novel ganglioside isolated from renal cell carcinoma. <i>Journal of Biological Chemistry</i> , 2001 , 276, 16695-703		46
926	Solution structure, backbone dynamics, and stability of a double mutant single-chain monellin. structural origin of sweetness. <i>Journal of Biological Chemistry</i> , 2001 , 276, 19624-30	5-4	31
925	DNA-XPA interactions: a (31)P NMR and molecular modeling study of dCCAATAACC association with the minimal DNA-binding domain (M98-F219) of the nucleotide excision repair protein XPA. 2001 , 29, 2635-43		22
924	An amino acid region at the N-terminus of rat hepatoma alpha1 \rightarrow 2 fucosyltransferase modulates enzyme activity and interaction with lipids: strong preference for glycosphingolipids containing terminal Galbeta1 \rightarrow 3GalNAc-structures. 2001 , 40, 5708-19		1
923	Lipid induced conformation of the tachykinin peptide Kassinin. 2001 , 18, 611-21, 623-5		18
922	Solution structure of p21(Waf1/Cip1/Sdi1) C-terminal domain bound to Cdk4. 2001 , 19, 419-27		4
921	The NMR-derived conformation of neuropeptide F from <i>Moniezia expansa</i> . 2002 , 19, 991-8		4
920	Lipopolysaccharides possessing two L-glycero-D-manno-heptopyranosyl-alpha-(1 \rightarrow 5)-3-deoxy-D-manno-oct-2-ulopyranosonic acid moieties in the core region. The structure of the core region of the lipopolysaccharides from <i>Burkholderia caryophylli</i> . <i>Journal of Biological Chemistry</i> , 2002 , 277, 10058-63	5-4	33

919	Structural and ICAM-1-docking properties of a cyclic peptide from the I-domain of LFA-1: an inhibitor of ICAM-1/LFA-1-mediated T-cell adhesion. 2002 , 19, 789-99		13
918	pH-dependent intramolecular binding and structure involving Cx43 cytoplasmic domains. <i>Journal of Biological Chemistry</i> , 2002 , 277, 36706-14	5.4	145
917	A new level of conotoxin diversity, a non-native disulfide bond connectivity in alpha-conotoxin AulB reduces structural definition but increases biological activity. <i>Journal of Biological Chemistry</i> , 2002 , 277, 48849-57	5.4	97
916	Solution structure of mu-conotoxin PIIIA, a preferential inhibitor of persistent tetrodotoxin-sensitive sodium channels. <i>Journal of Biological Chemistry</i> , 2002 , 277, 27247-55	5.4	66
915	Type I shorthorn sculpin antifreeze protein: recombinant synthesis, solution conformation, and ice growth inhibition studies. <i>Journal of Biological Chemistry</i> , 2002 , 277, 24073-80	5.4	25
914	Involvement of the second extracellular loop (E2) of the neurokinin-1 receptor in the binding of substance P. Photoaffinity labeling and modeling studies. <i>Journal of Biological Chemistry</i> , 2002 , 277, 22386-94	5.4	21
913	Structural characterizations of fusion peptide analogs of influenza virus hemagglutinin. Implication of the necessity of a helix-hinge-helix motif in fusion activity. <i>Journal of Biological Chemistry</i> , 2002 , 277, 22725-33	5.4	39
912	Solution structure of a SRP19 binding domain in human SRP RNA. 2002 , 132, 177-82		10
911	A comparison of the properties and the solution structure for RNA and DNA quadruplexes which contain two GGAGG sequences joined with a tetranucleotide linker. 2002 , 21, 785-801		13
910	NMR structural analysis of alpha-bungarotoxin and its complex with the principal alpha-neurotoxin-binding sequence on the alpha 7 subunit of a neuronal nicotinic acetylcholine receptor. <i>Journal of Biological Chemistry</i> , 2002 , 277, 12406-17	5.4	56
909	New binding site on common molecular scaffold provides HERG channel specificity of scorpion toxin BeKm-1. <i>Journal of Biological Chemistry</i> , 2002 , 277, 43104-9	5.4	51
908	Secondary structure of the third extracellular loop responsible for ligand selectivity of a mammalian gonadotropin-releasing hormone receptor. 2002 , 45, 1026-34		20
907	Structure of a novel P-superfamily spasmodic conotoxin reveals an inhibitory cystine knot motif. <i>Journal of Biological Chemistry</i> , 2002 , 277, 43033-40	5.4	32
906	Three-dimensional solution structure of the sodium channel agonist/antagonist delta-conotoxin TxVIA. <i>Journal of Biological Chemistry</i> , 2002 , 277, 36387-91	5.4	26
905	Solution structure and topology of the N-terminal membrane anchor domain of a microsomal cytochrome P450: prostaglandin I2 synthase. <i>Biochemical Journal</i> , 2002 , 368, 721-8	3.8	23
904	Structure of the antimicrobial peptide tachystatin A. <i>Journal of Biological Chemistry</i> , 2002 , 277, 23651-7	5.4	34
903	The three-dimensional structure of the gallium complex of azoverdin, a siderophore of <i>Azomonas macrocytogenes</i> ATCC 12334, determined by NMR using residual dipolar coupling constants. 2002 , 41, 12488-97		14
902	(1Z,3Z)-1,4-Di(pyridin-2-yl)buta-1,3-diene-2,3-diol: the planar highly conjugated symmetrical enediol with multiple intramolecular hydrogen bonds. 2002 , 67, 3339-45		29

901	Truncated hemoglobin from the cyanobacterium <i>Synechococcus</i> sp. PCC 7002: evidence for hexacoordination and covalent adduct formation in the ferric recombinant protein. 2002 , 41, 6902-10	62
900	Solution structure of ascidian trypsin inhibitor determined by nuclear magnetic resonance spectroscopy. 2002 , 41, 10657-64	10
899	Designing analogues of mini atrial natriuretic peptide based on structural analysis by NMR and restrained molecular dynamics. 2002 , 45, 881-7	3
898	Effect of glycosylation on MUC1 humoral immune recognition: NMR studies of MUC1 glycopeptide-antibody interactions. 2002 , 41, 9946-61	50
897	A Simple and Reliable Method for the Determination and Localization of Chitin in Abalone Nacre. 2002 , 14, 3252-3259	41
896	Stabilization of protein by replacement of a fluctuating loop: structural analysis of a chimera of bovine alpha-lactalbumin and equine lysozyme. 2002 , 41, 13807-13	12
895	Solution structure of a novel chromoprotein derived from apo-neocarzinostatin and a synthetic chromophore. 2002 , 41, 11731-9	35
894	Overexpression and structural study of the cathelicidin motif of the protegrin-3 precursor. 2002 , 41, 21-30	16
893	Spectroscopic Studies of Cyclodextrin Complexes with 2,5-Bis-(4-Methylphenyl)Oxazole. 2002 , 56, 1048-1054	2
892	Triterpenoid saponins from Caryophyllaceae family. 2002 , 26, 3-61	16
891	Isolation and quantitative analysis of phenolic antioxidants, free sugars, and polyols from mango (<i>Mangifera indica</i> L.) stem bark aqueous decoction used in Cuba as a nutritional supplement. 2002 , 50, 762-6	199
890	Antibiotic resistance peptides: interaction of peptides conferring macrolide and ketolide resistance with <i>Staphylococcus aureus</i> ribosomes: conformation of bound peptides as determined by transferred NOE experiments. 2002 , 41, 4218-29	22
889	Antagonistic effects of human cyclic MBP(87-99) altered peptide ligands in experimental allergic encephalomyelitis and human T-cell proliferation. 2002 , 45, 275-83	66
888	Solution structures of the cis- and trans-Pro30 isomers of a novel 38-residue toxin from the venom of <i>Hadronyche Infensa</i> sp. that contains a cystine-knot motif within its four disulfide bonds. 2002 , 41, 3294-301	36
887	Association of GM4 ganglioside with the membrane surrounding lipid droplets in shark liver. <i>Journal of Lipid Research</i> , 2002 , 43, 1019-25	6.3 15
886	Solution structure of betacellulin, a new member of EGF-family ligands. <i>Biochemical and Biophysical Research Communications</i> , 2002 , 294, 1040-6	3-4 17
885	Solution structure of moricin, an antibacterial peptide, isolated from the silkworm <i>Bombyx mori</i> . 2002 , 518, 33-8	34
884	D-Amino acid residue in the C-type natriuretic peptide from the venom of the mammal, <i>Ornithorhynchus anatinus</i> , the Australian platypus. 2002 , 524, 172-6	67

883	Protection against experimental malaria associated with AMA-1 peptide analogue structures. 2002 , 527, 95-100	29
882	Conformations of platypus venom C-type natriuretic peptide in aqueous solution and sodium dodecyl sulfate micelles. 2002 , 40, 711-9	24
881	¹ H-NMR structures of the Plasmodium falciparum 1758 erythrocyte binding peptide analogues and protection against malaria. 2002 , 71, 2773-85	19
880	Solution structure of omega-grammotoxin SIA, a gating modifier of P/Q and N-type Ca(2+) channel. 2002 , 321, 517-26	45
879	NMR structures of two variants of bovine pancreatic trypsin inhibitor (BPTI) reveal unexpected influence of mutations on protein structure and stability. 2002 , 321, 647-58	10
878	NMR structures of 36 and 73-residue fragments of the calreticulin P-domain. 2002 , 322, 773-84	49
877	A dimeric RNA quadruplex architecture comprised of two G:G(:A):G:G(:A) hexads, G:G:G:G tetrads and UUUU loops. 2002 , 322, 955-70	66
876	The solution structure of the recombinant hemoglobin from the cyanobacterium Synechocystis sp. PCC 6803 in its hemichrome state. 2002 , 324, 1015-29	53
875	Spatial structure of zervamicin IIB bound to DPC micelles: implications for voltage-gating. 2002 , 82, 762-71	40
874	A new zinc binding fold underlines the versatility of zinc binding modules in protein evolution. 2002 , 10, 639-48	19
873	Solution structure of paralytic peptide of silkworm, Bombyx mori. 2002 , 23, 2111-6	23
872	Activity and NMR structure of synthetic peptides of the bovine ATPase inhibitor protein, IF1. 2002 , 23, 2127-41	7
871	Structure-function studies of analogues of parathyroid hormone (PTH)-1-34 containing beta-amino acid residues in positions 11-13. 2002 , 41, 8162-75	17
870	Relationships between molecular properties and antimycobacterial activities of steroids. 2002 , 16, 107-13	15
869	Improvements in the Two-Dimensional Nuclear Magnetic Resonance Spectroscopy of Humic Substances. 2002 , 31, 388-392	8
868	The novel low temperature synthesis of nanocrystalline MgAl ₂ O ₄ spinel using gel precursors. 2002 , 37, 1041-1050	51
867	Structure and dynamics of the anticodon arm binding domain of Bacillus stearothermophilus Tyrosyl-tRNA synthetase. 2002 , 10, 311-7	16
866	Convergent Synthesis, NMR and Conformational Analysis of Tetra- and Pentasaccharide Haptens of the Shigella flexneri Serotype 5a O-Specific Polysaccharide. 2002 , 2002, 2486	14

865	¹ H, ¹³ C and ¹⁵ N NMR chemical shifts of substituted pyrazolo[1,5-a]pyrimidines. 2002 , 40, 480-482	5
864	Total assignment of ¹ H and ¹³ C NMR spectra of three triterpene saponins from roots of <i>Silene vulgaris</i> (Moench) Garcke. 2002 , 40, 618-621	17
863	Structural characterization of a cyclic dipeptidomimetic: the effect of ring size on a 1,2,5-trisubstituted-3-oxo-1,4-diazepine system. 2002 , 64, 16-25	3
862	Photomodulation of conformational states. III. Water-soluble bis-cysteinyl-peptides with (4-aminomethyl) phenylazobenzoic acid as backbone constituent. 2002 , 63, 382-93	35
861	Dimer structure of magainin 2 bound to phospholipid vesicles. 2002 , 64, 314-27	72
860	Studies of the receptor-bound conformation of alpha11beta3 antagonists by ¹⁵ N-edited NMR spectroscopy. 2002 , 66, 326-38	2
859	Solution structure of the unbound, oxidized Photosystem I subunit PsaC, containing [4Fe-4S] clusters F(A) and F(B): a conformational change occurs upon binding to photosystem I. 2002 , 7, 461-72	29
858	Synthesis and conformational analysis of a coumarinic acid-based cyclic prodrug of an opioid peptide with modified sensitivity to esterase-catalyzed bioconversion. 2002 , 59, 183-95	23
857	NMR structure of a minimum activity domain of human parathyroid peptide hormone: structural origin of receptor activation. 2002 , 60, 239-46	2
856	Functional expression of <i>Pseudomonas aeruginosa</i> GDP-4-keto-6-deoxy-D-mannose reductase which synthesizes GDP-rhamnose. <i>FEBS Journal</i> , 2002 , 269, 593-601	24
855	Conformational analysis by CD and NMR spectroscopy of a peptide encompassing the amphipathic domain of YopD from <i>Yersinia</i> . <i>FEBS Journal</i> , 2002 , 269, 3659-68	22
854	Solution NMR structure of five representative glycosylated polyene macrolide antibiotics with a sterol-dependent antifungal activity. <i>FEBS Journal</i> , 2002 , 269, 4533-41	30
853	Structure elucidation and conformational properties of eprosartan, a non peptide angiotensin II AT1 antagonist. 2002 , 28, 125-35	21
852	High resolution proton magnetic resonance spectroscopy of human cervical mucus. 2002 , 28, 827-40	7
851	Peptoid residues and beta-turn formation. 2002 , 8, 241-52	34
850	Synthesis and NMR characterization of 6-Phenyl-6-deoxy-2,3-di-O-methylcellulose. 2002 , 13, 413-427	3
849	Synthesis and NMR study of a linear pentasaccharide fragment of the <i>Shigella flexneri</i> 5a O-specific polysaccharide. 2002 , 58, 2593-2604	16
848	Selective conversion of O-succinimidyl carbamates to N-(O-carbamoyl)-succinmonoamides and ureas. 2002 , 43, 3443-3445	4

847	Structural analysis and chemical depolymerization of the capsular polysaccharide of <i>Streptococcus pneumoniae</i> type 1. 2002 , 337, 335-44	37
846	Conformational analysis of two xylose-containing N-glycans in aqueous solution by using ¹ H NMR ROESY and NOESY spectroscopy in combination with MD simulations. 2002 , 337, 2279-99	9
845	Rational structure-based design of a novel carboxypeptidase R inhibitor. 2002 , 9, 1129-39	13
844	High-resolution NMR structure of an AT-rich DNA sequence. 2002 , 22, 265-80	28
843	[NMR spectroscopy for studying structure and intramolecular dynamics of modified analogs of steroid hormones]. 2002 , 28, 220-35	9
842	Oligomerization of acidic fibroblast growth factor is not a prerequisite for its cell proliferation activity. 2002 , 11, 1050-61	31
841	Solution structure of a putative ribosome binding protein from <i>Mycoplasma pneumoniae</i> and comparison to a distant homolog. 2003 , 4, 235-43	13
840	[Catechol siderophore, produced by thermoresistant strain of <i>Bacillus licheniformis</i> VK21]. 2003 , 29, 597-604	25
839	The role of side chain conformational flexibility in surface recognition by <i>Tenebrio molitor</i> antifreeze protein. 2003 , 12, 1323-31	28
838	Multinuclear magnetic resonance, electrospray ionization time-of-flight mass spectral and molecular modelling characterization of lithocholic acid amide esters with various nitrogen heterocycles. 2003 , 41, 567-576	5
837	The role of cystine knots in collagen folding and stability, part I. Conformational properties of (Pro-Hyp-Gly) ₅ and (Pro-(4S)-FPro-Gly) ₅ model trimers with an artificial cystine knot. 2003 , 9, 3692-702	28
836	Spectroscopic and computational investigations of stable radical anions of triosmium benzoheterocycle clusters. 2003 , 9, 5749-56	31
835	The structural plasticity of the C terminus of p21Cip1 is a determinant for target protein recognition. 2003 , 4, 863-9	16
834	Conformational features of human melanin-concentrating hormone: an NMR and computational analysis. 2003 , 4, 73-81	7
833	NMR structure of the single QALGGH zinc finger domain from the <i>Arabidopsis thaliana</i> SUPERMAN protein. 2003 , 4, 171-80	36
832	Structure-function relationship studies of bovine parathyroid hormone [bPTH(1-34)] analogues containing alpha-amino-iso-butyric acid (Aib) residues. 2003 , 68, 437-57	8
831	The solution structures and activity of caerin 1.1 and caerin 1.4 in aqueous trifluoroethanol and dodecylphosphocholine micelles. 2003 , 69, 42-59	14
830	Synthesis, conformation, and immunosuppressive activity of CLX and its analogues. 2003 , 70, 497-511	5

829	Hyaluronate tetrasaccharide- Cu(II) interaction: a NMR study. 2003 , 70, 260-9	3
828	Conformational and biological characterization of human parathyroid hormone hPTH(1-34) analogues containing beta-amino acid residues in positions 17-19. 2003 , 70, 534-47	13
827	Isolation, structural characterization, and immunological evaluation of a high-molecular-weight exopolysaccharide from <i>Staphylococcus aureus</i> . 2003 , 338, 903-22	86
826	Structural elucidation of a novel core oligosaccharide backbone of the lipopolysaccharide from the new bacterial species <i>Agrobacterium larrymoorei</i> . 2003 , 338, 2721-30	14
825	Sine-modulated two-dimensional pure exchange solid-state NMR as a tool for characterizing dynamics in solids. 2003 , 377, 322-328	2
824	Structural characterization of β -pyridylaminocrotonoyl-2-pyridylamide by ESI-MS, NMR, single crystal X-ray analysis and ab initio methods. 2003 , 654, 61-69	7
823	¹ H-, ¹³ C-, and ¹⁵ N-NMR and ESI-TOF+ MS studies of a supramolecular complex of silver(I) and a cholaphane. 2003 , 668, 43-50	20
822	Conformational and biological studies for a pair of novel synthetic AT1 antagonists: stereoelectronic requirements for antihypertensive efficacy. 2003 , 31, 833-44	13
821	Structural elucidation and conformational properties of the toxin paralyisin beta-Ala-Tyr. 2003 , 31, 713-21	5
820	The structure of the antimicrobial peptide Ac-RRWWRF-NH ₂ bound to micelles and its interactions with phospholipid bilayers. 2003 , 61, 219-29	79
819	Structures of neuropeptide gamma from goldfish and mammalian neuropeptide gamma, as determined by ¹ H NMR spectroscopy. 2003 , 61, 274-85	9
818	Conformational studies of antimetastatic laminin-1 derived peptides in different solvent systems, using solution NMR spectroscopy. 2003 , 61, 24-39	3
817	nNOS inhibition, antimicrobial and anticancer activity of the amphibian skin peptide, citropin 1.1 and synthetic modifications. The solution structure of a modified citropin 1.1. <i>FEBS Journal</i> , 2003 , 270, 1141-53	60
816	Modified merozoite surface protein-1 peptides with short alpha helical regions are associated with inducing protection against malaria. <i>FEBS Journal</i> , 2003 , 270, 3946-52	27
815	Alpha helix shortening in 1522 MSP-1 conserved peptide analogs is associated with immunogenicity and protection against <i>P. falciparum</i> malaria. 2003 , 50, 400-9	26
814	Synthetic peptides mimicking the interleukin-6/gp130 interaction: a two-helix bundle system. Design and conformational studies. 2003 , 9, 90-105	1
813	Conformational features of a synthetic model of the first extracellular loop of the angiotensin II AT1A receptor. 2003 , 9, 229-43	9
812	Enantioselective binding of lambda- and delta-[Ru(bpy)(2)(HPIP)]Cl ₂ (HPIP=2-(2-hydroxyphenyl)imidazo[4,5-f][1,10]phenanthroline) to the hexanucleotide [d(5'-GTCGAC-3') ₂]. 2003 , 93, 221-34	14

811	Photocontrol of cell adhesion processes: model studies with cyclic azobenzene-RGD peptides. 2003 , 10, 487-90	57
810	Solution structure of Alo-3: a new knottin-type antifungal peptide from the insect <i>Acrocinus longimanus</i> . 2003 , 42, 14434-42	55
809	An alpha helix conformationally restricted peptide is recognized by cervical carcinoma patients' sera. 2003 , 46, 5389-94	25
808	PA1b, an insecticidal protein extracted from pea seeds (<i>Pisum sativum</i>): 1H-2-D NMR study and molecular modeling. 2003 , 42, 11915-23	43
807	Solution structure of the N-terminal amphitropic domain of <i>Escherichia coli</i> glucose-specific enzyme IIA in membrane-mimetic micelles. 2003 , 12, 1087-96	44
806	Solution structure of Pi4, a short four-disulfide-bridged scorpion toxin specific of potassium channels. 2003 , 12, 1844-54	12
805	Structural consequences of metallothionein dimerization: solution structure of the isolated Cd4-alpha-domain and comparison with the holoprotein dimer. 2003 , 42, 8403-10	17
804	A folded, secondary structure in step-growth oligomers from covalently linked, crowded aromatics. 2003 , 125, 4870-3	28
803	Introduction of a chemical constraint in a short peptide derived from human acidic fibroblast growth factor elicits mitogenic structural determinants. 2003 , 46, 2325-33	12
802	NMR solution structure of the glucagon antagonist [desHis1, desPhe6, Glu9]glucagon amide in the presence of perdeuterated dodecylphosphocholine micelles. 2003 , 42, 2825-35	19
801	Solution structure of a novel disintegrin, salmosin, from <i>Agkistrodon halys</i> venom. 2003 , 42, 14408-15	33
800	MSP-1 malaria pseudopeptide analogs: biological and immunological significance and three-dimensional structure. 2003 , 384, 71-82	10
799	Synthesis and structure-activity relationship of mannose-based peptidomimetics selectively blocking integrin alpha4beta7 binding to mucosal addressin cell adhesion molecule-1. 2003 , 46, 5752-62	18
798	An improved solution structure for psi-conotoxin PiiiE. 2003 , 42, 6347-52	14
797	Conformational studies of a bombolitin III-derived peptide mimicking the four-helix bundle structural motif of proteins. 2003 , 125, 15314-23	5
796	New probes of ligand flexibility in drug design: transferred (13)C CSA-dipolar cross-correlated relaxation at natural abundance. 2003 , 125, 11116-30	16
795	Structure of the nuclear factor ALY: insights into post-transcriptional regulatory and mRNA nuclear export processes. 2003 , 42, 7348-57	19
794	Distorting malaria peptide backbone structure to enable fitting into MHC class II molecules renders modified peptides immunogenic and protective. 2003 , 46, 2250-3	26

793	Characterization and three-dimensional structure determination of psi-conotoxin Piiiif, a novel noncompetitive antagonist of nicotinic acetylcholine receptors. 2003 , 42, 6353-62	38
792	Linearization of a naturally occurring circular protein maintains structure but eliminates hemolytic activity. 2003 , 42, 6688-95	102
791	NMR Studies of Structural Phase Transitions in Random Copolymers. 2003 , 36, 477-485	1
790	Structure and metal binding studies of the second copper binding domain of the Menkes ATPase. 2003 , 143, 209-18	27
789	The three-dimensional solution structure of NaD1, a new floral defensin from <i>Nicotiana glauca</i> and its application to a homology model of the crop defense protein alfAFP. 2003 , 325, 175-88	114
788	NMR structure of the HIV-1 regulatory protein VPR. 2003 , 327, 215-27	144
787	NMR solution structure of the non-RGD disintegrin obtustatin. 2003 , 329, 135-45	45
786	Disulfide pairings and secondary structure of porcine beta-microseminoprotein. 2003 , 541, 80-4	10
785	NMR studies of the fifth transmembrane segment of sarcoplasmic reticulum Ca ²⁺ -ATPase reveals a hinge close to the Ca ²⁺ -ligating residues. 2003 , 544, 50-6	8
784	Solution structure of epiregulin and the effect of its C-terminal domain for receptor binding affinity. 2003 , 553, 232-8	14
783	Contryphan-Vn: a modulator of Ca ²⁺ -dependent K ⁺ channels. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 303, 238-46	3-4 46
782	The solution structure of the oxidized bovine microsomal cytochrome b(5) mutant V61H. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 307, 600-9	3-4 2
781	Solution structure of the tachykinin peptide eledoisin. 2003 , 84, 655-64	22
780	Plasmodium falciparum SERA protein peptide analogues having short helical regions induce protection against malaria. 2003 , 85, 651-7	14
779	Solution structure of a peptide derived from the beta subunit of LFA-1. 2003 , 24, 827-35	5
778	Conformation of neuropeptide Y receptor antagonists: structural implications in receptor selectivity. 2003 , 24, 1035-43	9
777	6746 SERA peptide analogues immunogenicity and protective efficacy against malaria is associated with short alpha helix formation: malaria protection associated with peptides alpha helix shortening. 2003 , 24, 999-1006	23
776	Analysis of a Plasmodium falciparum EBA-175 peptide with high binding capacity to erythrocytes and their analogues using ¹ H NMR. 2003 , 141, 115-21	28

775	Structure of <i>Petunia hybrida</i> defensin 1, a novel plant defensin with five disulfide bonds. 2003 , 42, 8214-22	85
774	Novel sst(4)-selective somatostatin (SRIF) agonists. 4. Three-dimensional consensus structure by NMR. 2003 , 46, 5606-18	30
773	Structural characterization of hellethionins from <i>Helleborus purpurascens</i> . 2003 , 42, 2404-11	28
772	Presence of an unusual GM2 derivative, taurine-conjugated GM2, in Tay-Sachs brain. <i>Journal of Biological Chemistry</i> , 2003 , 278, 35286-91	5-4 24
771	Immunogenicity and protectivity of <i>Plasmodium falciparum</i> EBA-175 peptide and its analog is associated with alpha-helical region shortening and displacement. 2003 , 384, 1443-50	26
770	Isolation, structure, and activity of GID, a novel alpha 4/7-conotoxin with an extended N-terminal sequence. <i>Journal of Biological Chemistry</i> , 2003 , 278, 3137-44	5-4 110
769	Structural mimicry in the phage phi21 N peptide-boxB RNA complex. 2003 , 9, 663-76	24
768	Disulfide folding pathways of cystine knot proteins. Tying the knot within the circular backbone of the cyclotides. <i>Journal of Biological Chemistry</i> , 2003 , 278, 6314-22	5-4 103
767	Twists, knots, and rings in proteins. Structural definition of the cyclotide framework. <i>Journal of Biological Chemistry</i> , 2003 , 278, 8606-16	5-4 254
766	Solution structure of the recombinant penaeidin-3, a shrimp antimicrobial peptide. <i>Journal of Biological Chemistry</i> , 2003 , 278, 36859-67	5-4 46
765	A peptide derived from LFA-1 protein that modulates T-cell adhesion binds to soluble ICAM-1 protein. 2003 , 20, 635-44	7
764	Solution structures of stem-loop RNAs that bind to the two N-terminal RNA-binding domains of nucleolin. 2003 , 31, 6461-72	28
763	Solution structure of the <i>Mycobacterium tuberculosis</i> complex protein MPB70: from tuberculosis pathogenesis to inherited human corneal disease. <i>Journal of Biological Chemistry</i> , 2003 , 278, 43736-43	5-4 39
762	Roles of aromatic residues in the structure and biological activity of the small cytokine, growth-blocking peptide (GBP). <i>Journal of Biological Chemistry</i> , 2003 , 278, 10778-83	5-4 6
761	Conformational studies of the O-specific polysaccharide of <i>Shigella flexneri</i> 5a and of four related synthetic pentasaccharide fragments using NMR and molecular modeling. <i>Journal of Biological Chemistry</i> , 2003 , 278, 47928-36	5-4 31
760	Preorganization of molecular binding sites in designed diiron proteins. 2003 , 100, 3772-7	67
759	Structural study of DNA duplex containing an N-(2-deoxy-beta-D-erythro-pentofuranosyl) formamide frameshift by NMR and restrained molecular dynamics. 2003 , 31, 5930-40	9
758	Conformation of a bactericidal domain of puoroindoline a: structure and mechanism of action of a 13-residue antimicrobial peptide. 2003 , 185, 4938-47	101

757	NMR study on the structural changes of cytochrome P450cam upon the complex formation with putidaredoxin. Functional significance of the putidaredoxin-induced structural changes. <i>Journal of Biological Chemistry</i> , 2003 , 278, 39809-21	5-4	42
756	Structural basis for new pattern of conserved amino acid residues related to chitin-binding in the antifungal peptide from the coconut rhinoceros beetle <i>Oryctes rhinoceros</i> . <i>Journal of Biological Chemistry</i> , 2003 , 278, 22820-7	5-4	15
755	CCHX zinc finger derivatives retain the ability to bind Zn(II) and mediate protein-DNA interactions. <i>Journal of Biological Chemistry</i> , 2003 , 278, 28011-8	5-4	41
754	Cleavage of DNA without loss of genetic information by incorporation of a disaccharide nucleoside. 2003 , 31, 6758-69		4
753	Accommodation of alpha-substituted residues in the beta-peptide 12-helix: expanding the range of substitution patterns available to a foldamer scaffold. 2003 , 125, 8539-45		66
752	Identification of a novel family of proteins in snake venoms. Purification and structural characterization of nawaprin from <i>Naja nigricollis</i> snake venom. <i>Journal of Biological Chemistry</i> , 2003 , 278, 40097-104	5-4	49
751	X-ray Structures and Complete NMR Assignment by DFT Calculations of [Os(bpy) ₂ (CO)Cl]PF ₆ and [Os(bpy) ₂ (CO)H]PF ₆ Complexes. 2003 , 22, 4012-4019		26
750	The three-dimensional structural surface of two beta-sheet scorpion toxins mimics that of an alpha-helical dihydropyridine receptor segment. <i>Biochemical Journal</i> , 2003 , 370, 517-27	3-8	26
749	Activating the ryanodine receptor with dihydropyridine receptor II-III loop segments: size and charge do matter. 2004 , 9, 2860-72		7
748	Tissue-specific expression of head-to-tail cyclized miniproteins in Violaceae and structure determination of the root cyclotide <i>Viola hederacea</i> root cyclotide1. 2004 , 16, 2204-16		97
747	NMR solution structures of delta-conotoxin EVIA from <i>Conus ermineus</i> that selectively acts on vertebrate neuronal Na ⁺ channels. <i>Journal of Biological Chemistry</i> , 2004 , 279, 21356-66	5-4	33
746	Assignment of the four disulfides in the N-terminal somatomedin B domain of native vitronectin isolated from human plasma. <i>Journal of Biological Chemistry</i> , 2004 , 279, 35867-78	5-4	12
745	The solution structure of the N-terminal domain of human vitronectin: proximal sites that regulate fibrinolysis and cell migration. <i>Journal of Biological Chemistry</i> , 2004 , 279, 29359-66	5-4	31
744	Solution structure of spheniscin, a beta-defensin from the penguin stomach. <i>Journal of Biological Chemistry</i> , 2004 , 279, 30433-9	5-4	33
743	Full structural characterization of the lipid A components from the <i>Agrobacterium tumefaciens</i> strain C58 lipopolysaccharide fraction. 2004 , 14, 805-15		27
742	The 10th and 11th residues of short length N-terminal PTH(1-12) analogue are important for its optimum potency. 2004 , 64, 25-32		8
741	A novel type of highly negatively charged lipooligosaccharide from <i>Pseudomonas stutzeri</i> OX1 possessing two 4,6-O-(1-carboxy)-ethylidene residues in the outer core region. <i>FEBS Journal</i> , 2004 , 271, 2691-704		24
740	Design, structure and biological activity of beta-turn peptides of CD2 protein for inhibition of T-cell adhesion. <i>FEBS Journal</i> , 2004 , 271, 2873-86		13

739	Solution structure of 2',5' d(G4C4). Relevance to topological restrictions and nature's choice of phosphodiester links. <i>FEBS Journal</i> , 2004 , 271, 2956-66	13
738	Solution structure of IsTX. A male scorpion toxin from <i>Opisthacanthus madagascariensis</i> (Ischnuridae). <i>FEBS Journal</i> , 2004 , 271, 3855-64	17
737	Solution structure of the cyclotide palicourein: implications for the development of a pharmaceutical framework. 2004 , 12, 85-94	36
736	Solution structures of the C-terminal headpiece subdomains of human villin and advillin, evaluation of headpiece F-actin-binding requirements. 2004 , 13, 1276-87	40
735	The comparative study on the solution structures of the oxidized bovine microsomal cytochrome b5 and mutant V45H. 2004 , 13, 2161-9	10
734	Phenolic glycoside isolated from seeds of the greater plantain (<i>Plantago major</i> L.). 2004 , 396, 132-5	7
733	Characterization of the heme-histidine cross-link in cyanobacterial hemoglobins from <i>Synechocystis</i> sp. PCC 6803 and <i>Synechococcus</i> sp. PCC 7002. 2004 , 9, 183-94	25
732	Structural study of copper(I)-bleomycin. 2004 , 9, 323-34	7
731	In vivo 2D magnetic resonance spectroscopy of small animals. 2004 , 17, 317-38	16
730	Large scale enantiomeric synthesis, purification, and characterization of α -unsaturated amino acids via a Gly-Ni(II)-BPB-complex. 2004 , 60, 8233-8243	28
729	Conformational analysis of peptide analogues of silkworm chorion protein segments using CD, NMR and molecular modelling. 2004 , 10, 381-92	1
728	Solution conformational study of nociceptin and its 1-13 and 1-11 fragments using circular dichroism and two-dimensional NMR in conjunction with theoretical conformational analysis. 2004 , 10, 678-90	7
727	Influence of conformational flexibility on biological activity in cyclic astin analogues. 2004 , 76, 477-84	8
726	A synthetic peptide reproducing the mitochondrial targeting motif of AKAP121: a conformational study. 2004 , 76, 459-66	4
725	Peptide-chelating agent conjugate for selective targeting of somatostatin receptor type 1: synthesis and characterization. 2004 , 76, 527-34	
724	Synthesis and Characterization of the Diastereomers α and β $[\text{Ru}(\text{bpy})_2(\text{m-bpy-Gly-L-His-L-Lys})]\text{Cl}_2$ ^1H NMR Studies on Their Interactions with the Deoxynucleotide Duplex $d[(5'\text{-CGCGAATTCGCG-3}')_2]$. 2004 , 2004, 1427-1439	15
723	Elucidation of the Solution Conformations of Loloatin C by NMR Spectroscopy and Molecular Simulation. 2004 , 2004, 31-37	3
722	A Novel Core Region, Lacking Heptose and Phosphate, of the Lipopolysaccharide from the Gram-Negative Bacterium <i>Pseudomonas cichorii</i> (Pseudomonadaceae RNA Group 1). 2004 , 2004, 2427-2435	5

721	Structural Analysis of a Novel Polysaccharide of the Lipopolysaccharide-Deficient Extremophile Gram-Negative Bacterium <i>Thermus thermophilus</i> HB8. 2004 , 2004, 5047-5054		11
720	Chiral discrimination studies of (+)-(18-crown-6)-2,3,11,12-tetracarboxylic acid by high-performance liquid chromatography and NMR spectroscopy. 2004 , 42, 389-95		19
719	Binding mode of TMC-95A analogues to eukaryotic 20S proteasome. 2004 , 5, 1256-66		43
718	Synthesis and characterization of lithocholic acid derived dipyrromethanes: precursors for pyrrole-steroidal macrocycles. 2004 , 693, 81-86		5
717	Structural transitions as determinants of the action of the calcium-dependent antibiotic daptomycin. 2004 , 11, 949-57		208
716	Covariance NMR spectroscopy by singular value decomposition. <i>Journal of Magnetic Resonance</i> , 2004 , 171, 277-83	3	92
715	Novel avilamycin derivatives with improved polarity generated by targeted gene disruption. 2004 , 11, 1403-11		28
714	Structure-activity relationships for the beta-hairpin cationic antimicrobial peptide polyphemusin I. 2004 , 1698, 239-50		60
713	The complete structure of the lipooligosaccharide from the halophilic bacterium <i>Pseudoalteromonas issachenkonii</i> KMM 3549T. 2004 , 339, 1985-93		20
712	The structure of the phosphorylated carbohydrate backbone of the lipopolysaccharide of the phytopathogen bacterium <i>Pseudomonas tolaasii</i> . 2004 , 339, 2241-8		14
711	Structure of minor oligosaccharides from the lipopolysaccharide fraction from <i>Pseudomonas stutzeri</i> OX1. 2004 , 339, 2657-65		7
710	zFP538, a yellow fluorescent protein from coral, belongs to the DsRed subfamily of GFP-like proteins but possesses the unexpected site of fragmentation. 2004 , 43, 4764-72		19
709	Structure of the amodiaquine-FPIX mu oxo dimer solution complex at atomic resolution. 2004 , 43, 8078-84		35
708	Structural characterization of imazalil/beta-cyclodextrin inclusion complex. 2004 , 52, 1590-3		15
707	Three-dimensional structure of the mini-M conotoxin mr3a. 2004 , 43, 425-9		19
706	The Novel Synthesis of La _{0.8} Sr _{0.2} MnO ₃ Using the Michael-Addition Directed Hydrogelation of Acrylates for Materials Synthesis (MADHAMS) Method. 2004 , 16, 5336-5343		10
705	Conformational and Enantioselectivity in Host-Guest Chemistry: The Selective Binding of Cis Amides Examined by Free Energy Calculations. 2004 , 108, 17571-17582		4
704	KappaM-conotoxin RIIIK, structural and functional novelty in a K ⁺ channel antagonist. 2004 , 43, 8625-35		36

703	Solution structure and backbone dynamics of the K18G/R82E Alicyclobacillus acidocaldarius thioredoxin mutant: a molecular analysis of its reduced thermal stability. 2004 , 43, 6043-58		21
702	Cyanide binding to hexacoordinate cyanobacterial hemoglobins: hydrogen-bonding network and heme pocket rearrangement in ferric H117A Synechocystis hemoglobin. 2004 , 43, 12622-33		25
701	Design, synthesis, and biological evaluation of new cyclic disulfide decapeptides that inhibit the binding of AP-1 to DNA. 2004 , 47, 4239-46		18
700	Chemical diversity of polyene macrolides produced by Streptomyces noursei ATCC 11455 and recombinant strain ERD44 with genetically altered polyketide synthase NysC. 2004 , 48, 4120-9		49
699	Solution structure and functional characterization of SGTx1, a modifier of Kv2.1 channel gating. 2004 , 43, 890-7		95
698	Orientating peptide residues and increasing the distance between pockets to enable fitting into MHC-TCR complex determine protection against malaria. 2004 , 43, 6545-53		17
697	Structure and characterization of flavolipids, a novel class of biosurfactants produced by Flavobacterium sp. strain MTN11. 2004 , 70, 114-20		88
696	Solution structure of cryptdin-4, a mouse paneth cell alpha-defensin. 2004 , 43, 15759-66		32
695	Orientation and helical conformation of a tissue-specific hunter-killer peptide in micelles. 2004 , 13, 1988-96		8
694	NMR structure determination and calcium binding effects of lipopeptide antibiotic daptomycin. 2004 , 2, 1872-8		85
693	Structural Similarities of Micelle-bound Peptide YY (PYY) and Neuropeptide Y (NPY) are Related to their Affinity Profiles at the Y Receptors*1. 2004 ,		
692	Shortening and modifying the 1513 MSP-1 peptide's alpha-helical region induces protection against malaria. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 315, 418-27	3-4	25
691	Modifying RESA protein peptide 6671 to fit into HLA-DRbeta1* pockets induces protection against malaria. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 315, 1154-64	3-4	22
690	Changing ABRA protein peptide to fit into the HLA-DRbeta1*0301 molecule renders it protection-inducing. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 322, 119-25	3-4	15
689	Conformational preferences and activities of peptides from the catecholamine release-inhibitory (catestatin) region of chromogranin A. 2004 , 118, 75-87		21
688	Induction and displacement of an helix in the 6725 SERA peptide analogue confers protection against P. falciparum malaria. 2004 , 22, 1281-9		9
687	Presence of transient helical segments in the galanin-like peptide evident from (1)H NMR, circular dichroism, and prediction studies. 2004 , 146, 261-71		7
686	Mapping the anatomy of a Plasmodium falciparum MSP-1 epitope using pseudopeptide-induced mono- and polyclonal antibodies and CD and NMR conformation analysis. 2004 , 148, 110-22		10

685	NMR solution structure of the mitochondrial F1beta presequence from <i>Nicotiana plumbaginifolia</i> . 2004 , 336, 1129-40		45
684	Structural similarities of micelle-bound peptide YY (PYY) and neuropeptide Y (NPY) are related to their affinity profiles at the Y receptors. 2004 , 339, 1153-68		64
683	Mutational and structural analysis of stem-loop IIIc of the hepatitis C virus and GB virus B internal ribosome entry sites. 2004 , 343, 805-17		30
682	Synthesis and structure-activity relationships of 5,6,7,8-tetrahydro-4H-thieno[3,2-b]azepine derivatives: novel arginine vasopressin antagonists. 2004 , 47, 101-9		25
681	¹³ C NMR Study of the Self-Association of Chloroquine, Amodiaquine, and Quinine. 2004 , 108, 8505-8513		21
680	Mapping oxygen accessibility to ribonuclease A using high-resolution NMR relaxation spectroscopy. 2004 , 86, 1713-25		29
679	Molecular basis of the high-affinity activation of type 1 ryanodine receptors by imperatoxin A. <i>Biochemical Journal</i> , 2004 , 377, 385-94	3.8	32
678	Solution conformation of alpha-conotoxin G1C, a novel potent antagonist of alpha3beta2 nicotinic acetylcholine receptors. <i>Biochemical Journal</i> , 2004 , 380, 347-52	3.8	24
677	Solution structure of human orexin-A: regulator of appetite and wakefulness. 2004 , 37, 565-73		18
676	Cyclohexenyl nucleic acids: conformationally flexible oligonucleotides. 2005 , 33, 2452-63		25
675	The C-terminal domain of the HIV-1 regulatory protein Vpr adopts an antiparallel dimeric structure in solution via its leucine-zipper-like domain. <i>Biochemical Journal</i> , 2005 , 387, 333-41	3.8	41
674	Modulation of the catalytic mechanism of hen egg white lysozyme (HEWL) by photochromism of azobenzene. 2005 , 175, 100-107		7
673	¹ H, ¹³ C NMR spectral and single crystal structural studies of toxaphene congeners. Quantum chemical calculations for preferred conformers of 2,5-endo,6-exo,8,9,9,10,10-octachloro-2-bornene and their DFT/GIAO ¹³ C chemical shifts. 2005 , 754, 77-84		
672	Structural characterization of the carbohydrate backbone of the lipooligosaccharide of the marine bacterium <i>Arenibacter certesii</i> strain KMM 3941(T). 2005 , 340, 2540-9		19
671	Genes encoding enzymes responsible for biosynthesis of L-lyxose and attachment of eurenate during avilamycin biosynthesis. 2005 , 12, 1137-43		39
670	Isolation, gene expression and solution structure of a novel moricin analogue, antibacterial peptide from a lepidopteran insect, <i>Spodoptera litura</i> . 2005 , 1752, 83-92		24
669	Helical structure determined by NMR of the HIV-1 (345-392)Gag sequence, surrounding p2: implications for particle assembly and RNA packaging. 2005 , 14, 375-86		78
668	Relationship between temporary inhibition and structure of disulfide-linkage analogs of marinostatin, a natural ester-linked protein protease inhibitor. 2005 , 66, 49-58		8

667	Dynamic conformational changes of extracellular S5-P linkers in the hERG channel. 2005 , 569, 75-89	48
666	Chemical structure and biological activity of the <i>Caenorhabditis elegans</i> dauer-inducing pheromone. 2005 , 433, 541-5	279
665	Structure and different conformational states of native AMPA receptor complexes. 2005 , 433, 545-9	226
664	Solution structure of human proinsulin C-peptide. 2005 , 272, 4284-93	25
663	Complete Structural Elucidation of a Novel Lipooligosaccharide from the Outer Membrane of the Marine Bacterium <i>Shewanella pacifica</i> . 2005 , 2005, 2281-2291	19
662	Conformational studies of Aib-rich peptides containing lactam-bridged side chains: evidence of 3(10)-helix formation. 2005 , 80, 294-302	7
661	Quantitative analysis of helix-coil transition of block copolypeptide, Glu12-Ala12, by combined use of CD and NMR spectroscopy. 2005 , 80, 225-32	4
660	Photomodulation of conformational states. IV. Integrin-binding RGD-peptides with (4-aminomethyl)phenylazobenzoic acid as backbone constituent. 2005 , 77, 304-13	22
659	Solution structure of chi-conopeptide MrIA, a modulator of the human norepinephrine transporter. 2005 , 80, 815-23	38
658	Solution structure of delta-Am2766: a highly hydrophobic delta-conotoxin from <i>Conus amadis</i> that inhibits inactivation of neuronal voltage-gated sodium channels. 2005 , 2, 535-56	9
657	IP-COSY, a totally in-phase and sensitive COSY experiment. 2005 , 43, 372-9	33
656	A new ligand for immunoglobulin g subdomains by screening of a synthetic peptide library. 2005 , 6, 1242-53	38
655	The structure of the O-polysaccharide from <i>Pseudomonas stutzeri</i> OX1 containing two different 4-acetylamido-4,6-dideoxy-residues, tomosamine and perosamine. 2005 , 340, 651-6	12
654	The complete structure of the core carbohydrate backbone from the LPS of marine halophilic bacterium <i>Pseudoalteromonas carrageenovora</i> type strain IAM 12662T. 2005 , 340, 1475-82	12
653	The linkage between O-specific caryan and core region in the lipopolysaccharide of <i>Burkholderia caryophylli</i> is furnished by a primer monosaccharide. 2005 , 340, 1802-7	10
652	Synthesis and Structural Characterisation of Two Novel Diastereoisomeric Naproxen Appended β -Cyclodextrin Derivatives. 2005 , 51, 173-180	2
651	Structure of Circulin B and Implications for Antimicrobial Activity of the Cyclotides. 2005 , 11, 99-106	27
650	The Solution Structure of a Cyclic Analog of Neuropeptide Y with High Y1 Receptor Affinity by NMR, CD and MD Simulations. 2005 , 11, 143-152	

649	A dynamic N-capping motif in cytochrome b5: evidence for a pH-controlled conformational switch. 2006 , 63, 336-48		5
648	Solution structure and interaction of the antimicrobial polyphemusins with lipid membranes. 2005 , 44, 15504-13		87
647	Genes involved in formation and attachment of a two-carbon chain as a component of eurekaate, a branched-chain sugar moiety of avilamycin A. 2005 , 71, 400-6		19
646	Solution structure of synthetic penaeidin-4 with structural and functional comparisons with penaeidin-3. <i>Journal of Biological Chemistry</i> , 2005 , 280, 16009-18	5-4	34
645	Discovery, structural determination, and putative processing of the precursor protein that produces the cyclic trypsin inhibitor sunflower trypsin inhibitor 1. <i>Journal of Biological Chemistry</i> , 2005 , 280, 32245-53	5-4	30
644	Isolation and characterization of novel cyclotides from <i>Viola hederaceae</i> : solution structure and anti-HIV activity of vhl-1, a leaf-specific expressed cyclotide. <i>Journal of Biological Chemistry</i> , 2005 , 280, 22395-405	5-4	101
643	Complete structural characterization of the lipid A fraction of a clinical strain of <i>B. cepacia</i> genomovar I lipopolysaccharide. 2005 , 15, 561-70		53
642	Structural basis for phosphatidylinositol phosphate kinase type Iγ binding to talin at focal adhesions. <i>Journal of Biological Chemistry</i> , 2005 , 280, 8381-6	5-4	64
641	Identification and structural determination of the M(3) muscarinic acetylcholine receptor basolateral sorting signal. <i>Journal of Biological Chemistry</i> , 2005 , 280, 24568-75	5-4	17
640	Correlation of three-dimensional structures with the antibacterial activity of a group of peptides designed based on a nontoxic bacterial membrane anchor. <i>Journal of Biological Chemistry</i> , 2005 , 280, 5803-11	5-4	72
639	Solution structure and lipid membrane partitioning of VSTx1, an inhibitor of the KvAP potassium channel. 2005 , 44, 6015-23		88
638	Multiple loop conformations of peptides predicted by molecular dynamics simulations are compatible with nuclear magnetic resonance. 2005 , 44, 4829-40		13
637	Conformational studies of two bradykinin antagonists by using two-dimensional NMR techniques and molecular dynamics simulations. 2005 , 23, 125-34		2
636	NMR structures of double loops of an RNA aptamer against mammalian initiation factor 4A. 2005 , 33, 745-54		14
635	Solution structure of marinostatin, a natural ester-linked protein protease inhibitor. 2005 , 44, 2462-8		25
634	Isolation, solution structure, and insecticidal activity of kalata B2, a circular protein with a twist: do MBius strips exist in nature?. 2005 , 44, 851-60		199
633	Peptides inducing short-lived antibody responses against <i>Plasmodium falciparum</i> malaria have shorter structures and are read in a different MHC II functional register. 2005 , 44, 6745-54		22
632	Selective self-organization of guest molecules in self-assembled molecular boxes. 2005 , 127, 12697-708		69

631	A ribose sugar conformational switch in the LTR-retrotransposon Ty3 polypurine tract-containing RNA/DNA hybrid. 2005 , 127, 16344-5		13
630	N-terminal stathmin-like peptides bind tubulin and impede microtubule assembly. 2005 , 44, 14616-25		32
629	Disulfide bond mutagenesis and the structure and function of the head-to-tail macrocyclic trypsin inhibitor SFTI-1. 2005 , 44, 1145-53		50
628	Structural and DNA-binding studies on the bovine antimicrobial peptide, indolicidin: evidence for multiple conformations involved in binding to membranes and DNA. 2005 , 33, 4053-64		220
627	Structure-activity studies of peptides from the "hot-spot" region of human CD2 protein: development of peptides for immunomodulation. 2005 , 48, 6236-49		12
626	Novel solution structure of porcine beta-microseminoprotein. 2005 , 346, 1071-82		11
625	Elongating modified conserved peptides eliminates their immunogenicity and protective efficacy against <i>P. falciparum</i> malaria. 2005 , 150, 245-58		2
624	Structure and function of the potent cyclic and linear melanocortin analogues. 2005 , 150, 300-8		9
623	Protection against malaria induced by chirally modified <i>Plasmodium falciparum</i> 's MSP-1 42 pseudopeptides. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 329, 1053-66	3-4	6
622	Evidence that the tertiary structure of 20(S)-ginsenoside Rg(3) with tight hydrophobic packing near the chiral center is important for Na(+) channel regulation. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 333, 1194-201	3-4	37
621	Fitting modified HRP-I peptide analogue 3D structure into HLA-DR molecules induces protection against <i>Plasmodium falciparum</i> malaria. 2005 , 37, 336-49		6
620	Structural determinants of unexpected agonist activity in a retro-peptide analogue of the SDF-1alpha N-terminus. 2005 , 579, 5293-8		3
619	Automated NMR structure determination and disulfide bond identification of the myotoxin crostamine from <i>Crotalus durissus terrificus</i> . 2005 , 46, 759-67		66
618	Evidence supporting the hypothesis that specifically modifying a malaria peptide to fit into HLA-DRbeta1*03 molecules induces antibody production and protection. 2005 , 23, 1579-87		10
617	The elicitation of plant innate immunity by lipooligosaccharide of <i>Xanthomonas campestris</i> . <i>Journal of Biological Chemistry</i> , 2005 , 280, 33660-8	5-4	145
616	Probing the pH-dependent structural features of alpha-KTx12.1, a potassium channel blocker from the scorpion <i>Tityus serrulatus</i> . 2005 , 14, 1025-38		12
615	Structural and functional study of an <i>Anemonia</i> elastase inhibitor, a "nonclassical" Kazal-type inhibitor from <i>Anemonia sulcata</i> . 2005 , 44, 9626-36		30
614	The micelle-bound structure of an antimicrobial peptide derived from the alpha-chain of bovine hemoglobin isolated from the tick <i>Boophilus microplus</i> . 2005 , 44, 6440-51		34

613	Somatostatin receptor 1 selective analogues: 4. Three-dimensional consensus structure by NMR. 2005 , 48, 523-33	29
612	Synthesis, conformation, and stereodynamics of a salt of 2-[[2-(3,4-dichlorophenyl)-ethyl]propylamino]-1-pyridin-3-ylethanol. 2006 , 71, 792-5	4
611	Lariatins, antimycobacterial peptides produced by <i>Rhodococcus</i> sp. K01-B0171, have a lasso structure. 2006 , 128, 7486-91	113
610	Comparison of proposed putative active conformations of myelin basic protein epitope 87-99 linear altered peptide ligands by spectroscopic and modelling studies: the role of positions 91 and 96 in T-cell receptor activation. 2006 , 49, 6683-91	21
609	Base-displaced intercalated structure of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the <i>NarI</i> restriction enzyme, a hotspot for -2 bp deletions. 2006 , 128, 10085-95	57
608	A novel N14Y mutation in Connexin26 in keratitis-ichthyosis-deafness syndrome: analyses of altered gap junctional communication and molecular structure of N terminus of mutated Connexin26. 2006 , 169, 416-23	40
607	Comparison of penetratin and other homeodomain-derived cell-penetrating peptides: interaction in a membrane-mimicking environment and cellular uptake efficiency. 2006 , 45, 1408-20	67
606	NMR studies of the fifth transmembrane segment of Na ⁺ ,K ⁺ -ATPase reveals a non-helical ion-binding region. 2006 , 580, 4777-83	3
605	The PP-fold solution structure of human polypeptide YY and human PYY3-36 as determined by NMR. 2006 , 45, 8350-7	41
604	A conformational two-state peptide model system containing an ultrafast but soft light switch. 2006 , 90, 2099-108	24
603	The paperclip triplex: understanding the role of apex residues in tight turns. 2006 , 91, 2552-63	5
602	Formation of lactones from sialylated MUC1 glycopeptides. 2006 , 4, 713-20	17
601	Dermaseptin S9, an alpha-helical antimicrobial peptide with a hydrophobic core and cationic termini. 2006 , 45, 468-80	85
600	Solution structures of tetrahaem ferricytochrome c3 from <i>Desulfovibrio vulgaris</i> (Hildenborough) and its K45Q mutant: the molecular basis of cooperativity. 2006 , 1757, 143-53	7
599	Novel sst2-selective somatostatin agonists. Three-dimensional consensus structure by NMR. 2006 , 49, 4487-96	44
598	Fungicidal properties of two saponins from <i>Capsicum frutescens</i> and the relationship of structure and fungicidal activity. 2006 , 52, 336-42	22
597	Structural features in EIAV NCp11: a lentivirus nucleocapsid protein with a short linker. 2006 , 45, 5517-26	16
596	Solution structures of human LL-37 fragments and NMR-based identification of a minimal membrane-targeting antimicrobial and anticancer region. 2006 , 128, 5776-85	196

595	Solution structure of discrepin, a new K ⁺ -channel blocking peptide from the alpha-KTx15 subfamily. 2006 , 45, 1795-804		18
594	Oxygen accessibility to ribonuclease a: quantitative interpretation of nuclear spin relaxation induced by a freely diffusing paramagnet. 2006 , 110, 580-8		29
593	Current applications of bicelles in NMR studies of membrane-associated amphiphiles and proteins. 2006 , 45, 8453-65		208
592	Evolution of new function in the GTP cyclohydrolase II proteins of <i>Streptomyces coelicolor</i> . 2006 , 45, 12144-55		18
591	NMR studies of aurein 1.2 analogs. 2006 , 1758, 1203-14		40
590	Chemical synthesis and structure elucidation of bovine kappa-casein (1-44). <i>Biochemical and Biophysical Research Communications</i> , 2006 , 340, 1098-103	3-4	9
589	Protective cellular immunity against <i>P. falciparum</i> malaria merozoites is associated with a different P7 and P8 residue orientation in the MHC-peptide-TCR complex. 2006 , 88, 219-30		6
588	Discovery and characterization of a linear cyclotide from <i>Viola odorata</i> : implications for the processing of circular proteins. 2006 , 357, 1522-35		95
587	Structural analysis of peptides that interact with Newcastle disease virus. 2006 , 27, 1217-25		8
586	Two-, Three- and Four- Dimensional Nuclear Magnetic Resonance of Biomolecules. 2006 ,		
585	Kalata B8, a novel antiviral circular protein, exhibits conformational flexibility in the cystine knot motif. <i>Biochemical Journal</i> , 2006 , 393, 619-26	3.8	92
584	Small cationic protein from a marine turtle has beta-defensin-like fold and antibacterial and antiviral activity. 2006 , 64, 524-31		44
583	Conformational analysis of human calcitonin in solution. 2006 , 12, 51-7		4
582	Conformational analysis of a potent SSTR3-selective somatostatin analogue by NMR in water solution. 2006 , 12, 82-91		9
581	alpha-Conotoxin GI benzoylphenylalanine derivatives. (1)H-NMR structures and photoaffinity labeling of the <i>Torpedo californica</i> nicotinic acetylcholine receptor. 2006 , 273, 1373-88		13
580	Conformation and mode of membrane interaction in cyclotides. Spatial structure of kalata B1 bound to a dodecylphosphocholine micelle. 2006 , 273, 2658-72		111
579	Inhibitory properties and solution structure of a potent Bowman-Birk protease inhibitor from lentil (<i>Lens culinaris</i> , L) seeds. 2006 , 273, 4024-39		29
578	Structural evaluation of a novel pro-apoptotic peptide coupled to CNGRC tumor homing sequence by NMR. 2006 , 67, 417-24		6

577	Structural elucidation of the core-lipid A backbone from the lipopolysaccharide of <i>Acinetobacter radioresistens</i> S13, an organic solvent tolerant Gram-negative bacterium. 2006 , 341, 582-90	17
576	The structures of the cell wall teichoic acids from the thermophilic microorganism <i>Geobacillus thermoleovorans</i> strain Fango. 2006 , 341, 2613-8	3
575	Targeting retroviral Zn finger-DNA interactions: a small-molecule approach using the electrophilic nature of trans-platinum-nucleobase compounds. 2006 , 13, 539-48	55
574	Alternative O-GlcNAcylation/O-phosphorylation of Ser16 induce different conformational disturbances to the N terminus of murine estrogen receptor beta. 2006 , 13, 937-44	72
573	NMR spectroscopy of ribonucleic acids. 2006 , 48, 137-159	46
572	A putative bioactive conformation for the altered peptide ligand of myelin basic protein and inhibitor of experimental autoimmune encephalomyelitis [Arg91, Ala96] MBP87-99. 2006 , 25, 17-29	14
571	Structure elucidation and conformational study of V8: a novel synthetic non peptide AT(1) antagonist. 2006 , 40, 1097-104	15
570	¹ H, ¹³ C, ¹⁵ N NMR, ESI mass spectral and single crystal X-ray structural characterization of three spiro[pyrrolidine-2,3?-oxindoles]. 2006 , 800, 100-105	6
569	Synthesis and investigation of the spatial arrangement of the 17-acetoxy-7,8-dimethyl-3-methoxy-6-oxaestra-1,3,5(10),8(9)-tetraene. 2006 , 42, 198-205	2
568	New analogs of D-homoequilenine with substituents in the D ring. 2006 , 42, 1675-1682	
567	Contributions of the interdomain loop, amino terminus, and subunit interface to the ligand-facilitated dimerization of neurophysin: crystal structures and mutation studies of bovine neurophysin-I. 2007 , 16, 52-68	5
566	A nomenclature and data model to describe NMR experiments. 2006 , 36, 147-55	16
565	Spectroscopic Detection of Pseudo-Turns in Homodetic Cyclic Penta- and Hexapeptides Comprising Homoproline. 2006 , 12, 165-177	15
564	The influence of phosphorylation on the activity and structure of the neuronal IQ motif protein, PEP-19. 2006 , 1092, 16-27	11
563	Conformation analysis of aspartame-based sweeteners by NMR spectroscopy, molecular dynamics simulations, and X-ray diffraction studies. 2006 , 7, 377-87	6
562	Backbone dynamics of human parathyroid hormone (1-34): flexibility of the central region under different environmental conditions. 2006 , 84, 147-60	8
561	Solution structures of stomoxyn and spinigerin, two insect antimicrobial peptides with an alpha-helical conformation. 2006 , 81, 92-103	34
560	Structural Analysis of the Deep Rough Lipopolysaccharide from Gram Negative Bacterium <i>Alteromonas macleodii</i> ATCC 27126T: The First Finding of Kdo in the Inner Core of Lipopolysaccharides. 2006 , 2006, 4710-4716	15

559	Full Structural Characterisation of the Lipooligosaccharide of a Burkholderia pyrrocinia Clinical Isolate. 2006 , 2006, 4874-4883		24
558	Solution structures and behavior of trans-RuH(eta(1)-BH(4)) (binap)(1,2-diamine) complexes. 2006 , 44, 66-75		37
557	Structure and stereochemistry of products of hydroxylation of human steroid hormones by a housefly cytochrome P450 (CYP6A1). 2006 , 44, 467-74		15
556	¹ H, ¹³ C and ¹⁵ N NMR spectral characterization of twenty-seven 1,2-diaryl-(4E)-arylidene-2-imidazolin-5-ones. 2006 , 44, 895-900		1
555	The naturally occurring N6-threonyl adenine in anticodon loop of Schizosaccharomyces pombe tRNA _i causes formation of a unique U-turn motif. 2006 , 34, 2878-86		26
554	Structural and functional characterization of a mutant of Pseudocerastes persicus natriuretic peptide. <i>Protein and Peptide Letters</i> , 2006 , 13, 295-300	1.9	1
553	Structural Biology of Antimicrobial Peptides by NMR Spectroscopy. 2006 , 10, 569-581		27
552	Toward a better understanding of the basis of the molecular mimicry of polysaccharide antigens by peptides: the example of Shigella flexneri 5a. <i>Journal of Biological Chemistry</i> , 2006 , 281, 2317-32	5.4	34
551	The active conformation of avilamycin A is conferred by AviX12, a radical AdoMet enzyme. <i>Journal of Biological Chemistry</i> , 2006 , 281, 14756-63	5.4	21
550	The structures of glycolipids isolated from the highly thermophilic bacterium Thermus thermophilus Samu-SA1. 2006 , 16, 766-75		31
549	Probing the structure-function relationship of polyene macrolides: engineered biosynthesis of soluble nystatin analogues. 2006 , 49, 2431-9		32
548	Comparison of NMR structures and model-membrane interactions of 15-residue antimicrobial peptides derived from bovine lactoferricin. 2006 , 84, 312-26		31
547	Characterization of a defensin from the oyster Crassostrea gigas. Recombinant production, folding, solution structure, antimicrobial activities, and gene expression. <i>Journal of Biological Chemistry</i> , 2006 , 281, 313-23	5.4	147
546	Solution structure and antiestrogenic activity of the unique C-terminal, NR-box motif-containing region of MTA1s. <i>Journal of Biological Chemistry</i> , 2006 , 281, 25612-21	5.4	13
545	Chromatographic and Electrophoretic Separations Combined with Mass Spectrometry for Metabonomics. 2007 , 149-169		1
544	High-Resolution Magic Angle Spinning NMR Spectroscopy. 2007 , 113-147		6
543	Full structural characterization of Shigella flexneri M90T serotype 5 wild-type R-LPS and its delta galU mutant: glycine residue location in the inner core of the lipopolysaccharide. 2008 , 18, 260-9		18
542	Glycosylinositolphosphoceramides in Aspergillus fumigatus. 2008 , 18, 84-96		44

541	On the origin of some cubebene derivatives Diels-Alder adducts and the diene structures of solidago compounds. 2007 , 85, 996-1005	1
540	NMR structure and CD titration with metal cations of human prion alpha2-helix-related peptides. 2007 , 2007, 10720	7
539	A study of 7-deaza-2'-deoxyguanosine 2'-deoxycytidine base pairing in DNA. 2007 , 35, 6181-95	33
538	Structural modifications to a high-activity binding peptide located within the PfEMP1 NTS domain induce protection against <i>P. falciparum</i> malaria in Aotus monkeys. 2007 , 388, 25-36	10
537	RNA recognition mechanism of the minimal active domain of the human immunodeficiency virus type-2 nucleocapsid protein. 2007 , 141, 269-77	8
536	Pancreatic Trypsin Inhibitor. 2007 ,	
535	Peptides and Polypeptides. 2007 ,	
534	Ernst, Richard R.: The Success Story of Fourier Transformation in NMR. 2007 ,	
533	Synthetic vaccine update: applying lessons learned from recent SPf66 malarial vaccine physicochemical, structural and immunological characterization. 2007 , 25, 4487-501	10
532	Immunological and structural characterization of an epitope from the <i>Trypanosoma cruzi</i> KMP-11 protein. 2007 , 28, 1520-6	13
531	Structure-function studies of peptides for cell adhesion inhibition: identification of key residues by alanine mutation and peptide-truncation approach. 2007 , 28, 1498-508	5
530	Antibodies induced by <i>Plasmodium falciparum</i> merozoite surface antigen-2-designed pseudopeptides possess neutralizing properties of the in vitro malarial infection. 2007 , 28, 1954-65	5
529	EXPERIMENTAL 1H NMR METHODS. 2007 , 405-532	1
528	3D-Structure of the interior fusion peptide of HG/GBV-C by 1H NMR, CD and molecular dynamics studies. 2007 , 465, 187-96	4
527	NMR structural studies of the antibiotic lipopeptide daptomycin in DHPC micelles. 2007 , 1768, 3116-26	53
526	BTM-P1 polycationic peptide biological activity and 3D-dimensional structure. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 353, 908-14	3-4 17
525	New analogues of the potent cytotoxic saponin OSW-1. 2007 , 50, 3667-73	42
524	Solution structure of agelenin, an insecticidal peptide isolated from the spider <i>Agelena opulenta</i> , and its structural similarities to insect-specific calcium channel inhibitors. 2007 , 581, 3789-94	8

523	Proton-coupled electron transfer in a biomimetic peptide as a model of enzyme regulatory mechanisms. 2007 , 129, 4393-400	58
522	Principles and Applications of Multiple-Quantum Nmr. 2007 , 1-152	70
521	Structural characterization and biological evaluation of small interfering RNAs containing cyclohexenyl nucleosides. 2007 , 129, 9340-8	39
520	Structural analysis of immunotherapeutic peptides for autoimmune Myasthenia gravis. 2007 , 46, 14987-95	4
519	The structure of a novel insect peptide explains its Ca ²⁺ channel blocking and antifungal activities. 2007 , 46, 13733-41	26
518	Structures of pahayokolides A and B, cyclic peptides from a <i>Lyngbya</i> sp. 2007 , 70, 730-5	41
517	Stereochemistry modulates the stability of reduced interstrand cross-links arising from R- and S-alpha-CH ₃ -gamma-OH-1,N ₂ -propano-2'-deoxyguanosine in the 5'-CpG-3' DNA sequence. 2007 , 46, 2608-21	16
516	Structural approach to a novel tandem repeat DNA-binding domain, STPR, by CD and NMR. 2007 , 46, 1703-13	9
515	Structure and dynamics of surfactin studied by NMR in micellar media. 2007 , 129, 1968-77	40
514	DNA sequence modulates the conformation of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the <i>NarI</i> restriction enzyme. 2007 , 46, 8498-516	34
513	Laspartomycin, an acidic lipopeptide antibiotic with a unique peptide core. 2007 , 70, 443-6	26
512	Structure-based rational design of beta-hairpin peptides from discontinuous epitopes of cluster of differentiation 2 (CD2) protein to modulate cell adhesion interaction. 2007 , 50, 4038-47	16
511	Structure of the 1,N ₂ -etheno-2'-deoxyguanosine adduct in duplex DNA at pH 8.6. 2007 , 20, 1601-11	15
510	Structure and mechanism of action of the antimicrobial peptide piscidin. 2007 , 46, 1771-8	122
509	Retrocyclin-2: structural analysis of a potent anti-HIV theta-defensin. 2007 , 46, 9920-8	41
508	Common and divergent structural features of a series of corticotropin releasing factor-related peptides. 2007 , 129, 16102-14	46
507	Rational design, structure, and biological evaluation of cyclic peptides mimicking the vascular endothelial growth factor. 2007 , 50, 5135-46	31
506	Novel peptide inhibitors of human secretory phospholipase A ₂ with antiinflammatory activity: solution structure and molecular modeling. 2007 , 50, 5938-50	19

505	Determination of the conformation of the human VDAC1 N-terminal peptide, a protein moiety essential for the functional properties of the pore. 2007 , 8, 744-56	58
504	Structure of the alpha-homo-DNA:RNA duplex and the function of twist and slide to catalogue nucleic acid duplexes. 2007 , 13, 90-8	10
503	The complete structure and pro-inflammatory activity of the lipooligosaccharide of the highly epidemic and virulent gram-negative bacterium Burkholderia cenocepacia ET-12 (strain J2315). 2007 , 13, 3501-11	60
502	Biological and structural characterization of new linear gomesin analogues with improved therapeutic indices. 2007 , 88, 386-400	52
501	Role of Asn(2) and Glu(7) residues in the oxidative folding and on the conformation of the N-terminal loop of apamin. 2007 , 86, 447-62	9
500	Conformational solution studies of the SDS micelle-bound 11-28 fragment of two Alzheimer's beta-amyloid variants (E22K and A21G) using CD, NMR, and MD techniques. 2007 , 87, 23-39	15
499	Astressin-amide and stressin-acid are structurally different in dimethylsulfoxide. 2007 , 87, 196-205	7
498	Secondary structure inducing potential of beta-amino acids: torsion angle clustering facilitates comparison and analysis of the conformation during MD trajectories. 2007 , 88, 829-39	12
497	The Outer Membrane of the Marine Gram-Negative Bacterium Alteromonas addita is Composed of a Very Short-Chain Lipopolysaccharide with a High Negative Charge Density. 2007 , 2007, 1113-1122	12
496	Solution Structure and Behavior of Benzophenone-based Achiral Bisphosphine Ligands in Noyori-Type Ru(II)-Catalysts. 2007 , 25, 1163-1170	10
495	NMR structure of the peptidyl transferase RNA inhibitor antibiotic amicetin. 2007 , 45, 133-41	6
494	Chiral solvating properties of (S)-1-benzyl-6-methylpiperazine-2,5-dione. 2007 , 18, 464-475	41
493	Synthesis and biological evaluation of cyclic and branched peptide analogues as ligands for cholecystokinin type 1 receptor. 2007 , 15, 5845-53	6
492	The O-specific polysaccharide structure from the lipopolysaccharide of the Gram-negative bacterium Raoultella terrigena. 2007 , 342, 1514-8	15
491	Structural transition of a 15 amino acid residue peptide induced by GM1. 2007 , 342, 1895-903	12
490	Novel lithocholaphanes: Syntheses, NMR, MS, and molecular modeling studies. 2007 , 846, 65-73	21
489	Comparative NMR and IR spectral, X-ray structural and theoretical studies of eight 6-arylidenedibenzo[b,e]thiepin-11-one-5,5-dioxides. 2007 , 839, 94-98	6
488	Optimization of homonuclear 2D NMR for fast quantitative analysis: application to tropine-nortropine mixtures. 2007 , 43, 1243-8	36

487	NMR structure determination of a synthetic analogue of bacillomycin Lc reveals the strategic role of L-Asn1 in the natural iturinic antibiotics. 2007 , 67, 1374-81	30
486	Current analytical methods to study plant water extracts: the example of two mushrooms species, <i>Inonotus hispidus</i> and <i>Sparassis crispa</i> . 2007 , 18, 33-41	5
485	The solution structure of horseshoe crab antimicrobial peptide tachystatin B with an inhibitory cystine-knot motif. 2007 , 13, 269-79	18
484	Structure-function relationship studies of PTH(1-11) analogues containing sterically hindered dipeptide mimetics. 2007 , 13, 504-12	11
483	[Synthesis, structure, and biological properties of some 8 α -analogues of steroid estrogens with fluorine in position 2]. 2007 , 33, 315-23	6
482	[An NMR study of the conformational mobility of 7 α ,8 α analogues of steroid estrogens]. 2007 , 33, 324-31	8
481	Peptide synthesis: chemical or enzymatic. 2007 , 10, 0-0	124
480	GIAO/DFT ¹³ C NMR Chemical Shifts of 1,3,4-Thiadiazoles. 2007 , 182, 2217-2225	5
479	Divalent cation coordination and mode of membrane interaction in cyclotides: NMR spatial structure of ternary complex Kalata B7/Mn ²⁺ /DPC micelle. 2008 , 102, 1246-56	52
478	Investigation of the conformational transformations in some 7 β -methyl-6-oxa-14 β -analogues of steroidal estrogens by NMR spectroscopy. 2008 , 44, 654-665	2
477	Effect of divalent cations on the structure of the antibiotic daptomycin. 2008 , 37, 421-33	74
476	Complete hydrolysis of myo-inositol hexakisphosphate by a novel phytase from <i>Debaryomyces castellii</i> CBS 2923. 2008 , 78, 47-53	34
475	Early intermediates in the folding of dihydrofolate reductase from <i>Escherichia coli</i> detected by hydrogen exchange and NMR. 1995 , 4, 167-77	52
474	Comparative modeling of the three-dimensional structure of type II antifreeze protein. 1995 , 4, 460-71	39
473	The confirmation of the denatured structure of pyrrolidone carboxyl peptidase under non-denaturing conditions: difference in helix propensity of two synthetic peptides with single amino acid substitution. 2008 , 71, 737-42	1
472	Structural properties of the DNA-bound form of a novel tandem repeat DNA-binding domain, STPR. 2008 , 72, 414-26	8
471	Solution conformations of bradykinin antagonists modified with C α -C α cyclized nonaromatic residues. 2008 , 14, 819-29	2
470	The structure of S100A11 fragment explains a local structural change induced by phosphorylation. 2008 , 14, 1129-38	4

- 469 Conformational studies of vasopressin and mesotocin using NMR spectroscopy and molecular modelling methods. Part I: Studies in water. **2008**, 14, 76-84 18
- 468 Kissper, a kiwi fruit peptide with channel-like activity: structural and functional features. **2008**, 14, 742-54 32
- 467 Conformational analysis of a dermatan sulfate-derived tetrasaccharide by NMR, molecular modeling, and residual dipolar couplings. **2008**, 9, 240-52 31
- 466 The acylation and phosphorylation pattern of lipid A from *Xanthomonas campestris* strongly influence its ability to trigger the innate immune response in *Arabidopsis*. **2008**, 9, 896-904 49
- 465 Synthesis and biological properties of cylindramide derivatives: evidence for calcium-dependent cytotoxicity of tetramic acid lactams. **2008**, 9, 2474-86 26
- 464 Structural Elucidation with NMR Spectroscopy: Practical Strategies for Organic Chemists. **2008**, 2008, 2671-2688 80
- 463 The Structure of the O-Chain Polysaccharide from the Gram-Negative Endophytic Bacterium *Burkholderia phytofirmans* Strain PsJN. **2008**, 2008, 2303-2308 9
- 462 Structural propensities in the heme binding region of apocytochrome b5. I. Free peptides. **2008**, 90, 544-55 5
- 461 Three-dimensional consensus structure of sst2-selective somatostatin (SRIF) antagonists by NMR. **2008**, 89, 1077-87 10
- 460 The structure of the O-specific polysaccharide from the lipopolysaccharide of *Pseudomonas* sp. OX1 cultivated in the presence of the azo dye Orange II. **2008**, 343, 674-84 9
- 459 High-resolution NMR analysis of the conformations of native and base analog substituted retroviral and LTR-retrotransposon PPT primers. **2008**, 15, 254-62 15
- 458 Peptidoglycan and muropeptides from pathogens *Agrobacterium* and *Xanthomonas* elicit plant innate immunity: structure and activity. **2008**, 15, 438-48 113
- 457 Solid state structural studies of five bile acid derivatives. **2008**, 886, 197-206 12
- 456 Structural and functional consequences of the presence of a fourth disulfide bridge in the scorpion short toxins: solution structure of the potassium channel inhibitor HsTX1. **1999**, 8, 2672-85 48
- 455 NMR analysis of cleaved *Escherichia coli* thioredoxin (1-73/74-108) and its P76A variant: cis/trans peptide isomerization. **2000**, 9, 20-8 17
- 454 Conservative mutation Met8 --> Leu affects the folding process and structural stability of squash trypsin inhibitor CMTI-I. **2000**, 9, 273-9 5
- 453 Cloning, expression, purification, and preliminary characterization of a putative hemoglobin from the cyanobacterium *Synechocystis* sp. PCC 6803. **2000**, 9, 587-97 39
- 452 Elucidation of the solution structure of cardiotoxin analogue V from the Taiwan cobra (*Naja naja atra*)--identification of structural features important for the lethal action of snake venom cardiotoxins. **2000**, 9, 637-46 36

451	Conformational studies of highly potent 1-aminocyclohexane-1-carboxylic acid substituted V2 vasopressin agonists. 2008 , 66, 30-40	3
450	C-mannosylation in the hypertrehalosaemic hormone from the stick insect <i>Carausius morosus</i> . 2008 , 275, 1163-73	36
449	Purification and structural characterization of a D-amino acid-containing conopeptide, conomarphin, from <i>Conus marmoreus</i> . 2008 , 275, 1976-87	38
448	Bioactive N-terminal undecapeptides derived from parathyroid hormone: the role of alpha-helicity. 2005 , 65, 23-35	20
447	Conformational analysis by NMR and distance geometry techniques of a peptide mimetic of the third helix of the Antennapedia homeodomain. 2005 , 65, 200-8	
446	Structure-function studies of the functional and binding epitope of the human 37 kDa laminin receptor precursor protein. 2005 , 66, 9-18	18
445	¹ H Nuclear-Magnetic-Resonance Investigation of Oxidized Fe ₄ S ₄ Ferredoxin from <i>Thermotoga maritima</i> . <i>FEBS Journal</i> , 2008 , 229, 658-668	
444	Comparative Study on the Conformation of Phalloidin, Viroisin, and Related Derivatives in Aqueous Solution. <i>FEBS Journal</i> , 2008 , 232, 726-736	2
443	Structural Characterization of a Short Peptide Fragment that Mediates Estrogen-Receptor Dimerization. <i>FEBS Journal</i> , 2008 , 230, 879-885	
442	A novel beta-defensin structure: a potential strategy of big defensin for overcoming resistance by Gram-positive bacteria. 2008 , 47, 10611-9	30
441	Study Cases of Enzymatic Processes. 2008 , 253-378	3
440	Following fungal melanin biosynthesis with solid-state NMR: biopolymer molecular structures and possible connections to cell-wall polysaccharides. 2008 , 47, 4701-10	70
439	Enzyme Biocatalysis. 2008 ,	94
438	The structure of cyanine 5 terminally attached to double-stranded DNA: implications for FRET studies. 2008 , 47, 7857-62	85
437	The Arctic mutation alters helix length and type in the 11-28 beta-amyloid peptide monomer-CD, NMR and MD studies in an SDS micelle. 2008 , 164, 199-209	6
436	Syntheses and structural study of bile acid amidoalcohols. 2008 , 73, 1228-41	13
435	Structural characterisation of sporozoite components for a multistage, multi-epitope, anti-malarial vaccine. 2008 , 40, 543-57	20
434	HLA-DR allele reading register shifting is associated with immunity induced by SERA peptide analogues. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 372, 114-20	3-4 2

433	NMR structure of mussel mytilin, and antiviral-antibacterial activities of derived synthetic peptides. 2008 , 32, 227-38		56
432	Steroidal saponins. 2008 , 89, 45-141		26
431	RNA structure determination by NMR. <i>Methods in Molecular Biology</i> , 2008 , 452, 29-61	1.4	52
430	Three-dimensional structure of conotoxin tx3a: An m-1 branch peptide of the M-superfamily. 2008 , 47, 2826-32		14
429	Solution structure of a conformationally restricted fully active derivative of the human relaxin-like factor. 2008 , 47, 13308-17		4
428	3'-Intercalation of a N2-dG 1R-trans-anti-benzo[c]phenanthrene DNA adduct in an iterated (CG) ₃ repeat. 2008 , 21, 1348-58		9
427	Ring size in octreotide amide modulates differently agonist versus antagonist binding affinity and selectivity. 2008 , 51, 2676-81		20
426	The solution structure of a tetraheme cytochrome from <i>Shewanella frigidimarina</i> reveals a novel family structural motif. 2008 , 47, 11973-80		18
425	Design of a conformationally defined and proteolytically stable circular mimetic of brain-derived neurotrophic factor. <i>Journal of Biological Chemistry</i> , 2008 , 283, 33375-83	5.4	39
424	A new nerve growth factor-mimetic peptide active on neuropathic pain in rats. 2008 , 28, 2698-709		89
423	Structure-function analysis of the THAP zinc finger of THAP1, a large C2CH DNA-binding module linked to Rb/E2F pathways. <i>Journal of Biological Chemistry</i> , 2008 , 283, 4352-63	5.4	62
422	NMR of membrane-associated peptides and proteins. 2008 , 9, 50-69		52
421	Conversion shift of D-fructose to D-psicose for enzyme-catalyzed epimerization by addition of borate. 2008 , 74, 3008-13		53
420	Multidimensional Nuclear Magnetic Resonance of Biomolecules. 2008 ,		
419	Cell wall beta-(1,6)-glucan of <i>Saccharomyces cerevisiae</i> : structural characterization and in situ synthesis. <i>Journal of Biological Chemistry</i> , 2009 , 284, 13401-13412	5.4	92
418	C-terminal elongation of growth-blocking peptide enhances its biological activity and micelle binding affinity. <i>Journal of Biological Chemistry</i> , 2009 , 284, 29625-34	5.4	4
417	First structural characterization of <i>Burkholderia vietnamiensis</i> lipooligosaccharide from cystic fibrosis-associated lung transplantation strains. 2009 , 19, 1214-23		15
416	Structure-function relationship of Vpr: biological implications. 2009 , 7, 184-210		36

415	Conformational study of a somatostatin analogue by high-field n.m.r. spectroscopy, in aqueous solution. 1987 , 30, 541-7	4
414	Peptide conformation. 2009 , 31, 481-498	9
413	Rapid determination and NMR assignments of antiparallel sheets and helices of a scorpion and a cobra toxin. 1990 , 36, 227-30	10
412	Cyclic hexapeptides related to somatostatin. Conformational analysis employing ¹ H-NMR and molecular dynamics. 1990 , 36, 418-32	21
411	Determination of the structure of [Nle7]-endothelin by ¹ H NMR. 1991 , 37, 315-24	31
410	Conformation of a neurokinin antagonist in solution. 2009 , 39, 497-505	10
409	Pituitary adenylate cyclase activating polypeptide (PACAP) with 27 residues. 2009 , 40, 456-464	25
408	Solution conformation of endothelin and point mutants by nuclear magnetic resonance spectroscopy. 1992 , 40, 515-23	22
407	NMR study on solution structure of the site-specific mutant Leu48----Ala transforming growth factor alpha. 1992 , 39, 111-6	1
406	Reduced peptide bond cyclic somatostatin based opioid octapeptides. Synthesis, conformational properties and pharmacological characterization. 1992 , 39, 401-14	14
405	NMR and molecular modeling characterization of RGD containing peptides. 1992 , 39, 63-76	41
404	Spontaneous degradation via diketopiperazine formation of peptides containing a tetrahydroisoquinoline-3-carboxylic acid residue in the 2-position of the peptide sequence. 1993 , 41, 313-6	37
403	Solution conformation of Leu27 hGRF(1-32)NH ₂ and its deamidation products by 2D NMR. 1993 , 42, 24-32	13
402	Cyclic V3-loop-related HIV-1 conjugate vaccines. Synthesis, conformation and immunological properties. 1993 , 41, 455-66	33
401	Synthesis and solution conformation of c(D-Trp-D-Cys(SO ₃ -Na ⁺)-Pro-D-Val-Leu), a potent endothelin-A receptor antagonist. 1993 , 42, 194-203	8
400	Conformation of a biologically active C-terminal hexapeptide analog of the pheromone biosynthesis activating neuropeptide by NMR spectroscopy. 1994 , 43, 277-83	13
399	NMR spectra and restrained molecular dynamics of the mushroom toxin viroisin. 1994 , 43, 393-401	8
398	Solvent effects on the conformation of cyclo(-D-Trp-D-Asp-Pro-D-Val-Leu-). An NMR spectroscopy and molecular modeling study. 1994 , 43, 454-62	15

397	1H NMR analysis of fibril-forming peptide fragments of transthyretin. 1994 , 44, 388-98	26
396	The critical C-terminus of the small subunit of herpes simplex virus ribonucleotide reductase is mobile and conformationally similar to C-terminal peptides. 1994 , 44, 549-55	7
395	Conformational studies of a synthetic cyclic decapeptide fragment of rat transforming growth factor-alpha. 1995 , 46, 88-96	2
394	Synthesis, crystal structure and conformation in solution of four stereoisomeric cyclo-lanthionine derivatives. 1995 , 46, 290-301	13
393	Solution structure of a human calcitonin analog elucidated by NMR and distance geometry calculations. 1995 , 45, 305-11	8
392	Effect of trifluoroethanol on the solution structure and flexibility of desmopressin: a two-dimensional NMR study. 1995 , 45, 471-81	9
391	Solution structure of the B-chain of insulin as determined by 1H NMR spectroscopy. Comparison with the crystal structure of the insulin hexamer and with the solution structure of the insulin monomer. 1995 , 46, 424-33	25
390	1H and 13C NMR assignments and molecular modelling of a minor groove DNA-binding peptide from the HMG-I protein. 1995 , 45, 554-60	20
389	Conformational behaviour of a synthetic peptide of the C-terminus of villin that interacts with actin: an NMR, CD and stimulated annealing study. 1995 , 45, 574-86	7
388	Solution conformation of bovine lens alpha- and betaB2-crystallin terminal extensions. 1996 , 47, 9-19	3
387	Solution conformation of a peptide corresponding to residues 151-172 of HIV-1 integrase using NMR and CD spectroscopy. 1996 , 47, 117-22	7
386	Synthesis and conformational analysis of cyclic pentapeptide endothelin antagonists. 1996 , 48, 229-39	7
385	Backbone--side-chain interactions in serine. Synthesis, crystal structure and solution conformation of a linear model peptide N-Boc-L-Ser-L-Phe-OCH3. 1996 , 48, 299-303	2
384	Several polyhydroxymonoamide renin inhibitors assume similar conformations in the unbound and renin-bound states. 1996 , 48, 401-10	1
383	NMR solution structure of the receptor binding domain of Pseudomonas aeruginosa pilin strain P1. Identification of a beta-turn. 1996 , 48, 539-52	11
382	Evidence for a C-terminal turn in PBAN. An NMR and distance geometry study. 1996 , 47, 361-8	13
381	Conformational studies of an amphipathic peptide corresponding to human apolipoprotein A-II residues 18-30 with a C-terminal lipid binding motif EWLNS. 1996 , 48, 21-30	5
380	Lability of N-alkylated peptides towards TFA cleavage. 1996 , 47, 182-9	31

379	Solution structure of the phosphorylated sites of ribosomal protein S6 by 1H NMR spectroscopy. 1996 , 47, 282-8	1
378	Structure-function relationship studies of PTH(1-11) analogues containing D-amino acids. 2009 , 611, 1-7	12
377	Structural study and conformational behavior of the two different lipopolysaccharide O-antigens produced by the cystic fibrosis pathogen Burkholderia multivorans. 2009 , 15, 7156-66	18
376	A folded and functional synthetic PA1b: an interlocked entomotoxic miniprotein. 2009 , 92, 436-44	14
375	The structure of the O-specific polysaccharide from the lipopolysaccharide of Burkholderia anthina. 2009 , 344, 1697-700	11
374	The structure and purity of a reference dye standard used for quantification of C.I. Solvent Red 164 in fuels. 2009 , 82, 307-315	4
373	Synthesis, conformational analysis and immunological activity of beta3Phe-substituted Cyclolinopeptide A analogues. 2009 , 15, 166-74	7
372	Synthesis and receptor binding of opioid peptide analogues containing beta3-homo-amino acids. 2009 , 15, 777-82	14
371	Identification of a new antimicrobial lysine-rich cyclolipopeptide family from Xenorhabdus nematophila. 2009 , 62, 295-302	62
370	The structural and biological significance of the EAEEAE insert in the alpha-domain of human neuronal growth inhibitory factor. 2009 , 276, 3547-58	5
369	NMR and circular dichroism studies on the conformation of a 44-mer peptide from a CD4-binding domain of human immunodeficiency virus envelope glycoprotein. 1997 , 49, 432-43	2
368	Conformational analysis of cyclo(2,9)-Ac-QCRSVEGSCG-OH from the C-terminal loop of human growth hormone. 1997 , 49, 15-22	3
367	Inhibition of homotypic adhesion of T-cells: secondary structure of an ICAM-1-derived cyclic peptide. 1997 , 49, 517-26	19
366	Comparison of the solution conformations of a human immunodeficiency virus peptidomimetic and its retro-inverso isomer using 1H NMR spectroscopy. 1997 , 50, 421-35	14
365	Structural studies on endothelin receptor subtype B specific agonist IRL 1620 [suc-[Glu9, Ala11,15]ET-1(8-21)] and its analogs with dipalmitoyl phosphatidylcholine vesicles by NMR spectroscopy. 1998 , 51, 155-64	3
364	Solution conformational analysis of sodium complexed [Gly6]- and [Gly9]-antamanide analogs. 1998 , 51, 180-7	12
363	Chemical synthesis of dendrotoxin-I: revision of the reported structure. 1998 , 51, 355-64	13
362	Secondary interactions as driving force in heterocomplex formation of 2,7-disubstituted-1,8-naphthyridines: Quantum chemical, NMR and mass spectral investigations. 2009 , 931, 60-67	7

361	Characterization of glucuronic acid containing glycolipid in <i>Aspergillus fumigatus</i> mycelium. 2009 , 344, 1960-7	18
360	Structural basis for alpha-conotoxin potency and selectivity. 2009 , 17, 5894-9	9
359	Solution structure of Cn5, a crustacean toxin found in the venom of the scorpions <i>Centruroides noxius</i> and <i>Centruroides suffusus suffusus</i> . 2009 , 1794, 1591-8	9
358	Solution structure of a K(+) -channel blocker from the scorpion <i>Tityus cambridgei</i> . 2002 , 11, 390-400	11
357	Novel, potent, and radio-iodinatable somatostatin receptor 1 (sst1) selective analogues. 2009 , 52, 2733-46	35
356	An essential epitope of anti-MUC1 monoclonal antibody KL-6 revealed by focused glycopeptide library. 2009 , 131, 17102-9	65
355	Design of beta-hairpin peptides for modulation of cell adhesion by beta-turn constraint. 2009 , 52, 726-36	12
354	Conformational and dynamics simulation study of antimicrobial peptide hedistin-heterogeneity of its helix-turn-helix motif. 2009 , 1788, 2497-508	4
353	Structural characteristics of immunogenic liver-stage antigens derived from <i>P. falciparum</i> malarial proteins. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 384, 455-60	3-4 9
352	Solution structure of Hyp10Pro variant of conomorphin, a cysteine-free and D-amino-acid containing conopeptide. 2009 , 54, 153-60	13
351	Structure and function of a custom anticancer peptide, CB1a. 2009 , 30, 839-48	56
350	Synthesis and antimicrobial activity of truncated fragments and analogs of citropin 1.1: The solution structure of the SDS micelle-bound citropin-like peptides. 2009 , 168, 250-8	16
349	Correlations through the chemical bond I: Homonuclear shift correlation. 2009 , 27, 129-188	0
348	Model development for the viral Kcv potassium channel. 2009 , 96, 485-98	34
347	A de novo redesign of the WW domain. 2003 , 12, 2194-205	49
346	Chapter 1 NMR of Antimicrobial Peptides. 2009 , 65, 1-51	19
345	The stathmin-derived I19L peptide interacts with FtsZ and alters its bundling. 2009 , 48, 9734-44	10
344	chi-Conopeptide pharmacophore development: toward a novel class of norepinephrine transporter inhibitor (Xen2174) for pain. 2009 , 52, 6991-7002	63

343	A novel beta-defensin structure: big defensin changes its N-terminal structure to associate with the target membrane. 2009 , 48, 7629-35		14
342	Chapter 3 1D Double Quantum Filter NMR Studies. 2009 , 265-329		5
341	Screening of flavonoids as candidate antibiotics against <i>Enterococcus faecalis</i> . 2009 , 72, 719-24		54
340	Chapter 1 Quantitative 2D NMR Studies. 2009 , 66, 1-31		32
339	Functional neoglycopeptides: synthesis and characterization of a new class of MUC1 glycoprotein models having core 2-based O-glycan and complex-type N-glycan chains. 2009 , 48, 11117-33		32
338	Oligopyridine-ruthenium(II)-amino acid conjugates: synthesis, characterization, DNA binding properties and interactions with the oligonucleotide duplex d(5'-CGCGCG-3') ₂ . 2009 , 6403-15		16
337	Psi[CH(CF ₃)NH]Gly-peptides: synthesis and conformation analysis. 2009 , 7, 2286-96		39
336	Synthesis and structure of some 8 β -analogs of steroid estrogens. 2010 , 80, 1324-1330		5
335	Redox linked conformational changes in cytochrome c3 from <i>Desulfovibrio desulfuricans</i> ATCC 27774. 2010 , 49, 9620-9		15
334	Chiral eta(6)-arene/N-tosylethylenediamine-ruthenium(II) complexes: solution behavior and catalytic activity for asymmetric hydrogenation. 2010 , 5, 806-16		33
333	Solution structure of miRNA:mRNA complex. <i>Methods in Molecular Biology</i> , 2010 , 667, 251-65	1.4	1
332	A new non-natural arginine-like amino acid derivative with a sulfamoyl group in the side-chain. 2010 , 38, 691-700		21
331	Synthesis and structural studies of new analogues of PTH(1-11) containing C β -tetra-substituted amino acids in position 8. 2010 , 39, 1369-79		7
330	β -Helix peptides designed from EBV-gH protein display higher antigenicity and induction of monocyte apoptosis than the native peptide. 2010 , 39, 1507-19		
329	Structural Elucidation of a Novel <i>B. cenocepacia</i> ET-12 Lipooligosaccharide Isolated from a Cystic Fibrosis Patient after Lung Transplantation. 2010 , 2010, 1299-1306		6
328	A Urea-Linked Glucosamine Dimer as a Building Block for the Synthesis of Linear and Cyclic Neosaccharides. 2010 , 2010, 4062-4074		3
327	Full Structural Characterization of an Extracellular Polysaccharide Produced by the Freshwater Cyanobacterium <i>Oscillatoria planktothrix</i> FP1. 2010 , 2010, 5594-5600		13
326	Structure determination of the top-loop of the conserved 3'-terminal secondary structure in the genome of flaviviruses. 2010 , 11, 1404-12		4

325	Total synthesis, characterization, and conformational analysis of the naturally occurring hexadecapeptide integramide A and a diastereomer. 2010 , 16, 316-27	18
324	A doppel alpha-helix peptide fragment mimics the copper(II) interactions with the whole protein. 2010 , 16, 6212-23	24
323	Structural analysis of a helical peptide unfolding pathway. 2010 , 16, 5400-7	23
322	Heterocyclisations of 3-Trifluoroacetyl-1-Methyl-Lactams with 1,4-Bis-Nucleophiles. 2010 , 103, 243-255	5
321	SYNTHESIS OF 5-HYDROXY-3-TRIFLUOROMETHYL-PYRAZOLES BY RING OPENING OF 3-TRIFLUOROACETYL-BENZOLACTAMS. 2010 , 103, 655-664	3
320	New Trifluoromethylated Imidazolidines, Oxazolidines and Thiazolidines Starting from 3-(1,1-Dihydroxy-2,2,2-Trifluoroethyl)-1-Methyl-Tetrahydroazepin-2-one and β -(1,1-Dihydroxy-2,2,2-Trifluoroethyl)- β -Butyrolactone. 2010 , 104, 707-716	4
319	Configurational assignment of D- and L-isovalines in intact, natural, and synthetic peptides by 2D-NMR spectroscopy. 2010 , 7, 1612-24	9
318	Potential allosteric modulators of the proteasome activity. 2010 , 93, 481-95	22
317	Structural and biochemical characteristics of the cyclotide kalata B5 from <i>Oldenlandia affinis</i> . 2010 , 94, 647-58	22
316	Ultrafast 2D COSY with constant-time phase-modulated spatial encoding. <i>Journal of Magnetic Resonance</i> , 2010 , 204, 82-90	3 24
315	NMR spectroscopy and computational analysis of interaction between <i>Serratia marcescens</i> chitinase B and a dipeptide derived from natural-product cyclopentapeptide chitinase inhibitor argifin. 2010 , 18, 5835-44	8
314	NMR studies of new arginine vasopressin analogs modified with alpha-2-indanylglycine enantiomers at position 2 bound to sodium dodecyl sulfate micelles. 2010 , 151, 139-48	8
313	The structure of the carbohydrate backbone of the lipooligosaccharide from the halophilic bacterium <i>Arcobacter halophilus</i> . 2010 , 345, 850-3	10
312	Improved Resolution in Dipolar NMR Spectra Using Constant Time Evolution PISEMA Experiment. 2010 , 494, 104-110	8
311	Influence of bulky 3,3'-diphenylalanine enantiomers replacing position 2 of AVP analogues on their conformations: NMR and molecular modeling studies. 2010 , 45, 4065-73	8
310	Structure of a paralytic peptide from an insect, <i>Manduca sexta</i> . 1999 , 54, 256-61	25
309	Solution structure of LC5, the CCR5- derived peptide for HIV-1 inhibition. 2010 , 16, 165-70	10
308	Structure-activity relationship of a novel pentapeptide with cancer cell growth-inhibitory activity. 2010 , 16, 242-8	4

307	Peptide-peptoid hybrids based on (1-11)-parathyroid hormone analogs. 2010 , 16, 480-5	4
306	Synthesis, preferred conformation, and membrane activity of medium-length peptaibiotics: tylopeptin B. 2010 , 75, 169-81	15
305	Structure and activity of CPNGRC: a modified CD13/APN peptidic homing motif. 2010 , 75, 551-62	11
304	Against the rules: a marine bacterium, <i>Loktanella rosea</i> , possesses a unique lipopolysaccharide. 2010 , 20, 586-93	10
303	Structural characterisation of the natively unfolded enterocin EJ97. 2010 , 23, 507-18	7
302	Molecular requirements for the insecticidal activity of the plant peptide pea albumin 1 subunit b (PA1b). <i>Journal of Biological Chemistry</i> , 2010 , 285, 32689-32694	5-4 24
301	Simultaneous de novo identification of molecules in chemical mixtures by doubly indirect covariance NMR spectroscopy. 2010 , 132, 16922-7	28
300	Solution structure of GxTX-1E, a high-affinity tarantula toxin interacting with voltage sensors in Kv2.1 potassium channels. 2010 , 49, 5134-42	23
299	Selective removal of individual disulfide bonds within a potato type II serine proteinase inhibitor from <i>Nicotiana glauca</i> reveals differential stabilization of the reactive-site loop. 2010 , 395, 609-26	15
298	Structural and pharmacological characteristics of chimeric peptides derived from peptide E and beta-endorphin reveal the crucial role of the C-terminal YGGFL and YKKGE motifs in their analgesic properties. 2010 , 31, 962-72	2
297	3D structure determination of STARP peptides implicated in <i>P. falciparum</i> invasion of hepatic cells. 2010 , 28, 4989-96	7
296	Side chain cyclization based on serine residues: synthesis, structure, and activity of a novel cyclic analogue of the parathyroid hormone fragment 1-11. 2010 , 53, 8072-9	17
295	A conformationally constrained peptidomimetic binds to the extracellular region of HER2 protein. 2010 , 28, 289-308	37
294	The solution structure of (+)-spongistatin 1 in DMSO. 2010 , 12, 1788-91	20
293	Conformationally constrained peptides from CD2 to modulate protein-protein interactions between CD2 and CD58. 2011 , 54, 5307-19	32
292	Altered fatty acid metabolism in long duration road transport: An NMR-based metabolomics study in sheep. 2011 , 10, 1073-87	25
291	Ultrafast quantitative 2D NMR: an efficient tool for the measurement of specific isotopic enrichments in complex biological mixtures. 2011 , 83, 3112-9	56
290	Deconvolution of chemical mixtures with high complexity by NMR consensus trace clustering. 2011 , 83, 7412-7	48

289	Sørensen, Ole W.: Highlights from My Journey in NMR. 2011 ,		
288	Coherent-state transfer via highly mixed quantum spin chains. 2011 , 83,		48
287	Characterization of a designed vascular endothelial growth factor receptor antagonist helical peptide with antiangiogenic activity in vivo. 2011 , 54, 1391-400		34
286	Solution structure of LCI, a novel antimicrobial peptide from <i>Bacillus subtilis</i> . 2011 , 50, 3621-7		35
285	3D structure and immunogenicity of <i>Plasmodium falciparum</i> sporozoite induced associated protein peptides as components of fully-protective anti-malarial vaccine. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 416, 349-55	3-4	2
284	Structural determinants of the immunomodulatory properties of the C-terminal region of bovine β -casein. 2011 , 21, 770-776		15
283	NMR analysis of a kinetically trapped intermediate of a disulfide-deficient mutant of the starch-binding domain of glucoamylase. 2011 , 412, 304-15		1
282	Synthesis of both ionic species of ammonium dithiocarbamate derived cholic acid moieties. 2011 , 16, 6306-12		3
281	Solution structure of LC4 transmembrane segment of CCR5. <i>PLoS ONE</i> , 2011 , 6, e20452	3-7	11
280	NMR study of peplomycin in aqueous solution. Assignment of resonances by means of two-dimensional spectroscopy. 2011 , 64, 309-16		5
279	New analogs of steroid estrogens. 2011 , 47, 1234-1239		1
278	Identification, structure, and characterization of an exopolysaccharide produced by <i>Histophilus somni</i> during biofilm formation. 2011 , 11, 186		28
277	Synthesis, preferred conformation, protease stability, and membrane activity of heptaibin, a medium-length peptaibiotic. 2011 , 17, 585-94		28
276	Molecular Modeling Study of the Carbohydrate Region of the Endotoxin from <i>Burkholderia cenocepacia</i> ET-12. 2011 , 2011, 5114-5122		
275	Triple Hyp->Pro replacement in integramide A, a peptaib inhibitor of HIV-1 integrase: Effect on conformation and bioactivity. 2011 , 96, 49-59		3
274	Solution conformation of non-mammalian tachykinin physalaemin in lipid micelles by nuclear magnetic resonance. 2011 , 96, 252-9		3
273	Zinc to cadmium replacement in the <i>A. thaliana</i> SUPERMAN Cys[His]zinc finger induces structural rearrangements of typical DNA base determinant positions. 2011 , 95, 801-10		35
272	An efficient approach for the characterization of mucin-type glycopeptides: the effect of O-glycosylation on the conformation of synthetic mucin peptides. 2011 , 17, 2393-404		29

271	Targeting zinc finger domains with small molecules: solution structure and binding studies of the RanBP2-type zinc finger of RBM5. 2011 , 12, 2837-45		10
270	Design and application of robust rf pulses for toroid cavity NMR spectroscopy. <i>Journal of Magnetic Resonance</i> , 2011 , 209, 282-90	3	17
269	Deciphering the structural and biological properties of the lipid A moiety of lipopolysaccharides from Burkholderia cepacia strain ASP B 2D, in Arabidopsis thaliana. 2011 , 21, 184-94		29
268	Anti-group A streptococcal vaccine epitope: structure, stability, and its ability to interact with HLA class II molecules. <i>Journal of Biological Chemistry</i> , 2011 , 286, 6989-98	5-4	22
267	Disulfide-stabilized helical hairpin structure and activity of a novel antifungal peptide EcAMP1 from seeds of barnyard grass (Echinochloa crus-galli). <i>Journal of Biological Chemistry</i> , 2011 , 286, 25145-53	5-4	65
266	NMR structure of the A730 loop of the Neurospora VS ribozyme: insights into the formation of the active site. 2011 , 39, 4427-37		19
265	Possible structural role of the disaccharide unit in Fe-bleomycin before and after oxygen activation. 2012 , 65, 25-33		7
264	Giant DNA virus mimivirus encodes pathway for biosynthesis of unusual sugar 4-amino-4,6-dideoxy-D-glucose (Viosamine). <i>Journal of Biological Chemistry</i> , 2012 , 287, 3009-18	5-4	26
263	Decoding the functional roles of cationic side chains of the major antimicrobial region of human cathelicidin LL-37. 2012 , 56, 845-56		66
262	Three-dimensional solution structure of bottromycin A2: a potent antibiotic active against methicillin-resistant Staphylococcus aureus and vancomycin-resistant Enterococci. 2012 , 60, 169-71		15
261	Trichogin GA IV: a versatile template for the synthesis of novel peptaibiotics. 2012 , 10, 1285-99		42
260	Novel peptide foldameric motifs: a step forward in our understanding of the fully-extended conformation/3(10)-helix coexistence. 2012 , 10, 2413-21		24
259	Strategies for Structural Assignment of Marine Natural Products Through Advanced NMR-based Techniques. 2012 , 518-546		1
258	A novel β -conopeptide, CnIIIC, exerts potent and preferential inhibition of NaV1.2/1.4 channels and blocks neuronal nicotinic acetylcholine receptors. 2012 , 166, 1654-68		41
257	Solution structure of native and recombinant expressed toxin Cssl from the venom of the scorpion Centruroides suffusus suffusus, and their effects on Nav1.5 sodium channels. 2012 , 1824, 478-87		20
256	BASHD-J-resolved-COSY: a new method for measuring proton-proton spin coupling constants of multiplet signals. 2012 , 50, 713-6		3
255	Structural characterization of the cyclic cystine ladder motif of β -defensins. 2012 , 51, 9718-26		50
254	Structural basis for bifunctional zinc(II) macrocyclic complex recognition of thymine bulges in DNA. 2012 , 51, 5444-57		26

253	Structure, dynamics, and antimicrobial and immune modulatory activities of human LL-23 and its single-residue variants mutated on the basis of homologous primate cathelicidins. 2012 , 51, 653-64		44
252	Antimicrobial lipopeptaibol trichogin GA IV: role of the three Aib residues on conformation and bioactivity. 2012 , 43, 1761-77		24
251	¶Sulphate PNA (PNA S): highly selective DNA binding molecule showing promising antigene activity. <i>PLoS ONE</i> , 2012 , 7, e35774	3-7	33
250	Purification and structural characterization of siderophore (corynebactin) from <i>Corynebacterium diphtheriae</i> . <i>PLoS ONE</i> , 2012 , 7, e34591	3-7	24
249	Partial thioamide scan on the lipopeptaibiotic trichogin GA IV. Effects on folding and bioactivity. 2012 , 8, 1161-71		10
248	Structural Study of the Lipopolysaccharide O-Antigen Produced by the Emerging Cystic Fibrosis Pathogen <i>Pandora</i> sp. 2012 , 2012, 2243-2249		7
247	O-antigen and core carbohydrate of <i>Vibrio fischeri</i> lipopolysaccharide: composition and analysis of their role in <i>Euprymna scolopes</i> light organ colonization. <i>Journal of Biological Chemistry</i> , 2012 , 287, 8515-30	5-10	48
246	Protective immunity provided by a new modified SERA protein peptide: its immunogenetic characteristics and correlation with 3D structure. 2012 , 43, 183-94		4
245	Design, conformational studies and analysis of structure-function relationships of PTH (1-11) analogues: the essential role of Val in position 2. 2012 , 43, 207-18		4
244	Cyclic RGD peptides interfere with binding of the <i>Helicobacter pylori</i> protein CagL to integrins $\alpha_5\beta_1$ and $\alpha_6\beta_1$. 2012 , 43, 219-32		52
243	Structural characterization of two lipopolysaccharide O-antigens produced by the endofungal bacterium <i>Burkholderia</i> sp. HKI-402 (B4). 2012 , 347, 95-8		11
242	Conformational preferences of proline derivatives incorporated into vasopressin analogues: NMR and molecular modelling studies. 2012 , 79, 535-47		4
241	Highly potent antidiuretic antagonists: conformational studies of vasopressin analogues modified with 1-naphthylalanine enantiomers at position 2. 2012 , 79, 1033-42		3
240	Improving the double quantum filtered COSY experiment by "Moving Tube" NMR. <i>Journal of Magnetic Resonance</i> , 2012 , 219, 41-5	3	6
239	A solvent-dependent peptide spring unraveled by 2D-NMR. 2012 , 68, 4429-4433		15
238	Sanctolide A, a 14-membered PK-NRP hybrid macrolide from the cultured cyanobacterium <i>Oscillatoria sancta</i> (SAG 74.79). 2012 , 53, 3563-3567		12
237	Spectral simplification of proton homonuclear correlation experiments utilizing 1F_1 , 2F_1 fluorine decoupling. 2012 , 140, 104-106		2
236	Coordination chemistry and solution structure of Fe(II)-peplomycin. Two possible coordination geometries. 2012 , 111, 50-8		4

235	Aminoarabinose is essential for lipopolysaccharide export and intrinsic antimicrobial peptide resistance in <i>Burkholderia cenocepacia</i> (I). 2012 , 85, 962-74	79
234	Isovaline in naturally occurring peptides: A nondestructive methodology for configurational assignment. 2012 , 98, 36-49	17
233	Structural studies of seven homoisoflavonoids, six thiohomoisoflavonoids, and four structurally related compounds. 2012 , 23, 209-217	10
232	Therapeutic potential of the peptide leucine arginine as a new nonplant bowman-birk-like serine protease inhibitor. 2013 , 56, 6732-44	14
231	Micelle-bound conformations of neurohypophyseal hormone analogues modified with a C β -disubstituted residue: NMR and molecular modelling studies. 2013 , 31, 748-64	3
230	^1H and ^{13}C chemical shift assignment of the monomers that comprise carboxymethyl cellulose. <i>Carbohydrate Polymers</i> , 2013 , 97, 384-90	10.3 32
229	Molecular Biophysics for the Life Sciences. 2013 ,	1
228	Conformational preferences of zampanolide and dactylolide. 2013 , 15, 5246-9	12
227	Glycosaminoglycan Structural Characterization. 2013 ,	
226	Caulosegnins I-III: a highly diverse group of lasso peptides derived from a single biosynthetic gene cluster. 2013 , 135, 210-22	84
225	New bis-ferrocenyl end-capped peptides: synthesis and charge transfer properties. 2013 , 100, 14-24	12
224	Molecular insights into the AT1 antagonism based on biophysical and in silico studies of telmisartan. 2013 , 22, 4842-4857	8
223	Development of a ^{13}C -optimized 1.5-mm high temperature superconducting NMR probe. <i>Journal of Magnetic Resonance</i> , 2013 , 235, 58-65	3 57
222	Left-handed helical preference in an achiral peptide chain is induced by an L-amino acid in an N-terminal type II β -turn. 2013 , 78, 2248-55	35
221	Mitochondrial function in diabetes: novel methodology and new insight. 2013 , 62, 1833-42	27
220	The biolabile 2'-O-pivaloyloxymethyl modification in an RNA helix: an NMR solution structure. 2013 , 11, 2638-47	2
219	All-Thioamidated Homo- β -Peptides: Synthesis and Conformation. 2013 , 2013, 3455-3463	12
218	Chemical shift assignment of the complicated monomers comprising cellulose acetate by two-dimensional NMR spectroscopy. 2013 , 375, 136-44	16

217	NMR characterisation of a highly conserved secondary structural RNA motif of Halobacterium halobium 23S rRNA. 2013 , 11, 3382-92		2
216	The astexin-1 lasso peptides: biosynthesis, stability, and structural studies. 2013 , 20, 558-69		67
215	Basic NMR Experiments. 2013 , 53-82		
214	Spectroscopically labeled peptaibiotic analogs: the 4-nitrophenylalanine infrared absorption probe inserted at different positions into trichogin GA IV. 2013 , 19, 246-56		6
213	Chemistry and biology of the potent endotoxin from a Burkholderia dolosa clinical isolate from a cystic fibrosis patient. 2013 , 14, 1105-15		17
212	Functional evolution of scorpion venom peptides with an inhibitor cystine knot fold. 2013 , 33,		38
211	Fast and Ultrafast Quantitative 2D NMR. 2013 , 99-158		15
210	SGK196 is a glycosylation-specific O-mannose kinase required for dystroglycan function. 2013 , 341, 896-9		162
209	Novel inhibitor cystine knot peptides from Momordica charantia. <i>PLoS ONE</i> , 2013 , 8, e75334	3-7	12
208	Probing the residual structure in avian prion hexarepeats by CD, NMR and MD techniques. 2013 , 18, 11467-84		6
207	The biosynthesis of UDP-d-FucNAc-4N-(2)-oxoglutarate (UDP-Yelosamine) in Bacillus cereus ATCC 14579: Pat and Pyl, an aminotransferase and an ATP-dependent Grasp protein that ligates 2-oxoglutarate to UDP-4-amino-sugars. <i>Journal of Biological Chemistry</i> , 2014 , 289, 35620-32	5-4	13
206	Pure Shift NMR Spectroscopy. 2014 , 295-310		41
205	Concurrently optimized cooperative pulses in robust quantum control: application to broadband Ramsey-type pulse sequence elements. 2014 , 16, 115002		19
204	Communication: ultrafast homonuclear correlation spectroscopy with diagonal suppression. 2014 , 140, 231103		7
203	The 2.0 π helix in hetero-oligopeptides entirely composed of C α -disubstituted glycines with both side chains longer than methyls. 2014 , 102, 145-58		10
202	NMR based solvent exchange experiments to understand the conformational preference of intrinsically disordered proteins using FG-nucleoporin peptide as a model. 2014 , 102, 69-77		5
201	NMR Studies of Eye Lens Crystallins. 2014 , 139-152		3
200	Thermophiles as potential source of novel endotoxin antagonists: the full structure and bioactivity of the lipo-oligosaccharide from Thermomonas hydrothermalis. 2014 , 15, 2146-55		15

199	Conformational studies of neurohypophyseal hormones analogues with glycoconjugates by NMR spectroscopy. 2014 , 20, 406-14		3
198	Rapid pulsing artefacts in pulsed-field gradient double-quantum filtered COSY spectra. 2014 , 52, 329-32		4
197	Enhancement of the helical content and stability induced in a linear oligopeptide by an i, i+4 intramolecularly double stapled, overlapping, bicyclic [31, 22, 5]-(E)ene motif. 2014 , 102, 115-23		8
196	New 2-fluoro 8 β analogs of steroidal estrogens. 2014 , 50, 1520-1526		
195	Structural and conformational study of the O-polysaccharide produced by the metabolically versatile photosynthetic bacterium <i>Rhodospseudomonas palustris</i> strain BisA53. <i>Carbohydrate Polymers</i> , 2014 , 114, 384-391	10.3	10
194	Structural analysis and characterization of dextran produced by wild and mutant strains of <i>Leuconostoc mesenteroides</i> . <i>Carbohydrate Polymers</i> , 2014 , 99, 331-8	10.3	76
193	Structure and antimicrobial activity of platypus 'intermediate' defensin-like peptide. 2014 , 588, 1821-6		3
192	Base substituted 5'-O-(N-isoleucyl)sulfamoyl nucleoside analogues as potential antibacterial agents. 2014 , 22, 2875-86		14
191	Multidimensional approaches to NMR-based metabolomics. 2014 , 86, 47-57		97
190	Dietary fat, fatty acid saturation and mitochondrial bioenergetics. 2014 , 46, 33-44		37
189	NMR elucidation of the role of Mg ²⁺ in the structure and stability of the conserved RNA motifs of the EMCV IRES element. 2014 , 12, 1495-509		6
188	Covalently linked hopanoid-lipid A improves outer-membrane resistance of a <i>Bradyrhizobium</i> symbiont of legumes. 2014 , 5, 5106		67
187	Characterization of caulonodin lasso peptides revealed unprecedented N-terminal residues and a precursor motif essential for peptide maturation. 2014 , 5, 4032-4043		36
186	Synchronized and concurrent experiments in Moving Tube NMR: using separate sample volumes for different pulse sequences. <i>Journal of Magnetic Resonance</i> , 2014 , 247, 104-109		3
185	Structural dissection of the C-terminal sterile alpha motif (SAM) of human p73. 2014 , 558, 133-42		8
184	Customized metabolomics database for the analysis of NMR ¹ H- ¹ H TOCSY and ¹³ C- ¹ H HSQC-TOCSY spectra of complex mixtures. 2014 , 86, 5494-501		88
183	Chemical organization of the cell wall polysaccharide core of <i>Malassezia restricta</i> . <i>Journal of Biological Chemistry</i> , 2014 , 289, 12647-56	5-4	43
182	A novel α /7-conotoxin Lv1A from <i>Conus lividus</i> that selectively blocks α vs. β / β β nicotinic acetylcholine receptors. 2014 , 28, 1842-53		48

181	Nuclear magnetic resonance structure revealed that the human polyomavirus JC virus agnoprotein contains an α -helix encompassing the Leu/Ile/Phe-rich domain. <i>Journal of Virology</i> , 2014 , 88, 6556-75	6.6	17
180	Effect of head-to-tail cyclization on conformation of histatin-5. 2014 , 20, 952-7		4
179	Structural studies of the C-terminal 19-peptide of serum amyloid A and its Pro \rightarrow Ala variants interacting with human cystatin C. 2015 , 28, 413-26		5
178	GMDP: unusual physico-chemical and biological properties of the anomeric forms. 2015 , 21, 717-22		2
177	O-glycosylation effects on family 1 carbohydrate-binding module solution structures. 2015 , 282, 4341-56		15
176	Isolation and NMR Analysis of Antifungal Fengycin A and B from <i>Bacillus amyloliquefaciens</i> subsp. <i>plantarum</i> BC32-1. <i>Bulletin of the Korean Chemical Society</i> , 2015 , 36, 1316-1321	1.2	3
175	The Biosynthesis of UDP-D-QuiNAc in <i>Bacillus cereus</i> ATCC 14579. <i>PLoS ONE</i> , 2015 , 10, e0133790	3.7	9
174	Pushing the limits of signal resolution to make coupling measurement easier. 2015 , 51, 7939-42		22
173	Cysteine-Rich Peptide Family with Unusual Disulfide Connectivity from <i>Jasminum sambac</i> . 2015 , 78, 2791-9		10
172	Structural characterization and in vivo evaluation of β -hairpin peptidomimetics as specific CXCR4 imaging agents. 2015 , 12, 941-53		11
171	Investigating the Structural Variability and Binding Modes of the Glioma Targeting NFL-TBS.40-63 Peptide on Tubulin. 2015 , 54, 3660-9		7
170	Screening and assignment of phenylboronic acid and its anhydride formation by NMR spectroscopy. 2015 , 634, 95-97		4
169	Contribution of dihydrouridine in folding of the D-arm in tRNA. 2015 , 13, 4960-6		22
168	Fully resolved NMR correlation spectroscopy. 2015 , 21, 9044-7		43
167	^1H and ^2H NMR: Diagonal suppressed total correlation spectroscopy in spin echo mode. 2015 , 627, 58-63		2
166	Arginine-, D-arginine-vasopressin, and their inverso analogues in micellar and liposomic models of cell membrane: CD, NMR, and molecular dynamics studies. 2015 , 44, 727-43		7
165	NMR study on the interaction of the conserved CREX 'stem-loop' in the Hepatitis E virus genome with a naphthyridine-based ligand. 2015 , 13, 9665-72		1
164	Studies on the Chitin Binding Property of Novel Cysteine-Rich Peptides from <i>Alternanthera sessilis</i> . 2015 , 54, 6639-49		29

163	Synthesis and structural characterization of monomeric and dimeric peptide nucleic acids prepared by using microwave-promoted multicomponent reactions. 2015 , 13, 11052-71		5
162	Insect Gut Symbiont Susceptibility to Host Antimicrobial Peptides Caused by Alteration of the Bacterial Cell Envelope. <i>Journal of Biological Chemistry</i> , 2015 , 290, 21042-21053	5-4	32
161	Solution structure and base specificity of cytotoxic RC-RNase 2 from <i>Rana catesbeiana</i> . 2015 , 584, 70-8		2
160	Persistent cystic fibrosis isolate <i>Pseudomonas aeruginosa</i> strain RP73 exhibits an under-acylated LPS structure responsible of its low inflammatory activity. 2015 , 63, 166-75		20
159	Structural investigation of the lipopolysaccharide O-chain isolated from <i>Burkholderia fungorum</i> strain DSM 17061. 2016 , 433, 31-5		4
158	A Combined NMR and Computational Approach to Determine the RGD α 5 β Integrin Recognition Mode in Isolated Cell Membranes. 2016 , 22, 681-93		19
157	<i>Prevotella denticola</i> Lipopolysaccharide from a Cystic Fibrosis Isolate Possesses a Unique Chemical Structure. 2016 , 2016, 1732-1738		5
156	CLIP-COSY: A Clean In-Phase Experiment for the Rapid Acquisition of COSY-type Correlations. 2016 , 55, 7655-9		34
155	Multidimensional Nuclear Magnetic Resonance of Biomolecules. 2016 , 1-26		
154	Effects of the multiple O-glycosylation states on antibody recognition of the immunodominant motif in MUC1 extracellular tandem repeats. 2016 , 7, 1102-1122		26
153	Correlations Through the Chemical Bond I: Homonuclear Shift Correlation. 2016 , 203-241		1
152	CLIP-COSY: Reine Inphase-Signale und schnelle Akquisition COSY-artiger Korrelationen. 2016 , 128, 7785-7789		1
151	Structural and Conformational Study of the O-Antigenic Portion of the Lipopolysaccharide Isolated from <i>Burkholderia gladioli</i> pv. <i>cocovenenans</i> . 2016 , 2016, 748-755		3
150	Potent antidiuretic agonists, deamino-vasopressin and desmopressin, and their inverse analogs: NMR structure and interactions with micellar and liposomal models of cell membrane. 2016 , 106, 245-59		0
149	Synthetic Human NOTCH1 EGF Modules Unraveled Molecular Mechanisms for the Structural and Functional Roles of Calcium Ions and O-Glycans in the Ligand-Binding Region. 2016 , 55, 776-87		15
148	An unprecedented mechanism of nucleotide methylation in organisms containing thyX. 2016 , 351, 507-10		34
147	Relaxation Editing Using Long-Lived States and Coherences for Analysis of Mixtures. 2016 , 88, 3004-8		8
146	Rattusin structure reveals a novel defensin scaffold formed by intermolecular disulfide exchanges. <i>Scientific Reports</i> , 2017 , 7, 45282	4-9	12

145	Pure shift edited ultra high resolution NMR spectrum with complete eradication of axial peaks and unwanted couplings. <i>Journal of Magnetic Resonance</i> , 2017 , 279, 74-80	3	20
144	Differential Attenuation of NMR Signals by Complementary Ion-Exchange Resin Beads for De Novo Analysis of Complex Metabolomics Mixtures. 2017 , 23, 9239-9243		4
143	Insights into the anticancer properties of the first antimicrobial peptide from Archaea. 2017 , 1861, 2155-2164	27	
142	The Dual Activity Responsible for the Elongation and Branching of α (1,3)-Glucan in the Fungal Cell Wall. 2017 , 8,		57
141	The rare sugar -acetylated viosamine is a major component of Mimivirus fibers. <i>Journal of Biological Chemistry</i> , 2017 , 292, 7385-7394	5-4	9
140	Nuclear Magnetic Resonance Structure of the Human Polyoma JC Virus Agnoprotein. 2017 , 118, 3268-3280		7
139	The Lipid A from <i>Rhodopseudomonas palustris</i> Strain BisA53 LPS Possesses a Unique Structure and Low Immunostimulant Properties. 2017 , 23, 3637-3647		19
138	qNMR for profiling the production of fungal secondary metabolites. 2017 , 55, 670-676		7
137	The lipopolysaccharide core oligosaccharide of plays a critical role in maintaining a proper gut symbiosis with the bean bug. <i>Journal of Biological Chemistry</i> , 2017 , 292, 19226-19237	5-4	16
136	A discrete Fourier-encoded, diagonal-free experiment to simplify homonuclear 2D NMR correlations. 2017 , 147, 034201		2
135	Structure of the Lipopolysaccharide from the sp. ORS285 Mutant Strain. 2017 , 6, 541-553		9
134	Large-scale synthesis and structural analysis of a synthetic glycopeptide dendrimer as an anti-cancer vaccine candidate. 2016 , 15, 114-123		15
133	Effects of Aib residues insertion on the structural-functional properties of the frog skin-derived peptide esculentin-1a(1-21)NH. 2017 , 49, 139-150		12
132	Natural structural diversity within a conserved cyclic peptide scaffold. 2017 , 49, 103-116		11
131	Structure of O-Antigen and Hybrid Biosynthetic Locus in Clonal Variants Recovered from a Cystic Fibrosis Patient. 2017 , 8, 1027		9
130	The Deep-Sea Polyextremophile <i>Halobacteroides lacunaris</i> TB21 Rough-Type LPS: Structure and Inhibitory Activity towards Toxic LPS. 2017 , 15,		11
129	Conformational properties, membrane interaction, and antibacterial activity of the peptaibiotic chalciporin A: Multitechnique spectroscopic and biophysical investigations on the natural compound and labeled analogs. 2017 , 110, e23083		5
128	A novel peptide conformation: the β -bend ribbon. 2018 , 16, 7947-7958		5

127	Structural and Functional Assessment of mBjAMP1, an Antimicrobial Peptide from Branchiostoma japonicum, Revealed a Novel Hairpinin-like Scaffold with Membrane Permeable and DNA Binding Activity. 2018 , 61, 11101-11113		23
126	A Solution NMR Approach To Determine the Chemical Structures of Carbohydrates Using the Hydroxyl Groups as Starting Points. <i>ACS Omega</i> , 2018 , 3, 17957-17975	3.9	11
125	Non-Uniform and Absolute Minimal Sampling for High-Throughput Multidimensional NMR Applications. 2018 , 24, 11535-11544		11
124	DPFGSE-MDEC-J-resolved COSY: An efficient method for selectively measuring proton-proton spin coupling constants of the multiplet signals. 2018 , 56, 1183-1187		1
123	Specific PPII Turns Precede PPII Structures Binding to Allele-Specific HLA-DR PBRs in Fully-Protective Malaria Vaccine Components. 2018 , 6, 106		2
122	Structure and inflammatory activity of the LPS isolated from <i>Acetobacter pasteurianus</i> CIP103108. 2018 , 119, 1027-1035		14
121	Binding mode of AIF(370-394) peptide to CypA: insights from NMR, label-free and molecular docking studies. <i>Biochemical Journal</i> , 2018 , 475, 2377-2393	3.8	5
120	Single hydroxo-bridged group 13 metalloporphyrin dimers: Solution studies and solid-state structures. 2019 , 23, 969-989		3
119	Kinetic and thermodynamic effects of phosphorylation on p53 binding to MDM2. <i>Scientific Reports</i> , 2019 , 9, 693	4.9	15
118	Structural analysis and mode of action of BMAP-27, a cathelicidin-derived antimicrobial peptide. 2019 , 118, 170106		13
117	Cn29, a novel orphan peptide found in the venom of the scorpion <i>Centruroides noxius</i> : Structure and function. 2019 , 167, 184-191		3
116	Structural studies of the binding of an antagonistic cyclic peptide to the VEGFR1 domain 2. 2019 , 169, 65-75		6
115	Heterochiral Ala/(Me)Aze sequential oligopeptides: Synthesis and conformational study. 2019 , 25, e3165		1
114	Biomolecular NMR Spectroscopy and Structure Determination of DNA. 2019 , 421-469		2
113	DQF J-RES NMR: Suppressing the singlet signals for improving the J-RES spectra from complex mixtures. <i>Journal of Magnetic Resonance</i> , 2019 , 301, 19-29	3	2
112	Structure determination of UL49.5 transmembrane protein from bovine herpesvirus 1 by NMR spectroscopy and molecular dynamics. 2019 , 1861, 926-938		4
111	. 2019 ,		3
110	CRAGE enables rapid activation of biosynthetic gene clusters in undomesticated bacteria. 2019 , 4, 2498-2510		48

109	Epitranscriptomics. <i>Methods in Molecular Biology</i> , 2019 ,	1.4	1
108	Structure and antifungal activity of pelgipeptins from <i>Paenibacillus elgii</i> against phytopathogenic fungi. 2020 , 163, 154-163		11
107	The three-dimensional structure of the toxic peptide Cl13 from the scorpion <i>Centruroides limpidus</i> . 2020 , 184, 158-166		1
106	Pairing LPS Structure with Its Immunomodulatory Effects on Human Cellular Models. 2020 , 6, 1602-1616		23
105	Impaired O-Glycosylation at Consecutive Threonine TTX Motifs in Mucins Generates Conformationally Restricted Cancer Neoepitopes. 2020 , 59, 1221-1241		4
104	Salient Features of Monomeric Alpha-Synuclein Revealed by NMR Spectroscopy. 2020 , 10,		6
103	Imunofan-RDKVYR Peptide-Stimulates Skin Cell Proliferation and Promotes Tissue Repair. 2020 , 25,		3
102	Effect of Site-Specific O-Glycosylation on the Structural Behavior of NOTCH1 Receptor Extracellular EGF-like Domains 11 and 10. 2020 , 26, 12363-12372		3
101	Efficient Na triple-quantum signal imaging on clinical scanners: Cartesian imaging of single and triple-quantum Na (CRISTINA). 2020 , 84, 2412-2428		3
100	Structure and allosteric activity of a single-disulfide conopeptide from a human β and γ nicotinic acetylcholine receptors. <i>Journal of Biological Chemistry</i> , 2020 , 295, 7096-7112	5-4	3
99	Redirecting substrate regioselectivity using engineered N-GBD-CD2 branching sucrases for the production of pentasaccharide repeating units of <i>S. flexneri</i> 3a, 4a and 4b haptens. <i>Scientific Reports</i> , 2021 , 11, 2474	4-9	4
98	The muscle-relaxing C-terminal peptide from troponin I populates a nascent helix, facilitating binding to tropomyosin with a potent therapeutic effect. <i>Journal of Biological Chemistry</i> , 2021 , 296, 100278	5-4	1
97	Investigation of the Effects of Primary Structure Modifications within the RRE Motif on the Conformation of Synthetic Bovine Herpesvirus 1-Encoded UL49.5 Protein Fragments. 2021 , 18, e2000883		0
96	Diselenide crosslinks for enhanced and simplified oxidative protein folding. 2021 , 4,		5
95	Conformationally Constrained Peptides with High Affinity to the Vascular Endothelial Growth Factor. 2021 , 64, 10900-10907		1
94	Interactome and structural basis for targeting the human T-cell leukemia virus Tax oncoprotein.		
93	Probing the helical stability in a VEGF-mimetic peptide. 2021 , 116, 105379		0
92	Recent developments in biochemical nuclear magnetic resonance spectroscopy. 1985 , 31, 259-315		3

91	Characterization of Hydrogenated Fullerenes by NMR Spectroscopy. 2010 , 171-202		2
90	Analysis of complex carbohydrate primary and secondary structure via two-dimensional proton nuclear magnetic resonance spectroscopy. 1988 , 228, 759-84		3
89	Nuclear Magnetic Resonance Spectroscopy. 2013 , 113-173		2
88	Homonuclear Hartmann-Hahn Magnetization Transfer: New One- and Two-Dimensional NMR Methods for Structure Determination and Spectral Assignment. 1986 , 21-48		2
87	Three-Dimensional Structure Determination of Peptides Using Solution Nuclear Magnetic Resonance Spectroscopy. <i>Methods in Molecular Biology</i> , 2020 , 2068, 129-162	1.4	11
86	The Binding of Transition Metal Ions to DNA Oligonucleotides Studied by Nuclear Magnetic Resonance Spectroscopy. 1997 , 493-509		1
85	IDENTIFICATION OF FOLDED STRUCTURES IN IMMUNOGENIC PEPTIDES BY 2D NMR SPECTROSCOPY. 1987 , 477-480		2
84	Applications of 2D NMR to Biological Systems. 1988 , 435-488		1
83	NMR Methods Used to Derive Solution Conformations in Peptides. 1994 , 5, 423-430		1
82	Modern Nuclear Magnetic Resonance Spectroscopy of Peptides. 1985 , 437-473		4
81	¹ H NMR STRUCTURAL STUDIES OF A COMPLEX OF MELITTIN WITH PERDEUTERATED CALMODULIN. 1987 , 360-371		3
80	INVESTIGATION OF HELIX DIPOLE AND SALT BRIDGE STABILIZATION OF SYNTHETIC HELICAL PEPTIDES USING CD AND NMR. 1989 , 531-546		2
79	Cardiotoxin II from Taiwan cobra venom, <i>Naja naja atra</i> . Structure in solution and comparison among homologous cardiotoxins.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 23500-23508	5-4	25
78	Sialyl Lewis X mimics derived from a pharmacophore search are selectin inhibitors with anti-inflammatory activity.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 19663-19666	5-4	78
77	Functional and structural role of a tryptophan generally observed in protein-carbohydrate interaction. TRP-62 of hen egg white lysozyme.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 7070-7075	5-4	44
76	The tyrosine-containing internalization motif in the cytoplasmic domain of TGN38/41 lies within a nascent helix.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 7131-7136	5-4	27
75	Conformational change of adrenodoxin induced by reduction of iron-sulfur cluster. Proton nuclear magnetic resonance study.. <i>Journal of Biological Chemistry</i> , 1991 , 266, 6252-6258	5-4	26
74	Micelle-bound conformational preferences of a peptide derived from a murine major histocompatibility complex class I molecule.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 22830-22837	5-4	3

73	The structure of lipooligosaccharide produced by <i>Neisseria gonorrhoeae</i> , strain 15253, isolated from a patient with disseminated infection. Evidence for a new glycosylation pathway of the gonococcal lipooligosaccharide.. <i>Journal of Biological Chemistry</i> , 1994 , 269, 30345-30351	5-4	54
72	¹ H NMR study of [d(GCGATCGC)] ₂ and its interaction with minor groove binding 4',6-diamidino-2-phenylindole.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 3944-3951	5-4	38
71	Lipooligosaccharidic antigen containing a novel C4-branched 3,6-dideoxy-alpha-hexopyranose typifies <i>Mycobacterium gastrii</i> .. <i>Journal of Biological Chemistry</i> , 1993 , 268, 3168-3179	5-4	28
70	Identification of D-Sorbitol 3-Phosphate in the Normal and Diabetic Mammalian Lens. <i>Journal of Biological Chemistry</i> , 1989 , 264, 9278-9282	5-4	23
69	Structural Characterization of Nitrimyoglobin. <i>Journal of Biological Chemistry</i> , 1989 , 264, 6134-6145	5-4	36
68	Biosynthesis of Lipopolysaccharide in <i>Escherichia coli</i> . <i>Journal of Biological Chemistry</i> , 1989 , 264, 6956-6966	5-4	77
67	The flexible region of protein L12 from bacterial ribosomes studied by proton nuclear magnetic resonance. <i>Journal of Biological Chemistry</i> , 1989 , 264, 4498-4505	5-4	36
66	Mapping of neutralizing epitopes and the receptor binding site of human interleukin 1 beta. <i>Journal of Biological Chemistry</i> , 1993 , 268, 9771-9779	5-4	21
65	Import, processing, and two-dimensional NMR structure of a linker-deleted signal peptide of rat liver mitochondrial aldehyde dehydrogenase.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 19906-19914	5-4	62
64	Calcium plays distinctive structural roles in the N- and C-terminal domains of cardiac troponin C.. <i>Journal of Biological Chemistry</i> , 1993 , 268, 20966-20973	5-4	15
63	A proton nuclear magnetic resonance and molecular modeling study of cardiac troponin C. Calcium dependence and aromatic spectral assignments.. <i>Journal of Biological Chemistry</i> , 1990 , 265, 9754-9763	5-4	5
62	Conformational analysis of the tumor-associated carbohydrate antigen 19-9 and its Lea blood group antigen component as related to the specificity of monoclonal antibody CO19-9.. <i>Journal of Biological Chemistry</i> , 1990 , 265, 2028-2037	5-4	36
61	¹ H and ¹¹⁹ Sn magnetic resonance study of the SnIV protoporphyrin IX complex of equine myoglobin. Structure of the porphyrin-binding pocket. <i>Journal of Biological Chemistry</i> , 1991 , 266, 3728-3733	5-4	10
60	Novel O-Linked Carbohydrate Chains in the Cellulase Complex (Cellulosome) of <i>Clostridium thermocellum</i> . <i>Journal of Biological Chemistry</i> , 1989 , 264, 1027-1035	5-4	69
59	3-O-acetyl-sphingosine-series myelin glycolipids. <i>Journal of Lipid Research</i> , 2002 , 43, 751-761	6-3	23
58	Characterization of conformation and dynamics of CD4 Fragment (81-92) TYICEVEDQKEE and its benzylated derivative by ¹ H NMR spectroscopy and molecular modeling: Relevance of conformation to biological function. <i>Journal of Acquired Immune Deficiency Syndromes</i> , 1996 , 11, 222-32		2
57	Proton NMR study of the trimannosyl unit in a pentaantennary N-linked decasaccharide structure. Complete assignment of the proton resonances and conformational characterization. <i>FEBS Journal</i> , 1995 , 228, 822-9		11
56	¹ H, ¹³ C, ¹⁵ N-NMR resonance assignments of oxidized thioredoxin h from the eukaryotic green alga <i>Chlamydomonas reinhardtii</i> using new methods based on two-dimensional triple-resonance NMR spectroscopy and computer-assisted backbone assignment. <i>FEBS Journal</i> , 1995 , 229, 473-85		20

55	1H nuclear-magnetic-resonance investigation of oxidized Fe4S4 ferredoxin from <i>Thermotoga maritima</i> . Hyperfine-shifted resonances, sequence-specific assignments and secondary structure. <i>FEBS Journal</i> , 1995 , 229, 658-68		17
54	Elucidation of the primary structure of the lantibiotic epilancin K7 from <i>Staphylococcus epidermidis</i> K7. Cloning and characterisation of the epilancin-K7-encoding gene and NMR analysis of mature epilancin K7. <i>FEBS Journal</i> , 1995 , 230, 587-600		74
53	Structural characterization of a short peptide fragment that mediates estrogen-receptor dimerization. <i>FEBS Journal</i> , 1995 , 230, 879-85		4
52	Comparative Study on the Conformation of Phalloidin, Viroisin, and Related Derivatives in Aqueous Solution. <i>FEBS Journal</i> , 1995 , 232, 726-736		8
51	Mapping the anatomy of the immunodominant domain of the human immunodeficiency virus gp41 transmembrane protein: peptide conformation analysis using monoclonal antibodies and proton nuclear magnetic resonance spectroscopy. <i>Journal of Virology</i> , 1991 , 65, 1727-34	6.6	44
50	Nuclear magnetic resonance analysis of solution conformations in C4-V3 hybrid peptides derived from human immunodeficiency virus (HIV) type 1 gp120: relation to specificity of peptide-induced anti-HIV neutralizing antibodies. <i>Journal of Virology</i> , 1999 , 73, 746-50	6.6	11
49	Expression and Characterization of Stress Responsive Peptide-1; an Inducer of Antimicrobial Peptide Synthesis. 2019 , 4, 42-52		1
48	Altered metabolism of growth hormone receptor mutant mice: a combined NMR metabonomics and microarray study. <i>PLoS ONE</i> , 2008 , 3, e2764	3.7	42
47	Interplay between Structure and Charge as a Key to Allosteric Modulation of Human 20S Proteasome by the Basic Fragment of HIV-1 Tat Protein. <i>PLoS ONE</i> , 2015 , 10, e0143038	3.7	9
46	Solution Structure and Expression Profile of an Insect Cytokine: <i>Manduca sexta</i> Stress Response Peptide-2. <i>Protein and Peptide Letters</i> , 2017 , 24, 3-11	1.9	3
45	In vitro and in vivo effect of parathyroid hormone analogue (1-14) containing -amino-iso-butyric acid residue (Aib)1,3. <i>Yonsei Medical Journal</i> , 2006 , 47, 214-22	3	3
44	NMR Studies on Antitumor Drug Candidates, Berberine and Berberrubine. <i>Bulletin of the Korean Chemical Society</i> , 2002 , 23, 391-394	1.2	17
43	Tertiary Structure of Ginsenoside Re Studied by NMR Spectroscopy. <i>Bulletin of the Korean Chemical Society</i> , 2007 , 28, 2209-2213	1.2	4
42	Solution Structure of LXXLL-related Cofactor Peptide of Orphan Nuclear Receptor FTZ-F1. <i>Bulletin of the Korean Chemical Society</i> , 2012 , 33, 583-588	1.2	1
41	The glucuronyltransferase B4GAT1 is required for initiation of LARGE-mediated Hdyctroglycan functional glycosylation. <i>ELife</i> , 2014 , 3,	8.9	67
40	Computer-aided engineering of a branching sucrose for the glucodiversification of a tetrasaccharide precursor of <i>S. flexneri</i> antigenic oligosaccharides. <i>Scientific Reports</i> , 2021 , 11, 20294	4.9	1
39	Selection by site-directed antibodies of small regions of peptides which are ordered in water. <i>Novartis Foundation Symposium</i> , 1986 , 119, 58-75		1
38	NMR. 2009 , 333-374		

- 37 Introduction to Nuclear Magnetic Resonance. **2011**, 129-141
- 36 CALMODULIN FROM VARIOUS SOURCES AS STUDIED BY DIFFERENT NMR TECHNIQUES. **1987**, 397-399
- 35 Fundamentals of Structural Analysis of Glycosphingolipids by Proton Nuclear Magnetic Resonance Spectroscopy. **1987**, 49-70 1
- 34 Applications of Two-Dimensional NMR Experiments in Liquids. **1988**, 343-434
- 33 D. **1990**, 65-83
- 32 Protein structure determination by nuclear magnetic resonance spectroscopy. *Biotechnology*, **1990**, 14, 187-226
- 31 M. **1990**, 169-185
- 30 X. **1990**, 306-306 1
- 29 An ¹H NMR study of HTyr-d-Cys-Phe-d-PenOH (JOM-13), a highly selective ligand for the opioid receptor. **1991**, 105-109 1
- 28 ¹³C and ¹H resonance assignments and structural determination of an emerimicin peptide in DMSO. **1991**, 545-547
- 27 Toward the Structure of Mosaic Proteins: Expression, Purification and Structural Analysis of a Pair of Fibronectin Type1 Modules. **1993**, 623-631
- 26 Modern spectroscopic approaches to structure elucidation. *Biotechnology*, **1994**, 26, 389-444
- 25 Preferred Local Conformations of Peptides comprising the Entire Sequence of Bovine Pancreatic Trypsin Inhibitor (BPTI) and their Roles in Protein Folding. **1994**, 139-153 1
- 24 Computer-Supported Protein Structure Determination by NMR. *NATO ASI Series Series B: Physics*, **1994**, 197-207
- 23 Two-, three-, and four-dimensional nuclear magnetic resonance spectroscopy of protein pharmaceuticals. *Pharmaceutical Biotechnology*, **1995**, 7, 179-218
- 22 Multidimensional NMR of macromolecules. **1996**, 85-157
- 21 NMR, X-Ray and MS Investigations of Ethyl 2-Aroyl-and Ethyl 2-Arylcarbamoyl-4,5-dimethyl-1,2,3,6-tetrahydropyridazine-1- carboxylates: Flexible Multidentate Ligands. *Journal of Chemical Research*, **1999**, 23, 186-187 0.6
- 20 Synthesis and Evaluation of Novel Neamine-Nucleoside Conjugates as Potential Antibiotic Targets for Escherichia coli 16S Ribosomal RNA. *Methods in Molecular Biology*, **2019**, 1870, 151-163 1.4

19	Structural studies on Econotoxin SI. 2002 , 730-732		
18	Solution structure of the epidermal growth factor-like domain of heregulin-alpha, a ligand for p180erbB-4. <i>EMBO Journal</i> , 1994 , 13, 3517-23	13	10
17	Solution structure of human insulin-like growth factor II; recognition sites for receptors and binding proteins. <i>EMBO Journal</i> , 1994 , 13, 5590-7	13	12
16	Secondary structure determination for the Antennapedia homeodomain by nuclear magnetic resonance and evidence for a helix-turn-helix motif. <i>EMBO Journal</i> , 1988 , 7, 4305-9	13	40
15	The conformations of hirudin in solution: a study using nuclear magnetic resonance, distance geometry and restrained molecular dynamics. <i>EMBO Journal</i> , 1987 , 6, 529-37	13	11
14	Determination of the structure of the nucleocapsid protein NCp7 from the human immunodeficiency virus type 1 by 1H NMR. <i>EMBO Journal</i> , 1992 , 11, 3059-65	13	85
13	The internalization signal in the cytoplasmic tail of lysosomal acid phosphatase consists of the hexapeptide PGYRHV. <i>EMBO Journal</i> , 1992 , 11, 4391-9	13	19
12	Solution structure of a defensin-like peptide from platypus venom. <i>Biochemical Journal</i> , 1999 , 341 (Pt 3), 785-94	3.8	12
11	Defensin-like peptide-2 from platypus venom: member of a class of peptides with a distinct structural fold. <i>Biochemical Journal</i> , 2000 , 348 Pt 3, 649-56	3.8	18
10	Human T-lymphotrophic virus type I nucleocapsid protein NCp15: structural study and stability of the N-terminal zinc-finger. <i>Biochemical Journal</i> , 2000 , 352 Pt 2, 293-300	3.8	5
9	Conformational characterization of duplex DNA with solution-state NMR spectroscopy. <i>Journal of Magnetic Resonance Open</i> , 2022 , 10-11, 100035	0.4	1
8	Is Cys(MTSL) the Best α Amino Acid Residue to Electron Spin Labeling of Synthetically Accessible Peptide Molecules with Nitroxides?. <i>ACS Omega</i> , 2022 , 7, 5154-5165	3.9	1
7	Ecdysis triggering hormone peptide in the African malaria mosquito <i>Anopheles gambiae</i> : The peptide structure for receptor activation.. <i>Insect Science</i> , 2022 ,	3.6	0
6	Dynamic nuclear polarization-enhanced, double-quantum filtered C-C dipolar correlation spectroscopy of natural C abundant bone-tissue biomaterial.. <i>Journal of Magnetic Resonance</i> , 2022 , 335, 107144	3	0
5	Role of EPS in mitigation of plant abiotic stress: The case of <i>Methylobacterium extorquens</i> PA1. <i>Carbohydrate Polymers</i> , 2022 , 119863	10.3	1
4	<i>Bacteroides thetaiotaomicron</i> rough-type lipopolysaccharide: The chemical structure and the immunological activity. 2022 , 297, 120040		2
3	Therapeutic Peptides: Unravelling Conformational Dynamics by Systematic Application of Biophysical Techniques. 2022 , 23,		0
2	Structural and functional studies of scorpine: A channel blocker and cytolytic peptide. 2023 , 222, 106985		0

- 1 The Aharonov-Anandan phase and geometric double-quantum excitation in strongly coupled nuclear spin pairs. **2023**, 158, 124204

o