

CITATION REPORT

List of articles citing

A more meaningful scoring system for determining the severity of coronary heart disease

DOI: 10.1016/s0002-9149(83)80105-2
American Journal of Cardiology, 1983, 51, 606.

Source: <https://exaly.com/paper-pdf/16403526/citation-report.pdf>

Version: 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
1447	The ST segment/heart rate slope as a predictor of coronary artery disease: comparison with quantitative thallium imaging and conventional ST segment criteria. 1986 , 112, 296-304		51
1446	Plasma apoproteins and the severity of coronary artery disease. 1986 , 73, 978-86		61
1445	Plasma fibrinogen levels as an independent indicator of severity of coronary atherosclerosis. 1989 , 77, 209-13		112
1444	Alcohol consumption, serum lipids and severity of angiographically determined coronary artery disease. <i>American Journal of Cardiology</i> , 1990 , 65, 287-9	3	46
1443	Detection of human collateral circulation by vasodilation-thallium-201 tomography. <i>American Journal of Cardiology</i> , 1990 , 65, 991-8	3	8
1442	Progression and regression of minor coronary arterial narrowings by quantitative angiography after fenofibrate therapy. <i>American Journal of Cardiology</i> , 1991 , 67, 957-61	3	41
1441	Progression of carotid atherosclerosis in Japanese patients with coronary artery disease. 1992 , 23, 946-51		53
1440	Correlation of reciprocal ST-segment depression after acute myocardial infarction with coronary angiographic findings. 1992 , 36, 163-8		11
1439	Significance of R wave changes in exercise-induced supraventricular extrasystoles. Angiographic correlates. 1993 , 26, 197-206		1
1438	Effects of fenofibrate on angiographically examined coronary atherosclerosis and left ventricular function in hypercholesterolemic patients. 1993 , 98, 127-38		18
1437	Evidence that apolipoprotein(a) phenotype is a risk factor for coronary artery disease in men <i>American Journal of Cardiology</i> , 1994 , 74, 346-51	3	30
1436	Difference between women and men in left ventricular pump function during predischage exercise test after acute myocardial infarction. <i>American Journal of Cardiology</i> , 1994 , 73, 11-5	3	7
1435	Progression of carotid atherosclerosis in Japanese patients with coronary artery disease. 1994 , 3, 56-60		
1434	Lack of correlation between coronary risk factors and CAD severity. 1994 , 47, 37-43		20
1433	Origin and pathophysiological role of increased plasma endothelin-1 in patients with acute myocardial infarction. 1995 , 46, 557-65		6
1432	Sex differences in the anatomy of coronary artery disease. 1995 , 48, 723-30		19
1431	Serum glycoproteins and severity of coronary atherosclerosis. 1995 , 129, 234-8		67

1430	Band neutrophil count and the presence and severity of coronary atherosclerosis. 1996 , 132, 9-12		60
1429	Apolipoprotein(a) and pentanucleotide repeat polymorphisms are associated with the degree of atherosclerosis in coronary heart disease. 1996 , 123, 181-91		31
1428	Significance of angiotensin I-converting enzyme and angiotensin II type 1 receptor gene polymorphisms as risk factors for coronary heart disease. 1996 , 125, 161-9		87
1427	Paraoxonase gene polymorphism in Japanese subjects with coronary heart disease. 1996 , 57, 69-73		113
1426	Genotype distribution of estrogen receptor polymorphisms in men and postmenopausal women from healthy and coronary populations and its relation to serum lipid levels. 1997 , 17, 3006-12		87
1425	Quantitative comparison of angiographic characteristics of coronary artery disease in patients with noninsulin-dependent diabetes mellitus compared with matched nondiabetic control subjects. <i>American Journal of Cardiology</i> , 1997 , 80, 550-6	3	51
1424	Dyslipidemias have a detrimental effect on left ventricular systolic function in patients with a first acute myocardial infarction. <i>American Journal of Cardiology</i> , 1998 , 81, 531-7	3	34
1423	Pain intensity in nondiabetic patients with myocardial infarction or unstable angina. Its association with clinical and psychological features. 1998 , 67, 165-9		8
1422	ACE I/D gene polymorphism: presence of the ACE D allele increases the risk of coronary artery disease in younger individuals. 1998 , 139, 153-9		64
1421	Association of the insertion/deletion gene polymorphism of the apolipoprotein B signal peptide with myocardial infarction. 1998 , 141, 167-75		23
1420	Association of the Platelet Glycoprotein Ia C807T Gene Polymorphism With Nonfatal Myocardial Infarction in Younger Patients. 1999 , 93, 2449-2453		200
1419	A new promoter polymorphism in the gene of lipopolysaccharide receptor CD14 is associated with expired myocardial infarction in patients with low atherosclerotic risk profile. 1999 , 19, 932-8		145
1418	Angiotensin I converting enzyme gene polymorphism and insulin resistance in patients with angina pectoris. 1999 , 12, 291-7		11
1417	Angiotensin I-converting enzyme gene polymorphism in a low-risk European population for coronary artery disease. 1999 , 142, 211-6		33
1416	Angiotensinogen T174M and M235T gene polymorphisms are associated with the extent of coronary atherosclerosis. 1999 , 145, 309-14		52
1415	The p22 phox A640G gene polymorphism but not the C242T gene variation is associated with coronary heart disease in younger individuals. 1999 , 145, 315-23		92
1414	The effects of dyslipidemia on left ventricular systolic function in patients with stable angina pectoris. 1999 , 146, 117-24		27
1413	The graded effect of hyperhomocysteinemia on the severity and extent of coronary atherosclerosis. 1999 , 147, 379-86		57

1412	Insulin resistance and angiographical characteristics of coronary atherosclerosis. 1999 , 63, 666-73	16
1411	Descending aorta wall volume and coronary artery disease: a comparative study using enhanced computed tomography of the chest and coronary angiography. 2000 , 64, 842-7	5
1410	Circulating adhesion molecules and severity of coronary atherosclerosis. 2000 , 11, 77-81	51
1409	Preperitoneal fat thickness determined by ultrasonography is correlated with coronary stenosis and lipid disorders in non-obese male subjects. 2000 , 24, 502-7	31
1408	Susceptibility of apolipoprotein B-containing lipoproteins to oxidation and antioxidant status in acute coronary syndromes. 2000 , 23, 655-8	6
1407	Differences between coronary hyperresponsiveness to ergonovine and vasospastic angina. 2000 , 41, 257-68	3
1406	Increased chymase activity in internal thoracic artery of patients with hypercholesterolemia. 2000 , 35, 55-60	38
1405	Relation between green tea consumption and the severity of coronary atherosclerosis among Japanese men and women. 2000 , 10, 401-8	103
1404	The paraoxonase Leu-Met54 and Gln-Arg191 gene polymorphisms are not associated with the risk of coronary heart disease. 2000 , 152, 421-31	71
1403	Out-of-hospital ventricular fibrillation in patients with acute myocardial infarction: coronary angiographic determinants. 2000 , 35, 144-50	49
1402	Alcohol consumption and severity of angiographically determined coronary artery disease in Japanese men and women. 2001 , 156, 177-83	7
1401	Angiographic correlates of a positive troponin T test in patients with unstable angina. 2001 , 29, 1130-6	21
1400	A chymase gene variant is associated with atherosclerosis in venous coronary artery bypass grafts. 2001 , 12, 493-7	28
1399	Role of hypertension, dyslipidemia and diabetes mellitus in the development of coronary atherosclerosis in Japan. 2001 , 65, 731-7	25
1398	Apolipoprotein B as the best predictor of coronary artery disease in Iranian normolipidemic patients. 2001 , 34, 149-55	21
1397	Obesity, body fat distribution and coronary atherosclerosis among Japanese men and women. 2001 , 25, 191-7	21
1396	Extent of coronary atherosclerosis and homocysteine affect endothelial markers. 2001 , 52, 589-96	7
1395	Circadian variation in ischemic threshold in patients with stable angina: relation to plasma endothelin-1. 2002 , 53, 409-13	13

1394	Ultrasonic correlates of common carotid atherosclerosis in patients with coronary artery disease. 2002 , 53, 177-83		12
1393	Increased level of advanced oxidation protein products in patients with coronary artery disease. 2002 , 162, 221-5		141
1392	Activity and concentration of lipoprotein lipase in post-heparin plasma and the extent of coronary artery disease. 2002 , 163, 127-34		13
1391	Elevated level of plasma C-reactive protein in patients with unstable angina: its relations with coronary stenosis and lipid profile. 2002 , 53, 265-72		44
1390	Significance of small dense low-density lipoproteins and other risk factors in patients with various types of coronary heart disease. 2002 , 144, 1026-35		55
1389	Transcardiac gradient of soluble adhesion molecules predicts progression of coronary artery disease. 2002 , 84, 249-57		15
1388	Levels of soluble E-selectin and ICAM-1 in the coronary circulation of patients with stable coronary artery disease: association with the severity of coronary atherosclerosis. 2002 , 43, 93-101		24
1387	Does stent design affect the long-term outcome after coronary stenting?. 2002 , 56, 305-11		12
1386	Relation between type A behavior pattern and the extent of coronary atherosclerosis in Japanese women. 2002 , 9, 77-93		6
1385	Relation of erectile dysfunction to angiographic coronary artery disease. <i>American Journal of Cardiology</i> , 2003 , 91, 230-1	3	186
1384	Interaction between SREBP-1a and APOB polymorphisms influences total and low-density lipoprotein cholesterol levels in patients with coronary artery disease. 2003 , 63, 380-5		15
1383	The relation of QT dispersion and localized QT difference to coronary pathology in a population with unstable coronary artery disease. 2003 , 8, 22-9		3
1382	Endothelin-1 and vasopressin plasma levels are not associated with the insertion/deletion polymorphism of the human angiotensin I-converting enzyme gene in patients with coronary artery disease. 2003 , 17, 133-8		6
1381	The pre-existence of an acute coronary event predicts differences in biological parameters and clinical evolution among patients with longstanding stable angina. 2003 , 91, 193-200		3
1380	Ischemic preconditioning detected by treadmill exercise tests in patients with stable angina. 2003 , 54, 45-50		21
1379	Prognostic value of the carotid artery intima-media thickness for the presence and severity of coronary artery disease in type 2 diabetic patients. 2003 , 26, 3189-90		11
1378	Nocturnal oxygen desaturation correlates with the severity of coronary atherosclerosis in coronary artery disease. 2003 , 124, 936-41		80
1377	Impact of diabetes mellitus on worsening of the left ventricular ejection fraction in exercise-gated 201Tl myocardial single photon emission computed tomography in patients with coronary artery disease. 2003 , 67, 839-45		5

1376	Plasma levels of soluble thrombomodulin, C-reactive protein, and serum amyloid A protein in the atherosclerotic coronary circulation. 2003 , 44, 601-12	24
1375	Increased C-reactive protein level after renal stent implantation in patients with atherosclerotic renal stenosis. 2004 , 55, 479-84	16
1374	Waist-hip ratio and low HDL predict the risk of coronary artery disease in Pakistanis. 2004 , 20, 55-62	35
1373	Plasma vascular endothelial growth factor level is elevated in patients with multivessel coronary artery disease. 2004 , 27, 281-6	20
1372	Activation of nuclear factor-kappaB and correlation with elevated plasma c-reactive protein in patients with unstable angina. 2004 , 13, 173-8	29
1371	Time course of inflammatory response after renal artery stenting in patients with atherosclerotic renal stenosis. 2004 , 350, 115-21	30
1370	Atherosclerotic disease regression with statins: studies using vascular markers. 2004 , 96, 447-59	26
1369	Delayed intravascular catabolism of chylomicron-like emulsions is an independent predictor of coronary artery disease. 2004 , 176, 397-403	20
1368	Increases in lipids and immune cells in response to exercise and mental stress in patients with suspected coronary artery disease: effects of adjustment for shifts in plasma volume. 2004 , 65, 237-50	43
1367	Role of prehypertension in the development of coronary atherosclerosis in Japan. 2004 , 14, 57-62	43
1366	Plasma concentrations of monocyte chemoattractant protein 1 (MCP-1) and neopterin in the coronary circulation of patients with coronary artery disease. 2004 , 68, 114-20	16
1365	Elevated concentration of soluble vascular endothelial cadherin is associated with coronary atherosclerosis. 2004 , 68, 1-5	50
1364	Leg fat percentage correlates negatively with coronary atherosclerosis. 2004 , 68, 1173-8	23
1363	Endothelial function and baroreflex sensitivity according to the oral glucose tolerance test in patients with coronary artery disease and normal fasting glucose levels. 2005 , 109, 397-403	6
1362	N-terminal pro-B-type natriuretic Peptide predicts significant coronary artery lesion in the unstable angina patients with normal electrocardiogram, echocardiogram, and cardiac enzymes. 2005 , 69, 1472-6	18
1361	Relationship between myocardial injury and soluble P-selectin in non-ST elevation acute coronary syndromes. 2005 , 69, 530-5	7
1360	Lipoprotein(a) levels in patients with unstable angina and their relationship with atherothrombosis and myocardial damage. 2005 , 59, 150-5	2
1359	Relationship between leukocyte count and angiographical characteristics of coronary atherosclerosis. 2005 , 26, 1057-62	15

1358	P-wave dispersion in patients with stable coronary artery disease and its relationship with severity of the disease. 2005 , 38, 279-84	43
1357	Relationship between DNA damage, total antioxidant capacity and coronary artery disease. 2005 , 570, 197-203	76
1356	Changes in myocardial perfusion due to physical exercise in patients with stable coronary artery disease. 2005 , 32, 813-9	29
1355	Effects of the shape of coronary arteries on the presence, extent, and severity of their disease. 2005 , 20, 224-9	8
1354	Sp1-associated activation of macrophage inflammatory protein-2 promoter by CpG-oligodeoxynucleotide and lipopolysaccharide. 2005 , 62, 188-98	27
1353	Retinopathy is related to the angiographically detected severity and extent of coronary artery disease in patients with type 2 diabetes mellitus. 2005 , 46, 639-46	47
1352	Metabolic syndrome and risk of coronary heart disease in a Pakistani cohort. 2005 , 91, 1003-7	17
1351	Impaired renal function and atherosclerosis in a Pakistani cohort. 2005 , 21, 1201-7	3
1350	Gelatinase B C(-1562)T polymorphism in relation to ischaemic heart disease. 2005 , 65, 513-22	32
1349	Obesity, body fat and coronary atherosclerosis. 2005 , 98, 227-35	23
1348	Augmentation index, pulse pressure amplification and superoxide anion production in patients with coronary artery disease. 2005 , 99, 289-94	17
1347	Effects of xuezhikang, an extract of cholestin, on lipid profile and C-reactive protein: a short-term time course study in patients with stable angina. 2005 , 352, 217-24	43
1346	Angiotensinogen M235T polymorphism is associated with coronary artery disease severity. 2005 , 362, 176-81	37
1345	Protective effect of high diastolic blood pressure during exercise against exercise-induced myocardial ischemia. 2005 , 150, 790-5	15
1344	Erectile dysfunction: the need to be evaluated, the right to be treated. 2005 , 150, 620-6	14
1343	Coronary artery disease progression is associated with C-reactive protein and conventional risk factors but not soluble CD40 ligand. 2006 , 22, 691-6	13
1342	Ascending aortic blood pressure-derived indices are not correlated with the extent of coronary artery disease in patients with impaired left ventricular function. 2006 , 184, 370-6	10
1341	Significance of small dense low-density lipoprotein-cholesterol concentrations in relation to the severity of coronary heart diseases. 2006 , 189, 206-14	83

1340	The presence of metabolic syndrome is independently associated with elevated serum CD40 ligand and disease severity in patients with symptomatic coronary artery disease. 2006 , 55, 1029-34		25
1339	Significance of small dense low-density lipoprotein as a risk factor for coronary artery disease and acute coronary syndrome. 2006 , 47, 405-14		29
1338	Relationship between high-sensitivity C-reactive protein level and angiographical characteristics of coronary atherosclerosis. 2006 , 119, 319-323		0
1337	Time course of rapid C-reactive protein reduction by pravastatin in patients with stable angina. 2006 , 57, 1-7		17
1336	Predictors of long-term outcome following percutaneous coronary intervention in patients with type 2 diabetes mellitus. 2006 , 17, 131-8		3
1335	The association of QT dispersion and QT dispersion ratio with extent and severity of coronary artery disease. 2006 , 11, 43-51		10
1334	Paraoxonase and arylesterase activities in coronary artery disease. 2006 , 36, 779-87		63
1333	Relationship between true fasting plasma insulin level and angiographic characteristics of coronary atherosclerosis. 2006 , 29, 25-30		1
1332	Importance of blood cellular genomic profile in coronary heart disease. 2006 , 13, 17-26		10
1331	Comparison of presence and extent of coronary narrowing in patients with left bundle branch block without diabetes mellitus to patients with and without left bundle branch block but with diabetes mellitus. <i>American Journal of Cardiology</i> , 2006 , 97, 857-9	3	6
1330	Relationship between serum albumin level before initiating haemodialysis and angiographic severity of coronary atherosclerosis in end-stage renal disease patients. 2006 , 21, 1633-9		32
1329	Relationship between aortic valve sclerosis and the extent of coronary artery disease in patients undergoing diagnostic coronary angiography. 2006 , 106, 277-82		18
1328	Transcriptional profiling in coronary artery disease: indications for novel markers of coronary collateralization. 2006 , 114, 1811-20		45
1327	Effect of coronary angiography on oxidative DNA damage observed in circulating lymphocytes. 2007 , 58, 141-7		11
1326	Ascending aortic blood pressure waveform is related to coronary atherosclerosis in hypertensive as well as in normotensive subjects. 2007 , 16, 246-53		14
1325	Effects of percutaneous coronary intervention on peripheral venous blood circulating endothelial cells and plasma indices of endothelial damage/dysfunction. 2007 , 132, 1920-6		21
1324	Ability of ambulatory blood pressure monitoring and myocardial/carotid ultrasound to predict the location and the severity of coronary artery lesions in normotensive patients: a clinical study. 2007 , 30, 741-9		3
1323	Circulating endothelial progenitor cells, C-reactive protein and severity of coronary stenosis in Chinese patients with coronary artery disease. 2007 , 30, 133-41		36

1322	Linkage Analysis by a Transmission/Disequilibrium Test of Russian Sibling Pairs with Coronary Artery Disease. 2007 , 10, 43-54	5
1321	Lipoprotein(a) and LDL particle size are related to the severity of coronary artery disease. 2007 , 108, 282-9	34
1320	The relationship between plasma C-reactive protein levels and presence and severity of coronary stenosis in patients with stable angina. 2007 , 58, 657-62	6
1319	C-588T polymorphism of the human glutamate-cysteine ligase modifier subunit gene is not associated with the risk and extent of ischemic heart disease in a German cohort. 2007 , 45, 1416-8	3
1318	Echocardiographic epicardial fat thickness and coronary artery disease. 2007 , 71, 536-9	271
1317	Relationship between angiographically documented coronary artery disease and low bone mass in men. 2007 , 71, 1095-8	12
1316	Direct comparison of brain natriuretic peptide (BNP) and N-terminal pro-BNP secretion and extent of coronary artery stenosis in patients with stable coronary artery disease. 2007 , 71, 499-505	35
1315	The polymorphism in acetaldehyde dehydrogenase 2 gene, causing a substitution of Glu > Lys(504), is not associated with coronary atherosclerosis severity in Han Chinese. 2007 , 213, 215-20	15
1314	The relationship between Helicobacter pylori IgG titre and coronary atherosclerosis. 2007 , 62, 501-5	26
1313	Renal dysfunction is the most important predictor of the extent and severity of coronary artery disease in patients with diabetes mellitus. 2007 , 18, 463-9	18
1312	Association of RAGE gene polymorphisms with coronary artery disease in the Korean population. 2007 , 18, 1-8	19
1311	Association between Toll-like receptors and the extent and severity of coronary artery disease in patients with stable angina. 2007 , 18, 31-8	21
1310	Reduction of C-reactive protein by a single 80 mg of simvastatin in patients with unstable angina. 2007 , 376, 163-7	14
1309	A potential evidence to explain the reason behind the devastating prognosis of coronary artery disease in uraemic patients: renal insufficiency is associated with poor coronary collateral vessel development. 2007 , 115, 366-72	28
1308	Comparative predictive and diagnostic value of flow-mediated vasodilation in the brachial artery and intima media thickness of the carotid artery for assessment of coronary artery disease severity. 2007 , 117, 165-72	28
1307	Impaired nitrate-mediated dilatation could reflect nitrate tolerance in patients with coronary artery disease. 2007 , 120, 351-6	10
1306	Relationship of thyroid stimulating hormone with coronary atherosclerosis in angina patients. 2007 , 122, 56-60	24
1305	The relationship between carotid and coronary atherosclerotic damage in dialysis patients. 2007 , 30, 315-20	5

1304	Correlation among obstructive sleep apnea syndrome, coronary atherosclerosis and coronary heart disease. 2007 , 120, 1632-1634	4
1303	Incidence of changes and predictive factors for sexual function after coronary stenting. 2007 , 39, 16-21	9
1302	N-terminal pro-brain natriuretic peptide and coronary artery stenosis. 2007 , 37, 596; author reply 597	
1301	Serum sodium concentration is significantly associated with the angiographic characteristics of coronary atherosclerosis. 2007 , 28, 1136-42	7
1300	The role of paraoxonase (PON) enzyme in the extent and severity of the coronary artery disease in type-2 diabetic patients. 2007 , 22, 158-64	15
1299	DNA repair gene XRCC1 and XPD polymorphisms and their association with coronary artery disease risks and micronucleus frequency. 2007 , 22, 355-60	34
1298	Interleukin-18: a strong predictor of the extent of coronary artery disease in patients with unstable angina. 2007 , 22, 371-5	25
1297	Molecular scanning of the human carboxypeptidase E gene for mutations in Chinese subjects with coronary atherosclerosis. 2008 , 307, 31-9	4
1296	Level of CD14+ endothelial progenitor cells is not associated with coronary artery disease or cardiovascular risk factors. 2008 , 30, 319-26	6
1295	Investigating the association between K198N coding polymorphism in EDN1 and hypertension, lipoprotein levels, the metabolic syndrome and cardiovascular disease. 2008 , 123, 307-13	15
1294	Association of PARL rs3732581 genetic variant with insulin levels, metabolic syndrome and coronary artery disease. 2008 , 124, 263-70	5
1293	Determinants of coronary arterial flow-mediated dilatation following percutaneous coronary intervention. 2008 , 71, 327-32	2
1292	Correlation between lipoprotein(a) serum concentration and severity of coronary artery stenosis in an Iranian population according to Gensini score. 2008 , 41, 117-20	18
1291	Association of human carboxypeptidase E exon5 gene polymorphisms with angiographical characteristics of coronary atherosclerosis in a Chinese population. 2008 , 29, 736-44	6
1290	Determining the relationship between metabolic syndrome score and angiographic severity of coronary artery disease. 2008 , 62, 717-22	15
1289	Insulin resistance phenotypes and coronary artery disease in a native Pakistani cohort. 2008 , 62, 701-7	2
1288	Increased serum CXCL16 level is a marker for acute coronary syndromes. 2008 , 39, 332-7	15
1287	Circulating CXCL16 is related to the severity of coronary artery stenosis. 2008 , 39, 531-5	9

1286	Decreased circulating protective adiponectin level is associated with angiographic coronary disease progression in patients with angina pectoris. 2008 , 129, 76-80	28
1285	Waist circumference, metabolic syndrome and coronary artery disease in a Pakistani cohort. 2008 , 128, 77-82	7
1284	Circulating endothelial microparticles correlate inversely with endothelial function in patients with ischemic left ventricular dysfunction. 2008 , 14, 336-40	52
1283	Topography and severity of coronary artery disease in white-coat hypertension. 2008 , 19, 280-4	6
1282	Diabetes exacerbates angiographic coronary lesion progression in subjects with metabolic syndrome independent of CRP levels. 2008 , 388, 41-5	20
1281	Thoracic aorta atherosclerosis burden index predicts coronary artery disease in patients undergoing transesophageal echocardiography. 2008 , 197, 232-6	9
1280	The relationship between the metabolic syndrome defined by various criteria and the extent of coronary artery disease. 2008 , 197, 944-50	19
1279	The apolipoprotein All rs5082 variant is associated with reduced risk of coronary artery disease in an Australian male population. 2008 , 199, 333-9	23
1278	Key role of postchallenge hyperglycemia for the presence and extent of coronary atherosclerosis: an angiographic study. 2008 , 199, 317-22	36
1277	High triglyceride level is associated with severe coronary artery disease in hypertensive subjects. 2008 , 42, 146-52	6
1276	Relation of serum uric acid levels with the presence and severity of angiographic coronary artery disease. 2008 , 59, 166-71	20
1275	Racial differences in coronary artery lesions: a comparison of coronary artery lesions between mainland Chinese and Australian patients. 2008 , 59, 442-7	8
1274	Plasma C-reactive protein and 5-lipoxygenase-activating protein gene promoter poly-A polymorphism in patients with coronary artery disease. 2008 , 109, 25-32	3
1273	Proinsulin is an independent predictor of the angiographical characteristics of coronary atherosclerosis. 2008 , 110, 106-11	8
1272	Association of glycosylated haemoglobin level and diabetes mellitus duration with the severity of coronary artery disease. 2008 , 5, 184-9	22
1271	Truncal fat determined by dual-energy X-ray absorptiometry is an independent predictor of coronary artery disease extension. 2008 , 15, 428-33	4
1270	Diagnostic value of ischaemia-modified albumin for predicting myocardial ischaemia during myocardial perfusion scintigraphy. 2008 , 36, 147-51	1
1269	Plasma B-type natriuretic peptide levels reflect the presence and severity of stable coronary artery disease in chronic haemodialysis patients. 2009 , 24, 597-603	16

1268	The association of elasticity indexes of ascending aorta and the presence and the severity of coronary artery disease. 2008 , 19, 311-7		22
1267	The association of serum prolidase activity with the presence and severity of coronary artery disease. 2008 , 19, 319-25		45
1266	Impact of plasma adiponectin levels to the presence and severity of coronary artery disease in patients with metabolic syndrome. 2008 , 19, 79-84		17
1265	Small LDL-cholesterol is superior to LDL-cholesterol for determining severe coronary atherosclerosis. 2008 , 15, 250-60		68
1264	Echocardiographic Epicardial Fat Thickness on Short Term Prognosis in Patients with Acute Coronary Syndrome. 2008 , 16, 42		9
1263	Correlation between myocardial enzyme serum levels and markers of inflammation with severity of coronary artery disease and Gensini score: a hospital-based, prospective study in Greek patients. 2008 , 3, 699-710		23
1262	Vascular compliance is reduced in geriatric people with angiographic coronary atherosclerosis. 2009 , 37, 1443-9		0
1261	Correlation between pancreatic microcirculation and type 2 diabetes in patients with coronary artery disease: dynamic contrast-enhanced MR imaging. 2009 , 252, 704-11		18
1260	Reduction of circulating soluble fms-like tyrosine kinase-1 plays a significant role in renal dysfunction-associated aggravation of atherosclerosis. 2009 , 120, 2470-7		44
1259	Serum levels of remnant lipoprotein cholesterol and oxidized low-density lipoprotein in patients with coronary artery disease. 2009 , 53, 108-16		25
1258	Usefulness of serum cathepsin L as an independent biomarker in patients with coronary heart disease. <i>American Journal of Cardiology</i> , 2009 , 103, 476-81	3	41
1257	Association of neutrophil gelatinase-associated lipocalin with the severity of coronary artery disease. <i>American Journal of Cardiology</i> , 2009 , 104, 917-20	3	78
1256	Aortic and mitral valve atherosclerosis: Predictive factors and associations with coronary atherosclerosis using Gensini score. 2009 , 40, 124-7		11
1255	Potential link of microalbuminuria with metabolic syndrome in patients undergoing coronary angiography. 2009 , 40, 399-405		8
1254	Correlation among soluble markers and severity of disease in non-diabetic subjects with pre-mature coronary artery disease. 2009 , 330, 201-9		19
1253	Association of the mutation for the human carboxypeptidase E gene exon 4 with the severity of coronary artery atherosclerosis. 2009 , 36, 245-54		10
1252	Nonlinear association between serum testosterone levels and coronary artery disease in Iranian men. 2009 , 24, 297-306		15
1251	Coronary angiographic studies of impaired glucose regulation and coronary artery disease in Chinese nondiabetic subjects. 2009 , 36, 457-63		12

1250	Effects of chronic obstructive pulmonary disease on coronary atherosclerosis. 2009 , 24, 164-8	32
1249	Diagnostic value of plasma fibronectin level in predicting the presence and severity of coronary artery disease. 2009 , 88, 249-53	8
1248	Increased epicardial adipose tissue (EAT) volume in type 2 diabetes mellitus and association with metabolic syndrome and severity of coronary atherosclerosis. 2009 , 70, 876-82	163
1247	Association of serum sodium concentration with coronary atherosclerosis in China: follow-up study. 2009 , 30, 494-500	1
1246	Smoking and coronary atherosclerosis: follow-up study in China. 2009 , 36, 690-5	9
1245	Hepatocyte growth factor plays a critical role in the regulation of cytokine production and induction of endothelial progenitor cell mobilization: a pilot gene therapy study in patients with coronary heart disease. 2009 , 36, 790-6	26
1244	Evaluation of apolipoprotein M as a biomarker of coronary artery disease. 2009 , 42, 365-70	10
1243	Association of estrogen receptor alpha gene polymorphism with the presence of coronary artery disease documented by coronary angiography. 2009 , 42, 835-9	19
1242	Siglec-1 on monocytes is a potential risk marker for monitoring disease severity in coronary artery disease. 2009 , 42, 1057-63	25
1241	Overexpression of activated nuclear factor-kappa B in aorta of patients with coronary atherosclerosis. 2009 , 32, E42-7	15
1240	Total cholesterol content of erythrocyte membranes and coronary atherosclerosis: an intravascular ultrasound pilot study. 2009 , 60, 676-82	19
1239	Usefulness of brachial-ankle pulse wave velocity as a predictive marker of multiple coronary artery occlusive disease in Korean type 2 diabetes patients. 2009 , 85, 30-4	34
1238	Decreased ratio of high-molecular-weight to total adiponectin is associated with angiographic coronary atherosclerosis severity but not restenosis. 2009 , 405, 114-8	25
1237	Increased serum HMGB1 is related to the severity of coronary artery stenosis. 2009 , 406, 139-42	52
1236	Epicardial adipose tissue thickness by echocardiography is a marker for the presence and severity of coronary artery disease. 2009 , 19, 211-7	166
1235	Independent relationship of serum uric acid levels with leukocytes and coronary atherosclerotic burden. 2009 , 19, 729-35	16
1234	Functional genomics of blood cellular LXR-alpha gene in human coronary heart disease. 2009 , 46, 536-44	16
1233	Neopterin, CD4+CD28- lymphocytes and the extent and severity of coronary artery disease. 2009 , 135, 27-35	35

1232	The Syntax score predicts peri-procedural myocardial necrosis during percutaneous coronary intervention. 2009 , 135, 60-5	103
1231	Lipid profile and inflammatory markers associated with estrogen receptor alpha PvuII and XbaI gene polymorphisms. 2009 , 153, 288-95	6
1230	Is polymorphism within eNOS gene associated with the late onset of myocardial infarction? A pilot study. 2009 , 60, 588-95	18
1229	Helicobacter pylori infection is associated with decreased serum levels of high density lipoprotein, but not with the severity of coronary atherosclerosis. 2009 , 8, 59	45
1228	Inhibitory effects of Terminalia arjuna on platelet activation in vitro in healthy subjects and patients with coronary artery disease. 2009 , 20, 183-90	21
1227	Small arterial elasticity predicts the extent of coronary artery disease: Relationship with serum uric acid. 2009 , 202, 200-4	10
1226	Comparison of the long-term prognostic value of cystatin C to other indicators of renal function, markers of inflammation and systolic dysfunction among patients with acute coronary syndrome. 2009 , 207, 552-8	31
1225	Receptor for advanced glycation end products (RAGE) and its inflammatory ligand EN-RAGE in non-diabetic subjects with pre-mature coronary artery disease. 2009 , 207, 597-602	54
1224	Serial changes in circulating M30 antigen, a biomarker of apoptosis, in patients with acute coronary syndromes: relationship with the severity of coronary artery disease. 2009 , 20, 494-8	8
1223	Increased epicardial fat volume quantified by 64-multidetector computed tomography is associated with coronary atherosclerosis and totally occlusive lesions. 2009 , 73, 1927-33	87
1222	Serum choline levels in patients with stable angina and acute coronary syndromes: relation to the severity of coronary artery disease. 2010 , 21, 466-71	3
1221	The role of heart-type fatty acid-binding protein in predicting properties of coronary atherosclerosis in patients with acute coronary syndrome. 2010 , 21, 435-40	3
1220	Mycotic DNA in non-atherosclerotic aortic wall of coronary patients is associated with sICAM-1 expression. 2010 , 74, 749-53	1
1219	Coronary score adds prognostic information for patients with acute coronary syndrome. 2010 , 74, 490-5	39
1218	Plasma cystatin C concentration reflects the severity of coronary artery disease in patients without chronic kidney disease. 2010 , 74, 2441-7	22
1217	Cardio-ankle vascular index is independently associated with the severity of coronary atherosclerosis and left ventricular function in patients with ischemic heart disease. 2010 , 17, 249-58	68
1216	Anger rumination, social support, and cardiac symptoms in patients undergoing angiography. 2010 , 15, 841-57	11
1215	Growth differentiation factor 15 and coronary collateral formation. 2010 , 33, E1-5	7

1214	Apolipoprotein E polymorphism and the characteristics of diseased vessels in male Chinese patients with angiographic coronary artery disease: a case-case study. 2010 , 33, E30-4	7
1213	The association of circulating monocyte count with coronary collateral growth in patients with diabetes mellitus. 2010 , 47, 49-54	10
1212	Safety and effectiveness of percutaneous coronary intervention (PCI) in elderly patients. a 5-year consecutive study of 201 cases with PCI. 2010 , 51, 312-6	8
1211	The association between serum uric acid level and coronary artery disease. 2010 , 64, 900-7	39
1210	Abdominal visceral fat thickness measured by ultrasonography predicts the presence and severity of coronary artery disease. 2010 , 36, 1769-75	24
1209	Highly sensitive C-reactive protein and male gender are independently related to the severity of coronary disease in patients with metabolic syndrome and an acute coronary event. 2010 , 43, 297-302	11
1208	Impact of metabolic syndrome and its individual components on the presence and severity of angiographic coronary artery disease. 2010 , 51, 676-82	32
1207	The value of brachial-ankle pulse wave velocity as a predictor of coronary artery disease in high-risk patients. 2010 , 40, 224-9	21
1206	Correlates of the severity of coronary atherosclerosis in long-term kidney transplant patients. 2010 , 25, 706-11	2
1205	The relationship between glucose tolerance and severity of coronary artery disease using the Gensini score. 2010 , 61, 751-5	24
1204	Fibroblast growth factor 23 and fetuin A are independent predictors for the coronary artery disease extent in mild chronic kidney disease. 2010 , 5, 1780-6	65
1203	Metabolic syndrome and angiographic coronary artery disease prevalence in association with the Framingham risk score. 2010 , 8, 201-8	9
1202	Association of reduced zinc status with angiographically severe coronary atherosclerosis: a pilot study. 2010 , 61, 449-55	23
1201	Relationship between coronary artery disease and epicardial adipose tissue quantification at cardiac CT: comparison between automatic volumetric measurement and manual bidimensional estimation. 2010 , 17, 727-34	39
1200	Serum levels of apelin and ghrelin in patients with acute coronary syndromes and established coronary artery disease--KOZANI STUDY. 2010 , 155, 238-46	88
1199	The association of serum prolidase activity and erectile dysfunction. 2010 , 31, 146-54	19
1198	Relationship between the Gensini Score of Blood-Stasis Syndrome in Coronary Heart Disease and VEGF. 2010 , 12, 355-357	4
1197	The relationship between microalbuminuria and the presence and extent of coronary atherosclerosis. 2010 , 61, 184-91	9

1196	Non-diabetic hyperglycaemia correlates with angiographic coronary artery disease prevalence and severity. 2010 , 36, 402-8	16
1195	Serum adipocyte fatty acid binding proteins and adiponectin in patients with coronary artery disease: the significance of A-FABP/adiponectin ratio. 2010 , 411, 1761-5	17
1194	Serum levels of total p-cresylsulphate are associated with angiographic coronary atherosclerosis severity in stable angina patients with early stage of renal failure. 2010 , 211, 579-83	59
1193	Decreased serum levels of thioredoxin in patients with coronary artery disease plus hyperhomocysteinemia is strongly associated with the disease severity. 2010 , 212, 351-5	24
1192	Association of epicardial adipose tissue with coronary atherosclerosis is region-specific and independent of conventional risk factors and intra-abdominal adiposity. 2010 , 213, 279-87	76
1191	Visceral adiposity and the severity of coronary artery disease in middle-aged subjects with normal waist circumference and its relation with lipocalin-2 and MCP-1. 2010 , 213, 592-7	46
1190	Coronary heart disease is associated with mean platelet volume in type 2 diabetic patients. 2010 , 21, 368-72	47
1189	Microalbuminuria is not associated with endothelial dysfunction and coronary atherosclerosis in patients with acute coronary syndromes. 2010 , 32, 659-65	5
1188	The correlation between lipids ratio and degree of coronary artery stenosis. 2011 , 18, 53-6	16
1187	The same chromosome 9p21.3 locus is associated with type 2 diabetes and coronary artery disease in a Chinese Han population. 2011 , 60, 680-4	41
1186	Coronary atherosclerosis is associated with macrophage polarization in epicardial adipose tissue. 2011 , 58, 248-55	271
1185	The association of liver transaminase activity with presence and severity of premature coronary artery disease. 2011 , 62, 614-9	18
1184	Serum levels of vaspin and visfatin in patients with coronary artery disease-Kozani study. 2011 , 412, 48-52	83
1183	Increased plasma lipoprotein-associated phospholipase A2 (Lp-PLA2) levels are related to good collateral development in patients with isolated left coronary artery disease. 2011 , 148, 117-9	3
1182	Influence of body mass index on extent of coronary atherosclerosis and cardiac events in a cohort of patients at risk of coronary artery disease. 2011 , 21, 86-93	24
1181	Risk factors of cardiac troponin T elevation in patients with stable coronary artery disease after elective coronary drug-eluting stent implantation. 2011 , 34, 768-73	3
1180	Non-dimensional modeling in flow simulation studies of coronary arteries including side-branches: a novel diagnostic tool in coronary artery disease. 2011 , 216, 277-82	13
1179	Different association of circulating levels of adipocyte and epidermal fatty acid-binding proteins with metabolic syndrome and coronary atherosclerosis in Chinese adults. 2011 , 217, 194-200	13

1178	Study on the role of environmental parameters and HIF-1A gene polymorphism in coronary collateral formation among patients with ischemic heart disease. 2011 , 44, 1421-4		20
1177	Identification of ADAMTS7 as a novel locus for coronary atherosclerosis and association of ABO with myocardial infarction in the presence of coronary atherosclerosis: two genome-wide association studies. 2011 , 377, 383-92		399
1176	Procollagen III N-terminal peptide predicts short-term prognosis and cardiac remodeling in coronary heart disease patients with metabolic syndrome. 2011 , 341, 10-6		2
1175	Plasma neutrophil gelatinase-associated lipocalin levels in acute myocardial infarction and stable coronary artery disease. 2011 , 22, 333-8		47
1174	Time rate of blood pressure variation: a new factor associated with coronary atherosclerosis. 2011 , 29, 1109-14		29
1173	Circulating neuregulin-1 levels vary according to the angiographic severity of coronary artery disease and ischemia. 2011 , 22, 577-82		33
1172	Analysis of the correlation between non-high density lipoprotein cholesterol and coronary heart disease in elderly Chinese. 2011 , 50, 1279-85		3
1171	Severity of coronary atherosclerosis is an independent predictor of the left ventricular ejection fraction. 2011 , 38, 109-12		3
1170	Association of AGER gene G82S polymorphism with the severity of coronary artery disease in Chinese Han population. 2011 , 75, 470-4		4
1169	Understanding coronary atherosclerosis in relation to obesity: is getting the distribution of body fatness using dual-energy X-ray absorptiometry worth the effort? A novel perspective using Bayesian Networks. 2011 , 17, 32-9		3
1168	Elevated concentrations of pentraxin 3 are associated with coronary plaque vulnerability. 2011 , 58, 151-7		39
1167	B-type natriuretic peptide levels predict extent and severity of coronary disease in non-ST elevation coronary syndromes and normal left ventricular systolic function. 2011 , 167, 129-33		9
1166	High-sensitivity troponin T level and angiographic severity of coronary artery disease. <i>American Journal of Cardiology</i> , 2011 , 108, 639-43	3	42
1165	Comparison of the frequency of coronary artery disease in alcohol-related versus non-alcohol-related endstage liver disease. <i>American Journal of Cardiology</i> , 2011 , 108, 1552-5	3	40
1164	CagA-positive <i>Helicobacter pylori</i> strains enhanced coronary atherosclerosis by increasing serum OxLDL and HsCRP in patients with coronary heart disease. 2011 , 56, 109-14		37
1163	Relationship between major adverse cardiac events and angiographic findings in dialysis patients. 2011 , 43, 1171-8		6
1162	The severity of coronary atherosclerosis in diabetic and non-diabetic metabolic syndrome patients diagnosed according to different criteria and undergoing elective angiography. 2011 , 48, 21-7		13
1161	Plasma concentrations of soluble CD40 ligand in smokers with acute myocardial infarction: a pilot study. 2011 , 26, 131-7		12

1160	The S447X variant of lipoprotein lipase gene is inversely associated with severity of coronary artery disease. 2011 , 26, 457-63	8
1159	Lack of association between plasma myeloperoxidase levels and angiographic severity of coronary artery disease in patients with acute coronary syndrome. 2011 , 60, 137-42	7
1158	The relationship between serum levels of Zn and Cu and severity of coronary atherosclerosis. 2011 , 144, 436-44	19
1157	Coping mediates the association between Type D personality and perceived health in Chinese patients with coronary heart disease. 2011 , 18, 277-84	29
1156	Association of chemerin mRNA expression in human epicardial adipose tissue with coronary atherosclerosis. 2011 , 10, 87	61
1155	Multimodal cardiovascular magnetic resonance quantifies regional variation in vascular structure and function in patients with coronary artery disease: relationships with coronary disease severity. 2011 , 13, 61	9
1154	Pro/anti-inflammatory cytokines in the pathogenesis of premature coronary artery disease. 2011 , 31, 561-7	22
1153	Thymosin beta 4 is associated with collateral development in coronary artery disease. 2011 , 71, 625-30	9
1152	Angiotensinogen gene polymorphism in acute myocardial infarction patients. 2011 , 12, 42-7	15
1151	Inflammatory genes in epicardial fat contiguous with coronary atherosclerosis in the metabolic syndrome and type 2 diabetes: changes associated with pioglitazone. 2011 , 34, 730-3	87
1150	Correlation of femoral intima-media thickness and the severity of coronary artery disease. 2011 , 62, 134-9	23
1149	OX40 ligand levels and high-sensitivity C-reactive protein levels in blood from local coronary plaque and the femoral artery in patients with acute coronary syndrome or stable angina. 2011 , 39, 1275-83	6
1148	AGER gene -374T>A (rs1800624) polymorphism is not associated with the severity of non-diabetic coronary artery disease in Han Chinese. 2011 , 49, 547-9	1
1147	The relation of non-invasively and invasively assessed aortic pulsatile indices to the presence and severity of coronary artery disease. 2011 , 20, 309-16	5
1146	Uric acid and pentraxin-3 levels are independently associated with coronary artery disease risk in patients with stage 2 and 3 kidney disease. 2011 , 33, 325-31	39
1145	Using an abnormal increase in postexercise systolic blood pressure to diagnose coronary artery disease in gerontal patients. 2011 , 39, 637-46	2
1144	Depot-specific overexpression of proinflammatory, redox, endothelial cell, and angiogenic genes in epicardial fat adjacent to severe stable coronary atherosclerosis. 2011 , 9, 433-9	45
1143	Computational high-resolution heart phantoms for medical imaging and dosimetry simulations. 2011 , 56, 5845-64	12

1142	Relationship between total antioxidant capacity and the severity of coronary artery disease. 2012 , 3,	4
1141	Total cholesterol content of erythrocyte membranes is associated with the severity of coronary artery disease and the therapeutic effect of rosuvastatin. 2012 , 117, 390-8	21
1140	SYNTAX score: an independent predictor of long-term cardiac mortality in patients with acute ST-elevation myocardial infarction. 2012 , 23, 445-9	28
1139	Relation of aortic calcification, wall thickness, and distensibility with severity of coronary artery disease: evaluation with coronary CT angiography. 2012 , 53, 839-44	10
1138	Absence and Resolution of Fragmented QRS Predict Reversible Myocardial Ischemia With Higher Probability of ST Segment Resolution in Patients With ST Segment Elevation Myocardial Infarction. 2012 , 42, 674-83	13
1137	B-type natriuretic peptide as an independent predictor of coronary disease extension in non-ST elevation coronary syndromes with preserved systolic function. 2012 , 19, 366-73	8
1136	Microalbuminuria is associated with the severity of coronary artery disease independently of other cardiovascular risk factors. 2012 , 63, 457-60	13
1135	High levels of serum uric acid predict severity of coronary artery disease in patients with acute coronary syndrome. 2012 , 63, 448-52	38
1134	Associations between coronary heart disease and individual components of the metabolic syndrome according to glucose tolerance status. 2012 , 40, 934-42	1
1133	The association between different monocyte subsets and coronary collateral development. 2012 , 23, 16-21	8
1132	Association of coronary atherosclerotic burden with clinical presentation and prognosis in patients with stable and unstable coronary artery disease. 2012 , 101, 1003-11	29
1131	Association of serum calcitonin with coronary artery disease in individuals with and without chronic kidney disease. 2012 , 44, 1169-75	8
1130	Family history does not predict angiographic localization or severity of coronary artery disease. 2012 , 221, 451-7	4
1129	Regulatory T cells and IL-10 levels are reduced in patients with vulnerable coronary plaques. 2012 , 222, 519-23	63
1128	Do cardiac risk factors affect the homocysteine and asymmetric dimethylarginine relationship in patients with coronary artery diseases?. 2012 , 45, 1325-30	7
1127	Usefulness of sum of the thickness of plaque in the carotid artery for predicting the presence and the extent of the coronary artery disease in patients with type 2 diabetes mellitus without known coronary artery disease. 2012 , 96, 111-8	6
1126	Serum apolipoprotein A-V in patients with coronary artery disease and its association with triglyceride. 2012 , 6, 462-8	3
1125	Does metabolic syndrome predict significant angiographic coronary artery disease?. 2012 , 31, 769-778	2

1124	Does metabolic syndrome predict significant angiographic coronary artery disease?. 2012 , 31, 769-78	4
1123	Plasma ghrelin levels are closely associated with severity and morphology of angiographically-detected coronary atherosclerosis in Chinese patients with diabetes mellitus. 2012 , 33, 452-8	21
1122	Plasma adiponectin levels in relation to prognosis in patients with angiographic coronary artery disease. 2012 , 61, 1803-8	30
1121	Intérêt du dosage de la cystatine C chez des patients coronariens tunisiens. 2012 , 27, 20-27	
1120	Chitinase 3-like 1 gene-329G/A polymorphism, plasma concentration and risk of coronary heart disease in a Chinese population. 2012 , 499, 135-8	12
1119	Gender disparities in the association between epicardial adipose tissue volume and coronary atherosclerosis: a 3-dimensional cardiac computed tomography imaging study in Japanese subjects. 2012 , 11, 106	42
1118	Cystatin C and asymptomatic coronary artery disease in patients with metabolic syndrome and normal glomerular filtration rate. 2012 , 11, 108	27
1117	Insulin resistance predicts progression of de novo atherosclerotic plaques in patients with coronary heart disease: a one-year follow-up study. 2012 , 11, 71	44
1116	Apelin and vascular endothelial growth factor are associated with mobilization of endothelial progenitor cells after acute myocardial infarction. 2012 , 26, 400-9	13
1115	Adiponectin profile in Asian patients undergoing coronary revascularization and its association with plaque vulnerability: IDEAS-ADIPO study. 2012 , 20, 2451-7	2
1114	Coronary angiographic scoring systems: an evaluation of their equivalence and validity. 2012 , 164, 547-552.e1	129
1113	Correlation of coronary artery disease angiographic severity with intima-media thickness of carotid artery. 2012 , 2, 144-149	2
1112	Links between sleep disordered breathing, coronary atherosclerotic burden, and cardiac biomarkers in patients with stable coronary artery disease. 2012 , 60, 180-6	22
1111	Changes and significance of serum angiopoietin-2 levels in patients with coronary heart disease. 2012 , 17, 745-9	13
1110	Six lipoprotein lipase gene polymorphisms, lipid profile and coronary stenosis in a Tunisian population. 2012 , 39, 9893-901	18
1109	Plasma levels of lipometabolism-related miR-122 and miR-370 are increased in patients with hyperlipidemia and associated with coronary artery disease. 2012 , 11, 55	128
1108	High-density lipoprotein subclass and particle size in coronary heart disease patients with or without diabetes. 2012 , 11, 54	16
1107	Association of SNP rs6903956 on chromosome 6p24.1 with angiographical characteristics of coronary atherosclerosis in a Chinese population. 2012 , 7, e43732	17

1106	The relationship between the severity of coronary artery disease and epicardial adipose tissue depends on the left ventricular function. 2012 , 7, e48330		8
1105	Higher plasma thrombospondin-1 levels in patients with coronary artery disease and diabetes mellitus. 2012 , 42, 100-6		37
1104	The independent relationship of systemic inflammation with fragmented QRS complexes in patients with acute coronary syndromes. 2012 , 42, 449-57		13
1103	Is there any relationship between serum γ -glutamyltransferase levels and premature coronary artery disease?. 2012 , 2, 96-101		
1102	Polymorphisms of some increased cardiovascular risk related genes in acute coronary syndromes and their relation to coronary artery disease severity. 2012 , 1, 25-30		
1101	Comparison of aortic and carotid arterial stiffness parameters in patients with verified coronary artery disease. 2012 , 35, 26-31		25
1100	Angiotensin converting enzyme gene polymorphism is associated with severity of coronary artery disease in men with high total cholesterol levels. 2012 , 53, 175-82		16
1099	The association of serum chemerin level with risk of coronary artery disease in Chinese adults. 2012 , 41, 281-8		62
1098	Association of four apolipoprotein B polymorphisms with lipid profile and stenosis in Tunisian coronary patients. 2012 , 91, 75-9		16
1097	N-terminal pro-B-type natriuretic peptide is associated with severity of the coronary lesions in unstable angina patients with preserved left ventricular function. 2012 , 25, 126-31		8
1096	Prognostic usefulness of clinical and subclinical peripheral arterial disease in men with stable coronary heart disease. <i>American Journal of Cardiology</i> , 2012 , 110, 197-202	3	25
1095	Six-year prognosis of diabetic patients with coronary artery disease. 2012 , 42, 376-83		12
1094	Relationship of serum fetuin-A levels with coronary atherosclerotic burden and NAFLD in patients undergoing elective coronary angiography. 2013 , 11, 289-95		21
1093	The association between extracoronary calcification and coronary artery disease in patients with type 2 diabetes mellitus. 2013 , 28, 12-8		19
1092	Common variants in adiponectin gene are associated with coronary artery disease and angiographical severity of coronary atherosclerosis in type 2 diabetes. 2013 , 12, 67		21
1091	Changes of serum angiogenic factors concentrations in patients with diabetes and unstable angina pectoris. 2013 , 12, 34		22
1090	Prediction of coronary risk by SYNTAX and derived scores: synergy between percutaneous coronary intervention with taxus and cardiac surgery. 2013 , 62, 1219-1230		81
1089	Angiographic score assessment improves cardiovascular risk prediction: the clinical value of SYNTAX and Gensini application. 2013 , 102, 495-503		106

1088	Fasting plasma glucose correlates with angiographic coronary artery disease prevalence and severity in Chinese patients without known diabetes. 2013 , 50, 333-40	11
1087	Association between erectile dysfunction and coronary artery disease and its severity. 2013 , 65, 180-6	15
1086	Decreased plasma nesfatin-1 levels in patients with acute myocardial infarction. 2013 , 46, 167-71	30
1085	Leptin to adiponectin ratio as a useful predictor for cardiac syndrome X. 2013 , 18, 44-50	17
1084	Severity of coronary artery disease and echocardiographic parameters of ventricular diastolic function. 2014 , 31, 809-13	2
1083	Differential associations of angiographic extent and severity of coronary artery disease with asymmetric dimethylarginine but not insulin resistance in non-diabetic men with stable angina: a cross-sectional study. 2013 , 12, 145	10
1082	Usefulness of metabolic syndrome score in the prediction of angiographic coronary artery disease severity according to the presence of diabetes mellitus: relation with inflammatory markers and adipokines. 2013 , 12, 140	18
1081	Long-term prognostic impact of cystatin C on acute coronary syndrome octogenarians with diabetes mellitus. 2013 , 12, 157	11
1080	Correlation between the severity of periodontitis and coronary artery stenosis in a Chinese population. 2013 , 58, 333-8	8
1079	Mean platelet volume: an important predictor of coronary collateral development. 2013 , 24, 200-4	13
1078	Association of two variants in the interleukin-6 receptor gene and premature coronary heart disease in a Chinese Han population. 2013 , 40, 1021-6	5
1077	Association between nonalcoholic fatty liver disease and coronary artery disease. 2013 , 59, 290-7	11
1076	Low HDL-C predicts risk and PCI outcomes in the Han Chinese population. 2013 , 226, 193-7	8
1075	Metabolically obese status with normal weight is associated with both the prevalence and severity of angiographic coronary artery disease. 2013 , 62, 952-60	38
1074	Serum osteoprotegerin and osteopontin levels are associated with arterial stiffness and the presence and severity of coronary artery disease. 2013 , 167, 1924-8	84
1073	Non-invasively measured aortic wave reflection and pulse pressure amplification are related to the severity of coronary artery disease. 2013 , 62, 131-7	21
1072	Prehospital statin therapy and one-year mortality in patients with stable coronary artery disease undergoing percutaneous coronary intervention. 2013 , 24, 145-50	2
1071	Sagittal abdominal diameter to triceps skinfold thickness ratio: a novel anthropometric index to predict premature coronary atherosclerosis. 2013 , 227, 329-33	8

1070	NLRP3 inflammasomes show high expression in aorta of patients with atherosclerosis. 2013 , 22, 746-50	111
1069	Hemoglobin A1C in non-diabetic patients: an independent predictor of coronary artery disease and its severity. 2013 , 102, 225-32	25
1068	Adiponectin and long-term mortality in coronary artery disease participants and controls. 2013 , 33, e19-29	30
1067	Assessment of the 9p21.3 locus in severity of coronary artery disease in the presence and absence of type 2 diabetes. 2013 , 14, 11	21
1066	The prognostic significance of SYNTAX score after early percutaneous transluminal coronary angioplasty for acute ST elevation myocardial infarction. 2013 , 22, 341-5	10
1065	Effects of opium consumption on cardiometabolic diseases. 2013 , 10, 733-40	37
1064	Carotid intima-media thickness and carotid plaques improves prediction of obstructive angiographic coronary artery disease in women. 2013 , 64, 57-63	17
1063	New echocardiographic parameter of aortic stiffness and atherosclerosis in patients with coronary artery disease: aortic propagation velocity. 2013 , 62, 236-40	13
1062	Relation between mean platelet volume and severity of atherosclerosis in patients with acute coronary syndromes. 2013 , 64, 131-6	41
1061	Frequency and predictors of renal artery stenosis in hypertensive patients undergoing coronary angiography. 2013 , 64, 385-90	5
1060	Reduced paraoxonase 1 activity as a marker for severe coronary artery disease. 2013 , 35, 97-103	32
1059	Relationship of high sensitivity C-reactive protein with presence and severity of coronary artery disease. 2013 , 29, 1425-9	11
1058	The association of ABO blood groups with extent of coronary atherosclerosis in Croatian patients suffering from chronic coronary artery disease. 2013 , 23, 351-9	15
1057	Sonographic evaluation for predicting the presence and severity of coronary artery disease. 2013 , 29, 125-30	13
1056	Anxiety and adverse coronary artery disease outcomes in Chinese patients. 2013 , 75, 530-6	17
1055	Monocyte heterogeneity in myocardial infarction with and without ST elevation and its association with angiographic findings. 2013 , 24, 404-11	6
1054	Association between nonalcoholic fatty liver disease and coronary artery disease complexity in patients with acute coronary syndrome: a pilot study. 2013 , 64, 604-8	25
1053	Disturbed coronary hemodynamics in vessels with intermediate stenoses evaluated with fractional flow reserve: a combined analysis of epicardial and microcirculatory involvement in ischemic heart disease. 2013 , 128, 2557-66	110

1052	De Gruyter. 2013 , 21,	1
1051	Myocardial ischemia during mental stress: role of coronary artery disease burden and vasomotion. 2013 , 2, e000321	66
1050	Adaptive immunity is related to coronary artery disease severity after acute coronary syndrome in subjects with metabolic syndrome. 2013 , 10, 32-9	11
1049	Effect of transradial coronary angiography procedure on vascular diameter and vasodilator functions in the access site. 2013 , 64, 515-21	17
1048	A Pro 12 Ala substitution in the PPAR α polymorphism may decrease the number of diseased vessels and the severity of angiographic coronary artery. 2013 , 24, 347-51	9
1047	Central diastolic blood pressure is associated with the degree of coronary collateral development. 2013 , 64, 546-52	6
1046	Relationship between epicardial fat measured by echocardiography and coronary atherosclerosis: a single-blind historical cohort study. 2013 , 30, 505-11	32
1045	The effect of smoking on myocardial performance index in middle-aged males after first acute myocardial infarction. 2013 , 30, 155-63	2
1044	Cell-Free circulating DNA: a new biomarker for the acute coronary syndrome. 2013 , 124, 76-84	38
1043	Serum pentraxin 3 levels are associated with the complexity and severity of coronary artery disease in patients with stable angina pectoris. 2013 , 61, 278-85	32
1042	Presence of angina pectoris is related to extensive coronary artery disease in diabetic patients. 2013 , 36, 475-9	4
1041	Serum uric Acid is not an independent risk factor for premature coronary artery disease. 2013 , 3, 246-53	10
1040	Serum surfactant protein D is associated with the prognosis in patients with chronic kidney disease. 2013 , 14, 461-5	9
1039	The association of insulin-like growth factor-1 with severity of coronary artery disease. 2013 , 14, 416-20	7
1038	Aortic valve calcium score is associated with coronary calcified plaque burden. 2013 , 54, 355-61	6
1037	Association between serum inhibin-B levels and coronary artery disease in aging males. 2013 , 9, 796-801	3
1036	Decreased renal function is an independent predictor of severity of coronary artery disease: an application of Gensini score. 2013 , 28, 1615-21	12
1035	Maximal Exercise-Corrected QT as a Predictor of Coronary Artery Disease: Comparison of Simpler Heart Rate Corrections. 2013 , 43, 655-63	5

1034	Serum levels of fibroblast growth factor 19 are inversely associated with coronary artery disease in chinese individuals. 2013 , 8, e72345	24
1033	Usefulness of the neutrophil-to-lymphocyte ratio in predicting the severity of coronary artery disease: a Gensini score assessment. 2014 , 21, 1271-82	39
1032	Combined effects of admission serum creatinine concentration with age and gender on the prognostic significance of subjects with acute ST-elevation myocardial infarction in China. 2014 , 9, e108986	3
1031	Higher fibrinogen level is independently linked with the presence and severity of new-onset coronary atherosclerosis among Han Chinese population. 2014 , 9, e113460	23
1030	Elevated small dense low-density lipoprotein cholesterol as a predictor for future cardiovascular events in patients with stable coronary artery disease. 2014 , 21, 755-67	53
1029	A novel predictor of infarct-related artery patency before percutaneous intervention and in-hospital outcomes for ST-segment elevation myocardial infarction patients: serum bilirubin level. 2014 , 10, 91-7	6
1028	NLRP3 and downstream cytokine expression elevated in the monocytes of patients with coronary artery disease. 2014 , 10, 791-800	39
1027	The associations of epicardial adipose tissue with coronary artery disease and coronary atherosclerosis. 2014 , 55, 197-203	31
1026	Role of insulin-like growth factor 1 in stent thrombosis under effective dual antiplatelet therapy. 2014 , 10, 242-9	1
1025	Usefulness of the Doppler flow of the ophthalmic artery in the evaluation of carotid and coronary atherosclerosis. 2014 , 44, 406-14	9
1024	The role of insulin-like growth factor-1 in development of coronary no-reflow and severity of coronary artery disease in patients with acute myocardial infarction. 2014 , 10, 12-7	3
1023	Apelin: a potential marker of coronary artery stenosis and atherosclerotic plaque stability in ACS patients. 2014 , 55, 204-12	27
1022	Low serum adropin is associated with coronary atherosclerosis in type 2 diabetic and non-diabetic patients. 2014 , 52, 751-8	71
1021	Visceral adiposity index score indicated the severity of coronary heart disease in Chinese adults. 2014 , 6, 143	18
1020	The relation between neutrophil-to-lymphocyte ratio and coronary chronic total occlusions. 2014 , 14, 130	10
1019	Chemerin is a novel biomarker of acute coronary syndrome but not of stable angina pectoris. 2014 , 13, 145	27
1018	Elevated plasma IL-37, IL-18, and IL-18BP concentrations in patients with acute coronary syndrome. 2014 , 2014, 165742	68
1017	Association of fibrinogen with severity of stable coronary artery disease in patients with type 2 diabetic mellitus. 2014 , 2014, 485687	21

1016	An association study between genetic polymorphism in the interleukin-6 receptor gene and coronary heart disease. 2014 , 2014, 504727	8
1015	Central pulse pressure amplification is associated with more extensive and severe coronary artery disease. 2014 , 48, 167-75	5
1014	Levels of acylation stimulating protein and the complement component 3 precursor are associated with the occurrence and development of coronary heart disease. 2014 , 8, 1861-1866	6
1013	A genetic polymorphism in RBP4 is associated with coronary artery disease. 2014 , 15, 22309-19	9
1012	Clinical implications of tenascin-C and OX40 ligand in patients with acute coronary syndrome. 2014 , 2, 132-136	9
1011	Effects of atorvastatin loading prior to primary percutaneous coronary intervention on endothelial function and inflammatory factors in patients with ST-segment elevation myocardial infarction. 2014 , 7, 316-322	6
1010	The chromosome 9p21 variant not predicting long-term cardiovascular mortality in Chinese with established coronary artery disease: an eleven-year follow-up study. 2014 , 2014, 626907	6
1009	Impact of coronary collateral circulation and severity of coronary artery disease in the development of postoperative atrial fibrillation. 2014 , 19, 394-7	4
1008	Non-invasive assessment of functionally significant coronary stenoses through mathematical analysis of spectral ECG components. 2014 , 1, e000144	4
1007	Correlation between comprehensive evaluation of coronary artery lesion severity and long-term clinical outcomes in Chinese octogenarians with acute coronary syndrome. 2014 , 23, 1125-31	3
1006	Does estimated glomerular filtration rate have an effect on left ventricular function after ST-elevation myocardial infarction?. 2014 , 31, 318-24	4
1005	Mitral annulus displacement measured by two-dimensional speckle tracking imaging to assess the left ventricular longitudinal systolic function in coronary heart disease. 2014 , 42, 544-9	2
1004	Clinical utility of serum cystatin C in predicting coronary artery disease in patients without chronic kidney disease. 2014 , 28, 191-7	18
1003	Implication of plasma intermedin levels in patients who underwent first-time diagnostic coronary angiography: a single centre, cross-sectional study. 2014 , 14, 182	2
1002	The relationship of plasma neuropeptide Y levels with coronary collateral development. 2014 , 25, 73-8	4
1001	microRNA-155 is inversely associated with severity of coronary stenotic lesions calculated by the Gensini score. 2014 , 25, 304-10	37
1000	Association of vitamin D status with mental stress-induced myocardial ischemia in patients with coronary artery disease. 2014 , 76, 569-75	4
999	Sex differences in mental stress-induced myocardial ischemia in young survivors of an acute myocardial infarction. 2014 , 76, 171-80	75

998	Correlation between the serum alkaline phosphatase level and the severity of coronary artery disease. 2014 , 25, 349-52	11
997	Relationship between toll-like receptor 4 levels in aorta and severity of atherosclerosis. 2014 , 42, 958-65	8
996	The association study of plasma levels of pigment epithelium-derived factor with acute coronary syndrome in the chinese han population. 2014 , 127, 31-7	9
995	Validation of a novel clinical prediction score for severe coronary artery diseases before elective coronary angiography. 2014 , 9, e94493	12
994	Hand and heart, hand in hand: is radiological hand osteoarthritis associated with atherosclerosis?. 2014 , 17, 299-303	12
993	The severity of coronary artery disease and reversible ischemia revealed by N-terminal pro-brain natriuretic peptide in patients with unstable angina and preserved left ventricular function. 2014 , 52, 143-8	2
992	High-density lipoprotein cholesterol and risk of cardiovascular events in octogenarian patients with acute coronary syndrome: long-term follow-up study. 2014 , 174, 133-4	2
991	B-type natriuretic peptide levels predict extent and severity of coronary artery disease in non-ST elevation acute coronary syndrome and normal left ventricular function. 2014 , 66, 183-7	7
990	The association between Factor V Leiden with the presence and severity of coronary artery disease. 2014 , 47, 356-60	11
989	Association of epicardial fat thickness with TIMI risk score in NSTEMI/USAP patients. 2014 , 39, 755-60	17
988	Comment on: Triglyceride-to-HDL cholesterol ratio: Predictive value for CHD severity and new-onset heart failure. 2014 , 39, 356	
987	Framingham risk score and severity of coronary artery disease. 2014 , 39, 638-43	10
986	The role of bilirubin and its protective function against coronary heart disease. 2014 , 39, 711-5	9
985	Renal impairment and coronary collaterals in patients with acute coronary syndrome. 2014 , 39, 379-83	2
984	Triglyceride-to-HDL cholesterol ratio. Predictive value for CHD severity and new-onset heart failure. 2014 , 39, 105-10	21
983	Soluble TWEAK levels are independently associated with coronary artery disease severity in patients with stage 2-3 kidney disease. 2014 , 46, 411-5	5
982	Association between haptoglobin 2-2 genotype and coronary artery disease and its severity in a tunisian population. 2014 , 52, 269-82	7
981	Plasma Urotensin II Act as a Diagnostic Biomarker for Acute Coronary Syndromes. 2014 , 20, 145-151	

980	Human coronary heart disease: importance of blood cellular miR-2909 RNomics. 2014 , 392, 49-63	19
979	Relation between the GRACE score and severity of atherosclerosis in acute coronary syndrome. 2014 , 63, 24-8	24
978	Correlation of the coronary severity scores and mean platelet volume in diabetes mellitus. 2014 , 31, 140-8	11
977	Relation of ABO blood groups to the severity of coronary atherosclerosis: an Gensini score assessment. 2014 , 237, 748-53	32
976	Effects of thyroid dysfunction on the severity of coronary artery lesions and its prognosis. 2014 , 64, 496-500	16
975	Carotid intima media thickness and coronary atherosclerosis linkage in symptomatic intermediate risk patients evaluated by coronary computed tomography angiography. 2014 , 176, 988-93	27
974	Increased serum concentrations of pentosidine are related to presence and severity of coronary artery disease. 2014 , 134, 633-8	27
973	Glomerular filtration rate is associated with burden of coronary atherosclerosis in patients with acute coronary syndrome. 2014 , 65, 350-6	9
972	Lower vitamin D level is associated with poor coronary collateral circulation. 2014 , 48, 278-83	7
971	Associations between retinal microvascular structure and the severity and extent of coronary artery disease. 2014 , 236, 25-30	32
970	Relationship between increased serum resistin level and severity of coronary artery disease. 2014 , 65, 239-42	16
969	Serum total p-cresylsulfate level is associated with abnormal QTc interval in stable angina patients with early stage of renal failure. 2014 , 437, 25-30	11
968	Serum cystatin C levels are associated with coronary artery disease and its severity. 2014 , 47, 176-81	18
967	Association between neutrophil to lymphocyte ratio and severity of coronary artery disease. 2014 , 20, 50-4	83
966	Association between serum chemokine CC-motif ligand 17 and coronary artery disease. 2014 , 176, 541-2	3
965	The association between body mass index and coronary artery disease severity: a comparison of black and white patients. 2014 , 167, 514-20	11
964	Correlation of chronic renal dysfunction and albuminuria with severity of coronary artery lesions in patients with coronary artery disease. 2014 , 69, 55-9	2
963	Correlation between brachial-ankle pulse wave velocity, carotid artery intima-media thickness, ankle-brachial index, and the severity of coronary lesions. 2014 , 70, 1205-11	13

962	Impact of completeness of revascularization by coronary intervention on exercise capacity early after acute ST-elevation myocardial infarction. 2014 , 9, 50		4
961	Comparison by computed tomographic angiography-the presence and extent of coronary arterial atherosclerosis in South Asians versus Caucasians with diabetes mellitus. <i>American Journal of Cardiology</i> , 2014 , 113, 1782-7	3	16
960	The roles of a novel anti-inflammatory factor, milk fat globule-epidermal growth factor 8, in patients with coronary atherosclerotic heart disease. 2014 , 233, 661-665		25
959	Prediction of coronary artery disease severity using CHADS2 and CHA2DS2-VASc scores and a newly defined CHA2DS2-VASc-HS score. <i>American Journal of Cardiology</i> , 2014 , 113, 950-6	3	59
958	Plasma PCSK9 levels are associated with the severity of coronary stenosis in patients with atherosclerosis. 2014 , 174, 863-4		36
957	Overexpression of caspase-1 in aorta of patients with coronary atherosclerosis. 2014 , 23, 1070-4		9
956	Association between HMW adiponectin, HMW-total adiponectin ratio and early-onset coronary artery disease in Chinese population. 2014 , 235, 392-7		21
955	Relation of leukocytes and its subsets counts with the severity of stable coronary artery disease in patients with diabetic mellitus. 2014 , 9, e90663		15
954	Increased arterial stiffness after coronary artery revascularization correlates with serious coronary artery lesions and poor clinical outcomes in patients with chronic kidney disease. 2014 , 4, 280-9		1
953	Angiographic coronary artery disease and high-sensitivity troponin-T in a native Pakistani cohort presenting with chronic chest pain. 2014 , 68, 1473-7		
952	Angiographic evaluation of the effects of glucose metabolic status on progression of coronary artery lesions in patients with coronary artery disease. 2014 , 6, 541-6		4
951	Coronary Artery Disease Severity and Cardiovascular Biomarkers in Patients with Peripheral Artery Disease. 2015 , 24, 278-82		9
950	Detection of circulating IgG antibodies to apolipoprotein B100 in acute myocardial infarction. 2015 , 5, 712-6		5
949	Late percutaneous coronary intervention prevents left ventricular remodeling and improves clinical outcomes in patients with ST-elevation myocardial infarction. 2015 , 38, 82-91		6
948	Cyclophilin A in cardiovascular homeostasis and diseases. 2015 , 235, 1-15		35
947	Interaction between microRNA expression and classical risk factors in the risk of coronary heart disease. 2015 , 5, 14925		30
946	South Asians and Anglo Australians with heart disease in Australia. 2015 , 39, 568-576		2
945	Associations of lipid levels susceptibility loci with coronary artery disease in Chinese population. 2015 , 14, 80		9

944	Association of circulating angiogenesis inhibitors and asymmetric dimethyl arginine with coronary plaque burden. 2015 , 8, 13	8
943	Circulating adipokines and insulin resistance in subjects with combined cardiac and metabolic syndrome X. 2015 , 7, 83	7
942	Relationship between serum vitamin D levels and angiographic severity and extent of coronary artery disease. 2015 , 45, 940-8	13
941	Pro-protein convertase subtilisin/kexin 9 concentrations correlate with coronary artery disease atheroma burden in a Pakistani cohort with chronic chest pain. 2015 , 69, 738-42	8
940	Risk Factors, Coronary Severity, Outcome and ABO Blood Group: A Large Chinese Han Cohort Study. 2015 , 94, e1708	10
939	The Pivotal Role of a Novel Biomarker of Reactive Oxygen Species in Chronic Kidney Disease. 2015 , 94, e1040	14
938	Relationship between inter-arm blood pressure differences and future cardiovascular events in coronary artery disease. 2015 , 33, 1780-9; discussion 1790	14
937	The Combined Effect of Ear Lobe Crease and Conventional Risk Factor in the Diagnosis of Angiographically Diagnosed Coronary Artery Disease and the Short-Term Prognosis in Patients Who Underwent Coronary Stents. 2015 , 94, e815	7
936	Proprotein Convertase Subtilisin/Kexin type 9, C-Reactive Protein, Coronary Severity, and Outcomes in Patients With Stable Coronary Artery Disease: A Prospective Observational Cohort Study. 2015 , 94, e2426	35
935	A combination of the neutrophil-to-lymphocyte ratio and the GRACE risk score better predicts PCI outcomes in Chinese Han patients with acute coronary syndrome. 2015 , 15, 995-1001	14
934	The association between platelet/lymphocyte ratio and coronary artery disease severity. 2015 , 15, 640-7	59
933	A randomized comparison study assessing the impact of cilostazol on the heart rate and arrhythmias by 24-hour ambulatory holter electrocardiographic monitoring after drug-eluting stent implantation for coronary artery disease. 2015 , 22, 152-64	7
932	Association between Serum Fibroblast Growth Factor 21 and Coronary Artery Disease in Patients with Type 2 Diabetes. 2015 , 30, 586-90	18
931	Associations between Carotid Artery Plaque Score, Carotid Hemodynamics and Coronary Heart Disease. 2015 , 12, 14275-84	9
930	Hyperhomocysteinemia is an independent predictor of long-term clinical outcomes in Chinese octogenarians with acute coronary syndrome. 2015 , 10, 1467-74	15
929	The association between triglyceride/high-density lipoprotein cholesterol ratio and all-cause mortality in acute coronary syndrome after coronary revascularization. 2015 , 10, e0123521	41
928	Association of SNP Rs9943582 in APLNR with Left Ventricle Systolic Dysfunction in Patients with Coronary Artery Disease in a Chinese Han GeneID Population. 2015 , 10, e0125926	10
927	Correlations between Phlegm Syndrome of Chinese Medicine and Coronary Angiography: A Systematic Review and Meta-Analysis. 2015 , 2015, 751743	4

926	Polymorphism on Chromosome 9p21.3 Is Associated with Severity and Early-Onset CAD in Type 2 Diabetic Tunisian Population. 2015 , 2015, 792679	3
925	Association of Aortic Diameters with Coronary Artery Disease Severity and Albumin Excretion. 2015 , 2015, 857628	2
924	Association of Plasma Myeloperoxidase Level with Risk of Coronary Artery Disease in Patients with Type 2 Diabetes. 2015 , 2015, 761939	16
923	The relationship between neutrophil to lymphocyte ratio, platelet to lymphocyte ratio and thrombolysis in myocardial infarction risk score in patients with ST elevation acute myocardial infarction before primary coronary intervention. 2015 , 11, 126-35	8
922	Hemoglobin A1c Level Is Not Related to the Severity of Atherosclerosis in Patients with Acute Coronary Syndrome. 2015 , 2015, 192108	5
921	Relation between Apolipoprotein E Gene Polymorphism and Severity of Coronary Artery Disease in Acute Myocardial Infarction. 2015 , 2015, 363458	7
920	Serum Galectin-9 Levels Are Associated with Coronary Artery Disease in Chinese Individuals. 2015 , 2015, 457167	13
919	Thyroid Function, Prevalent Coronary Heart Disease, and Severity of Coronary Atherosclerosis in Patients Undergoing Coronary Angiography. 2015 , 2015, 708272	11
918	Elevated Concentration of Interferon-Inducible Protein of 10 kD (IP-10) Is Associated With Coronary Atherosclerosis. 2015 , 56, 269-72	19
917	Clinical utility of the ratio between circulating fibrinogen and fibrin (ogen) degradation products for evaluating coronary artery disease in type 2 diabetic patients. 2015 , 128, 727-32	
916	Fractional Flow Reserve Guided Percutaneous Coronary Intervention Improves Clinical Outcome with Reduced Cost in Contemporary Clinical Practice. 2015 , 128, 2000-5	4
915	Increased serum homocitrulline concentrations are associated with the severity of coronary artery disease. 2015 , 53, 103-10	20
914	Cardiac valve calcification is associated with presence and severity of coronary artery disease in patients with pre-dialysis chronic kidney disease. 2015 , 19, 1090-7	15
913	Coronary Artery Bypass Surgery and Percutaneous Coronary Revascularization: Impact on Morbidity and Mortality in Patients with Coronary Artery Disease. 2015 , 683-726	2
912	Role of ultrasound in the diagnosis and treatment of nonalcoholic fatty liver disease and its complications. 2015 , 9, 603-27	79
911	Inflammatory bowel disease, liver diseases and endothelial function: is there a linkage?. 2015 , 16, 11-21	20
910	The relationship of TIMI risk index with SYNTAX and Gensini risk scores in predicting the extent and severity of coronary artery disease in patients with STEMI undergoing primary percutaneous coronary intervention. 2015 , 9, 257-66	9
909	Red Blood Cell Distribution Width is Independently Correlated With Diurnal QTc Variation in Patients With Coronary Heart Disease. 2015 , 94, e822	3

908	Tissue Factor Pathway Inhibitor-2 Gene Polymorphisms Associate With Coronary Atherosclerosis in Chinese Population. 2015 , 94, e1675		3
907	Comparison of Angiographic Burden of Coronary Artery Disease in Patients With Versus Without Hepatitis C Infection. <i>American Journal of Cardiology</i> , 2015 , 116, 1041-4	3	16
906	Increased Platelet Distribution Width Is Associated With Severity of Coronary Artery Disease in Patients With Acute Coronary Syndrome. 2015 , 66, 638-43		25
905	Correlation of ascending aorta elasticity and the severity of coronary artery stenosis in hypertensive patients with coronary heart disease assessed by M-mode and tissue Doppler echocardiography. 2015 , 71, 785-8		7
904	Association of polymorphisms in long non-coding RNA H19 with coronary artery disease risk in a Chinese population. 2015 , 772, 15-22		99
903	High-density lipoprotein subfractions in relation with the severity of coronary artery disease: A Gensini score assessment. 2015 , 9, 26-34		31
902	Association of serum total bilirubin level with severity of coronary atherosclerosis is linked to systemic inflammation. 2015 , 240, 110-4		87
901	Severity of coronary artery disease is independently associated with the frequency of early age-related macular degeneration. 2015 , 99, 365-70		14
900	MiR-17-5p as circulating biomarkers for the severity of coronary atherosclerosis in coronary artery disease. 2015 , 197, 123-4		29
899	Lipoprotein-associated Phospholipase A2 is Associated with Angiographic Coronary Artery Disease and Coronary Artery Risk Factors in the Elderly. 2015 , 9, 82-86		2
898	Expression of NLRP3 in subcutaneous adipose tissue is associated with coronary atherosclerosis. 2015 , 242, 407-14		39
897	Clinical significance of pulse pressure in patients with coronary artery disease. 2015 , 190, 299-301		10
896	Evaluation of association between common genetic variants on chromosome 9p21 and coronary artery disease in Turkish population. 2015 , 15, 196-203		14
895	The prevalence of erectile dysfunction among angiographically detected patients of coronary artery disease in North-India. 2015 , 5, 272-276		
894	Epicardial adipose tissue thickness is associated with myocardial infarction and impaired coronary perfusion. 2015 , 15, 224-31		13
893	Different levels of glycosylated hemoglobin influence severity and long-term prognosis of coronary heart disease patients with stent implantation. 2015 , 9, 361-366		2
892	Relationship of renin-angiotensin-aldosterone system polymorphisms and phenotypes to mortality in Chinese coronary atherosclerosis patients. 2014 , 4, 4600		7
891	Epicardial adipose tissue thickness and acute coronary syndrome: a matter of how much or how?. 2015 , 199, 8-9		5

890	Serum chemokine CCL17/thymus activation and regulated chemokine is correlated with coronary artery diseases. 2015 , 238, 365-9	12
889	Angina and mental stress-induced myocardial ischemia. 2015 , 78, 433-437	19
888	The atherosclerosis burden score (ABS): a convenient ultrasound-based score of peripheral atherosclerosis for coronary artery disease prediction. 2015 , 8, 138-47	8
887	Association of male pattern baldness with angiographic coronary artery disease severity and collateral development. 2015 , 23, 265-74	10
886	Coronary severity score and C-reactive protein predict major adverse cardiovascular events in patients with stable coronary artery disease (from the Taichung CAD study). 2015 , 445, 93-100	17
885	A study of IgG antibodies to the ApoB protein in non-ST segment elevation acute coronary syndrome. 2015 , 49, 136-41	2
884	Clonal restriction and predominance of regulatory T cells in coronary thrombi of patients with acute coronary syndromes. 2015 , 36, 1041-8	30
883	Association of hydrogen sulfide with alterations of monocyte chemokine receptors, CCR2 and CX3CR1 in patients with coronary artery disease. 2015 , 64, 627-35	23
882	Association between the ABCA1-565C/T gene promoter polymorphism and coronary heart disease severity and cholesterol efflux in the Chinese Han population. 2015 , 19, 347-52	7
881	Hematological parameters can predict the extent of coronary artery disease in patients with end-stage renal disease. 2015 , 47, 1719-25	11
880	Associations between pentraxin 3 and severity of coronary artery disease. 2015 , 5, e007123	16
879	The roles of a novel inflammatory neopterin in subjects with coronary atherosclerotic heart disease. 2015 , 24, 169-172	19
878	Testosterone and high-sensitive C-reactive protein in coronary artery disease patients awaiting coronary artery bypass graft. 2015 , 47, 493-8	4
877	Association between anger and mental stress-induced myocardial ischemia. 2015 , 169, 115-21.e2	33
876	Treg/Th17 balance in stable CAD patients with different stages of coronary atherosclerosis. 2015 , 238, 17-21	59
875	Impact of waist circumference on hospital outcome and coronary angiographic findings of patients with acute ST-segment elevation myocardial infarction. 2015 , 67, 159-165	3
874	Link between angiographic extent and severity of coronary artery disease and degree of sensorineural hearing loss. 2015 , 40, 481-6	10
873	Pericardial fat volume is an independent risk factor for the severity of coronary artery disease in patients with preserved ejection fraction. 2015 , 65, 37-41	18

872	Serum Endocan Level and the Severity of Coronary Artery Disease: A Pilot Study. 2015 , 66, 727-31	60
871	The clinical, angiographic and prognosis characteristics of elderly patients with acute ST-segment elevation myocardial infarction--the first elderly STEMI population study in northwest of China. 2015 , 179, 326-8	3
870	Apolipoprotein B/apolipoprotein A1 ratio and non-high-density lipoprotein cholesterol. Predictive value for CHD severity and prognostic utility in CHD patients. 2015 , 40 Suppl 1, 1-7	16
869	Effects of Percutaneous Coronary Intervention on Serum Angiopoietin-2 in Patients with Coronary Heart Disease. 2016 , 129, 631-5	9
868	Relationship between osteopenic syndrome and severity of coronary artery disease detected with coronary angiography and Gensini score in men. 2016 , 11, 377-82	6
867	The Association between Platelet/Lymphocyte Ratio and Coronary Artery Disease Severity in Asymptomatic Low Ejection Fraction Patients. 2016 , 46, 821-826	15
866	The Correlation between Serum Level of N-Terminal Pro-B-type Natriuretic Peptide and Gensini Score in Patients with Acute Coronary Syndrome. 2016 , 1, 36-40	
865	The Relationship between P & QT Dispersions and Presence & Severity of Stable Coronary Artery Disease. 2016 , 46, 522-9	3
864	Association of rs10757274 and rs2383206 Polymorphisms on 9p21 locus with Coronary Artery Disease in Turkish Population. 2016 , 46, 615-621	5
863	Reduced T-Cell Thymic Export Reflected by sj-TREC in Patients with Coronary Artery Disease. 2016 , 23, 632-43	11
862	Elevated Levels of Plasma Superoxide Dismutases 1 and 2 in Patients with Coronary Artery Disease. 2016 , 2016, 3708905	13
861	The NLRP3 rs10754558 Polymorphism Is Associated with the Occurrence and Prognosis of Coronary Artery Disease in the Chinese Han Population. 2016 , 2016, 3185397	23
860	The Correlation of Serum Myeloid-Related Protein-8/14 and Eosinophil Cationic Protein in Patients with Coronary Artery Disease. 2016 , 2016, 4980251	10
859	Prevalence of Hepatitis C Virus Seropositivity and Its Impact on Coronary Artery Disease among Egyptian Patients Referred for Coronary Angiography. 2016 , 2016, 1623197	4
858	Putative consequences of exposure to Helicobacter pylori infection in patients with coronary heart disease in terms of humoral immune response and inflammation. 2016 , 12, 45-54	15
857	Human serum pre-β ₂ -high density lipoprotein levels are independently and negatively associated with coronary artery diseases. 2016 , 13, 36	12
856	Correlation between plasma angiopoietin-1, angiopoietin-2 and matrix metalloproteinase-2 in coronary heart disease. 2016 , 12, 1214-1219	8
855	Blood count parameters can predict the severity of coronary artery disease. 2016 , 31, 1093-1100	20

854	Associations between parameters of flow-mediated vasodilatation obtained by continuous measurement approaches and the presence of coronary artery disease and the severity of coronary atherosclerosis. 2016 , 38, 443-50	2
853	SYNTAX score-0 patients: risk stratification in nonobstructive coronary artery disease. 2016 , 105, 901-911	9
852	Increased Serum Level of Growth Differentiation Factor 15 (GDF-15) is Associated with Coronary Artery Disease. 2016 , 34, 138-43	14
851	Non-acute percutaneous coronary intervention versus medical therapy in patients with ischaemic heart disease. 2016 ,	1
850	Does a Reduction in the Glomerular Filtration Rate Increase the Overall Severity of Coronary Artery Stenosis?. 2016 , 55, 871-7	1
849	Relationship between diagonal earlobe creases and coronary artery disease as determined via angiography. 2016 , 6, e008558	15
848	Lower coronary plaque burden in patients with HIV presenting with acute coronary syndrome. 2016 , 3, e000511	9
847	Drug-eluting stents versus bare-metal stents for stable ischaemic heart disease. 2016 ,	1
846	IL-34 is associated with the presence and severity of renal dysfunction and coronary artery disease in patients with heart failure. 2016 , 6, 39324	21
845	Drug-eluting stents versus bare-metal stents for acute coronary syndrome. 2016 ,	2
844	Cataract surgery is more prevalent and occurs at an earlier age in a high cardiovascular risk cohort: Comparison with the Blue Mountains Eye Study. 2016 , 212, 72-5	4
843	The relationship between coronary atherosclerosis and body fat distribution measured using dual energy X-ray absorptiometry. 2016 , 248, 190-5	4
842	Polymorphisms of TGF β 1 and TGFBR2 in relation to coronary artery disease in a Chinese population. 2016 , 49, 873-8	11
841	Serum IgE levels are associated with coronary artery disease severity. 2016 , 251, 355-360	24
840	Aortic pulsatility assessed by an oscillometric method is associated with coronary atherosclerosis in elderly people. 2016 , 25, 373-380	6
839	Fragmented QRS complexes have predictive value of imperfect ST-segment resolution in patients with STEMI after primary percutaneous coronary intervention. 2016 , 34, 398-402	8
838	Hypertension is associated with narrower retinal arteriolar calibre in persons with and without coronary artery disease. 2016 , 30, 761-765	4
837	Association of combined genetic variations in PPAR α and LXRE with coronary artery disease and severity in Thai population. 2016 , 248, 140-8	10

836	Is monocyte to HDL ratio superior to monocyte count in predicting the cardiovascular outcomes: evidence from a large cohort of Chinese patients undergoing coronary angiography. 2016 , 48, 305-12	28
835	Large HDL Subfraction But Not HDL-C Is Closely Linked With Risk Factors, Coronary Severity and Outcomes in a Cohort of Nontreated Patients With Stable Coronary Artery Disease: A Prospective Observational Study. 2016 , 95, e2600	24
834	Plasma d-Dimer as a Useful Marker Predicts Severity of Atherosclerotic Lesion and Short-Term Outcome in Patients With Coronary Artery Disease. 2016 , 22, 633-40	11
833	Testosterone as a marker of coronary artery disease severity in middle aged males. 2016 , 68 Suppl 3, S16-S20	10
832	High-density lipoprotein subclass profile and mortality in patients with coronary artery disease: Results from the GENES study. 2016 , 109, 607-617	6
831	The APOA5 rs662799 polymorphism is associated with dyslipidemia and the severity of coronary heart disease in Chinese women. 2016 , 15, 170	14
830	Increased expression of the adipocytokine omentin in the epicardial adipose tissue of coronary artery disease patients. 2016 , 251, 299-304	31
829	Variants in 9p21 Predicts Severity of Coronary Artery Disease in a Chinese Han Population. 2016 , 80, 274-81	6
828	Usefulness of carotid plaque (sum and maximum of plaque thickness) in combination with intima-media thickness for the detection of coronary artery disease in asymptomatic patients with diabetes. 2016 , 7, 396-403	24
827	Age at Menopause and Extent of Coronary Artery Disease Among Postmenopausal Women with Acute Coronary Syndromes. 2016 , 129, 1205-1212	16
826	Non-HDL-C is a Better Predictor for the Severity of Coronary Atherosclerosis Compared with LDL-C. 2016 , 25, 975-81	30
825	Coronary Computed Tomography (CT) Angiography as a Predictor of Cardiac and Noncardiac Vascular Events in Asymptomatic Type 2 Diabetics: A 7-Year Population-Based Cohort Study. 2016 , 5,	16
824	Cardiac involvement in patients with cirrhosis: a focus on clinical features and diagnosis. 2016 , 17, 26-36	6
823	Time course of various cell origin circulating microparticles in ST-segment elevation myocardial infarction patients undergoing percutaneous transluminal coronary intervention. 2016 , 11, 1481-1486	7
822	Editorial : Scoring and Other Assessments of Plaque Morphology and Pathology. 2016 , 67, 889-893	
821	Analysis of the genetic association between IL27 variants and coronary artery disease in a Chinese Han population. 2016 , 6, 25782	5
820	Serum levels of mitochondrial inhibitory factor 1 are independently associated with long-term prognosis in coronary artery disease: the GENES Study. 2016 , 14, 125	13
819	The roles of the GPX1 Pro198Leu and OGG1 Ser326Cys variants in coronary artery disease / GPX1 geni Pro198Leu ve OGG1 geni Ser326Cys varyantları koroner arter hastalığındaki rolleri. 2016 , 41,	

818	Lipoprotein(a) and lipid profiles of patients awaiting coronary artery bypass graft; a cross sectional study. 2016 , 16, 213	1
817	Association of decreased serum sTREM-1 level with the severity of coronary artery disease: Inhibitory effect of sTREM-1 on TNF- α and oxLDL-induced inflammatory reactions in endothelial cells. 2016 , 95, e4693	6
816	Modification of the association between smoking status and severity of coronary stenosis by vitamin D in patients suspected of coronary heart disease. 2016 , 95, e4817	3
815	Urine proteomics in the diagnosis of stable angina. 2016 , 16, 70	16
814	The rs5888 single nucleotide polymorphism in scavenger receptor class B type 1 (SCARB1) gene and the risk of premature coronary artery disease: a case-control study. 2016 , 15, 7	17
813	Description of a Heart Team approach to coronary revascularization and its beneficial long-term effect on clinical events after PCI. 2016 , 105, 388-400	21
812	Transradial catheterization may decrease the radial artery luminal diameter and impair the vasodilatation response in the access site at late term: an observational study. 2016 , 31, 482-9	13
811	Relationship Between Hematologic Indices and Global Registry of Acute Coronary Events Risk Score in Patients With ST-Segment Elevation Myocardial Infarction. 2016 , 22, 60-8	36
810	Fibrinogen and the Severity of Coronary Atherosclerosis among Adults with and without Statin Treatment: Lipid as a mediator. 2016 , 25, 558-67	4
809	Serum Cystatin C and the Risk of Coronary Heart Disease in Ethnic Chinese Patients With Normal Renal Function. 2016 , 47, 13-9	4
808	Metabolic syndrome and retinal microvascular calibre in a high cardiovascular disease risk cohort. 2016 , 100, 1041-6	8
807	Evaluation of galectin-3 levels in acute coronary syndrome. 2016 , 65, 26-30	19
806	Association between vaspin level and coronary artery disease in patients with type 2 diabetes. 2016 , 113, 26-32	22
805	Circadian Rhythm of Infarct Size and Left Ventricular Function Evaluated with Tissue Doppler Echocardiography in ST Elevation Myocardial Infarction. 2016 , 25, 250-6	9
804	Platelet-to-Lymphocyte Ratio Improves the Predictive Power of GRACE Risk Score for Long-Term Cardiovascular Events in Patients with Acute Coronary Syndrome. 2016 , 134, 39-46	10
803	Epicardial fat thickness: A surrogate marker of coronary artery disease - Assessment by echocardiography. 2016 , 68, 336-41	29
802	Is galectin-3 a biomarker, a player-or both-in the presence of coronary atherosclerosis?. 2016 , 64, 764-70	23
801	In-hospital free fatty acids levels predict the severity of myocardial ischemia of acute coronary syndrome. 2016 , 16, 29	11

800	Predictive and prognostic value of admission neutrophil-to-lymphocyte ratio in patients with CHD. 2016 , 41, 605-613	19
799	Coronary Flow Reserve and Microcirculatory Resistance in Patients With Intermediate Coronary Stenosis. 2016 , 67, 1158-1169	175
798	Human Endothelial Cell-Specific Molecule-1 (Endocan) and Coronary Artery Disease and Microvascular Angina. 2016 , 67, 846-53	23
797	Correlation between epicardial adipose tissue and severity of coronary artery stenosis evaluated by 64-MDCT. 2016 , 40, 477-80	3
796	Hepatosteatosi s and carotid intima-media thickness in patients with myocardial infarction. 2016 , 43, 77-82	6
795	Volume elastic modulus of the brachial artery and coronary artery stenosis in patients with suspected stable coronary artery disease. 2016 , 31, 1467-75	1
794	Serum Trimethylamine-N-Oxide is Elevated in CKD and Correlates with Coronary Atherosclerosis Burden. 2016 , 27, 305-13	241
793	Correlation of NLRP3 with severity and prognosis of coronary atherosclerosis in acute coronary syndrome patients. 2016 , 31, 1218-29	45
792	Paraoxanase as an indicator of myocardial ischemia and its utility in determining extension of ischemia. 2016 , 34, 45-8	3
791	Young women post-MI have higher plasma concentrations of interleukin-6 before and after stress testing. 2016 , 51, 92-98	20
790	Aberrant serum polyunsaturated fatty acids profile is relevant with acute coronary syndrome. 2016 , 31, 1209-17	8
789	Increased Glycated Hemoglobin Level is Associated With SYNTAX Score II in Patients With Type 2 Diabetes Mellitus. 2016 , 67, 384-90	6
788	Association of Platelet to Lymphocyte Ratio With Inflammation and Severity of Coronary Atherosclerosis in Patients With Stable Coronary Artery Disease. 2016 , 67, 89-95	79
787	NQO1 C609T Polymorphism is Associated with Coronary Artery Disease in a Gender-Dependent Manner. 2017 , 17, 35-41	4
786	Renal acoustic radiation force impulse elastography in the evaluation of coronary artery disease. 2017 , 58, 156-163	9
785	Ratios of leptin to insulin and adiponectin to endothelin are sex-dependently associated with extent of coronary atherosclerosis. 2017 , 22, 239-245	10
784	Serum Endothelial Cell-Specific Molecule 1 (Endocan) Levels in Patients With Acute Myocardial Infarction and Its Clinical Significance. 2017 , 68, 354-359	21
783	Effects of antioxidant system on coronary artery lesions in patients with abnormal glucose metabolism. 2017 , 29, 141-146	8

782	Severity of coronary atherosclerosis in patients with COPD. 2017 , 11, 751-756	13
781	The prognostic value of autoantibodies against β -adrenoceptor and cardiac troponin-I for clinical outcomes in STEMI. 2017 , 18, 34-41	6
780	High serum YKL-40 level positively correlates with coronary artery disease. 2017 , 11, 133-139	8
779	SYNTAX Score and Long-Term Outcomes: The BARI-2D Trial. 2017 , 69, 395-403	38
778	Impact of metabolic syndrome on angiographic severity in Egyptian patients with acute coronary syndrome. 2017 , 7, 13-16	
777	Depression and chest pain in patients with coronary artery disease. 2017 , 230, 420-426	26
776	The relation between coronary artery disease severity and fragmented QRS complex in patients with left bundle branch block. 2017 , 69, 119-126	1
775	ApoB is superior to LDL-C or non-HDL-C as a lipid marker for predicting the presence and severity of atherosclerosis in female patients with myocardial infarction. 2017 , 58, 223-225	2
774	Plasma free fatty acids in relation with the severity of coronary artery disease in non-diabetics: A Gensini score assessment. 2017 , 14, 48-52	4
773	Predictive value of non-fasting remnant cholesterol for short-term outcome of diabetics with new-onset stable coronary artery disease. 2017 , 16, 7	10
772	Association between peripheral blood cells mitochondrial DNA content and severity of coronary heart disease. 2017 , 261, 105-110	31
771	Interaction Between Vitamin D and Lipoprotein (a) on the Presence and Extent of Coronary Heart Disease. 2017 , 26, 1079-1084	3
770	Impact of multi-vessel therapy to the risk of periprocedural myocardial injury after elective coronary intervention: exploratory study. 2017 , 17, 69	7
769	Serum CXCL10 and CXCL12 chemokine levels are associated with the severity of coronary artery disease and coronary artery occlusion. 2017 , 233, 23-28	37
768	High levels of serum β -microglobulin predict severity of coronary artery disease. 2017 , 17, 71	14
767	Relationship between inter-arm blood pressure difference and severity of coronary atherosclerosis. 2017 , 263, 171-176	13
766	Association between matrix metalloproteinases 2-1306C/T polymorphism and the risk of coronary artery disease in Iranian population. 2017 , 24, 185-189	2
765	Association between circulating microRNA-208a and severity of coronary heart disease. 2017 , 77, 379-384	30

764	High-density lipoprotein cholesterol levels are associated with coronary severity but not with outcomes in new-onset patients with stable coronary artery disease. 2017 , 263, 104-111	7
763	Significance of reciprocal ST segment depression in ST elevation myocardial infarction. 2017 , 5, 23-27	1
762	Associations between P2RY12 gene polymorphisms and risks of clopidogrel resistance and adverse cardiovascular events after PCI in patients with acute coronary syndrome. 2017 , 96, e6553	7
761	Novel and traditional lipid-related biomarkers and their combinations in predicting coronary severity. 2017 , 7, 360	18
760	Platelet to lymphocyte ratio is associated with the severity of coronary artery disease and clinical outcomes of percutaneous coronary intervention in the Chinese Han population. 2017 , 13, 731-738	20
759	Gender differences in the prevalence of coronary artery tortuosity and its association with coronary artery disease. 2017 , 14, 23-27	15
758	Relationship between two blood stasis syndromes and inflammatory factors in patients with acute coronary syndrome. 2017 , 23, 845-849	8
757	Neutrophil-derived microparticles are released into the coronary circulation following percutaneous coronary intervention in acute coronary syndrome patients. 2017 , 37,	19
756	Telomere Shortening, Regenerative Capacity, and Cardiovascular Outcomes. 2017 , 120, 1130-1138	46
755	Postprandial Blood Glucose Outweighs Fasting Blood Glucose and HbA1c in screening Coronary Heart Disease. 2017 , 7, 14212	16
754	Plasma interleukin-27 levels in patients with coronary artery disease. 2017 , 96, e8260	4
753	Genetic Variation at the Locus is Associated With Reduced Severity of Coronary Artery Disease. 2017 , 6,	17
752	Cardiac Computed Tomography Angiographic Findings as Predictors of Late Heart Failure in an Asymptomatic Diabetic Cohort: An 8-Year Prospective Follow-Up Study. 2017 , 138, 218-227	3
751	Increased plasma levels of lncRNA H19 and LIPCAR are associated with increased risk of coronary artery disease in a Chinese population. 2017 , 7, 7491	107
750	The association of the Syntax score II with carotid intima media thickness and epicardial fat tissue. 2017 , 69, 752-756	5
749	Extent of coronary artery disease in patients undergoing angiography for stable or acute coronary syndromes. 2017 , 58, 115-121	9
748	Polymorphism in ERCC1 confers susceptibility of coronary artery disease and severity of coronary artery atherosclerosis in a Chinese Han population. 2017 , 7, 6407	2
747	The clinical value of inflammatory biomarkers in coronary artery disease: PTX3 as a new inflammatory marker. 2017 , 97, 64-67	13

746	Discrepancy between fractional flow reserve and instantaneous wave-free ratio: Clinical and angiographic characteristics. 2017 , 245, 63-68	36
745	Interleukin-18, matrix metalloproteinase-22 and -29 are independent risk factors of human coronary heart disease. 2017 , 18, 685-695	4
744	Three-Vessel Assessment of Coronary Microvascular Dysfunction in Patients With Clinical Suspicion of Ischemia: Prospective Observational Study With the Index of Microcirculatory Resistance. 2017 , 10,	13
743	Sex-dependent effects of diabetes mellitus on the revascularization rate in mid-term follow up of young patients with coronary artery disease. 2017 , 31, 1686-1690	1
742	The relationship between S-adenosylhomocysteine and coronary artery lesions: A case control study. 2017 , 471, 314-320	11
741	The Relationship between Pulse Wave Velocity and Coronary Artery Stenosis and Percutaneous Coronary Intervention: a retrospective observational study. 2017 , 17, 45	4
740	Macrophage migration inhibitory factor promoter polymorphisms (-794 CATT5-8): Relationship with soluble MIF levels in coronary atherosclerotic disease subjects. 2017 , 17, 144	8
739	Elevated levels of pre- β -high-density lipoprotein are associated with cholesterol ester transfer protein, the presence and severity of coronary artery disease. 2017 , 16, 4	8
738	CHADS-VASc-HSF score - New predictor of severity of coronary artery disease in 2976 patients. 2017 , 228, 1002-1006	14
737	Diurnal glycemic fluctuation is associated with severity of coronary artery disease in prediabetic patients: Possible role of nitrotyrosine and glyceraldehyde-derived advanced glycation end products. 2017 , 69, 625-631	13
736	Coronary Artery Aneurysm Differs From Coronary Artery Ectasia: Angiographic Characteristics and Cardiovascular Risk Factor Analysis in Patients Referred for Coronary Angiography. 2017 , 68, 823-830	31
735	Endocan: Endothelial Dysfunction, Inflammation, or Both?. 2017 , 68, 80	1
734	Relationship between arterial stiffness parameters and the extent and severity of coronary artery disease. 2017 , 59, e134-e141	1
733	Endocan: a novel biomarker associated with well-developed coronary collateral circulation in patients with stable angina and chronic total occlusion. 2017 , 43, 60-67	8
732	The impact of vascular endothelial growth factor +405 C/G polymorphism on long-term outcome and severity of coronary artery disease. 2017 , 31,	8
731	Cystatin (C) as a predictor of coronary artery disease in subclinical atherosclerotic hypertensive patients. 2017 , 7, 42-47	
730	Netrin-1 is associated with macrophage infiltration and polarization in human epicardial adipose tissue in coronary artery disease. 2017 , 69, 851-858	15
729	The Impact of Chronic Repetitive Leg Ischemia on Left Ventricular Function and Severity of Coronary Atherosclerosis in Patients with Acute Coronary Syndrome. 2017 , 26, 447-450	

728	MTHFR C677T gene polymorphism and the severity of coronary lesions in acute coronary syndrome. 2017 , 96, e9044	5
727	Coronary Artery Disease in Premenopausal and Postmenopausal Women. 2017 , 58, 174-179	9
726	Clinical and biochemical factors associated with acute myocardial infarction: Risk factors for acute myocardial infarction. 2017 ,	
725	Risk of Cardiovascular Disease Due to Chronic Hepatitis C Infection: A Review. 2017 , 5, 343-362	44
724	The relationship between apnoea hypopnoea index and Gensini score in patients with acute myocardial infarction undergoing emergency primary percutaneous coronary intervention. 2017 , 9, 2476-2483 ⁵	
723	infection and severity of coronary atherosclerosis in patients with chronic coronary artery disease. 2017 , 13, 933-938	10
722	Association Between Oxidative Stress and Peripheral Leukocyte Telomere Length in Patients with Premature Coronary Artery Disease. 2017 , 23, 4382-4390	18
721	Genome-Wide Association Analysis for Severity of Coronary Artery Disease Using the Gensini Scoring System. 2017 , 4, 57	11
720	Serum Cytokine Profile in Relation to the Severity of Coronary Artery Disease. 2017 , 2017, 4013685	36
719	The Relationship between Body Mass Index and the Severity of Coronary Artery Disease in Patients Referred for Coronary Angiography. 2017 , 2017, 5481671	6
718	Gender-linked impact of epicardial adipose tissue volume in patients who underwent coronary artery bypass graft surgery or non-coronary valve surgery. 2017 , 12, e0177170	7
717	Plasma Omentin-1 Level as a Predictor of Good Coronary Collateral Circulation. 2017 , 24, 940-948	17
716	Association of plasma fatty acid alteration with the severity of coronary artery disease lesions in Tunisian patients. 2017 , 16, 154	9
715	Serum Tartrate-resistant Acid Phosphatase-5b Levels are Associated with the Severity and Extent of Coronary Atherosclerosis in Patients with Coronary Artery Disease. 2017 , 24, 1058-1068	8
714	Association of Serum Prealbumin with Angiographic Severity in Patients with Acute Coronary Syndrome. 2017 , 23, 4041-4049	13
713	Effect of Metabolic Syndrome Score, Metabolic Syndrome, and Its Individual Components on the Prevalence and Severity of Angiographic Coronary Artery Disease. 2017 , 130, 669-677	12
712	Identification of Risk Factors Influencing In-Stent Restenosis with Acute Coronary Syndrome Presentation. 2017 , 53, 203-210	3
711	Hyponatremia Is Associated with Worse Outcomes from Fall Injuries in the Elderly. 2017 , 14,	12

710	gene polymorphism is associated with increased susceptibility to coronary artery disease in Chinese Han population: a case-control study. 2017 , 8, 89055-89063	7
709	The relationship between the bilirubin and the severity of coronary artery disease (CAD) in patients with chronic stable angina. 2018 , 8, 43-50	
708	Correlation of plasma 25-hydroxy-vitamin D level with angiographic severity in coronary artery disease. 2018 , 8, 57-60	1
707	Age at menopause, extent of coronary artery disease and outcome among postmenopausal women with acute coronary syndromes. 2018 , 259, 8-13	11
706	Relationship between Syntax Score and prognostic localization of coronary artery lesions with conventional risk factors, plasma profile markers, and carotid atherosclerosis (CAPP Study 2). 2018 , 257, 306-311	6
705	Myocardial flow reserve derived by dynamic perfusion single-photon emission computed tomography reflects the severity of coronary atherosclerosis. 2018 , 34, 1493-1501	7
704	Protease-Activated Receptor-2 Plays a Critical Role in Vascular Inflammation and Atherosclerosis in Apolipoprotein E-Deficient Mice. 2018 , 138, 1706-1719	41
703	IL-13 may be involved in the development of CAD via different mechanisms under different conditions in a Chinese Han population. 2018 , 8, 6182	6
702	Coronary microvascular dysfunction in patients with stable coronary artery disease: The CE-MARC 2 coronary physiology sub-study. 2018 , 266, 7-14	22
701	Circulating pro-angiogenic micro-ribonucleic acid in patients with coronary heart disease. 2018 , 27, 336-342	9
700	Lipoprotein(a) level associates with coronary artery disease rather than carotid lesions in patients with familial hypercholesterolemia. 2018 , 32, e22442	9
699	Serum albumin level and abnormal corrected QT interval in patients with coronary artery disease and chronic kidney disease. 2018 , 48, 1242-1251	1
698	Uric acid levels are associated with endothelial dysfunction and severity of coronary atherosclerosis during a first episode of acute coronary syndrome. 2018 , 14, 191-199	24
697	Assessment of left and right atrial geometrical changes in patients with stable coronary artery disease: Left and right atrial strain and strain rate imaging study. 2018 , 70, 101-106	3
696	Incremental prognostic value of coronary and systemic atherosclerosis after myocardial infarction. 2018 , 261, 6-11	9
695	Nonculprit Plaque Characteristics in Patients With Acute Coronary Syndrome Caused by Plaque Erosion vs Plaque Rupture: A 3-Vessel Optical Coherence Tomography Study. 2018 , 3, 207-214	41
694	History of vasomotor symptoms, extent of coronary artery disease, and clinical outcomes after acute coronary syndrome in postmenopausal women. 2018 , 25, 635-640	4
693	Alteration in microRNA-155 level correspond to severity of coronary heart disease. 2018 , 78, 219-223	26

692	Evaluation of serum cysteine-rich protein 61 levels in patients with coronary artery disease. 2018 , 12, 329-339	10
691	Risk of discontinuation of clopidogrel after 1 month following bare-metal stents: a propensity-score adjusted comparison with continued administration of clopidogrel after drug-eluting stents. 2018 , 45, 432-439	1
690	Adverse outcome has a U-shaped relation with acute phase change in insulin sensitivity after ST-Elevation Myocardial Infarction. 2018 , 254, 16-22	1
689	Effect modification of hypertension on the association of vitamin D deficiency with severity of coronary stenosis. 2018 , 27, 134-140	1
688	Prognostic significance of arterial stiffness and osteoprotegerin in patients with stable coronary artery disease. 2018 , 48, e12890	17
687	Predictive value of big endothelin-1 on outcomes in patients with myocardial infarction younger than 35 years old. 2018 , 15, 25-33	2
686	Transcoronary gradients of HDL-associated MicroRNAs in unstable coronary artery disease. 2018 , 253, 138-144	14
685	Reduced Forced Expiratory Volume in 1 Second Percentage Predicted Is Associated With Diffuse Coronary Atherosclerosis in Hospitalized Patients Undergoing Coronary Angiography. 2018 , 355, 307-313	1
684	Noncoding RNAs in Cardiovascular Disease: Pathological Relevance and Emerging Role as Biomarkers and Therapeutics. 2018 , 31, 150-165	58
683	Epicardial adipose tissue thickness, carotid intima media thickness and total cholesterol/HDL ratio: a combined cut off for detecting coronary artery disease. 2018 , 8, 73-79	1
682	Elevated lipoprotein(a) and low-density lipoprotein cholesterol as predictors of the severity and complexity of angiographic lesions in patients with premature coronary artery disease. 2018 , 12, 1019-1026	19
681	Local Thickness of Epicardial Adipose Tissue Surrounding the Left Anterior Descending Artery Is a Simple Predictor of Coronary Artery Disease - New Prediction Model in Combination With Framingham Risk Score. 2018 , 82, 1369-1378	12
680	Soluble glycoprotein 130 is inversely related to severity of coronary atherosclerosis. 2018 , 23, 527-532	10
679	Influence of sleep-disordered breathing assessed by pulse oximetry on long-term clinical outcomes in patients who underwent percutaneous coronary intervention. 2018 , 107, 711-718	4
678	Higher levels of circulating naïve CD8CD45RA cells are associated with lower extent of coronary atherosclerosis and vascular dysfunction. 2018 , 259, 26-30	6
677	Delayed flow-mediated vasodilation and critical coronary stenosis. 2018 , 66, 1-7	14
676	Relationship between high-density lipoprotein cholesterol and the red cell distribution width in patients with coronary artery disease. 2018 , 17, 53	9
675	Implication of coronary CT angiography combined with four-dimensional speckle tracking echocardiography for predicting major adverse cardiac events. 2018 , 34, 1287-1293	2

674	The significance of post-stress decrease in left ventricular ejection fraction in patients undergoing regadenoson stress gated SPECT myocardial perfusion imaging. 2018 , 25, 1313-1323	14
673	Mental Stress, Exercise, and Other Determinants of Elevation in High-Sensitivity Troponin Levels: A Call for Standardization of Laboratory Protocols. 2018 , 11, 612-615	
672	The Comparison of Angiographic Scoring Systems With the Predictors of Atherosclerosis. 2018 , 69, 158-163	7
671	Association of bone metabolism markers with coronary atherosclerosis and coronary artery disease in postmenopausal women. 2018 , 36, 352-363	5
670	Western Dietary Pattern Is Associated With Severe Coronary Artery Disease. 2018 , 69, 339-346	18
669	A spectrum of retinal vasculature measures and coronary artery disease. 2018 , 268, 215-224	29
668	Association between coronary artery disease severity and overactive bladder in geriatric patients. 2018 , 36, 35-40	4
667	The association between red cell distribution width, erythropoietin levels, and coronary artery disease. 2018 , 29, 74-80	4
666	Application of pressure-derived myocardial fractional flow reserve in chronic hemodialysis patients. 2018 , 71, 52-58	8
665	Pentraxin-3 and coronary artery disease. 2018 , 102, 1-2	2
664	Ethnic Differences in the Risk Factors and Severity of Coronary Artery Disease: a Patient-Based Study in Iran. 2018 , 5, 623-631	6
663	Pericardial fat volume and coronary calcifications for prediction of coronary artery disease extent in patients with type 2 diabetes mellitus. 2018 , 49, 971-975	
662	Correlations of DAPT score and PRECISE-DAPT score with the extent of coronary stenosis in acute coronary syndrome. 2018 , 97, e12531	12
661	Association between blood neutrophil-to-lymphocyte ratio and severity of coronary artery disease: Evidence from 17 observational studies involving 7017 cases. 2018 , 97, e12432	17
660	Automatic Severity Classification of Coronary Artery Disease via Recurrent Capsule Network. 2018 ,	4
659	Association of lymphocyte to monocyte ratio with severity of coronary artery disease. 2018 , 97, e12813	30
658	Inflammation-based Glasgow Prognostic Score in patients with acute ST-segment elevation myocardial infarction: A prospective cohort study. 2018 , 97, e13615	14
657	Baseline levels of serum high sensitivity C reactive protein and lipids in predicting the residual risk of cardiovascular events in Chinese population with stable coronary artery disease: a prospective cohort study. 2018 , 17, 273	5

656	Reduced Plasma Kallistatin Is Associated With the Severity of Coronary Artery Disease, and Kallistatin Treatment Attenuates Atherosclerotic Plaque Formation in Mice. 2018 , 7, e009562	10
655	Young Women With Coronary Artery Disease Exhibit Higher Concentrations of Interleukin-6 at Baseline and in Response to Mental Stress. 2018 , 7, e010329	18
654	Predictive value of combining the level of lipoprotein-associated phospholipase A2 and antithrombin III for acute coronary syndrome risk. 2018 , 9, 517-522	5
653	Systemic microvascular dysfunction in microvascular and vasospastic angina. 2018 , 39, 4086-4097	83
652	GRACE and TIMI risk scores in predicting the angiographic severity of non-ST elevation acute coronary syndrome. 2018 , 70 Suppl 3, S250-S253	9
651	Association of TAFI gene polymorphisms with severity of coronary stenosis in stable coronary artery disease. 2018 , 171, 171-176	4
650	The relationship of Charlson comorbidity index with stent restenosis and extent of coronary artery disease. 2018 , 10, 70-75	1
649	Associations between circulating IgG antibodies to Apolipoprotein B-derived peptide antigens and acute coronary syndrome in a Chinese Han population. 2018 , 38,	2
648	Free triiodothyronine and global registry of acute coronary events risk score on predicting long-term major adverse cardiac events in STEMI patients undergoing primary PCI. 2018 , 17, 234	6
647	The atheroprotective role of lipoxin A prevents oxLDL-induced apoptotic signaling in macrophages via JNK pathway. 2018 , 278, 259-268	16
646	The role of retinol-binding protein 4 and its relationship with sex hormones in coronary artery disease. 2018 , 506, 204-210	9
645	Relationship between the endothelial glycocalyx and the extent of coronary atherosclerosis. 2018 , 25, e12504	7
644	Triglyceride glucose and haemoglobin glycation index for predicting outcomes in diabetes patients with new-onset, stable coronary artery disease: a nested case-control study. 2018 , 50, 576-586	32
643	Associations between retinal arteriolar and venular calibre with the prevalence of impaired fasting glucose and diabetes mellitus: A cross-sectional study. 2018 , 13, e0189627	5
642	Impact of right coronary dominance on triple-vessel coronary artery disease: A cross-sectional study. 2018 , 97, e11685	2
641	Elevated plasma lipoprotein(a) levels were associated with increased risk of cardiovascular events in Chinese patients with stable coronary artery disease. 2018 , 8, 7726	21
640	The relationship of plasma miR-503 and coronary collateral circulation in patients with coronary artery disease. 2018 , 207, 145-151	15
639	Epicardial fat thickness can predict severity and multivessel distribution in Egyptian patients with atherosclerotic coronary artery stenosis. 2018 , 70, 323-327	4

638	Lipoprotein(a) and apolipoprotein(a) isoform size: Associations with angiographic extent and severity of coronary artery disease, and carotid artery plaque. 2018 , 275, 232-238	13
637	Association between cardiac autonomic nervous dysfunction and the severity of coronary lesions in patients with stable coronary artery disease. 2018 , 46, 3729-3740	6
636	Plasma Chemerin Levels Are Increased in ST Elevation Myocardial Infarction Patients with High Thrombus Burden. 2018 , 2018, 5812704	4
635	The relationship between fragmented QRS complexes and syntax II scores in patients with ST-segment elevation myocardial infarction. 2018 , 51, 825-829	3
634	Roles of Perivascular Adipose Tissue in the Pathogenesis of Atherosclerosis. 2018 , 9, 3	38
633	Change of Inflammatory Factors in Patients with Acute Coronary Syndrome. 2018 , 131, 1444-1449	15
632	Increased Serum Interleukin-34 Levels Are Related to the Presence and Severity of Cardiac Dysfunction in Patients With Ischemic Cardiomyopathy. 2018 , 9, 904	7
631	Associations between circulating full-length angiopoietin-like protein 8 levels and severity of coronary artery disease in Chinese non-diabetic patients: a case-control study. 2018 , 17, 92	13
630	A study of the association of rs12040273 with susceptibility and severity of coronary artery disease in a Chinese Han population. 2018 , 18, 10	3
629	Clinical analysis of lectin-like oxidized low-density lipoprotein receptor-1 in patients with in-stent restenosis after percutaneous coronary intervention. 2018 , 97, e0366	8
628	Atherogenic index of plasma (AIP): a novel predictive indicator for the coronary artery disease in postmenopausal women. 2018 , 17, 197	87
627	Association between cognitive function and parameters of echocardiography and coronary artery angiography. 2018 , 76, 225-230	2
626	The different relations of PCSK9 and Lp(a) to the presence and severity of atherosclerotic lesions in patients with familial hypercholesterolemia. 2018 , 277, 7-14	15
625	Cardiac Phase Space Tomography: A novel method of assessing coronary artery disease utilizing machine learning. 2018 , 13, e0198603	9
624	Effect of the Epicardial Adipose Tissue Volume on the Prevalence of Paroxysmal and Persistent Atrial Fibrillation. 2018 , 82, 1778-1787	16
623	Association between elevated plasma microRNA-223 content and severity of coronary heart disease. 2018 , 78, 373-378	16
622	Erectile dysfunction severity as a predictor of left main and/or three-vessel disease in acute coronary syndrome patients. 2018 , 70 Suppl 3, S56-S59	5
621	Familial hypercholesterolemia in very young myocardial infarction. 2018 , 8, 8861	9

620	Association Between the Presence or Severity of Coronary Artery Disease and Pericardial Fat, Paracardial Fat, Epicardial Fat, Visceral Fat, and Subcutaneous Fat as Assessed by Multi-Detector Row Computed Tomography. 2018 , 59, 695-704	18
619	Tenascin C: A Potential Biomarker for Predicting the Severity of Coronary Atherosclerosis. 2019 , 26, 31-38	10
618	Association of serum uric acid levels with SYNTAX score II and long term mortality in the patients with stable angina pectoris who undergo percutaneous coronary interventions due to multivessel and/or unprotected left main disease. 2019 , 35, 1-7	6
617	Plasma Lipoprotein(a) Concentration Is Associated With the Coronary Severity but Not With Events in Stable Coronary Artery Disease Patients: A Chinese Cohort Study. 2019 , 28, 1009-1017	10
616	New non-invasive indexes of arterial stiffness are significantly correlated with severity and complexity of coronary atherosclerosis. 2018 , 1-7	1
615	Early Versus Late New-Onset Atrial Fibrillation in Acute Myocardial Infarction: Differences in Clinical Characteristics and Predictors. 2019 , 70, 921-928	12
614	Left ventricular dyssynchrony in coronary artery disease patients without regional wall-motion abnormality: Correlation with Gensini score. 2019 , 36, 1689-1697	2
613	Expression of B2 Receptor on Circulating CD34-Positive Cells and Outcomes of Myocardial Infarction. 2019 , 2019, 7816438	0
612	Circulating ESM-1 levels are correlated with the presence of coronary artery disease in patients with obstructive sleep apnea. 2019 , 20, 188	4
611	Association between the No-Reflow Phenomenon and Soluble CD40 Ligand Level in Patients with Acute ST-Segment Elevation Myocardial Infarction. 2019 , 55,	2
610	Lipoprotein(a) and coronary artery disease in Chinese postmenopausal female patients: a large cross-sectional cohort study. 2019 , 95, 534-540	4
609	Applying Pulse Spectrum Analysis to Facilitate the Diagnosis of Coronary Artery Disease. 2019 , 2019, 2709486	6
608	Clinical characteristics of patients with premature acute coronary syndrome and adverse cardiovascular events after PCI. 2019 , 18, 793-801	5
607	Serum sLYVE-1 is not associated with coronary disease but with renal dysfunction: a retrospective study. 2019 , 9, 10816	2
606	Plasma fatty acid-binding protein 4 (FABP4) level is associated with abnormal QTc interval in patients with stable angina and chronic kidney disease. 2019 , 19, 153	2
605	Impact of free fatty acids on prognosis in coronary artery disease patients under different glucose metabolism status. 2019 , 18, 134	11
604	Association between Nonalcoholic Fatty Liver and Gensini Score in Patients with Coronary Heart Disease: A Cross-Sectional Study. 2019 , 144, 90-96	1
603	DNA methylation of antisense noncoding RNA in the INK locus (ANRIL) is associated with coronary artery disease in a Chinese population. 2019 , 9, 15340	4

602	Tumour necrosis factor like cytokine 1A levels and lesion complexity in non-smoking patients with coronary artery disease. 2019 , 24, 764-770	0
601	Correlation between coronary stenosis and Toll-like receptors 2 and 4 levels in Chinese Zhuang patients with coronary heart disease. 2019 , 18, 2346-2352	2
600	The Relationship Between Mean Platelet Volume and In-Hospital Mortality in Geriatric Patients with ST Segment Elevation Myocardial Infarction Who Underwent Primary Percutaneous Coronary Intervention. 2019 , 4, 135-141	3
599	Clinical Significance of Increased Cardiac Troponin T in Patients with Chronic Hemodialysis and Cardiovascular Disease: Comparison to B-Type Natriuretic Peptide and A-Type Natriuretic Peptide Increase. 2019 , 44, 1050-1062	6
598	Very low-density lipoprotein cholesterol is associated with extent and severity of coronary artery disease in patients with type 2 diabetes mellitus. 2019 , 7, 2050312119871786	0
597	Role of adipokine zinc-lycoprotein in coronary heart disease. 2019 , 317, E1055-E1062	3
596	Associations of SNPs of the Gene with Serum Adiponectin Levels, Unstable Angina, and Coronary Artery Disease. 2019 , 9,	4
595	Biobanking for discovery of novel cardiovascular biomarkers using imaging-quantified disease burden: protocol for the longitudinal, prospective, BioHEART-CT cohort study. 2019 , 9, e028649	20
594	Low expression of PRMT5 in peripheral blood may serve as a potential independent risk factor in assessments of the risk of stable CAD and AMI. 2019 , 19, 31	8
593	sSema4D levels are increased in coronary heart disease and associated with the extent of coronary artery stenosis. 2019 , 219, 329-335	9
592	Association of common polymorphisms in IL-6 and IL6ST genes with levels of inflammatory markers and coronary stenosis. 2019 , 21, 100593	3
591	High-sensitivity C-reactive protein and hypertension: combined effects on coronary severity and cardiovascular outcomes. 2019 , 42, 1783-1793	5
590	Cell cycle arrest in peripheral blood mononuclear cells: A non-invasive method for diagnosis of coronary artery disease. 2019 , 84, 153-160	3
589	Association of circulating BMP9 with coronary heart disease and hypertension in Chinese populations. 2019 , 19, 131	4
588	Eosinophils count in peripheral circulation is associated with coronary artery disease. 2019 , 286, 128-134	7
587	Association of circulating growth differentiation factor-15, Kröpel-like factor 4 and growth arrest-specific 6 with coronary artery disease. 2019 , 495, 630-636	3
586	Circulating microRNA-378 levels serve as a novel biomarker for assessing the severity of coronary stenosis in patients with coronary artery disease. 2019 , 39,	5
585	Comparison of autonomic stress reactivity in young healthy versus aging subjects with heart disease. 2019 , 14, e0216278	7

584	Circulating MIF Levels Predict Clinical Outcomes in Patients With ST-Elevation Myocardial Infarction After Percutaneous Coronary Intervention. 2019 , 35, 1366-1376	9
583	Cutaneous microvascular function in patients with obstructive or non-obstructive coronary artery disease evaluated by laser speckle contrast imaging. 2019 , 125, 103880	1
582	Lipoprotein(a) and Cardiovascular Outcomes in Patients With Coronary Artery Disease and Prediabetes or Diabetes. 2019 , 42, 1312-1318	50
581	A guide for Gensini Score calculation. 2019 , 287, 181-183	51
580	Non-HDL-c/TC: A Novel Lipid-Related Marker in the Assessment of Severity of Coronary Artery Lesions and Cardiovascular Outcomes. 2019 , 2019, 5931975	5
579	Using path analysis to investigate the relationships between depression, anxiety, and health-related quality of life among patients with coronary artery disease. 2019 , 28, 2695-2704	8
578	Correlation between Gensini Score and Duration of Diabetes in Patients Undergoing Coronary Angiography. 2019 , 11, e4010	2
577	High-sensitive cardiac Troponin T and exercise stress test for evaluation of angiographically significant coronary disease. 2019 , 287, 1-6	4
576	Association of Aortic Valve Sclerosis and Clinical Factors in Patients With Acute Myocardial Infarction. 2019 , 357, 474-482	3
575	Gender-specific associations between atherogenic index of plasma and the presence and severity of acute coronary syndrome in very young adults: a hospital-based observational study. 2019 , 18, 99	26
574	Alterations in the gut microbiome and metabolism with coronary artery disease severity. 2019 , 7, 68	93
573	Renal haemodynamics and coronary atherosclerotic burden are associated in patients with hypertension and mild coronary artery disease. 2019 , 17, 3255-3263	4
572	Impaired mitochondrial oxidative phosphorylation capacity in epicardial adipose tissue is associated with decreased concentration of adiponectin and severity of coronary atherosclerosis. 2019 , 9, 3535	9
571	LncRNA MALAT1 gene polymorphisms in coronary artery disease: a case-control study in a Chinese population. 2019 , 39,	13
570	Potential associations of circulating growth differentiation factor-15 with sex hormones in male patients with coronary artery disease. 2019 , 114, 108792	8
569	A multi-stage association study of plasma cytokines identifies osteopontin as a biomarker for acute coronary syndrome risk and severity. 2019 , 9, 5121	0
568	Angiographic features of patients with coronary plaque erosion. 2019 , 288, 12-16	14
567	DNA methylation and hydroxymethylation are associated with the degree of coronary atherosclerosis in elderly patients with coronary heart disease. 2019 , 224, 241-248	22

566	Effect of CD14 polymorphisms on the risk of cardiovascular disease: evidence from a meta-analysis. 2019 , 18, 74	6
565	The impact of preoperative fibrinogen-albumin ratio on mortality in patients with acute ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2019 , 493, 8-13	25
564	Secretory vimentin is associated with coronary artery disease in patients and induces atherogenesis in ApoE mice. 2019 , 283, 9-16	11
563	Biomarkers enhance the long-term predictive ability of the KAMIR risk score in Chinese patients with ST-elevation myocardial infarction. 2019 , 132, 30-41	6
562	Correlation between the serum procalcitonin level and the extension and severity of coronary artery disease in patients with non-ST-segment elevation myocardial infarction. 2019 , 8, 62-66	1
561	Obstructive sleep apnoea in acute coronary syndrome. 2019 , 28,	10
560	ADAMTS-5 Decreases in Coronary Arteries and Plasma from Patients with Coronary Artery Disease. 2019 , 2019, 6129748	4
559	Triglyceride to high-density lipoprotein cholesterol ratio as a risk factor of repeat revascularization among patients with acute coronary syndrome after first-time percutaneous coronary intervention. 2019 , 11, 5087-5095	2
558	Relationship between Random Blood Glucose, Fasting Blood Glucose, and Gensini Score in Patients with Acute Myocardial Infarction. 2019 , 2019, 9707513	7
557	Interleukin-34 Levels Were Associated with Prognosis in Patients with Acute Myocardial Infarction. 2019 , 60, 1259-1267	6
556	Plasma levels of receptor interacting protein kinase-3 correlated with coronary artery disease. 2019 , 132, 1400-1405	9
555	Comparison of Apolipoprotein B/A1 ratio, Framingham risk score and TC/HDL-c for predicting clinical outcomes in patients undergoing percutaneous coronary intervention. 2019 , 18, 202	12
554	Association of lipoprotein(a) and coronary artery disease in 1003 patients with stage 3-5 chronic kidney disease undergoing coronary angiography. 2019 , 30, 137-142	1
553	Impact of Non-Alcoholic Fatty Liver Disease on Cardiovascular Outcomes in Patients With Stable Coronary Artery Disease: A Matched Case-Control Study. 2019 , 10, e00011	12
552	Sex-Specific Association Between Coronary Artery Disease Severity and Myocardial Ischemia Induced by Mental Stress. 2019 , 81, 57-66	9
551	Relationship between serum homocysteine levels and long-term outcomes in patients with ST-segment elevation myocardial infarction. 2019 , 132, 1028-1036	3
550	Low expression of PIK3C2A gene: A potential biomarker to predict the risk of acute myocardial infarction. 2019 , 98, e15061	7
549	Monocyte/lymphocyte ratio is related to the severity of coronary artery disease and clinical outcome in patients with non-ST-elevation myocardial infarction. 2019 , 98, e16267	31

548	Effect of coronary collateral circulation on the prognosis of elderly patients with acute ST-segment elevation myocardial infarction treated with underwent primary percutaneous coronary intervention. 2019 , 98, e16502	3
547	Posttraumatic stress disorder is associated with enhanced interleukin-6 response to mental stress in subjects with a recent myocardial infarction. 2019 , 75, 26-33	31
546	Severity of obstructive sleep apnea and extension of coronary artery disease. 2019 , 23, 747-752	8
545	The role of CD27-CD70 signaling in myocardial infarction and cardiac remodeling. 2019 , 278, 210-216	1
544	Serum Metrnl is associated with the presence and severity of coronary artery disease. 2019 , 23, 271-280	22
543	Usefulness of platelet to lymphocyte ratio for predicting recurrence of atrial fibrillation after direct current cardioversion. 2019 , 24, e12616	4
542	3D-speckle tracking echocardiography for assessment of coronary artery disease severity in stable angina pectoris. 2019 , 36, 320-327	8
541	Overexpression of GAS5 inhibits abnormal activation of Wnt/ β -catenin signaling pathway in myocardial tissues of rats with coronary artery disease. 2019 , 234, 11348-11359	19
540	Predictive value of fasting blood glucose for serious coronary atherosclerosis in non-diabetic patients. 2019 , 47, 152-158	3
539	Clinical and genetic characteristics of familial hypercholesterolemia patients with type 2 diabetes. 2019 , 42, 591-598	6
538	Beta 2-Microglobulin and the Severity of Coronary Stenosis in Patients With Acute Coronary Syndrome. 2019 , 28, 575-582	5
537	Association of Prediabetes With Higher Coronary Atherosclerotic Burden Among Patients With First Diagnosed Acute Coronary Syndrome. 2019 , 70, 174-180	23
536	Predictive Value of the Acute-to-Chronic Glycemic Ratio for In-Hospital Outcomes in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Percutaneous Coronary Intervention. 2020 , 71, 38-47	11
535	Associations between specific plasma ceramides and severity of coronary-artery stenosis assessed by coronary angiography. 2020 , 46, 150-157	16
534	Association of Genetic Variations in NRF2, NQO1, HMOX1, and MT with Severity of Coronary Artery Disease and Related Risk Factors. 2020 , 20, 176-189	10
533	Comparison of MESA of and Framingham risk scores in the prediction of coronary artery disease severity. 2020 , 45, 139-144	0
532	Clinical value of ARG1 in acute myocardial infarction patients: Bioinformatics-based approach. 2020 , 121, 109590	8
531	Lipoprotein(a) as a marker for predicting coronary collateral circulation in patients with acute myocardial infarction. 2020 , 17, 67-78	2

530	Risk Factors and Outcomes of Very Young Adults Who Experience Myocardial Infarction: The Partners YOUNG-MI Registry. 2020 , 133, 605-612.e1		27
529	Upregulation of microRNA-218 reduces cardiac microvascular endothelial cells injury induced by coronary artery disease through the inhibition of HMGB1. 2020 , 235, 3079-3095		11
528	Lipoprotein(a) Is Associated with the Presence and Severity of New-Onset Coronary Artery Disease in Postmenopausal Women. 2020 , 29, 503-510		1
527	The independent and incremental value of ultrasound carotid plaque length to predict the presence and severity of coronary artery disease: analysis from the carotid plaque length prospective registry. 2020 , 21, 389-396		4
526	Long-term Prognostic Impact of Physical Activity in Patients With Stable Coronary Heart Disease. <i>American Journal of Cardiology</i> , 2020 , 125, 176-181	3	1
525	Lipoprotein(a) levels are associated with coronary severity but not with outcomes in Chinese patients underwent percutaneous coronary intervention. 2020 , 30, 265-273		8
524	Predictive value of the combination of age, creatinine, and ejection fraction score and diabetes in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention. 2020 , 31, 109-117		8
523	Clinical and procedural characteristics of persons living with HIV presenting with acute coronary syndrome. 2020 , 34, 81-90		3
522	Fasting serum total bile acid level is associated with coronary artery disease, myocardial infarction and severity of coronary lesions. 2020 , 292, 193-200		19
521	Fasting Blood Glucose and HbA Correlate With Severity of Coronary Artery Disease in Elective PCI Patients With HbA 5.7% to 6.4. 2020 , 71, 167-174		8
520	High cardiovascular risk of patients with type 2 diabetes is only partially attributed to angiographic burden of atherosclerosis. 2020 , 17, 1479164120953612		4
519	Association of high-sensitivity troponin T and I with the severity of stable coronary artery disease in patients with chronic kidney disease. 2020 , 313, 81-87		4
518	Long-term prognostic utility of low-density lipoprotein (LDL) triglyceride in real-world patients with coronary artery disease and diabetes or prediabetes. 2020 , 19, 152		5
517	Naive CD8 T Cells Expressing CD95 Increase Human Cardiovascular Disease Severity. 2020 , 40, 2845-2859		5
516	Sex differences in the inflammatory response to stress and risk of adverse cardiovascular outcomes among patients with coronary heart disease. 2020 , 90, 294-302		13
515	Association of plasma free fatty acids levels with the presence and severity of coronary and carotid atherosclerotic plaque in patients with type 2 diabetes mellitus. 2020 , 20, 156		8
514	Prognostic Value of Braden Scale in Patients With Acute Myocardial Infarction: From the Retrospective Multicenter Study for Early Evaluation of Acute Chest Pain. 2020 , 35, E53-E61		3
513	Study of correlation between epicardial fat thickness and severity of coronary artery disease. 2020 , 72, 445-447		1

512	Management of multivessel coronary artery disease in patients with non-ST-elevation myocardial infarction: a complex path to precision medicine. 2020 , 11, 2040622320938527	8
511	CircRNA ZNF609 in peripheral blood leukocytes acts as a protective factor and a potential biomarker for coronary artery disease. 2020 , 8, 741	15
510	Associations between the psoas major muscle index and the presence and severity of coronary artery disease. 2020 , 99, e21086	1
509	Impact of Smoking Status on Growth Differentiation Factor 15 and Mortality in Patients With Suspected or Known Coronary Artery Disease: The ANOX Study. 2020 , 9, e018217	2
508	Predictive efficacy of neutrophil-to-lymphocyte ratio for long-term prognosis in new onset acute coronary syndrome: a retrospective cohort study. 2020 , 20, 500	4
507	On-pump versus off-pump coronary artery bypass surgery for multi-vessel coronary revascularization. 2020 , 12, 5639-5646	5
506	Associations of myeloperoxidase, interleukin-17A and heparin-binding EGF-like growth factor levels with in-stent restenosis after percutaneous coronary intervention: a single-centre case-control study in China. 2020 , 10, e039405	6
505	Circulating Levels of CILP2 Are Elevated in Coronary Heart Disease and Associated with Atherosclerosis. 2020 , 2020, 1871984	1
504	Serum exosomal miR-122-5p is a new biomarker for both acute coronary syndrome and underlying coronary artery stenosis. 2020 , 25, 539-547	5
503	Overexpression of MicroRNA-16 Alleviates Atherosclerosis by Inhibition of Inflammatory Pathways. 2020 , 2020, 8504238	11
502	The effects of percutaneous coronary intervention on mortality in elderly patients with non-ST-segment elevation myocardial infarction undergoing coronary angiography. 2020 , 65, 81-88	1
501	Impact of increased serum 8-hydroxy-2Ndeoxyguanosine levels on extent of coronary artery lesions in elderly patients with type 2 diabetes. 2020 , 48, 300060520934653	1
500	Activation of NLRP3 inflammasome assembly is associated with smoking status of patients with coronary artery disease. 2020 , 87, 106820	4
499	The relationship of body composition indices with the significance, extension and severity of coronary artery disease. 2020 , 30, 2279-2285	1
498	Cohort Profile: The Leipzig Research Center for Civilization Diseases-Heart Study (LIFE-Heart). 2020 , 49, 1439-1440h	8
497	Interaction between a haptoglobin genetic variant and coronary artery disease (CAD) risk factors on CAD severity in Singaporean Chinese population. 2020 , 8, e1450	1
496	IL-35 polymorphisms and cognitive decline did not show any association in patients with coronary heart disease over a 2-year period: A retrospective observational study (STROBE compliant). 2020 , 99, e21390	1
495	Barthel Index as a Predictor of Mortality in Patients with Acute Coronary Syndrome: Better Activities of Daily Living, Better Prognosis. 2020 , 15, 1951-1961	6

494	Short report: depression and anxiety symptoms as predictors of adverse cardiovascular events in Chinese patients after percutaneous coronary intervention. 2021 , 26, 1126-1133	3
493	Association between Obstructive Sleep Apnea and SYNTAX Score. 2020 , 9,	1
492	Predictive value of cardiopulmonary fitness parameters in the prognosis of patients with acute coronary syndrome after percutaneous coronary intervention. 2020 , 48, 300060520949081	0
491	Relation of Fibrinogen-to-Albumin Ratio to Severity of Coronary Artery Disease and Long-Term Prognosis in Patients with Non-ST Elevation Acute Coronary Syndrome. 2020 , 2020, 1860268	9
490	Correlation between the Serum Platelet-Derived Growth Factor, Angiopoietin-1, and Severity of Coronary Heart Disease. 2020 , 2020, 3602608	3
489	Association of Genes Related to Oxidative Stress with the Extent of Coronary Atherosclerosis. 2020 , 10,	1
488	Free Thiol β -GPI (β -Glycoprotein-I) Provides a Link Between Inflammation and Oxidative Stress in Atherosclerotic Coronary Artery Disease. 2020 , 40, 2794-2804	2
487	The fasting plasma glucose is correlated to the prevalence and severity of coronary artery disease in population without diabetes history in Xinjiang, China. 2020 , 13, 683-690	
486	Relative Fat Mass Index can be solution for obesity paradox in coronary artery disease severity prediction calculated by SYNTAX Score. 2021 , 97, 434-441	0
485	Association between plasma leptin/adiponectin ratios with the extent and severity of coronary artery disease. 2020 , 20, 474	5
484	Incomplete protective effect of coronary collateral circulation for acute myocardial infarction patients. 2020 , 99, e22750	0
483	Methylation of CpG sites in C1QTNF1 (C1q and tumor necrosis factor related protein 1) differs by gender in acute coronary syndrome in Han population: a case-control study. 2020 , 42, 681-689	2
482	Clinical significance of serum concentrations of neuregulin-4, in acute coronary syndrome. 2020 , 10, 5797	5
481	Associations between the IL-6-neutralizing sIL-6R-sgp130 buffer system and coronary artery disease in postmenopausal women. 2020 , 8, 379	1
480	Relationship between Long Noncoding RNA H19 Polymorphisms and Risk of Coronary Artery Disease in a Chinese Population: A Case-Control Study. 2020 , 2020, 9839612	5
479	Genome-wide analysis of carotid plaque burden suggests a role of IL5 in men. 2020 , 15, e0233728	4
478	Whole blood viscosity in microvascular angina and coronary artery disease: Significance and utility. 2020 , 39, 17-23	
477	Circulating metabolites from the choline pathway and acute coronary syndromes in a Chinese case-control study. 2020 , 17, 39	0

476	Serum Exosomal MicroRNA-21, MicroRNA-126, and PTEN Are Novel Biomarkers for Diagnosis of Acute Coronary Syndrome. 2020 , 11, 654	26
475	Predicting Chronic Myocardial Ischemia Using CCTA-Based Radiomics Machine Learning Nomogram. 2020 , 1	4
474	Whole blood viscosity in microvascular angina and coronary artery disease: Significance and utility. 2020 , 39, 17-23	7
473	Elevated Plasma Adiponectin Levels Are Associated with Abnormal Corrected QT Interval in Patients with Stable Angina. 2020 , 61, 29-38	3
472	Association of Risk Factors with Coronary Angiographic Findings of Female Patients with Acute Myocardial Infarction at a Tertiary Hospital of Bangladesh. 2020 , 34, 100-110	
471	Association between the Non-high-Density Lipoprotein Cholesterol to High-Density Lipoprotein Cholesterol Ratio and the Risk of Coronary Artery Disease. 2020 , 2020, 7146028	3
470	Abnormal P-wave axis and myocardial ischemia development during mental stress. 2020 , 60, 3-7	2
469	Residual Gensini Score Is Associated With Long-Term Cardiac Mortality in Patients With Heart Failure After Percutaneous Coronary Intervention. 2020 , 2, 89-94	4
468	Predictive value of leukocyte telomere length for the severity of coronary artery disease. 2020 , 17, 175-183	0
467	Plasma P-selectin level is associated with severity of coronary heart disease in Chinese Han population. 2020 , 48, 300060519896437	1
466	Nitrate Esters Alleviated Coronary Atherosclerosis Through Inhibition of NF- κ B-Regulated Macrophage Polarization Shift in Epicardial Adipose Tissue. 2020 , 75, 475-482	4
465	Predictive values of heart rate variability, deceleration and acceleration capacity of heart rate in post-infarction patients with LVEF \geq 5. 2020 , 25, e12771	4
464	Serum level of HDL particles are independently associated with long-term prognosis in patients with coronary artery disease: The GENES study. 2020 , 10, 8138	9
463	Association of the methylene-tetrahydrofolate reductase gene rs1801133 C677T variant with serum homocysteine levels, and the severity of coronary artery disease. 2020 , 10, 10064	4
462	Association of platelet to lymphocyte ratio with in-hospital major adverse cardiovascular events and the severity of coronary artery disease assessed by the Gensini score in patients with acute myocardial infarction. 2020 , 133, 415-423	8
461	Lower Gensini Score in Navy Personnel Compared to Civilians, rather than for Non-Commissioned Officers. 2020 , 441, 012177	
460	Analysis on the polymorphisms of site RS4977574, and RS1333045 in region 9p21 and the susceptibility of coronary heart disease in Chinese population. 2020 , 21, 36	3
459	Extracellular ubiquitin levels are increased in coronary heart disease and associated with the severity of the disease. 2020 , 80, 256-264	4

458	Up-regulation of long non-coding RNA THRIL in coronary heart disease: Prediction for disease risk, correlation with inflammation, coronary artery stenosis, and major adverse cardiovascular events. 2020 , 34, e23196	13
457	Association between impaired cutaneous microvascular endothelial function and lectin-like oxidized low-density lipoprotein receptor-1 in patients with coronary slow flow. 2020 , 129, 103984	1
456	Elevated 1-h postload plasma glucose levels identify coronary heart disease patients with greater severity of coronary artery lesions and higher risk of 1-year re-admission. 2020 , 17, 1479164119896978	2
455	Distinct Characteristics of VEGF-D and VEGF-C to Predict Mortality in Patients With Suspected or Known Coronary Artery Disease. 2020 , 9, e015761	12
454	Serum VEGF: Diagnostic Value of Acute Coronary Syndrome from Stable Angina Pectoris and Prognostic Value of Coronary Artery Disease. 2020 , 2020, 6786302	5
453	Clinical utility of serum cystatin C for prediction of multi-vessel disease by coronary angiography in type 2 diabetes mellitus patients with normal renal function. 2020 , 20, 183	1
452	Association of Posttraumatic Stress Disorder With Mental Stress-Induced Myocardial Ischemia in Adults After Myocardial Infarction. 2020 , 3, e202734	12
451	Contradictions between DAPT and PRECISE-DAPT scores with the severity of coronary lesion in acute coronary syndrome. 2020 , 99, e19699	1
450	Association of small dense low-density lipoprotein with cardiovascular outcome in patients with coronary artery disease and diabetes: a prospective, observational cohort study. 2020 , 19, 45	15
449	Drug-eluting stents versus bare-metal stents for acute coronary syndrome. 2017 , 8, CD012481	10
448	Multiple Non-coding ANRIL Transcripts Are Associated with Risk of Coronary Artery Disease: a Promising Circulating Biomarker. 2021 , 14, 229-237	3
447	Cardiac Rehabilitation with Targeted Intensity Improves Cardiopulmonary Functions Accompanying with Reduced Copeptin Level in Patients with Coronary Artery Disease. 2021 , 14, 317-326	3
446	Relation Between Globulin, Fibrinogen, and Albumin With the Presence and Severity of Coronary Artery Disease. 2021 , 72, 174-180	3
445	Early risk stratification of acute myocardial infarction using a simple physiological prognostic scoring system: insights from the REACP study. 2021 , 20, 147-159	3
444	Rapidly and Accurately Detecting Significant Coronary Artery Stenosis in Patients with Suspected Stable Coronary Artery Disease and Normal Segmental Wall Motion by Speckle Tracking Automated Functional Imaging. 2021 , 47, 546-555	1
443	Differential associations of cardiac troponin T and cardiac troponin I with coronary artery pathology and dynamics in response to short-duration exercise. 2021 , 88, 23-29	3
442	Associations between smoking habits and major adverse cardiovascular events in patients who underwent coronary computed tomography angiography as screening for coronary artery disease. 2021 , 36, 483-491	0
441	Serum lipoprotein(a) and risk of periprocedural myocardial injury in patients undergoing percutaneous coronary intervention. 2020 , 44, 176	4

440	Association study of hsa_circ_0001946, hsa-miR-7-5p and PARP1 in coronary atherosclerotic heart disease. 2021 , 328, 1-7	5
439	Colchicine Inhibits Neutrophil Extracellular Trap Formation in Patients With Acute Coronary Syndrome After Percutaneous Coronary Intervention. 2021 , 10, e018993	28
438	The role of the plasma glycosylated hemoglobin A1c/Apolipoprotein A-I ratio in predicting cardiovascular outcomes in acute coronary syndrome. 2021 , 31, 570-578	4
437	The potential value of Copeptin and Pentraxin3 for evaluating the severity of coronary stenosis in patients with coronary artery disease. 2021 , 87, 32-38	2
436	Presence and severity of coronary artery disease in patients who achieved intensive blood pressure reduction at the time of coronary computed tomography angiography. 2021 , 44, 206-214	0
435	Relationship between P wave peak time and coronary artery disease severity in non-ST elevation acute coronary syndrome. 2021 , 46, 188-194	
434	Plasma proprotein convertase subtilisin/kexin type 9 concentration and recurrent cardiovascular events in patients with familial hypercholesterolemia. 2019 , 2047487319880985	6
433	Atherogenic dyslipidaemia and cardiovascular events in patients with diabetes or pre-diabetes and stable coronary artery disease: a prospective, cohort study. 2021 , 11, e037340	1
432	A risk biomarker for obstructive sleep apnea in patients with coronary artery disease: monocyte to high-density lipoprotein ratio. 2021 , 25, 1519-1526	
431	FABP4 and omentin-1 gene expression in epicardial adipose tissue from coronary artery disease patients. 2021 , 44, e20200441	0
430	Nutritional status and severity of coronary artery disease. 2021 , 32, 644-649	1
429	Higher serum level of Cystatin C: An additional risk factor of CAD. 2021 , 100, e24269	2
428	Thrombospondin-1 Levels in Patients with Coronary Heart Disease. 2021 , 11, 277-291	
427	Hypertrophy and Insulin Resistance of Epicardial Adipose Tissue Adipocytes: Association with the Coronary Artery Disease Severity. 2021 , 9,	11
426	Correlation between Carotid Intima-Media Thickness and Coronary Heart Disease in Patients with Hyperhomocysteinemia and Predictive Value of Revascularization. 2021 , 11, 2341-2349	
425	The Significance of Platelet/Lymphocyte Ratio in Relation to Coronary Angiographic Findings in Patients with Coronary Artery Disease. 2021 , 3, 183-190	
424	Circulating PGLYRP1 Levels as a Potential Biomarker for Coronary Artery Disease and Heart Failure. 2021 , 77, 578-585	2
423	[Optical coherence tomography angiography and cardiovascular diseases. An overview of the current knowledge]. 2021 , 118, 1119-1127	1

422	Prognostic Value of the Nutritional Risk Screening 2002 Scale in Patients With Acute Myocardial Infarction: Insights From the Retrospective Multicenter Study for Early Evaluation of Acute Chest Pain. 2021 , 36, 546-555	1
421	Retinal microvasculature impairments in patients with coronary artery disease: An optical coherence tomography angiography study. 2021 ,	3
420	Gensini Score and Thrombus Burden Add Predictive Value to the SYNTAX Score in Detecting No-Reflow after Myocardial Infarction. 2021 , 116, 466-472	0
419	Association between <i>Toxoplasma gondii</i> infection and coronary atherosclerosis. 2021 , 115, 1190-1197	
418	Liver fibrosis scores and coronary atherosclerosis: novel findings in patients with stable coronary artery disease. 2021 , 15, 413-423	6
417	Impact of diabetes on coronary severity and cardiovascular outcomes in patients with heterozygous familial hypercholesterolaemia. 2021 ,	3
416	Polymorphism of Interleukin-1 Gene Cluster in Polish Patients with Acute Coronary Syndrome. 2021 , 10,	1
415	Effects of sleep apnea hypopnea syndromes on cardiovascular events: a systematic review and meta-analysis. 2021 , 1	
414	Long non-coding RNA MALAT1 and its target microRNA-125b associate with disease risk, severity, and major adverse cardiovascular event of coronary heart disease. 2021 , 35, e23593	6
413	Perivascular Adipose Tissue Inflammation in Ischemic Heart Disease. 2021 , 41, 1239-1250	5
412	Association between high-density lipoprotein cholesterol levels and major adverse cardiovascular events in patients who underwent coronary computed tomography angiography: FU-CCTA registry. 2021 , 36, 1457-1465	0
411	Systemic immune-inflammation index predicts the severity of coronary stenosis in patients with coronary heart disease. 2021 , 32, 715-720	10
410	Blood urea nitrogen is associated with long-term all-cause mortality in stable angina pectoris patients: 8-year follow-up results. 2021 , 61, 66-70	1
409	Effects of Bivalirudin and Unfractionated Heparin on Liver and Renal Function in Chinese Patients with Coronary Artery Disease Undergoing Coronary Angiography with/without Percutaneous Coronary Intervention. 2021 , 9, 477-483	
408	Stenosis-DetNet: Sequence consistency-based stenosis detection for X-ray coronary angiography. 2021 , 89, 101900	5
407	Assessing the severity of cardiovascular disease in 213 088 patients with coronary heart disease: a retrospective cohort study. 2021 , 8,	
406	CD36+/CD61+ Microparticles Correlate with the Risk of Percutaneous Cardiac Interventions in Coronary Artery Disease Patients and the Effects of Ticagrelor. 2021 , 1	
405	Relationship between the Severity of Coronary Artery Disease and Cardiovascular Risk Factors in Acute Coronary Syndrome: Based on Tehran Heart Center's Data Registry. 2020 , 15, 165-170	1

404	Low-Density Lipoprotein Cholesterol 4: The Notable Risk Factor of Coronary Artery Disease Development. 2021 , 8, 619386	2
403	Periprocedural myocardial injury and coronary artery disease in patients undergoing transcatheter aortic valve replacement. 2021 , 35, 8-8	1
402	Smoking, alcohol and opioids effect on coronary microcirculation: an update overview. 2021 , 21, 185	7
401	Nutritional Risk Screening 2002 was associated with acute kidney injury and mortality in patients with acute coronary syndrome: Insight from the REACP study. 2021 , 31, 1121-1128	2
400	Association of Serum Level and DNA Methylation Status of Brain-Derived Neurotrophic Factor with the Severity of Coronary Artery Disease.. 2022 , 37, 159-168	0
399	Metabolic Signatures in Coronary Artery Disease: Results from the BioHEART-CT Study. 2021 , 10,	2
398	Impaired Peripheral Microvascular Function and Risk of Major Adverse Cardiovascular Events in Patients With Coronary Artery Disease. 2021 , 41, 1801-1809	7
397	Uric acid in chronic coronary syndromes: Relationship with coronary artery disease severity and left ventricular diastolic parameter. 2021 , 31, 1501-1508	7
396	Gut microbiota dysbiosis in stable coronary artery disease combined with type 2 diabetes mellitus influences cardiovascular prognosis. 2021 , 31, 1454-1466	2
395	Analysis of risk factors for different subtypes of acute coronary syndrome. 2021 , 49, 3000605211008326	3
394	Correlation between echocardiographic calcium score and coronary artery lesion severity on invasive coronary angiography. 2021 , 73, 307-312	
393	Prognosis of Coronary Atherosclerotic Burden in Non-Ischemic Dilated Cardiomyopathies. 2021 , 10,	0
392	The role of markers of endothelial dysfunction, oxidative and cellular stress in the prediction of myocardial infarction in comorbid patients with stable coronary heart disease. 2021 , 12, 23-27	
391	Risk stratification of ST-segment elevation myocardial infarction (STEMI) patients using machine learning based on lipid profiles. 2021 , 20, 48	4
390	Plaque erosion and acute coronary syndromes: phenotype, molecular characteristics and future directions. 2021 , 18, 724-734	24
389	Lack of association between serum IL-25 levels and acute coronary syndrome: a preliminary study. 2021 , 61, 60-65	
388	Cholesterol-related gene variants are associated with diabetes in coronary artery disease patients. 2021 , 48, 3945-3954	0
387	Coronary lesion complexity in patients with heterozygous familial hypercholesterolemia hospitalized for acute myocardial infarction: data from the RICO survey. 2021 , 20, 45	1

- 386 Relationship of epicardial fat volume with coronary plaque characteristics, coronary artery calcification score, coronary stenosis, and CT-FFR for lesion-specific ischemia in patients with known or suspected coronary artery disease. **2021**, 332, 8-14 3
- 385 Associations of Lipoprotein(a) With Coronary Atherosclerotic Burden and All-Cause Mortality in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. **2021**, 8, 638679 0
- 384 Value of galectin-3 in the diagnosis of acute coronary syndrome and the assessment of coronary artery lesions. **2021**, 15, 647-657
- 383 The Relationship Between Gensini Score and Erectile Dysfunction in Patients with Chronic Coronary Syndrome. **2021**, 9, 100376
- 382 Accuracy of triggering receptor expressed on myeloid cells 1 in diagnosis and prognosis of acute myocardial infarction: a prospective cohort study. **2021**, 9, e11655 2
- 381 Major dietary patterns and metabolic syndrome associated with severity of coronary artery disease: A structural equation modeling. **2021**, 2601060211020655 0
- 380 Serum levels of IL-32 in patients with coronary artery disease and its relationship with the serum levels of IL-6 and TNF- α . **2021**, 48, 4263-4271 3
- 379 A Study of Associations Between rs9349379 (PHACTR1), rs2891168 (CDKN2B-AS), rs11838776 (COL4A2) and rs4880 (SOD2) Polymorphic Variants and Coronary Artery Disease in Iranian Population. **2021**, 1 1
- 378 Patients with New-Onset Tumor of Severe Coronary Artery Disease May be at a Higher Risk of Arrhythmia. **2021**, 2021, 1948624
- 377 Coronary artery disease burden relation with the presentation of acute cardiac events and ventricular fibrillation. **2021**, 0
- 376 Novel Insights Into the Interaction Between the Autonomic Nervous System and Inflammation on Coronary Physiology: A Quantitative Flow Ratio Study. **2021**, 8, 700943 3
- 375 Sexual Dysfunction and the Impact of Beta-Blockers in Young Males With Coronary Artery Disease. **2021**, 8, 708200 1
- 374 CD200 Limits Monopoiesis and Monocyte Recruitment in Atherosclerosis. **2021**, 129, 280-295 3
- 373 The clinical role of combined serum C1q and hsCRP in predicting coronary artery disease. **2021**, 93, 50-58 0
- 372 Association between rs3088440 (G > A) polymorphism at 9p21.3 locus with the occurrence and severity of coronary artery disease in an Iranian population. **2021**, 48, 5905-5912
- 371 Circulating FoxP3+ T-lymphocytes in chronic coronary artery disease: Associations with the severity of atherosclerosis and lipid metabolism. **2021**, 36, 45-51
- 370 Increased circulating erythrocyte-derived microparticles in patients with acute coronary syndromes. **2021**, 15, 741-751 0
- 369 Non-mercaptalbumin is significantly associated with the coronary plaque burden and the severity of coronary artery disease. **2021**, 11, 14242 1

368	Prediction of presence and severity of coronary artery disease using prediction for atherosclerotic cardiovascular disease risk in China scoring system. 2021 , 9, 5453-5461	0
367	Effect of periodontal disease on oxidative stress markers in patients with atherosclerosis. 2021 , 1	1
366	Association Between Androgenic Alopecia and Coronary Artery Disease: A Cross-Sectional Study of Han Chinese Male Population. 2021 , 14, 4809-4818	0
365	Positive correlations between plasma BPI level and MPO-DNA and S100A8/A9 in myocardial infarction. 2021 , 1-9	0
364	Characteristics of circular RNAs expression of peripheral blood mononuclear cells in humans with coronary artery disease. 2021 , 53, 349-357	2
363	Relationship between frontal QRS-T duration and the severity of coronary artery disease in who were non-diabetic and had stable angina pectoris. 2021 , 25, 572-578	1
362	Reduction in GLP-1 secretory capacity may be a novel independent risk factor of coronary artery stenosis. 2021 , 11, 15578	
361	Coronary artery disease burden in women poorly explained by traditional risk factors: Sex disaggregated analyses from the BioHEART-CT study. 2021 , 333, 100-107	0
360	The Association of Thyroid Hormones with Cardiogenic Shock and Prognosis in Patients with ST Segment Elevation Myocardial Infarction (STEMI) Treated with Primary PCI. 2021 ,	1
359	Sortilin and Homocysteine as Potential Biomarkers for Coronary Artery Diseases. 2021 , 14, 6167-6176	0
358	Upregulation of long non-coding RNA myocardial infarction-associated transcription is correlated with coronary artery stenosis and elevated inflammation in patients with coronary atherosclerotic heart disease. 2021 , 37, 1038-1047	0
357	Increased levels of VCAM-1 is associated with higher occurrence of coronary artery disease in adults with moderate to severe obstructive sleep apnea. 2021 , 85, 131-137	1
356	Sleep profile and the risk of cardiovascular events in patients undergoing percutaneous coronary intervention. 2021 , 1-13	1
355	Gender Differences in the Relationship of Waist Circumference to Coronary Artery Lesions and One-Year Re-Admission Among Coronary Artery Disease Patients with Normal Body Mass Index. 2021 , 14, 4097-4107	3
354	Comprehensive Analysis of mRNA Expression Profiling and Identification of Potential Diagnostic Biomarkers in Coronary Artery Disease. 2021 , 6, 24016-24026	1
353	Serum glutamate and glutamine-to-glutamate ratio are associated with coronary angiography defined coronary artery disease.. 2022 , 32, 186-194	1
352	Patient Endothelial Colony-Forming Cells to Model Coronary Artery Disease Susceptibility and Unravel the Role of Dysregulated Mitochondrial Redox Signalling. 2021 , 10,	1
351	Risk Stratification of Patients with Acute Coronary Syndrome. 2021 , 10,	2

350	Long non-coding RNA THRIL is upregulated in coronary heart disease and binds to microRNA-424 to upregulate TXNIP in mice. 2021 , 138, 104215	1
349	Predictive role of Frontal QRS-T angle and Selvester QRS Score in determining angiographic slow flow phenomenon following percutaneous coronary intervention in patients with Non-ST elevation myocardial infarction. 2021 , 69, 20-26	
348	Expression of neutrophil elastase and myeloperoxidase genes in coronary atherosclerosis. 2021 , 25, 101336	1
347	Inflammatory markers in acute myocardial infarction and the correlation with the severity of coronary heart disease. 2021 , 53, 1041-1047	5
346	Relation between Baseline Coronary Atherosclerotic Status, Cardiovascular Events, and Malignancies in Type 2 Diabetics: A Long-Term Prospective Cohort Study. 2021 , 146, 419-425	2
345	Multi-omics study reveals that statin therapy is associated with restoration of gut microbiota homeostasis and improvement in outcomes in patients with acute coronary syndrome. 2021 , 11, 5778-5793	10
344	Association Between the Level of Low-Density Lipoprotein Cholesterol and Coronary Atherosclerosis in Patients Who Have Undergone Coronary Computed Tomography Angiography. 2021 , 12, 10-15	0
343	Comparison Between Small Dense LDL-Cholesterol and LDL-Cholesterol to Predict Coronary Events in Stable Coronary Heart Disease. 2008 , 147-153	0
342	Cyclophilin A: Novel Biomarker for Oxidative Stress and Cardiovascular Diseases. 2015 , 405-438	1
341	The Clinical Role of Angiotensin-Like Protein 3 in Evaluating Coronary Artery Disease in Patients with Obstructive Sleep Apnea. 2020 , 34, 773-780	4
340	Trans and conjugated fatty acids in dairy products: Cause for concern?. 2020 , 87-120	3
339	Recovery of Left Ventricular Systolic Function and Clinical Outcomes in Young Adults With Myocardial Infarction. 2020 , 75, 2804-2815	12
338	Association between mitochondrial DNA haplogroup variation and coronary artery disease. 2020 , 30, 960-966	2
337	Testosterone is negatively associated with the severity of coronary atherosclerosis in men. 2012 , 14, 875-8	33
336	Colchicine inhibits neutrophil extracellular trap formation in acute coronary syndrome patients after percutaneous coronary intervention.	2
335	Gene polymorphism but not catalytic activity of angiotensin I-converting enzyme is associated with coronary artery disease and myocardial infarction in low-risk patients. 1995 , 92, 2796-9	65
334	Coronary artery disease and polymorphisms in a receptor mediating shear stress-dependent platelet activation. 1997 , 96, 3281-6	72
333	Coronary disease is not associated with robust alterations in inflammatory gene expression in human epicardial fat. 2019 , 4,	7

332	Selective Topical Cooling for Myocardial Protection. 1993 , 1, 176-181	2
331	Association of the Platelet Glycoprotein Ia C807T Gene Polymorphism With Nonfatal Myocardial Infarction in Younger Patients. 1999 , 93, 2449-2453	5
330	Correlation between prediabetes and coronary artery disease severity in patients undergoing elective coronary angiography. 2019 , 71, 34	6
329	Prevalence of type 2 diabetes mellitus and association of HbA1c with severity of coronary artery disease in patients presenting as non-diabetic acute coronary syndrome. 2020 , 72, 66	2
328	Is the ratio of apoB/apoA-1 the best predictor for the severity of coronary artery lesions in Chinese diabetics with stable angina pectoris? An assessment based on Gensini scores. 2015 , 12, 402-9	9
327	Relationship of lipid and lipoprotein ratios with coronary severity in patients with new on-set coronary artery disease complicated with type 2 diabetics. 2016 , 13, 685-692	5
326	Plasma ghrelin and pro-inflammatory markers in patients with obstructive sleep apnea and stable coronary heart disease. 2013 , 19, 251-6	19
325	A New Potential Predictor of Coronary Artery Disease: The Ratio of Mitral Peak Filling Velocity to Mitral Annular Velocity in Early Diastole. 2017 , 23, 1180-1189	2
324	Correlation Between Pigment Epithelium-Derived Factor (PEDF) level and Degree of Coronary Angiography and Severity of Coronary Artery Disease in a Chinese Population. 2018 , 24, 1751-1758	3
323	Relationship of Autophagy and Apoptosis with Total Occlusion of Coronary Arteries. 2018 , 24, 6984-6988	7
322	R/S ratio in lead II, and the prognostic significance of red cell distribution width in acute coronary syndrome. 2019 , 7, 2217-2226	3
321	Retinal and choroidal vascular changes in coronary heart disease: an optical coherence tomography angiography study. 2019 , 10, 1532-1544	44
320	Association of NT-proBNP and multiple biomarkers with severity of angiographic coronary artery disease in diabetic and pre-diabetic Chinese patients. 2011 , 6, e22563	5
319	Imbalanced frequencies of Th17 and Treg cells in acute coronary syndromes are mediated by IL-6-STAT3 signaling. 2013 , 8, e72804	18
318	Depressive symptoms are associated with mental stress-induced myocardial ischemia after acute myocardial infarction. 2014 , 9, e102986	30
317	SDF1 Polymorphisms Influence Outcome in Patients with Symptomatic Cardiovascular Disease. 2016 , 11, e0161933	9
316	Prediction of Coronary Artery Disease Extent and Severity Using Pulse Wave Velocity. 2016 , 11, e0168598	13
315	Oral glucose tolerance test predicts increased carotid plaque burden in patients with acute coronary syndrome. 2017 , 12, e0183839	2

314	Epicardial Adipose Tissue Thickness Is an Independent Predictor of Critical and Complex Coronary Artery Disease by Gensini and Syntax Scores. 2016 , 43, 29-37	15
313	Association of Heart Rate Recovery With Microalbuminuria in Non-Obstructive Coronary Artery Disease. 2017 , 8, 206-213	2
312	Associations Between High Levels of High-Density Lipoprotein Cholesterol and the Presence and Severity of Coronary Artery Disease in Patients Who Have Undergone Coronary Computed Tomography Angiography. 2020 , 12, 734-739	1
311	The role of oxidative DNA damage and GSTM1, GSTT1, and hOGG1 gene polymorphisms in coronary artery disease risk. 2016 , 16, 931-938	11
310	Assessment of atrial conduction times in prediabetic patients with coronary artery disease. 2017 , 17, 374-380	3
309	The value of real-time myocardial contrast echocardiography for detecting coronary microcirculation function in coronary artery disease patients. 2018 , 19, 27-33	2
308	The significant coronary tortuosity and atherosclerotic coronary artery disease; What is the relation?. 2018 , 10, 209-213	3
307	Genetic variations of Mrf-2/ARID5B confer risk of coronary atherosclerosis in the Japanese population. 2008 , 49, 313-27	7
306	N-terminal pro-B-type natriuretic peptide level is depressed in patients with significant coronary artery disease who have high body mass index. 2008 , 49, 403-12	5
305	Brachial arterial stiffness predicts coronary atherosclerosis in patients at risk for cardiovascular diseases. 2004 , 45, 231-42	20
304	[Combined markers of initial stages of coronary atherosclerosis]. 2017 , 63, 272-277	3
303	DNA hydroxymethylation combined with carotid plaques as a novel biomarker for coronary atherosclerosis. 2019 , 11, 3170-3181	9
302	Serum retinol-binding protein 4 is associated with the presence and severity of coronary artery disease in patients with subclinical hypothyroidism. 2019 , 11, 4510-4520	7
301	Toll-like receptor 4 rs11536889 is associated with angiographic extent and severity of coronary artery disease in a Chinese population. 2017 , 8, 2025-2033	7
300	Association between fibrinogen level and the severity of coronary stenosis in 418 male patients with myocardial infarction younger than 35 years old. 2017 , 8, 81361-81368	13
299	Combined value of red blood cell distribution width and global registry of acute coronary events risk score on predicting long-term major adverse cardiac events in STEMI patients undergoing primary PCI. 2018 , 9, 13971-13980	7
298	Increased levels of total P-Cresylsulphate and indoxyl sulphate are associated with coronary artery disease in patients with diabetic nephropathy. 2010 , 7, 275-84	50
297	Acute Phase Response: Implication in ST-segment Elevation Myocardial Infarction. 2014 , 8, 44-51	5

296	Association of vitamin D deficiency and premature coronary artery disease. 2019 , 10, 80-85	2
295	Obstructive sleep apnea and severity of coronary artery disease. 2018 , 9, 276-282	6
294	Correlation of heart rate and blood pressure variability as well as hs-CRP with the burden of stable coronary artery disease. 2020 , 68, 376-382	2
293	Serum Cortisol Level as a Predictor of In-Hospital Mortality in Patients Undergoing Primary Percutaneous Intervention for ST Segment Elevation Myocardial Infarction. 2019 , 4, 41-44	0
292	Red Blood Cell Distribution Width (RDW) and its Association with Coronary Atherosclerotic Burden in Patients with Stable Angina Pectoris. 2012 , 9,	3
291	Importance of Reciprocal ST Segment Depression in the Extensive Coronary Artery Disease. 2010 , 7,	1
290	Assessment of the Correlation between Severity of Coronary Artery Disease and Waist-Hip Ratio. 2017 , 5, 929-933	5
289	Relationship between peripheral leukocyte count and the severity of stable angina determined by coronary angiography. 2004 , 19, 99-103	1
288	An increased monocyte count predicts coronary artery spasm in patients with resting chest pain and insignificant coronary artery stenosis. 2006 , 21, 97-102	12
287	Correlation between serum free fatty acids levels and Gensini score in elderly patients with coronary heart disease. 2014 , 11, 57-62	7
286	Kinetics of plasma microRNA-499 expression in acute myocardial infarction. 2015 , 7, 890-6	39
285	Plasma α -antitrypsin: a neglected predictor of angiographic severity in patients with stable angina pectoris. 2015 , 128, 755-61	3
284	Independent Prognostic Value of High-sensitivity C-reactive Protein in Patients with Coronary Artery Ectasia. 2016 , 129, 2582-2588	6
283	Serum sex hormone and growth arrest-specific protein 6 levels in male patients with coronary heart disease. 2016 , 18, 644-9	7
282	The association between fasting serum insulin, apo-lipoproteins level, and severity of coronary artery involvement in non-diabetic patients. 2014 , 3, 192	2
281	Association between prolonged PR intervals and significant coronary artery disease in patients with non-ST elevation myocardial infarction and unstable angina. 2020 , 9, 6	1
280	Assessment of long-term mortality in patients with complex coronary artery disease undergoing percutaneous intervention: comparison of multiple anatomical and clinical prognostic risk scores. 2017 , 13, 1177-1184	6
279	Predictors of high-sensitivity cardiac troponin in stable patients undergoing coronary angiography. 2016 , 12, 329-36	3

278	Subtle myocardial damage associated with diagnostic coronary angiography alone. 2010 , 6, 388-93	2
277	A Survey of Game Theory in Wireless Sensor Networks Security. 2011 , 6,	37
276	Plasma tissue factor pathway inhibitor levels in angiographically defined coronary artery disease among saudis. 2013 , 28, 191-4	2
275	Serum neutrophil gelatinase-associated lipocalin levels are correlated with the complexity and the severity of atherosclerosis in acute coronary syndrome. 2015 , 15, 450-5	14
274	Novel predictors of infarct-related artery patency for ST-segment elevation myocardial infarction: Platelet-to-lymphocyte ratio, uric acid, and neutrophil-to-lymphocyte ratio. 2015 , 15, 648-56	17
273	Altered heart rate variability depend on the characteristics of coronary lesions in stable angina pectoris. 2015 , 15, 496-501	11
272	Relationship between red cell distribution width and long-term mortality in patients with non-ST elevation acute coronary syndrome. 2015 , 15, 634-9	18
271	The relation between intensity and complexity of coronary artery lesion and oxidative stress in patients with acute coronary syndrome. 2015 , 15, 795-800	20
270	High levels of HB-EGF and interleukin-18 are associated with a high risk of in-stent restenosis. 2015 , 15, 907-12	10
269	The association of paraoxonase 1 gene L55M polymorphism with the extent and severity of coronary artery disease in the Turkish population and its dependence on gender. 2016 , 16, 175-82	14
268	Relationship of Plasma Adiponectin and Waist-hip Ratio with Coronary Artery Disease. 2016 , 70, 413-418	6
267	Decreased beta cell function and insulin sensitivity contributed to coronary artery disease in patients with normal glucose tolerance. 2012 , 19, 806-13	10
266	Insulin response to oral glucose load is associated with coronary artery disease in subjects with normal glucose tolerance. 2008 , 15, 6-12	10
265	Hyperuricemia and severity of coronary artery disease: An observational study in adults 35 years of age and younger with acute coronary syndrome. 2019 , 26, 275-282	8
264	Red cell distribution width is correlated with extensive coronary artery disease in patients with diabetes mellitus. 2017 , 28, 319-323	1
263	Assessment of the left atrial volume index and plasma NT-proANP level in patients with acute ST-elevation myocardial infarction. 2013 , 68, 997-1003	3
262	Clinical application of simultaneous detection of cystatin C, cathepsin S, and IL-1 in classification of coronary artery disease. 2017 , 31, 315-320	3
261	Association of Thyroid Function with Severity of Coronary Artery Disease in Euthyroid Patients. 2015 , 9, OC10-3	8

260 Classification of Coronary Artery Lesions Based on XGBoost. **2021**, 178-187

259 SUBCELLULAR LOCALIZATION OF FoxP3 TRANSCRIPTION FACTOR IN PATIENTS WITH ACUTE CORONARY SYNDROME: COMPARATIVE ANALYSIS AND PROSPECTIVE OBSERVATION. **2021**, 23, 731-736

258 Levels of miR-130b-5p in peripheral blood are associated with severity of coronary artery disease. **2021**, 48, 7719-7732

1

257 Association Between Major Adverse Cardiovascular Events and the Ratio of Subcutaneous Fat Area to Visceral Fat Area in Patients Who Have Undergone Multidetector Row Computed Tomography. **2021**, 3, 674-681

0

256 C Deletion at the re74650330 Locus of the Gene (rs74650330) Increases the Risk of Coronary Artery Disease in Individuals with Low-Density Lipoprotein Cholesterol Levels. **2021**, 25, 660-667

255 Effects of sleep habits on acute myocardial infarction risk and severity of coronary artery disease in Chinese population. **2021**, 21, 481

1

254 Association of Global Coagulation Profiles With Cardiovascular Risk Factors and Atherosclerosis: A Sex Disaggregated Analysis From the BioHEART-CT Study. **2021**, 10, e020604

0

253 Genetic and epigenetic associations of ANRIL with coronary artery disease and risk factors. **2021**, 14, 240

1

252 Carotid Endarterectomy in Patients with Ischemic Heart Disease : Evaluation of Our Patients and Review of Literature. **2003**, 12, 723-728

6

251 Association between coronary endothelial dysfunction and local inflammation of atherosclerotic coronary arteries. **2003**, 67-73

250 Coronary Artery Bypass Surgery and Percutaneous Coronary Revascularization: Impact on Morbidity and Mortality in Patients with Coronary Artery Disease. **2007**, 1073-1111

0

249 Elevated Small Dense Low-density Lipoprotein Cholesterol Concentration as a Promising Risk Marker for Severe Stable Coronary Heart Disease, Independently of Adiponectin and C-reactive Protein. **2007**, 19, 81-93

248 Do Japanese diagnostic criteria for metabolic syndrome (MS) reflect the severity on coronary arteriography?. **2007**, 53, 131-137

247 Determination of plasma copeptin level in patients with coronary heart disease and its diagnostic value for severity of coronary atherosclerosis. **2009**, 29, 1154-1157

246 Development of quantitative ELISA for determination of serum adiponectin and its clinical application. **2010**, 29, 850-853

245 Blood cellular mutant LXR- β protein stability governs initiation of coronary heart disease. **2013**, 5, 305-12

1

244 MCV and MCH Values in Coronary Artery Patients with Positive Gensini Score. **2013**, 10,

243 Cyclophilin A: Novel Biomarker for Oxidative Stress and Cardiovascular Diseases. **2014**, 1-27

242	The Cardiac Catheterization Laboratory for Coronary Angioplasty. 1986 , 61-92	
241	Techniques of Cardiac Catheterization and Coronary Angiography. 1989 , 10-21	
240	Different Lipoprotein Patterns in Men and Women Examined by Coronary Angiography. 1993 , 125-129	
239	Predictors of development of cardiovascular events after stenting of coronary arteries at patients with obesity. 2014 , 5, 5-9	
238	[Markers of visceral adipose tissue metabolic abnormalities in patients with coronary atherosclerosis in relation to the presence of type 2 diabetes mellitus]. 2015 , 87, 31-36	
237	Association of Angiotensin-Converting Enzyme Genotype, Insertion/Deletion Polymorphism and Saphenous Vein Graft Atherosclerosis in Iranian Patients. 2015 , 30, 557-61	2
236	Adiponectin as Biomarker in Coronary Artery Disease. 2015 , 1-17	
235	EPICARDIAL OBESITY AND A RANGE OF METABOLIC RISK FACTORS RELATION WITH THE PREVALENCE INDEX OF CORONARY ATHEROSCLEROSIS. 2015 , 14, 35-40	1
234	Analysis of the Clinical Characteristics of Patients with Acute Coronary Syndrome in Different States of Renal Function. 2015 , 64, 357-61	1
233	Serum cystatin C and neutrophil gelatinase-associated lipocalin in predicting the severity of coronary artery disease in diabetic patients. 2016 , 16, 756-761	1
232	Relationship between calculated total antioxidant status and atherosclerotic coronary artery disease. 2016 , 16, 689-95	5
231	Association between parathyroid hormone levels and the extensiveness of coronary artery disease. 2016 , 16, 839-843	2
230	Correlation between Serum Dehydroepiandrosterone Sulfate and Coronary Artery Disease in Middle-Aged and Elderly Men. 2016 , 66, 111-116	
229	Cholesteryl ester transfer protein gene polymorphism (I405V) and premature coronary artery disease in an Iranian population. 2016 , 16, 114-20	6
228	Adiponectin as Biomarker in Coronary Artery Disease. 2016 , 635-651	1
227	The association between aspirin resistance and extent and severity of coronary atherosclerosis. 2018 , 5, 323-328	2
226	Homocysteine enhances the predictive value of the GRACE risk score in patients with ST-elevation myocardial infarction. 2017 , 18, 182-193	5
225	Association of Multiple Genetic Variants with the Extension and Severity of Coronary Artery Disease. 2018 , 110, 16-23	2

- 224 Relationship between myocardial performance index and severity of coronary artery disease in patients with non-ST-segment elevation acute coronary syndrome. **2017**, 28, 4-7
- 223 Relationship Between Morning Hypertension and T-Peak to T-End Interval in Patients with Suspected Coronary Heart Disease. **2017**, 2, 407-414
- 222 Similarity of Adipocytokines Level in Radial and Coronary Artery Associated with Epicardial Adipose Tissue Thickness. **2017**, 17, 107-116
- 221 Association of a disintegrin and metalloproteinase with a thrombospondin type 1 motif member 13 polymorphisms with severity of coronary stenosis in type 2 diabetes mellitus. **2018**, 23, 59 3
- 220 Possible Roles of Epicardial Adipose Tissue in the Pathogenesis of Coronary Atherosclerosis. **2018**, 4, 5-10 1
- 219 Relationship between Aortic Valve Sclerosis and the Severity of Coronary Artery Disease in Patients Undergoing Diagnostic Coronary Angiography. **2018**, 11,
- 218 Fragmente QRS kompleksleri koroner anjiyografi uygulanan stabil koroner arter hastalarında daha yüksek Gensini skorunu içermektedir. **2018**, 57, 71-74 1
- 217 MULTIMARKER DIAGNOSTIC PANELS FOR ATHEROSCLEROSIS. **2018**, 65-73 4
- 216 Correlation of Resting Heart Rate with the Severity and Complexity of Coronary Artery Disease: A Single-Center Retrospective Study. **2019**, 10, 104 3
- 215 Serum level of High-density lipoprotein particles are independently associated with long-term prognosis in patients with coronary artery disease: The GENES study.
- 214 Diagnostic value of chaperone activity of heat shock proteins 70 in coronary atherosclerosis. **2019**, 10, 60-64
- 213 Relevance of Syntax Score as a Marker of Complex Coronary Artery Disease and Plaque Burden in Young Coronary Artery Disease - A Single Centre Multifactorial Analysis. **2019**, 8, 3039-3043
- 212 Akut Koroner Sendromlu Hastalarda Plazma von Willebrand Faktör Düzeyi ile Koroner Arter Hastalığı Ciddiyeti Arasındaki İlişi.
- 211 Gensini skoru pompa destekli koroner arter baypas greftleme sonrası gelişen atriyal fibrilasyonun değerlendirilmesi.
- 210 A Coronary Artery Calcium Score of Zero in Patients Who Have Undergone Coronary Computed Tomography Angiography Is Associated With Freedom From Major Adverse Cardiovascular Events. **2020**, 12, 662-667 0
- 209 Correlation of glycosylated hemoglobin level with the severity of coronary artery disease in diabetic patients. **2020**, 4, 001-005
- 208 Association between coronary artery disease severity and videocapillaroscopic findings of nail fold capillary circulation.
- 207 Lipoprotein(a) as a predictor for the presence and severity of premature coronary artery disease: a cross-sectional analysis of 2433 patients. **2021**, 32, 78-83 1

- 206 New Opportunities for Determining the Terms of Carrying out the Control Coronarangiography after Percutaneous Coronary Intervention. **2020**, 5, 51-59
- 205 Non-Alcoholic Fatty Liver Disease as a Coronary Heart Disease Severity Predictor. **2020**, 11, 182-192 1
- 204 Discordance of Low-Density Lipoprotein Cholesterol and Non-High-Density Lipoprotein Cholesterol and Coronary Artery Disease Severity. **2020**, 114, 469-475 1
- 203 MELD-XI Score and Coronary Artery Disease Prevalence and Extent Among In-Hospital Patients With End-Stage Liver Failure Awaiting Transplant. **2020**, 18, 88-92 0
- 202 The Relationship between Lipoprotein-Associated Phospholipase A2 and Coronary Heart Disease. **2020**, 10, 1229-1238
- 201 Spiritual well-being and coronary artery diseases severity: Mediating effects of anger rumination and worry. **2021**, 80, 501-512 1
- 200 Elevated serum miR-3129-5p contributes to the progression of coronary heart disease via targeting mTOR. **2021**, 37, 314-323 0
- 199 Validation of combined markers of the presence and severity of coronary atherosclerosis. **2020**, 23, 65
- 198 Validation of combined markers of the presence and severity of coronary atherosclerosis. **2020**, 23, 94
- 197 Association between Endothelial Nitric Oxide Synthase Gene Polymorphism (Glu298Asp) and Coronary No-Reflow Phenomenon in Acute Myocardial Infarction. **2019**, 73, 529-535 0
- 196 Role of Urotensin-II in Saphenous Vein Graft Disease. **2020**, 35, 675-682 1
- 195 Frailty Significantly Associated with a Risk for Mid-term Outcomes in Elderly Chronic Coronary Syndrome Patients: a Prospective Study. **2020**, 35, 897-905 1
- 194 Elevated Plasma von Willebrand Factor Antigen and Activity Levels Are Associated With the Severity of Coronary Stenosis. **2020**, 26, 1076029619900552 0
- 193 Multivariate Analysis of Coronary Artery Lesions in Patients with Coronary Heart Disease. **2020**, 10, 2727-2732 1
- 192 Low profilin 1 serum levels are associated with diabetes, family history and multivessel lesions in patients with coronary artery disease. **2021**, 17, 305-308
- 191 Kalp Tipi Yağ Asid Bağlayıcı Proteinin Akut Koroner Sendrom Risk Skorları ile İlişkisi [tr] / The Relationship between Cardiac Fatty Acid Binding Protein and Acute Coronary Syndrome Risk Scores [en]. 165-175
- 190 Factors affecting brain structure in smoking-related diseases: Chronic Obstructive Pulmonary Disease (COPD) and coronary artery disease. **2021**, 16, e0259375 0
- 189 Cardiac CT scanning in coronary artery disease: Epicardial fat volume and its correlation with coronary artery lesions and left ventricular function. **2020**, 20, 2961-2968 1

188	Impact of obstructive sleep apnea on left ventricular mass index in men with coronary artery disease. 2020 , 16, 1675-1682	3
187	T786C eNOS polymorphism as risk factor for recurrent myocardial infarction in young and middleaged patients. 2020 , 17-21	
186	Exercise testing score for myocardial ischemia gradation. 2007 , 7, 61-72	
185	Variation in high-sensitivity C-reactive protein levels over 24 hours in patients with stable coronary artery disease. 2010 , 37, 42-8	20
184	Comparing angiography features of inferior versus anterior myocardial infarction regarding severity and extension in a cohort of Iranian patients. 2011 , 16, 484-9	2
183	Angiotensin- converting enzyme insertion/deletion polymorphism and its association with coronary artery disease in an Iranian population. 2013 , 8, 89-94	2
182	Comparative evaluation of the diagnostic accuracy of (99m)Tc-sestamibi gated SPECT using five different sets of image acquisitions at stress and rest phases for the diagnosis of coronary artery disease. 2013 , 4, 10-6	1
181	Relationship between Renal Artery Stenosis and Severity of Coronary Artery Disease in Patients with Coronary Atherosclerotic Disease. 2012 , 6, 84-7	3
180	The relationship between level of androgenic hormones and coronary artery disease in men. 2007 , 18, 362-6	12
179	Impact of estimated HDL particle size via the ratio of HDL-C and apoprotein A-I on short-term prognosis of diabetic patients with stable coronary artery disease. 2014 , 11, 245-52	3
178	The relationship between regional brain volumes and the extent of coronary artery disease in mild cognitive impairment. 2014 , 19, 739-45	12
177	Coronary artery disease and plasma apolipoprotein E4 in mild cognitive impairment. 2014 , 10, 244-51	2
176	Association of serum chemerin levels with the severity of coronary artery disease in patients with metabolic syndrome. 2014 , 7, 5461-8	8
175	Rheumatoid factor, anti-nuclear antibody in ischemic heart disease: Acute versus chronic patients. 2014 , 10, 305-10	1
174	Analysis of the risk factors and characteristics of coronary artery disease of Han, Uygur and Kazak patients with acute myocardial infarction in Xinjiang district. 2015 , 8, 2831-8	2
173	Association analysis of GWAS and candidate gene loci in a Chinese population with coronary heart disease. 2015 , 8, 7497-506	3
172	5,10-methylene tetrahydrofolate reductase C677T gene polymorphism, homocysteine concentration and the extent of premature coronary artery disease in southern Iran. 2013 , 12, 437-48	6
171	Obstructive sleep apnea, diagnosed by the Berlin questionnaire and association with coronary artery disease severity. 2015 , 11, 275-80	5

170	Association between left ventricular end-diastolic pressure and coronary artery disease as well as its extent and severity. 2015 , 8, 18673-80	3
169	Correlation analysis between serum lipoprotein (a) and the incidence of aortic valve sclerosis. 2015 , 8, 19318-24	2
168	Synergistic effect of hypertension with diabetes mellitus and gender on severity of coronary atherosclerosis: Findings from Tehran Heart Center registry. 2015 , 11, 317-22	7
167	Lack of Association between High-Density Lipoprotein Cholesterol and Angiographic Coronary Lesion Severity in Chinese Patients with Low Background Low-Density Lipoprotein Cholesterol. 2015 , 31, 528-35	4
166	The Association between Androgenic Hormone Levels and the Risk of Developing Coronary Artery Disease (CAD). 2016 , 45, 14-9	6
165	Analyzing Gensini Score as a Semi-Continuous Outcome. 2016 , 11, 55-61	6
164	Evaluation of Echocardiographic Epicardial Fat Thickness as a Sign of Cardiovascular Risk in Positive Exercise Test Patients. 2016 , 32, 684-689	3
163	Relationship between R-R interval and left ventricular systolic synchrony in subjects with coronary artery disease determined using angiography. 2016 , 6, 175-184	
162	Prevalence of non-alcoholic fatty liver disease in patients with coronary artery disease. 2016 , 12, 201-205	6
161	Free triiodothyronine in relation to coronary severity at different ages: Gensini score assessment in 4206 euthyroid patients. 2016 , 13, 978-983	5
160	Altered serum level of cartilage oligomeric matrix protein and its association with coronary calcification in patients with coronary heart disease. 2017 , 14, 87-92	6
159	The correlation between plasma human neutrophil peptide 1-3 levels and severity of coronary artery disease. 2016 , 1, e133-e138	2
158	Predictors of Impaired Reperfusion after Percutaneous Coronary Intervention in Patients with In-Hospital Acute Stent Thrombosis: A Retrospective Analyses of 5 Years of Data. 2017 , 33, 384-392	3
157	Association of invasive treatment and lower mortality of patients \geq 80 years with acute myocardial infarction: a propensity-matched analysis. 2018 , 15, 666-674	2
156	Corrected QT Interval Prolongations in Patients with Non-ST-Elevation Acute Coronary Syndrome. 2018 , 13, 173-179	1
155	Association of Epicardial Adipose Tissue Thickness with Extent and Complexity of Coronary Artery Disease in Patients with Acute Coronary Syndrome. 2019 , 35, 459-467	2
154	[Correlation of lipoprotein(a) with clinical stability and severity of coronary artery lesions in patients with coronary artery disease]. 2019 , 39, 235-240	1
153	Association between Vascular Endothelial Growth Factor Plasma Levels and rs699947 Polymorphism and Coronary Collateral Vessel Formation. 2019 , 14, 121-127	1

- 152 [Serum annexin A2 level is significantly elevated in patients with coronary heart disease]. **2020**, 40, 382-387
- 151 Prothrombin Gene G20210A Variant in Angiographically Documented Patients with Coronary Artery Stenosis. **2019**, 14, 150-155
- 150 Association between α -1-antitrypsin and acute coronary syndrome. **2020**, 20, 119
- 149 [Correlation of plasma N-acetyl-neuraminic acid level with TIMI risk stratification and clinical outcomes in patients with acute coronary syndrome]. **2020**, 40, 1253-1258
- 148 Inflammation is associated with the presence and severity of chronic coronary syndrome through soluble CD40 ligand. **2020**, 10, 329-339 1
- 147 Dyslipidemia, Not Inflammatory Markers or Adipokines, Contributes Significantly to a Higher SYNTAX Score in Stable Coronary Artery Disease (from the Taichung CAD Study). **2021**, 37, 232-238
- 146 [Correlation of serum ADAMTS13 and TSP1 levels with myocardial injury and prognosis in patients with acute coronary syndrome]. **2021**, 41, 710-715
- 145 Major Dietary Patterns Relationship with Severity of Coronary Artery Disease in Gaza-Strip, Palestine: A Cross-Sectional Study. **2021**, 31, 599-610
- 144 Using the Gensini score to estimate severity of STEMI, NSTEMI, unstable angina, and anginal syndrome. **2021**, 100, e27331 1
- 143 Using the Gensini score to estimate severity of STEMI, NSTEMI, unstable angina, and anginal syndrome. **2021**, 100, e27331 1
- 142 Aorta unfolding index as an indicator for coronary artery disease severity. **2021**, 2841851211057277
- 141 The relation of circulating cell division cycle 42 expression with Th1, Th2, and Th17 cells, adhesion molecules, and biochemical indexes in coronary heart disease patients. **2021**, 1 5
- 140 Androgen-dependent tissue factor pathway inhibitor regulating protein: a review of its peripheral actions and association with cardiometabolic diseases. **2021**, 1 0
- 139 Relationship Between Plasma Total Homocysteine Levels and Mean Corrected TIMI Frame Count in Patients with Acute Myocardial Infarction. **2021**, 14, 8161-8172 0
- 138 MicroRNA-34a in coronary heart disease: Correlation with disease risk, blood lipid, stenosis degree, inflammatory cytokines, and cell adhesion molecules. **2021**, 36, e24138 1
- 137 Telomere Length and Oxidative Stress in Patients with ST-Segment Elevation and Non-ST-Segment Elevation Myocardial Infarction. **2021**, 183
- 136 Relation of circulating lncRNA GAS5 and miR-21 with biochemical indexes, stenosis severity, and inflammatory cytokines in coronary heart disease patients.. **2022**, e24202 2
- 135 Association between α -1-antitrypsin and acute coronary syndrome. **2020**, 20, 1-1 1

134	Prevalence, risk factors, and coronary angiographic profile in patients with tortuous coronary artery. 2021 , 63, 547-554	
133	Predictive Value of Gensini Score in the Long-Term Outcomes of Patients With Coronary Artery Disease Who Underwent PCI.. 2021 , 8, 778615	2
132	The Association of Thyroid Hormones with Coronary Atherosclerotic Severity in Euthyroid Patients.. 2022 , 54, 12-19	0
131	Relationship between Braden Scale scores and acute kidney injury among patients with acute coronary syndrome: a multicentre retrospective cohort study.. 2022 , 12, e049102	
130	Alterations of Gut Microbiome and Serum Metabolome in Coronary Artery Disease Patients Complicated With Non-alcoholic Fatty Liver Disease Are Associated With Adverse Cardiovascular Outcomes.. 2021 , 8, 805812	1
129	Immunoglobulin E Sensitization to Mammalian Oligosaccharide Galactose-a-1,3 (EGal) Is Associated With Noncalcified Plaque, Obstructive Coronary Artery Disease, and ST-Segment-Elevated Myocardial Infarction.. 2022 , ATVBAHA121316878	2
128	Early life stress and autonomic response to acute mental stress in individuals with coronary heart disease.. 2022 ,	
127	Impact of serum uric acid levels on the clinical prognosis and severity of coronary artery disease in patients with acute coronary syndrome and hypertension after percutaneous coronary intervention: a prospective cohort study.. 2022 , 12, e052031	1
126	The interplay of long noncoding RNA HULC with microRNA-128-3p and their correlations with lipid level, stenosis degree, inflammatory cytokines, and cell adhesion molecules in coronary heart disease patients.. 2022 , 1	
125	Preexisting Cardiovascular Risk Factors and Coronary Artery Atherosclerosis in Patients with and without Cancer.. 2022 , 2022, 4570926	0
124	Impact of Chronic Kidney Disease on the Associations of Cardiovascular Biomarkers With Adverse Outcomes in Patients With Suspected or Known Coronary Artery Disease: The EXCEED-J Study.. 2022 , e023464	0
123	A Novel Integrated Biomarker for Evaluation of Risk and Severity of Coronary Atherosclerosis, and Its Validation.. 2022 , 12,	0
122	Carotid intima-media thickness predicted the presence but not the severity or complexity of coronary artery disease in a South Asian population.. 2022 ,	
121	Retinal Microvasculature Changes in Patients With Coronary Total Occlusion on Optical Coherence Tomography Angiography.. 2021 , 8, 708491	1
120	Association between coronary endothelial dysfunction and local inflammation of atherosclerotic coronary arteries. 2003 , 249, 67-73	5
119	Diagnostic performance of coronary computed tomography angiography-derived fractional flow reserve in lesion-specific ischemia patients with different Gensini score levels.. 2022 , 10, 412	0
118	Relationship between Apolipoprotein E Genotype and Lipoprotein Profile in Patients with Coronary Heart Disease.. 2022 , 27,	
117	Prognostic Value of N-Terminal Pro-B-Type Natriuretic Peptide and High-Sensitivity C-Reactive Protein in Patients With Previous Myocardial Infarction.. 2022 , 9, 797297	0

116	Albumin to globulin ratio was associated with in-stent restenosis and revascularization events after percutaneous coronary intervention.. 2022,	1
115	Predictive value of ACEF II score for adverse prognosis in patients with coronary heart disease after percutaneous coronary intervention.. 2022,	1
114	Comprehensive Metabolic Profiling of Inflammation Indicated Key Roles of Glycerophospholipid and Arginine Metabolism in Coronary Artery Disease.. 2022, 13, 829425	0
113	Association between Non-alcoholic Fatty Liver Disease and the Severity of Coronary Artery Stenosis in Eastern Chinese Population. 2022, 21,	0
112	Association of serum cardiac troponin I and severity of coronary stenosis in patients with varied renal functions: a retrospective cohort study.. 2022, 12, e054722	
111	Late Percutaneous Coronary Intervention is Associated with Better Prognosis of Patients with Acute Myocardial Infarction.. 2022, 15, 2621-2627	0
110	Serum miR-497-5p serves as a diagnostic biomarker for acute coronary syndrome and predicts the occurrence of major adverse cardiovascular events after percutaneous coronary intervention.. 2022 , 13, 8266-8276	0
109	Association Between Insulin Resistance Estimated by Triglyceride Glucose Index and In-Stent Restenosis in Non-Diabetic Patients. 2022, 10, 12-17	
108	Increased Neutrophil elastase and proteinase 3 are closely associated with occurrence and severity of stroke and acute myocardial infarction in patients with type 2 diabetes mellitus.. 2022, 186, 109853	0
107	Circulating FAM19A5 level is associated with the presence and severity of coronary artery disease.. 2022,	
106	Nonalcoholic Fatty Liver Disease-Associated Liver Fibrosis Is Linked with the Severity of Coronary Artery Disease Mediated by Systemic Inflammation.. 2021, 2021, 6591784	2
105	Metabolic Impairment in Coronary Artery Disease: Elevated Serum Acylcarnitines Under the Spotlights.. 2021, 8, 792350	1
104	The gut microbiome and microbial metabolites in acute myocardial infarction.. 2021,	1
103	Serum Levels of lncRNA CCHE1 and TCF21 in Patients with Coronary Artery Disease and Their Clinical Significances.. 2021, 2021, 8526144	
102	Relations of physical signs to genotype, lipid and inflammatory markers, coronary stenosis or calcification, and outcomes in patients with heterozygous familial hypercholesterolemia. 2021, 19, 498	0
101	Apolipoprotein B/A1 Ratio Is Associated with Severity of Coronary Artery Stenosis in CAD Patients but Not in Non-CAD Patients Undergoing Percutaneous Coronary Intervention.. 2021, 2021, 8959019	1
100	Coronary ostial assessment of myeloperoxidase level correlated to severity of coronary artery disease. 2021, 63, 646-650	
99	Analysis of Coronary Artery Lesion Degree and Related Risk Factors in Patients with Coronary Heart Disease Based on Computer-Aided Diagnosis of Coronary Angiography.. 2021, 2021, 2370496	1

98	Interleukin-35 Levels in Patients with Stable Coronary Artery Disease.. 2022 , 118, 400-408	0
97	Association between fibrinogen-to-albumin ratio and the presence and severity of coronary artery disease in patients with acute coronary syndrome. 2021 , 21, 588	4
96	Association of Serum β -Hydroxybutyrate and Coronary Artery Disease in an Urban Chinese Population.. 2022 , 9, 828824	0
95	High Resting Heart Rate and High BMI Predicted Severe Coronary Atherosclerosis Burden in Patients With Stable Angina Pectoris by SYNTAX Score. 2018 , 69, 380-386	2
94	Relationship between Severity and Complexity of Coronary Artery Involvement and Obstructive Sleep Apnea Based on STOP-BANG Questionnaire.. 2022 , 13, 34	
93	Features of atherosclerosis in patients with angina and no obstructive coronary artery disease.. 2022 ,	0
92	The correlation between plasma human neutrophil peptide 1-3 levels and severity of coronary artery disease. 2016 , 1, 133-138	3
91	D-dimer to Creatinine Ratio: A Novel Biomarker Associated with Gensini Score in ST-Segment Elevation Myocardial Infarction Patients.. 2022 , 28, 10760296221099938	
90	Diabetic Retinopathy as a Predictor of Angiographic Coronary Atherosclerosis Severity in Patients with Type 2 Diabetes Mellitus.. 2022 , 15, 1485-1494	
89	Clinical value and expression of Homer 1, homocysteine, S-adenosyl-l-homocysteine, fibroblast growth factors 23 in coronary heart disease.. 2022 , 22, 215	0
88	Association of serum sclerostin and osteoprotegerin levels with the presence, severity and prognosis in patients with acute myocardial infarction.. 2022 , 22, 213	0
87	Exercise-induced change in circulating NT-proBNP could not distinguish between patients with and without coronary artery disease: the CADENCE study.. 2022 , 56, 107-113	
86	Sex differences in coronary artery disease and diabetes revealed by scRNA-Seq and CITE-Seq of human CD4+ T cells.	0
85	Lipoprotein(a) and Cardiovascular Outcomes in Patients With Coronary Artery Disease and Different Metabolic Phenotypes. 2022 , 9,	1
84	The Ratio of Red Blood Cell Distribution Width to Albumin Is Correlated With All-Cause Mortality of Patients After Percutaneous Coronary Intervention A Retrospective Cohort Study. 2022 , 9,	0
83	Serum levels of IL-37 and correlation with inflammatory cytokines and clinical outcomes in patients with coronary artery disease. jim-2021-002134	0
82	Heterogeneity in coronary heart disease risk. 2022 , 12,	0
81	Prognostic Value of Combining Apelin-12 and Estimated Glomerular Filtration Rate in Patients with ST-Segment Elevation Myocardial Infarction. 2022 , 2022, 1-9	

- 80 Long-term Prognostic Significance of Pentraxin-3 in Patients with Non-ST Elevation Myocardial Infarction and Coronary Stenting. **2022**, 10, 175-180
- 79 Prediabetes in acute coronary syndrome. **2022**, 11, 138
- 78 The Importance of Osmolarity in the Prognosis Prediction of ST-elevation and Depression in aVR Derivation of Patients with Acute Coronary Syndrome. **2022**, 10, 281-289
- 77 Assessment of Clinical Features in HIV-Infected Patients with Acute Coronary Syndromes Undergoing Percutaneous Coronary Intervention in China. **2022**, 2022, 1-8
- 76 Immunodominant MHC-II Restricted Epitopes in Human Apolipoprotein B. 0
- 75 MHR and NHR but not LHR were associated with coronary artery disease in patients with chest pain with controlled LDL-C. jim-2021-002314
- 74 Lipoprotein Subfractions as Markers for Predicting the Presence and Severity of Coronary Artery Disease in Patients Undergoing Coronary Angiography. 000331972211121
- 73 Association of High-Sensitivity Troponin T and I Blood Concentrations With All-Cause Mortality and Cardiovascular Outcome in Stable Patients Results From the INTERCATH Cohort. 2
- 72 Gensini score values for predicting periprocedural myocardial infarction: An observational study analysis. **2022**, 101, e29491
- 71 Association between Neutrophil to Lymphocyte Ratio and the Extent of Coronary Artery Disease in Patients with STEMI Versus Patients With Stable Angina Undergoing PCI: An Idea for Investigation of Other Inflammatory Diseases. **2021**, 9, 124-130
- 70 Carotid intima-media thickness predicted the presence but not the severity or complexity of coronary artery disease in a South Asian population. **2022**, 34, 183-192
- 69 Comparison of three non-insulin-based insulin resistance indexes in predicting the presence and severity of coronary artery disease. 9, 1
- 68 Predictors of Stent Restenosis in Han and Uygur Patients with Coronary Heart Disease after PCI in the Xinjiang Region. **2022**, 2022, 1-8
- 67 Triglyceride-glucose index in the prediction of adverse cardiovascular events in patients with premature coronary artery disease: a retrospective cohort study. **2022**, 21, 0
- 66 Long-term prognostic value of macrophage migration inhibitory factor in ST-segment elevation myocardial infarction patients with metabolic syndrome after percutaneous coronary intervention. 9,
- 65 Esheets in serum protein are independent risk factors for coronary lesions besides LDL-C in coronary heart disease patients. 9,
- 64 The prevalence and survival of pulmonary hypertension due to left heart failure: A retrospective analysis of a multicenter prospective cohort study. 9,
- 63 Inter-arm blood pressure difference as a tool for predicting coronary artery disease severity. **2022**, 9, e002063 1

62	Production of Reactive Oxygen Species by Epicardial Adipocytes Is Associated with an Increase in Postprandial Glycemia, Postprandial Insulin, and a Decrease in Serum Adiponectin in Patients with Severe Coronary Atherosclerosis. 2022 , 10, 2054	1
61	The constricting effect of reduced coronary artery compliance on the left ventricle is an important cause of reduced diastolic function in patients with coronary heart disease. 2022 , 22,	
60	The relationship between Vaspin, Nesfatin-1 plasma levels and presence of fragmented QRS with the severity of coronary atherosclerosis. 2022 , 67, 298-303	
59	Hypertension and clinical outcomes in patients with familial hypercholesterolemia. 2022 , 40, 1713-1721	
58	Association between circulating oxidized OxLDL/LDL-C ratio and the severity of coronary atherosclerosis, along with other emerging biomarkers of cardiovascular disease in patients with type 2 diabetes. 2022 , 191, 110040	0
57	Prediction of microvascular obstruction by coronary artery angiography score after acute ST-segment elevation myocardial infarction: a single-center retrospective observational study. 2022 , 22,	0
56	Application of an antibody microarray for serum protein profiling of coronary artery stenosis. 2022 , 631, 55-63	1
55	Visfatin is negatively associated with coronary artery lesions in subjects with impaired fasting glucose. 2022 , 17, 1405-1411	0
54	Ratio of transmitral early filling velocity to early diastolic strain rate predicts obstructive coronary stenosis. 2022 , 35, 371	0
53	A retrospective study on the relationship between fibrosis-4 index and all-cause mortality in patients with acute myocardial infarction. 2022 , 24,	0
52	Sex Differences in Coronary Artery Disease and Diabetes Revealed by scRNA-Seq and CITE-Seq of Human CD4+ T Cells. 2022 , 23, 9875	0
51	Clinical Characteristics Predicting Worse Long-Term Outcomes in Patients with Myocardial Infarction and Non-Obstructive Coronary Arteries (MINOCA). 2022 , 9, 286	0
50	Effects of gut microbiota and gut-derived lipopolysaccharide on patients with coronary heart disease.	0
49	Association between major adverse cardiovascular events and pentraxin-3 in patients who have undergone coronary computed tomography angiography: from the FU-CCTA registry.	0
48	Clinical characteristics and risk factors of in-hospital gastrointestinal bleeding in patients with acute myocardial infarction. 9,	0
47	Association of Nonalcoholic Fatty Liver Disease with Ventricular Tachycardia and Sinus Arrest in Patients with Non-ST-Segment Elevation Myocardial Infarction. 2022 , 63, 814-820	1
46	Nonlinear association of 1,5-anhydroglucitol with the prevalence and severity of coronary artery disease in chinese patients undergoing coronary angiography. 13,	0
45	Fibroblast growth factor 21 is associated with widening QRS complex and prolonged corrected QT interval in patients with stable angina. 2022 , 22,	0

- 44 The correlation of pericoronary adipose tissue with coronary artery disease and left ventricular function. **2022**, 22, ○
- 43 ANGPTL3 and Cardiovascular Outcomes in Patients With Acute Coronary Syndrome and Obstructive Sleep Apnea. **2022**, 11, ○
- 42 Circulating long non-coding RNA Coromarker expression correlated with inflammation, coronary artery stenosis, and plaque vulnerability in patients with coronary artery disease. ○
- 41 The relationship between soluble lymphocyte activation gene-3 and coronary artery disease. 9, ○
- 40 Features of Combined Cardiac and Cerebral Angiography in Patients with Cerebral Artery Stenosis and Coronary Artery Stenosis. **2022**, 12, 8907-8913 ○
- 39 Association of severity of coronary and carotid atherosclerosis and functional condition of a liver at patients with stable stenocardia and obesity. **2022**, 54-60 ○
- 38 Association of the severity of vascular damage with discordance between the fractional flow reserve and non-hyperemic pressure ratios. **2022**, ○
- 37 Circulating miR-143 and miR-145 as promising biomarkers for evaluating severity of coronary artery stenosis in patients with acute coronary syndrome. **2022**, ○
- 36 Positive Association of Plasma Trimethylamine-N-Oxide and Atherosclerosis in Patient with Acute Coronary Syndrome. **2022**, 2022, 1-9 2
- 35 Increased long non-coding RNA NORAD reflects serious cardiovascular stenosis, aggravated inflammation status, and higher lipid level in coronary heart disease. ○
- 34 Identification of circular RNA BTBD7_hsa_circ_0000563 as a novel biomarker for coronary artery disease and the functional discovery of BTBD7_hsa_circ_0000563 based on peripheral blood mononuclear cells: a case control study. **2022**, 19, 1
- 33 Association between neutrophil-to-lymphocyte ratio and severity of coronary artery disease. **2022**, 0 ○
- 32 Trajectories and determinants of left ventricular ejection fraction after the first myocardial infarction in the current era of primary coronary interventions. 9, ○
- 31 Small LDL subfractions are associated with coronary atherosclerosis despite no differences in conventional lipids. ○
- 30 Serum gamma glutamiltransferaz seviyeleri stabil koroner arter hastalarıda koroner kollateral geliiminin belirleyicisi olabilir mi?. ○
- 29 Blood group B and decreased risk of coronary artery disease in hypertensive elderly.. **2022**, 63, 388-399 ○
- 28 Data-independent acquisition proteomics reveals circulating biomarkers of coronary chronic total occlusion in humans. 9, ○
- 27 miR194 hypomethylation regulates coronary artery disease pathogenesis. **2022**, 15, ○

26 Association of Coronary Artery Severity and Late In-Stent Restenosis: An Angiographic Imaging Study. 000331972211509 ○

25 The association of the paraoxonase 1 Q192R polymorphism with coronary artery disease (CAD) and cardiometabolic risk factors in Iranian patients suspected of CAD. 9, ○

24 Association of Epicardial Adipose Tissue Adipocytes Hypertrophy with Biomarkers of Low-Grade Inflammation and Extracellular Matrix Remodeling in Patients with Coronary Artery Disease. **2023**, 11, 241 ○

23 Association of lipoprotein(a) with coronary severity in patients with new-onset acute myocardial infarction: A large cross-sectional study. **2023**, 540, 117220 ○

22 Prognostic value of gut microbiota-derived metabolites in patients with ST-segment elevation myocardial infarction. **2022**, ○

21 The association of glycated hemoglobin A1c with coronary artery disease, myocardial infarction, and severity of coronary lesions. 108155892211405 ○

20 Thyroid-stimulating hormone within the normal reference range has a U-shaped association with the severity of coronary artery disease in nondiabetic patients but is diluted in diabetic patients. 108155892211491 ○

19 CETP Taq1B polymorphism and its association with cardiovascular risk factors in an Iranian population under angiography, A Cross-Sectional Study. ○

18 Evaluation of cases with early repolarization on electrocardiogram and normal population in terms of laboratory and clinical results. **2023**, 6, 433-440 ○

17 Risk factors and inflammatory markers in acute coronary syndrome-ST elevation myocardial infarction (STEMI). **2022**, ○

16 The Caprini Risk Score for Early Prediction of Mortality in Patients With Acute Coronary Syndrome. Publish Ahead of Print, ○

15 Joint effects of paraoxonase 1 rs662 polymorphism and vitamins C/E intake on coronary artery disease severity (Gensini and SYNTAX scores) and lipid profile in patients undergoing coronary angiography. 9, ○

14 Circulating Total Bilirubin and Long-Term Prognosis in Patients With Previous Myocardial Infarction. **2023**, 3, 242-251 ○

13 Coronary ectasia in different scenarios, primarily in myocardial infarction with nonobstructive coronary artery disease. **2023**, 24, 167-171 ○

12 Alterations of NMR-Based Lipoprotein Profile Distinguish Unstable Angina Patients with Different Severity of Coronary Lesions. **2023**, 13, 273 ○

11 Proteomic profiling of epicardial fat in heart failure with preserved versus reduced and mildly reduced ejection fraction. **2023**, 27, 727-735 ○

10 Sex Differences in Vascular Response to Mental Stress and Adverse Cardiovascular Events Among Patients With Ischemic Heart Disease. **2023**, 43, ○

9 HbA1c is a predictive factor of severe coronary stenosis and major adverse cardiovascular events in patients with both type 2 diabetes and coronary heart disease. **2023**, 15, ○

- 8 Earlobe Creases as a Marker of the Risk for Coronary Atherosclerosis Before Angiography in Elderly and Non-elderly Patients. **2023**, ○
- 7 Quantitation of diffuse myocardial ischemia with mental stress and its association with cardiovascular events in individuals with recent myocardial infarction. ○
- 6 Correlation of elevated serum uric acid with coronary artery disease in Xinjiang, China: A retrospective case-control study. **2023**, 102, e33256 ○
- 5 Chinese herbal prescription Qing-Xin-Jie-Yu granule combined with conventional treatment for intermediate coronary lesions: study protocol for a randomized, double-blinded, placebo-controlled trial. ○
- 4 Association Between Plasma Ceramides and One-Year Mortality in Patients with Acute Coronary Syndrome: Insight from the PEACP Study. Volume 18, 571-584 ○
- 3 The association of serum uric acid/albumin ratio with the development of coronary collateral circulation in patients with chronic total occluded coronary arteries. **2023**, 15, 14-21 ○
- 2 Long- and very long-chain ceramides are predictors of acute kidney injury in patients with acute coronary syndrome: the PEACP study. **2023**, 22, ○
- 1 Role of combined use of mean platelet volume-to-lymphocyte ratio and monocyte to high-density lipoprotein cholesterol ratio in predicting patients with acute myocardial infarction. **2023**, 18, ○