
CITATION REPORT List of articles citing



DOI: 10.1016/s0735-1097(83)80100-4 Journal of the American College of Cardiology, 1983, 1, 994-10

Source: https://exaly.com/paper-pdf/16409559/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
317	Radionuclide Exercise Testing for Coronary Artery Disease. 1984 , 2, 367-378		6
316	One-Vessel Coronary Artery Disease. 1984 , 144, 1233		3
315	Diagnostic methods in the assessment of the patient with chest pain. <i>Journal of the American College of Cardiology</i> , 1984 , 4, 290-1	15.1	4
314	Prognostic implications of rest and exercise radionuclide ventriculography in patients with suspected or proven coronary heart disease. 1984 , 6, 707-18		26
313	Prevalence of high-risk thallium-201 scintigraphic findings in left main coronary artery stenosis: comparison with patients with multiple- and single-vessel coronary artery disease. 1984 , 53, 462-9		88
312	Prognostic indicators from radionuclide angiography in medically treated patients with coronary artery disease. 1984 , 53, 18-22		147
311	Thallium-201 myocardial perfusion imaging without exercise. Rest and dipyridamole studies. 1984 , 86, 159-61		13
310	Angina pectoris. 1984 , 311, 1121-3		
309	Left ventricular dilatation. Prognostic value in severe left ventricular dysfunction secondary to coronary artery disease. 1985 , 88, 215-20		20
308	Nuclear cardiology: current indications and clinical usefulness. 1985 , 10, 1-76		15
307	Determination of cardiac risk by dipyridamole-thallium imaging before peripheral vascular surgery. 1985 , 312, 389-94		622
306	Detection of high risk coronary artery disease by thallium imaging. <i>Heart</i> , 1985 , 53, 616-23	5.1	15
305	Prognostic significance of normal quantitative planar thallium-201 stress scintigraphy in patients with chest pain. <i>Journal of the American College of Cardiology</i> , 1985 , 6, 27-30	15.1	122
304	Selection of patients for preoperative coronary angiography: Use of dipyridamole-stressEhallium myocardial imaging. 1985 , 2, 504-510		
303	The persistent defect on exercise thallium imaging and its fate after myocardial revascularization: does it represent scar or ischemia?. 1985 , 110, 996-1001		168
302	Thallium-201 myocardial scintigraphy. 1985 , 109, 113-29		58
301	The use of rest and exercise radionuclide ventriculography in risk stratification in patients with suspected coronary artery disease. 1985 , 110, 864-72		50

(1986-1985)

300	coronary artery disease. 1985 , 110, 135-43	115	
299	Accuracy of left ventricular ejection fraction using the nuclear stethoscope in left ventricular aneurysm. 1985 , 55, 177-80	10	
298	Prognosis with chest pain and normal thallium-201 exercise scintigrams. 1985 , 55, 920-6	127	
297	Comparison of exercise electrocardiography and quantitative thallium imaging for one-vessel coronary artery disease. 1985 , 56, 257-61	39	
296	Comparison of stress digital ventriculography, stress thallium scintigraphy, and digital fluoroscopy in the diagnosis of coronary artery disease in subjects without prior myocardial infarction. 1985 , 56, 434-40	26	
295	Increased thallium lung uptake after exercise in isolated left anterior descending coronary artery disease. 1985 , 55, 1469-73	35	
294	Prognostic significance of regional myocardial ischemia of the ventricular septum assessed by thallium-201 exercise testing. 1986 , 58, 359-60	7	
293	Cardiac cinefluoroscopy as an inexpensive aid in the diagnosis of coronary artery disease. 1986 , 57, 1041-6	45	
292	Assessment of regional myocardial perfusion with thallium imaging during transient left anterior descending coronary arterial occlusion during angioplasty. 1986 , 57, 1083-7	8	
291	Monkey business. 1986 , 57, 471-5	59	
291 290	Monkey business. 1986, 57, 471-5 Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986, 57, 503-8	59 62	
	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the		
290	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986 , 57, 503-8 Quantitative thallium imaging findings in patients with normal coronary angiographic findings and	62 54	
290 289	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986 , 57, 503-8 Quantitative thallium imaging findings in patients with normal coronary angiographic findings and in clinically normal subjects. 1986 , 57, 509-12 Thallium-201 myocardial imaging during pharmacologic coronary vasodilation: comparison of oral and intravenous administration of dipyridamole. <i>Journal of the American College of Cardiology</i> , 1986	62 54 1 94	
290 289 288	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986, 57, 503-8 Quantitative thallium imaging findings in patients with normal coronary angiographic findings and in clinically normal subjects. 1986, 57, 509-12 Thallium-201 myocardial imaging during pharmacologic coronary vasodilation: comparison of oral and intravenous administration of dipyridamole. <i>Journal of the American College of Cardiology</i> , 1986, 8, 76-83 Guidelines for clinical use of cardiac radionuclide imaging, December 1986. <i>Journal of the American</i>	62 54 1 94	
290 289 288 287	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986, 57, 503-8 Quantitative thallium imaging findings in patients with normal coronary angiographic findings and in clinically normal subjects. 1986, 57, 509-12 Thallium-201 myocardial imaging during pharmacologic coronary vasodilation: comparison of oral and intravenous administration of dipyridamole. <i>Journal of the American College of Cardiology</i> , 1986, 8, 76-83 Guidelines for clinical use of cardiac radionuclide imaging, December 1986. <i>Journal of the American College of Cardiology</i> , 1986, 8, 1471-1483 Extent and severity of myocardial hypoperfusion as predictors of prognosis in patients with	62 54 1 94 1 33 1 359	
290 289 288 287 286	Usefulness of oral dipyridamole suspension for stress thallium imaging without exercise in the detection of coronary artery disease. 1986, 57, 503-8 Quantitative thallium imaging findings in patients with normal coronary angiographic findings and in clinically normal subjects. 1986, 57, 509-12 Thallium-201 myocardial imaging during pharmacologic coronary vasodilation: comparison of oral and intravenous administration of dipyridamole. <i>Journal of the American College of Cardiology</i> , 1986, 8, 76-83 Guidelines for clinical use of cardiac radionuclide imaging, December 1986. <i>Journal of the American College of Cardiology</i> , 1986, 8, 1471-1483 Extent and severity of myocardial hypoperfusion as predictors of prognosis in patients with suspected coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1986, 7, 464-71 Determination of the quantitative thallium imaging variables that optimize detection of coronary	62 54 1 94 1 33 1 359	

282	Patients with suspected Lassa fever in London during 1984. 1986 , 292, 58-58		1
281	Patients with suspected Lassa fever in London during 1984. 1986 , 292, 57-8		1
280	Patients with suspected Lassa fever in London during 1984. 1986 , 292, 58-58		
279	Simultaneous measurement of left ventricular function and myocardial perfusion during a single exercise test: dual isotope imaging with gold-195 m and thallium-201. 1986 , 7, 493-500		5
278	The prevalence and clinical significance of residual myocardial ischemia 2 weeks after uncomplicated non-Q wave infarction: a prospective natural history study. 1986 , 73, 1186-98		220
277	Exercise Thallium Imaging in Patients With Diabetes Mellitus. 1987 , 147, 313		32
276	Nuclear magnetic resonance imaging: current applications and future prospects. 1987 , 75, 88-95		23
275	Lung uptake of thallium as a prognostic indicator. 1987 , 317, 1532-4		9
274	Thallium-201 stress imaging in hypertensive patients. 1987 , 10, 16-21		53
273	Prognostic importance of thallium uptake by the lungs during exercise in coronary artery disease. 1987 , 317, 1485-9		218
272	The efficacy of cardiovascular nuclear medicine exercise studies. 1987, 17, 104-20		55
271	The role of radionuclide cardiac imaging in coronary artery bypass surgery. 1987 , 113, 163-70		3
270	Computer versus visual analysis of exercise thallium-201 images: a critical appraisal in 325 patients with chest pain. 1987 , 114, 1129-37		17
269	Silent myocardial ischemia: II. Prognosis and implications for the clinical assessment of patients with coronary artery disease. 1987 , 114, 627-38		31
268	Prognostic value of computer-quantitated exercise thallium imaging early after percutaneous transluminal coronary angioplasty. <i>Journal of the American College of Cardiology</i> , 1987 , 10, 275-83	15.1	41
267	Reply. Journal of the American College of Cardiology, 1987 , 10, 1365-1366	15.1	
266	Quantitative analysis of dipyridamole-thallium images for the detection of coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1987 , 10, 142-9	15.1	43
265	Limited Prognostic Value of Early Rest Thallium-201 Imaging in Patients Admitted for Suspected Acute Myocardial Infarction. 1987 , 1, 129-133		2

264	Prognostic Implications of Normal Exercise Thallium Images. 1987 , 1, 209-212	30
263	Angina pectoris: diagnosis, treatment, and prognosis. 1987 , 12, 213-81	1
262	Role of nuclear cardiology in evaluating the total ischemic burden in coronary artery disease. 1987 , 59, 31C-38C	8
261	Incremental prognostic power of clinical history, exercise electrocardiography and myocardial perfusion scintigraphy in suspected coronary artery disease. 1987 , 59, 270-7	130
260	Usefulness of residual ischemic myocardium within prior infarct zone for identifying patients at high risk late after acute myocardial infarction. 1987 , 60, 15-9	118
259	Role of exercise thallium-201 myocardial perfusion scintigraphy in predicting prognosis in suspected coronary artery disease. 1987 , 59, 531-4	53
258	Risk stratification in stable angina pectoris. 1987 , 59, 695-7	10
257	Sensitivity, specificity, and prognostic significance of noninvasive testing for occult or known coronary disease. 1987 , 29, 241-70	56
256	Exercise thallium scans: patterns of use and impact on management of patients with known or suspected coronary artery disease. 1987 , 59, 50-5	28
255	Myocardial scintigraphy25 years after start. 1988 , 13, 619-36	14
254	Myocardial perfusion imaging with 99mTc-methoxy-isobutyl-isonitrile (MIBI): comparison of short and long time intervals between rest and stress injections. Preliminary results. 1988 , 13, 515-22	57
253	Usefulness of exercise electrocardiography and thallium scintigraphy in unstable angina pectoris in predicting the extent and severity of coronary artery disease. 1988 , 62, 1164-70	27
252	Comparative accuracy of clinical tests for diagnosis and prognosis of coronary artery disease. 1988 , 62, 896-900	17
251	Use of exercise thallium-201 imaging for risk stratification of elderly patients with coronary artery disease. 1988 , 61, 269-72	66
250	The clinical and pathophysiologic implications of pain, ST abnormalities, and scintigraphic changes induced during dipyridamole infusion: their relationships to the peripheral hemodynamic response. 1988 , 116, 1071-80	32
249	Factors affecting sensitivity and specificity of a diagnostic test: the exercise thallium scintigram. 1988 , 84, 699-710	103
248	Exercise echocardiography: ready or not?. <i>Journal of the American College of Cardiology</i> , 1988 , 11, 1355-& _{5.1}	31
247	Superiority of quantitative exercise thallium-201 variables in determining long-term prognosis in ambulatory patients with chest pain: a comparison with cardiac catheterization. <i>Journal of the</i> 15.1 American College of Cardiology, 1988 , 12, 25-34	146

246	Thallium-201 scintigraphy in risk assessment for ambulatory patients with chest pain: does everyone need catheterization?. <i>Journal of the American College of Cardiology</i> , 1988 , 12, 35-6	15.1	4
245	Prognostic value and limitations of exercise radionuclide angiography in medically treated coronary artery disease. 1988 , 63, 573-82		20
244	DOBUTAMINE STRESS TEST. 1988 , 332, 1347-1348		2
243	Prognostic utility of the exercise thallium-201 test in ambulatory patients with chest pain: comparison with cardiac catheterization. 1988 , 77, 745-58		146
242	Methodologic Problems in Exercise Testing Research. 1988 , 148, 1289		13
241	Prognosis and thallium-201 scintigraphy in patients admitted with chest pain without confirmed acute myocardial infarction. <i>Heart</i> , 1988 , 59, 184-9	5.1	15
240	Cine NMR: An Emerging Technology. 1988, 5, 373-382		5
239	The clinical use of nuclear exercise tests. 1988 , 4, 613-22		2
238	Noninvasive evaluation of angina pectoris. 1988 , 23, 95-101, 105-7, 111-5		2
237	After the Myocardial Infarction: A Review and Approach to Risk Stratification. 1988 , 6, 153-163		
236	Non-invasive data provide independent prognostic information in patients with chest pain without previous myocardial infarction: findings in male patients who have had cardiac catheterization. 1988 , 9, 418-26		24
235	Exercise thallium-201 scintigraphy and prognosis in typical angina pectoris and negative exercise electrocardiography. 1989 , 64, 282-7		40
234	Prognostic utility of predischarge dipyridamole-thallium imaging compared to predischarge submaximal exercise electrocardiography and maximal exercise thallium imaging after uncomplicated acute myocardial infarction. 1989 , 64, 1243-8		69
233	Noninvasive cardiac risk stratification of diabetic and nondiabetic uremic renal allograft candidates using dipyridamole-thallium-201 imaging and radionuclide ventriculography. 1989 , 64, 1017-21		71
232	Comparison of myocardial washout rate of thallium-201 between rest, dipyridamole with and without aminophylline, and exercise states in normal subjects. 1989 , 64, 1022-8		12
231	A look at 15 years of planar thallium-201 imaging. 1989 , 118, 581-601		41
230	Prognostic importance of silent myocardial ischemia detected by intravenous dipyridamole thallium myocardial imaging in asymptomatic patients with coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 1635-41	15.1	90
229	Dipyridamole thallium-201 scintigraphy: an excellent alternative to exercise scintigraphy. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 1642-4	15.1	26

[1990-1989]

228	Technetium-99m isonitrile myocardial uptake at rest. I. Relation to severity of coronary artery stenosis. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 1673-7	41
227	Importance of epidemiology and biostatistics in deciding clinical strategies for using diagnostic tests: a simplified approach using examples from coronary artery disease. <i>Journal of the American</i> 15.1 <i>College of Cardiology</i> , 1989 , 13, 1653-65	70
226	Task Force I: Determination of prognosis in patients with ischemic heart disease. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 1016-25	24
225	Comparison of planar and tomographic exercise thallium-201 imaging methods for the evaluation of coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1989 , 13, 613-6	33
224	Perfusion imaging. 1989 , 45, 896-921	1
223	Cardiac evaluation of the patient with stroke. 1990 , 21, 14-23	49
222	Prognostic value of radionuclide angiography in medically treated patients with coronary artery disease. A comparison with clinical and catheterization variables. 1990 , 82, 1705-17	161
221	Ability of dipyridamole-thallium-201 imaging one to four days after acute myocardial infarction to predict in-hospital and late recurrent myocardial ischemic events. 1990 , 65, 160-7	98
220	Comparison of thallium redistribution with rest "reinjection" imaging for the detection of viable myocardium. 1990 , 66, 158-63	132
219	Prevalence and correlates of increased lung/heart ratio of thallium-201 during dipyridamole stress imaging for suspected coronary artery disease. 1990 , 66, 1324-8	29
218	Assessment of post-infarction jeopardized myocardium by vasodilationthallium-201 tomography: impact on risk stratification. 1990 , 11, 1093-100	2
217	The value of thallium-201 imaging. 1990 , 323, 190-2	8
216	Prevalence and prognostic significance of exercise-induced silent myocardial ischemia detected by thallium scintigraphy and electrocardiography in asymptomatic volunteers. 1990 , 81, 428-36	218
215	Altered myocardial perfusion in patients with angina pectoris or silent ischemia during exercise as assessed by quantitative thallium-201 single-photon emission computed tomography. 1990 , 82, 1305-15	48
214	Relations of the myocardial imaging agents 99mTc-MIBI and 201T1 to myocardial blood flow in a canine model of myocardial ischemic insult. 1990 , 81, 289-96	53
213	Additional prognostic value of exercise testing and thallium-201 scintigraphy in catheterized patients without previous myocardial infarction. 1990 , 27, 235-43	3
212	Prognostic value of a normal exercise echocardiogram. 1990 , 120, 49-55	144
211	Appraisal of clinical models based on results of stress nuclear imaging in risk stratification. 1990 , 120, 1487-90	6

210	Prognostic value of dipyridamole thallium scintigraphy for evaluation of ischemic heart disease. <i>Journal of the American College of Cardiology</i> , 1990 , 15, 109-16	15.1	122
209	The variable extent of jeopardized myocardium in patients with single vessel coronary artery disease: quantification by thallium-201 single photon emission computed tomography. <i>Journal of the American College of Cardiology</i> , 1991 , 17, 355-62	15.1	82
208	Thallium scintigraphy for diagnosis and risk assessment of coronary artery disease. 1991 , 338, 786-788		4
207	Ribose infusion accelerates thallium redistribution with early imaging compared with late 24-hour imaging without ribose. <i>Journal of the American College of Cardiology</i> , 1991 , 18, 1671-81	15.1	26
206	Dobutamine thallium myocardial perfusion tomography. <i>Journal of the American College of Cardiology</i> , 1991 , 18, 1471-9	15.1	128
205	Stress Thallium Imaging. 1991 , 5, 173-184		6
204	Prognostic value of 201Tl myocardial scintigraphy 1991 , 84, 2203-2204		
203	Prediction of death and myocardial infarction by radionuclide angiocardiography in patients with suspected coronary artery disease. 1991 , 67, 919-26		20
202	Silent versus symptomatic ischemia during a thallium-201 exercise test. 1991 , 68, 1600-8		17
201	Current status of nuclear cardiology techniques. 1991 , 16, 451-535		7
200	Exercise thallium-201 perfusion scintigraphy in the assessment of coronary artery disease. 1991 , 67, 2D-11D		66
199	Comparison of exercise performance in left main and three-vessel coronary artery disease. 1991 , 22, 14-20		7
198	Technetium 99m SESTAMIBI myocardial perfusion imaging: comparison between treadmill, dipyridamole and trans-oesophageal atrial pacing "stress" tests in normal subjects. 1991 , 18, 247-51		18
197	Prognostic value of thallium-201 myocardial perfusion imaging. A diagnostic tool comes of age. 1991 , 83, 363-81		305
196	The relationship between intermittent claudication and coronary artery disease-is it more than wethink?. 1991 , vmr-2, 137-146		8
195	Nuclear medicine. 1992 , 68, 82-105		2
194	Independent and incremental prognostic value of tests performed in hierarchical order to evaluate patients with suspected coronary artery disease. Validation of models based on these tests. 1992 , 85, 237-48		116
193	Viable myocardium and reinjection of thallium. <i>Heart</i> , 1992 , 68, 537-9	5.1	2

192	Coronary artery disease and cardiac events with asymptomatic and symptomatic cerebrovascular disease. 1992 , 23, 939-45		67
191	Cardiac imaging after acute myocardial infarction. Identification of patients at continued risk. 1992 , 92, 93-6, 99-100		
190	Prognostic significance of silent exertional myocardial ischaemia in symptomatic men without previous myocardial infarction. 1992 , 13, 183-7		13
189	Prognostic value of dipyridamole thallium-201 imaging in elderly patients. <i>Journal of the American College of Cardiology</i> , 1992 , 19, 1390-8	15.1	64
188	Safety and diagnostic accuracy of adenosine thallium-201 scintigraphy in patients unable to exercise and those with left bundle branch block. 1992 , 124, 614-21		48
187	Influence of exercise intensity on the presence, distribution, and size of thallium-201 defects. 1992 , 123, 909-16		54
186	Risk stratification in coronary disease: a contrary viewpoint. 1992 , 116, 927-36		17
185	Long-Term Prognostic Value of Thallium Scintigraphy and Coronary Angiography after Myocardial Infarction in Young Men. 1992 , 6, 291-303		
184	Prognostic value of thallium-201 myocardial perfusion imaging in three primary patient populations. 1992 , 70, 23E-29E		6
183	Technetium-99m sestamibi myocardial imaging at rest for assessment of myocardial infarction and first-pass ejection fraction. Multicenter Cardiolite Study Group. 1992 , 69, 22-7		25
182	A new method for evaluating lung uptake of thallium-201 during stress myocardial scintigraphy. 1992 , 70, 940-4		3
181	Prevalence and prognostic significance of exercise-induced silent myocardial ischemia in apparently healthy subjects. 1992 , 69, 14B-18B		14
180	Noninvasive identification of severe coronary artery disease using exercise tomographic thallium-201 imaging. 1992 , 70, 14-20		169
179	Prognostic implications of transient asymptomatic myocardial ischemia as detected by ambulatory electrocardiographic monitoring. 1992 , 35, 77-96		4
178	Patients with a normal exercise thallium-201 myocardial scintigram: always a good prognosis?. 1993 , 20, 151-8		9
177	Prognostic significance from 10-year follow-up of a qualitatively normal planar exercise thallium test in suspected coronary artery disease. 1993 , 71, 1270-3		38
176	Comparison of myocardial perfusion imaging with technetium-99m tetrofosmin versus thallium-201 in coronary artery disease. 1993 , 72, 1015-9		78
175	Prognosis of patients with an isolated fixed thallium-201 defect and no prior myocardial infarction. 1993 , 72, 1199-201		6

174	Prognostic implications of normal exercise SPECT thallium images in patients with strongly positive exercise electrocardiograms. 1993 , 72, 1201-3		33
173	Minimal residual coronary obstructions in patients who suffered a first myocardial infarction. A prospective study comparing coronary angiography and exercise thallium scintigraphy. 1993 , 16, 879-8	2	4
172	Sequential internal mammary artery grafts for coronary artery bypass. 1993 , 56, 1136-40		16
171	Variables associated with a poor prognosis in patients with an ischemic thallium-201 exercise test. 1993 , 125, 335-44		34
170	Nuclear cardiology (1). 1993 , 329, 775-83		113
169	Independent and incremental prognostic value of exercise single-photon emission computed tomographic (SPECT) thallium imaging in coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1993 , 22, 665-70	15.1	294
168	Factors affecting uniformity in interpretation of planar thallium-201 imaging in a multicenter trial. The Multicenter Study on Silent Myocardial Ischemia (MSSMI) Thallium-201 Investigators. <i>Journal of the American College of Cardiology</i> , 1993 , 21, 1064-74	15.1	47
167	Dobutamine thallium-201 tomography for evaluating patients with suspected coronary artery disease unable to undergo exercise or vasodilator pharmacologic stress testing. <i>Journal of the American College of Cardiology</i> , 1993 , 21, 1583-90	15.1	129
166	Prognostic value of an increase in fluorine-18 deoxyglucose uptake in patients with myocardial infarction: comparison with stress thallium imaging. <i>Journal of the American College of Cardiology</i> , 1993 , 22, 1621-7	15.1	114
165	Quantitative adenosine 201Tl single-photon emission computed tomography for the early assessment of patients surviving acute myocardial infarction. 1993 , 87, 1197-210		70
164	Coronary artery disease after heart transplantation: non-invasive evaluation with exercise thallium scintigraphy. 1993 , 14, 226-9		47
163	Exercise Thallium-201 Imaging Characteristics and Prognostic Value in Patients with Left Bundle Branch Block. 1994 , 8, 58-62		
162	Prognostic Value of Left Ventricular Cavity Dilatation in Patients with Increased Thallium Lung Uptake during Exercise Thallium-201 Testing. 1994 , 8, 381-385		
161	Cardiac stress testing with thallium-201 imaging reveals silent ischemia in individuals with paraplegia. 1994 , 75, 946-950		74
160	Myocardial Perfusion Imaging During Pharmacologic Stress Testing. 1994 , 12, 223-245		36
159	Prognosis of Acute and Chronic Coronary Artery Disease by Myocardial Perfusion Imaging. 1994 , 12, 271-287		14
158	Exercise technetium-99m sestamibi tomography for cardiac risk stratification of patients with stable chest pain. 1994 , 89, 615-22		119
157	Prognostic value of myocardial hypoperfusion indexes in patients with suspected or known coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 1994 , 1, 325-37	2.1	5

156	Prognostic value of adenosine single-photon emission computed tomographic thallium imaging in medically treated patients with angiographic evidence of coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 1994 , 1, 254-61	2.1	33
155	Myocardial technetium 99m-labeled sestamibi single-photon emission computed tomographic imaging in the detection of coronary artery disease: comparison between early (15 minutes) and delayed (60 minutes) imaging. <i>Journal of Nuclear Cardiology</i> , 1994 , 1, 441-8	2.1	12
154	Myocardial single-photon emission computed tomographic imaging with technetium 99m tetrofosmin: stress-rest imaging with same-day and separate-day rest imaging. <i>Journal of Nuclear Cardiology</i> , 1994 , 1, 138-43	2.1	25
153	Correlation of resting electrocardiographic ST-T wave abnormalities with exercise thallium-201 stress testing in patients with known or suspected coronary artery disease. 1994 , 74, 211-5		6
152	Prognostic significance of a fixed thallium defect one to six months after onset of acute myocardial infarction or unstable angina. Multicenter Myocardial Ischemia Research Group. 1994 , 74, 1196-200		17
151	Prognostic value of thallium-201 single-photon emission computed tomographic myocardial perfusion imaging according to extent of myocardial defect. Study in 1,926 patients with follow-up at 33 months. <i>Journal of the American College of Cardiology</i> , 1994 , 23, 1096-106	15.1	159
150	Does myocardial perfusion scintigraphy demonstrate clinical usefulness in patients with markedly positive exercise tests? An assessment of the method in a high-risk subset. 1994 , 127, 805-816		12
149	Exercise tomographic thallium-201 imaging in patients with severe coronary artery disease and normal electrocardiograms. 1994 , 121, 825-32		65
148	Diagnostic and prognostic value of dipyridamole echocardiography in patients with suspected coronary artery disease. Comparison with exercise electrocardiography. 1994 , 89, 1160-73		150
147	Cardiac prognosis of patients with carotid stenosis and no history of coronary artery disease. Veterans Affairs Cooperative Study Group 167. 1994 , 25, 759-65		79
146	Stress imaging. Current clinical options for the diagnosis, localization, and evaluation of coronary artery disease. 1995 , 79, 1025-61		6
145	Enhanced hepatic uptake of thallium-201 in patients with severe narrowing of the right coronary artery. 1995 , 76, 1020-4		1
144	Postexercise systolic blood pressure response: association with the presence and extent of perfusion abnormalities on thallium-201 scintigraphy. 1995 , 129, 227-34		15
143	Incremental value of prognostic testing in patients with known or suspected ischemic heart disease: a basis for optimal utilization of exercise technetium-99m sestamibi myocardial perfusion single-photon emission computed tomography. <i>Journal of the American College of Cardiology</i> , 1995 ,	15.1	457
142	Long-term prediction of major ischemic events by exercise thallium-201 single-photon emission computed tomography. Incremental prognostic value compared with clinical, exercise testing, catheterization and radionuclide angiographic data. <i>Journal of the American College of Cardiology</i> ,	15.1	84
141	Independent prognostic value of intravenous dipyridamole with technetium-99m sestamibi tomographic imaging in predicting cardiac events and cardiac-related hospital admissions. <i>Journal of the American College of Cardiology</i> , 1995 , 26, 1202-8	15.1	79
140	Guidelines for clinical use of cardiac radionuclide imaging. Report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Committee on Radionuclide Imaging), developed in collaboration with the American Society of Nuclear Cardiology, Journal of the American College of Cardiology, 1995, 25, 56	15.1 21-47	196
139	Reply. Journal of the American College of Cardiology, 1995 , 26, 1560-1562	15.1	1

138	Identification of severe and extensive coronary artery disease by automatic measurement of transient ischemic dilation of the left ventricle in dual-isotope myocardial perfusion SPECT. <i>Journal of the American College of Cardiology</i> , 1996 , 27, 1612-20	15.1	195
137	Prognostic value of dobutamine-atropine stress technetium-99m sestamibi perfusion scintigraphy in patients with chest pain. <i>Journal of the American College of Cardiology</i> , 1996 , 28, 447-54	15.1	35
136	Predictors of prognosis by quantitative assessment of coronary angiography, single photon emission computed tomography thallium imaging, and treadmill exercise testing. 1996 , 131, 582-90		8
135	Prognostic value of iodine-123 labelled BMIPP fatty acid analogue imaging in patients with myocardial infarction. 1996 , 23, 272-9		41
134	Prognostic value of stress dobutamine technetium-99m sestamibi single-photon emission computed tomography (SPECT) in patients with suspected coronary artery disease. 1996 , 78, 1092-1096	6	17
133	Relation between perfusion defects on stress technetium-99m sestamibi SPECT scintigraphy and the location of a subsequent acute myocardial infarction. 1996 , 78, 26-30		9
132	Clinical and angiographic significance of a normal thallium-201 tomographic study in patients with a strongly positive exercise electrocardiogram. 1996 , 78, 638-41		7
131	The site of acute myocardial infarction is related to the coronary territory of transient defects on prior myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, 382-8	2.1	8
130	Risk assessment in patients with stable coronary artery disease: incremental value of nuclear imaging. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, S41-9	2.1	16
129	Myocardial perfusion stress protocols. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, G11-G15	2.1	3
128	American Society of Nuclear Cardiology project on myocardial perfusion imaging: measuring outcomes in response to emerging guidelines. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, 436-42	2.1	13
127	Prognostic utility of increased pulmonary thallium uptake in patients without ischemia. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, 301-7	2.1	6
126	Comparative diagnostic accuracy of Tl-201 and Tc-99m sestamibi SPECT imaging (perfusion and ECG-gated SPECT) in detecting coronary artery disease in women. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 69-77	15.1	316
125	Importance of estimated functional capacity as a predictor of all-cause mortality among patients referred for exercise thallium single-photon emission computed tomography: report of 3,400 patients from a single center. <i>Journal of the American College of Cardiology</i> , 1997 , 30, 641-8	15.1	151
124	Evaluation of exercise thallium scintigraphy versus exercise electrocardiography in predicting survival outcomes and morbid cardiac events in patients with single- and double-vessel disease. Findings from the Angioplasty Compared to Medicine (ACME) Study. <i>Journal of the American</i>	15.1	59
123	College of Cardiology, 1997, 30, 1256-63 GanzkEperverteilung von Tc-99m-Sestamibi nach pharmakologischer Belastung mit Arbutamin und Dipyridamol im Vergleich zum Ruhezustand. 1997, 36, 19-22		1
122	Prognostic value of dobutamine stress echocardiography in patients undergoing diagnostic coronary arteriography. 1997 , 79, 1610-4		18
121	Incremental value of rubidium-82 positron emission tomography for prognostic assessment of known or suspected coronary artery disease. 1997 , 80, 865-70		109

120	Prognostic value of persistent thallium-201 defects that become reversible after reinjection in patients with chronic myocardial infarction. <i>Journal of Nuclear Cardiology</i> , 1997 , 4, 195-201	2.1	4
119	Prognosis in patients with left bundle branch block and normal dipyridamole thallium-201 scintigraphy. <i>Journal of Nuclear Cardiology</i> , 1998 , 5, 414-7	2.1	17
118	Prognostic value of stress echocardiography in the evaluation of atypical chest pain patients without known coronary artery disease. 1998 , 81, 545-51		34
117	Predictive value of planar 18F-fluorodeoxyglucose imaging for cardiac events in patients after acute myocardial infarction. 1998 , 81, 1072-7		26
116	The usefulness of positron emission tomography. 1998 , 23, 69-120		12
115	Prognostic value of exercise thallium-201 imaging performed within 2 years of coronary artery bypass graft surgery. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 848-54	15.1	50
114	Thallium reinjection versus standard stress/delay redistribution imaging for prediction of cardiac events. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 1280-5	15.1	14
113	Functional assessment of coronary stenoses. <i>Journal of the American College of Cardiology</i> , 1998 , 32, 1134-5	15.1	1
112	Correspondence. Journal of the American College of Cardiology, 1998 , 32, 1134	15.1	3
111	Correspondence. Journal of the American College of Cardiology, 1998 , 32, 1133-1134	15.1	1
110	Prognostic value of exercise thallium-201 imaging in a community population. 1998 , 135, 663-70		11
109	Long-term prognostic value of exercise echocardiography compared with exercise 201Tl, ECG, and clinical variables in patients evaluated for coronary artery disease. 1998 , 98, 2679-86		121
108	Investigation and management of stable angina: revised guidelines 1998. <i>Heart</i> , 1999 , 81, 546-555	5.1	20
107	Long-term additive prognostic value of thallium-201 myocardial perfusion imaging over clinical and exercise stress test in low to intermediate risk patients: study in 1137 patients with 6-year follow-up. 1999 , 100, 1521-7		142
106	Wintergreen panel summaries. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 93-155	2.1	21
105	Diagnostic and Prognostic Use of Stress Echocardiography and Radionuclide Scintigraphy. 1999 , 16, 85	7-877	1
104	GuBs de actuaciB clBica de la Sociedad EspaBla de CardiologB. CardiologB nuclear: bases tBnicas y aplicaciones clBicas. 1999 , 52, 957-989		8
103	ACC/AHA/ACP-ASIM guidelines for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Patients With Chronic Stable Angina). <i>Journal of the</i>	15.1	482

102	50th anniversary historical article. Perfusion imaging. <i>Journal of the American College of Cardiology</i> , 1999 , 34, 9-11	5.1	4
101	Diagnostic and prognostic value of myocardial perfusion imaging in patients with known or suspected stable coronary artery disease. 2000 , 17, 587-95		4
100	Stress nuclear myocardial perfusion imaging versus stress echocardiography: prognostic comparisons. 2000 , 43, 231-44		5
99	An overview of radiotracers in nuclear cardiology. <i>Journal of Nuclear Cardiology</i> , 2000 , 7, 701-7 2	.1	20
98	Contributions of nuclear cardiology to diagnosis and prognosis of patients with coronary artery disease. 2000 , 101, 1465-78		291
97	[Nuclear cardiology: technical bases and clinical applications]. 2000 , 19, 29-64		O
96	Perfusion imaging. <i>Journal of the American College of Cardiology</i> , 2000 , 35, 30B-31B	5.1	
95	Identification of chest pain patients appropriate for an emergency department observation unit. 2001 , 19, 35-66		25
94	Exercise echocardiography for the prognostic stratification of patients with low pretest probability of coronary artery disease. 2001 , 111, 18-23		35
93	Cardiac nuclear medicine in monitoring patients with coronary heart disease. 2001 , 31, 223-37		21
92	Long-term prognostic value of 201Tl single-photon emission computed tomographic myocardial perfusion imaging after coronary stenting. 2001 , 141, 999-1006		34
91	Cardiac imaging. 2001 , 13, 155-163		О
90	Technetium-99m-labeled myocardial perfusion agents: Are they better than thallium-201?. 2001 , 9, 160-7	2	22
89	Post-myocardial infarction risk stratification with stress nuclear myocardial perfusion imaging versus echocardiography: separate but not equal. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 215-8	.1	1
88	Comparison of Tl-201 with Tc-99m-labeled myocardial perfusion agents: technical, physiologic, and clinical issues. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 482-98	.1	42
87	Incidence of major cardiovascular events in black patients with normal myocardial stress perfusion study results. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 541-7	.1	12
86	Principal uses of myocardial perfusion scintigraphy in the management of patients with known or suspected coronary artery disease. 2001 , 43, 281-302		22
85	Non-invasive investigations. 2001 , 59, 29-44		2

84	Assessment of transient left ventricular dilatation on rest and exercise on Tc-99m tetrofosmin myocardial SPECT. 2002 , 27, 34-9	13
83	Can the estimation of ejection fraction during gated single photon emission computed tomography at rest add information to the cardiac perfusion study?. 2002 , 23, 899-906	1
82	Exercise echocardiography and thallium-201 single-photon emission computed tomography stress test for 5- and 10-year prognosis of mortality and specific cardiac events. 2002 , 15, 1326-34	13
81	Stress echocardiography for risk stratification of patients with chest pain and normal or slightly narrowed coronary arteries. 2002 , 15, 1285-9	9
80	Validation of an algorithm for predicting cardiac events in renal transplant candidates. 2002, 89, 847-50	49
79	Predictive value of myocardial tomoscintigraphy in asymptomatic diabetic patients after percutaneous coronary intervention. 2003 , 90, 165-73	8
78	Comparison of the predischarge exercise thallium-201 perfusion defect after myocardial infarction with myocardium at risk measured during acute infarction with technetium-99m sestamibi imaging. 2003 , 145, 357-63	О
77	[Clinical predictors of the presence of perfusion defects in more than than one territory in myocardial scintigraphy with GATED-SPECT]. 2003 , 22, 20-5	
76	Medium- to long-term prognostic impact of dipyridamole thallum-201 myocardial single-photon emission computed tomography in elderly patients. 2003 , 67, 913-7	10
75	Outcome prediction in patients at high risk for coronary artery disease: comparison between 99mTc tetrofosmin and 99mTc sestamibi. 2004 , 232, 58-65	17
74	In retroSPECT: the prognostic value of nuclear cardiology-past, present, and future. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 237-8	
73	Prognostic value of normal stress echocardiogram in patients with suspected coronary artery diseasea British general hospital experience. 2004 , 94, 181-6	21
72	Incremental prognostic value of myocardial perfusion 99m-technetium-sestamibi SPECT in the elderly. 2004 , 93, 137-43	29
71	The prognostic value of myocardial perfusion imaging in patients with suspected or known coronary artery disease. 2004 , 25, 217-20	1
70	Radionuclide-based insights into the pathophysiology of ischemic heart disease: beyond diagnosis. 2005 , 53, 176-91	5
69	Prognostic value of quantitative stress myocardial perfusion imaging in unstable angina patients with negative cardiac enzymes and no new ischemic ECG changes. <i>Journal of Nuclear Cardiology</i> , 2.1 2005 , 12, 32-6	8
68	The clinical importance of electrocardiographic changes during pharmacologic stress testing with radionuclide myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 466-72	9
67	Potential indications for coronary angiography by computed tomography. 2005 , 3, 161-6, 174	4

66	Prediction of mortality in patients with coronary artery disease undergoing vasodilator stress testing: a comparison between 99mTc-tetrofosmin and 99mTc-sestamibi. 2007 , 28, 457-63		1
65	Quantification of left ventricular volumes and ejection fraction from gated 99mTc-MIBI SPECT: validation of an elastic surface model approach in comparison to cardiac magnetic resonance imaging, 4D-MSPECT and QGS. 2007 , 34, 900-9		27
64	Three dimensional first-pass myocardial perfusion imaging at 3T: feasibility study. 2008, 10, 57		45
63	PronEtico de la enfermedad coronaria crilica mediante gated-SPECT de perfusili miocidica. 2008 , 8, 25B-34B		
62	Radiopharmaceutical agents for myocardial perfusion imaging. 2008, 118, 1668-74		69
61	Stress/Rest Myocardial Perfusion Abnormalities by Gated SPECT: Still the Best Predictor of Cardiac Events in Stable Ischemic Heart Disease. 2009 , 50, 546-53		60
60	Normal myocardial perfusion scan portends a benign prognosis independent from the pretest probability of coronary artery disease. Sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2009 , 54, 93-100	3	7
59	New protocol of myocardial SPECT imaging with technetium-99m sestamibi for reducing the time interval between rest and adenosine stress phases. 2009 , 2, 70-6		2
58	Coronary artery phenotypes in subjects with negative myocardial perfusion imaging and typical angina pectoris. 2010 , 340, 350-5		
57	Abnormal myocardial perfusion in the absence of anatomically significant coronary artery disease: implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12	2.1	2
57 56		2.1	6
	implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12 Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> ,		
56	implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12 Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2010 , 56, 51-8 Systolic 3D first-pass myocardial perfusion MRI: Comparison with diastolic imaging in healthy		6
56 55	implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12 Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2010 , 56, 51-8 Systolic 3D first-pass myocardial perfusion MRI: Comparison with diastolic imaging in healthy subjects. 2010 , 63, 858-64		6
56 55 54	implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12 Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2010 , 56, 51-8 Systolic 3D first-pass myocardial perfusion MRI: Comparison with diastolic imaging in healthy subjects. 2010 , 63, 858-64 Multimodality cardiovascular molecular imaging technology. 2010 , 51 Suppl 1, 38S-50S Ninth annual Mario S. Verani memorial lecture: testing our tests: the evidence bar is rising. Regulatory considerations in the development of cardiac imaging agents. <i>Journal of Nuclear</i>	3	6 18
56 55 54 53	implications and clinical significance. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 8-12 Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2010 , 56, 51-8 Systolic 3D first-pass myocardial perfusion MRI: Comparison with diastolic imaging in healthy subjects. 2010 , 63, 858-64 Multimodality cardiovascular molecular imaging technology. 2010 , 51 Suppl 1, 38S-50S Ninth annual Mario S. Verani memorial lecture: testing our tests: the evidence bar is rising. Regulatory considerations in the development of cardiac imaging agents. <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 547-60 Accelerated, high spatial resolution cardiovascular magnetic resonance myocardial perfusion	2.1	6 18 15
56 55 54 53 52	Relation between prognosis and myocardial perfusion imaging from the difference of end-point criterion for exercise stress testing: a sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2010, 56, 51-8 Systolic 3D first-pass myocardial perfusion MRI: Comparison with diastolic imaging in healthy subjects. 2010, 63, 858-64 Multimodality cardiovascular molecular imaging technology. 2010, 51 Suppl 1, 38S-50S Ninth annual Mario S. Verani memorial lecture: testing our tests: the evidence bar is rising. Regulatory considerations in the development of cardiac imaging agents. <i>Journal of Nuclear Cardiology</i> , 2011, 18, 547-60 Accelerated, high spatial resolution cardiovascular magnetic resonance myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2011, 18, 952-8	2.1	6 18 15

(2000-2012)

48	The complementary roles of radionuclide myocardial perfusion imaging and cardiac computed tomography. 2012 , 47, 228-39	3
47	Three-dimensional first-pass myocardial perfusion MRI using a stack-of-spirals acquisition. 2013 , 69, 839-44	33
46	Advanced cardiovascular magnetic resonance myocardial perfusion imaging: high-spatial resolution versus 3-dimensional whole-heart coverage. 2013 , 6, 339-48	31
45	Radioiodinated acebutolol as a new highly selective radiotracer for myocardial perfusion imaging. 2014 , 57, 593-9	18
44	Ischemic burden by 3-dimensional myocardial perfusion cardiovascular magnetic resonance: comparison with myocardial perfusion scintigraphy. 2014 , 7, 647-54	31
43	Role of coronary physiology in the contemporary management of coronary artery disease. 2015 , 3, 148-55	2
42	Prediction Models for Cardiac Risk Classification with Nuclear Cardiology Techniques. 2016 , 9, 1	
41	The elusive role of myocardial perfusion imaging in stable ischemic heart disease: Is ISCHEMIA the answer?. <i>Journal of Nuclear Cardiology</i> , 2017 , 24, 1610-1618	2
40	Myocardial perfusion scintigraphy for risk stratification of patients with coronary artery disease: the AMICO registry. 2020 ,	О
39	Perfusion Measurements of the Myocardium. 2015 , 1279-1354	1
39	Perfusion Measurements of the Myocardium. 2015 , 1279-1354 Cardiac Rehabilitation. 2011 , 713-740	1
38	Cardiac Rehabilitation. 2011 , 713-740	1
38	Cardiac Rehabilitation. 2011 , 713-740 State-of-the-Art Myocardial Perfusion Imaging. 1994 , 12, 199-222	1 28
38 37 36	Cardiac Rehabilitation. 2011, 713-740 State-of-the-Art Myocardial Perfusion Imaging. 1994, 12, 199-222 The Prognosis in Stable and Unstable Angina. 1991, 9, 27-38	1 28 25
38 37 36 35	Cardiac Rehabilitation. 2011, 713-740 State-of-the-Art Myocardial Perfusion Imaging. 1994, 12, 199-222 The Prognosis in Stable and Unstable Angina. 1991, 9, 27-38 Risk Stratification in Angina Pectoris. 1991, 9, 39-47 Comparative Analysis of the Diagnostic and Prognostic Value of Exercise ECG and Thallium-201	1 28 25 3
38 37 36 35 34	Cardiac Rehabilitation. 2011, 713-740 State-of-the-Art Myocardial Perfusion Imaging. 1994, 12, 199-222 The Prognosis in Stable and Unstable Angina. 1991, 9, 27-38 Risk Stratification in Angina Pectoris. 1991, 9, 39-47 Comparative Analysis of the Diagnostic and Prognostic Value of Exercise ECG and Thallium-201 Scintigraphic Markers of Myocardial Ischemia in Asymptomatic and Symptomatic Patients. 1989, 7, 565-575 Exercise and rest technetium-99m-tetrofosmin lung uptake: correlation with left ventricular	1 28 25 3

Myocardial Blood Flow and Perfusion: Radionuclide Techniques. 2002, 555-577 30 Application of stress echocardiography to the prediction of outcomes. Developments in 29 Cardiovascular Medicine, 2003, 167-205 28 Nuclear Cardiology Techniques in Acute Coronary Syndromes. 2011, 148-162 Perfusion Measurements of the Myocardium: Radionuclide Methods and Related Techniques. 2014, 1-89 27 26 Impact of Radionuclide Techniques on Clinical Cardiology. 1985, 381-399 The Quantification of Residual Ischemia in Patients Recovering from Acute Myocardial Infarction by 25 Exercise Thallium-201 Scintigraphy. 1985, 483-494 The frequency, pathophysiology, and prognosis of exercise-induced silent ischemia. 1987, 96-106 24 4 Radionuclide Imaging of the Cardiovascular System. 1988, 103-113 23 The Diagnostic and Prognostic Value of Exercise ECG and Radionuclide Markers of Ischemia in 2.2 Asymptomatic and Symptomatic Populations. Developments in Cardiovascular Medicine, 1988, 25-38 Chest Pain in a Young Man. 1988, 82-94 21 Radionuclide imaging of the heart. 1991, 1-40 20 Antianginal treatment and 201Tl scintigraphy.. 1991, 84, 2203-2204 19 18 Clinical Applications. 1992, 207-214 Planar Thallium-201 imaging: Useful in clinical cardiology?. 1992, 65-78 17 Prognostic evaluation and follow-up of chronic coronary artery disease. Developments in 16 1 Cardiovascular Medicine, 1994, 216-240 Imaging to justify no intervention. Developments in Cardiovascular Medicine, 1996, 75-92 State-of-the-Art Diagnosis in Myocardial Ischema. Developments in Cardiovascular Medicine, 1997, 83-101 14 Nuclear cardiology: which clinical questions can be answered now and in the future?. Developments 13 in Cardiovascular Medicine, 1998, 261-278

CITATION REPORT

Serial stress myocardial perfusion imaging in patients with percutaneous coronary intervention: a retrospective single-center cohort study. *Journal of the Japanese Coronary Association*, **2015**, 21, 20-27

11	Cardiovascular event risk estimated by myocardial perfusion SPECT combined with clinical data. <i>Journal of Cardiology</i> , 2021 ,	3	1
10	Advances in Cardiac Diagnosis. <i>Clinics in Sports Medicine</i> , 1984 , 3, 417-424	2.6	
9	Radioisotope Techniques in the Evaluation of Patients with Coronary Artery Disease. <i>Primary Care - Clinics in Office Practice</i> , 1985 , 12, 67-76	2.2	
8	Mental stress and myocardial ischemia. Correlates and potential interventions. <i>Texas Heart Institute Journal</i> , 1993 , 20, 152-7	0.8	9
7	Investigation and management of stable angina: revised guidelines 1998. Joint Working Party of the British Cardiac Society and Royal College of Physicians of London. <i>Heart</i> , 1999 , 81, 546-55	5.1	20
6	Comparative evaluation of the diagnostic accuracy of (99m)Tc-sestamibi gated SPECT using five different sets of image acquisitions at stress and rest phases for the diagnosis of coronary artery disease. <i>American Journal of Nuclear Medicine and Molecular Imaging</i> , 2013 , 4, 10-6	2.2	1
5	Chest Pain with Normal Thallium-201 Myocardial Perfusion Image - Is It Really Normal?. <i>Acta Cardiologica Sinica</i> , 2016 , 32, 328-36	1.1	
4	Imaging guidelines for nuclear cardiology procedures. American Society of Nuclear Cardiology. Myocardial perfusion stress protocols. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, G11-5	2.1	1
3	Proceedings of the 4th Invitational Wintergreen Conference. Wintergreen, Virginia, USA. July 12-14, 1998. Abstracts. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 93-155	2.1	9
2	Assessing the Adequacy of Myocardial Perfusion in Man: Anatomic and Functional Techniques. 1985 , 23, 589-596		О
1	Myocardial Ischemic Disease. 2013 , 464-489		O