

Response of Monocotyledons to BAS 9052 OH

Weed Science

32, 28-32

DOI: [10.1017/s004317450005846x](https://doi.org/10.1017/s004317450005846x)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Sethoxydim Metabolism in Monocotyledonous and Dicotyledonous Plants. <i>Weed Science</i> , 1985, 33, 771-773.	1.5	25
2	Tolerance of Red Fescue (<i>Festuca rubra</i>) and Bentgrass (<i>Agrostis</i> spp.) to Sethoxydim. <i>Weed Science</i> , 1986, 34, 457-461.	1.5	16
3	Effects of Sublethal Concentrations of Bentazon, Fluazifop, Haloxyfop, and Sethoxydim on Corn (<i>Zea mays</i>). <i>Weed Science</i> , 1986, 34, 171-174.	1.5	11
4	Physiological influences of fenoxaprop on corn (<i>Zea mays</i>). <i>Pesticide Biochemistry and Physiology</i> , 1987, 28, 333-340.	3.6	6
5	Influence of Environment on Corn (<i>Zea mays</i>) Tolerance to Sethoxydim. <i>Weed Science</i> , 1987, 35, 568-575.	1.5	23
6	Rice (<i>Oryza sativa</i>) Tolerance to Fenoxaprop. <i>Weed Science</i> , 1987, 35, 401-406.	1.5	18
7	Susceptibility of Rice (<i>Oryza sativa</i>) to Various Postemergence Grass Herbicides. <i>Weed Science</i> , 1987, 35, 686-690.	1.5	7
8	Retention, absorption, translocation and distribution of sethoxydim in monocotyledonous and dicotyledonous plants*. <i>Weed Research</i> , 1987, 27, 179-186.	1.7	10
9	Moisture Stress Effects on Absorption and Translocation of Four Foliar-Applied Herbicides. <i>Weed Technology</i> , 1988, 2, 437-441.	0.9	24
10	Four Foliar Pathogenic Fungi for Controlling Seedling Johnsongrass (<i>Sorghum halepense</i>). <i>Weed Science</i> , 1989, 37, 802-809.	1.5	9
11	Tolerance of Five Perennial Cool-Season Grasses to Fluazifop. <i>Weed Technology</i> , 1989, 3, 385-388.	0.9	4
12	Effect of Sethoxydim and Haloxyfop on Acetyl-Coenzyme a Carboxylase Activity in <i>Festuca</i> Species. <i>Weed Science</i> , 1989, 37, 512-516.	1.5	71
13	Herbicides for Postemergence Control of Annual Grass Weeds in Seedling Forage Grasses. <i>Weed Science</i> , 1989, 37, 375-379.	1.5	15
14	Tolerance of Corn (<i>Zea mays</i>) to Sethoxydim Applied With Precision Postemergence-directed Sprayer Equipment. <i>Weed Technology</i> , 1989, 3, 663-667.	0.9	7
15	The Effect of Sethoxydim on Corn (<i>Zea mays</i>) and Giant Foxtail (<i>Setaria faberi</i>). <i>Weed Science</i> , 1989, 37, 600-603.	1.5	6
16	Control of Seedling Grasses with Postemergence Grass Herbicides. <i>Weed Technology</i> , 1989, 3, 39-43.	0.9	24
17	Inhibition of corn acetyl-CoA carboxylase by cyclohexanedione and aryloxyphenoxypropionate herbicides. <i>Pesticide Biochemistry and Physiology</i> , 1989, 34, 76-85.	3.6	108
18	Compatibility of Sethoxydim with Five Postemergence Broadleaf Herbicides. <i>Weed Technology</i> , 1990, 4, 128-133.	0.9	70

#	ARTICLE	IF	CITATIONS
19	Improving the Efficiency of Sethoxydim in Flax. <i>Weed Technology</i> , 1990, 4, 749-753.	0.9	4
20	Kinetics of inhibition of acetyl-coenzyme A carboxylase by sethoxydim and haloxyfop. <i>Pesticide Biochemistry and Physiology</i> , 1991, 39, 100-109.	3.6	73
21	Control of Downy Brome (<i>Bromus tectorum</i>) in Conservation Fallow Systems. <i>Weed Technology</i> , 1991, 5, 557-562.	0.9	10
22	A Chlorophyll Assay to Assess Sethoxydim Activity in Two Grass Species. <i>Weed Technology</i> , 1991, 5, 355-362.	0.9	1
23	Lipid Biosynthesis Inhibitors. <i>Weed Science</i> , 1991, 39, 435-449.	1.5	118
24	Effects of Various Adjuvants on Sethoxydim Activity. <i>Weed Technology</i> , 1992, 6, 865-870.	0.9	24
25	Ammonium Sulfate Increases Efficacy of Sethoxydim Through Increased Absorption and Translocation. <i>Weed Science</i> , 1992, 40, 351-358.	1.5	20
26	Herbicide Tolerance and Weed Control in Sethoxydim-Tolerant Corn (<i>Zea mays</i>). <i>Weed Science</i> , 1993, 41, 213-217.	1.5	27
27	Resistance of Selected Ornamental Grasses to Graminicides. <i>Weed Technology</i> , 1993, 7, 326-330.	0.9	6
28	Rotation Affects Downy Brome (<i>Bromus tectorum</i>) in Winter Wheat (<i>Triticum aestivum</i>). <i>Weed Technology</i> , 1994, 8, 728-732.	0.9	62
29	Weed management in conservation tillage systems for wheat production in North and South America. <i>Crop Protection</i> , 1994, 13, 243-259.	2.1	93
30	Interactions of Sethoxydim and Corn (<i>Zea mays</i>) Postemergence Broadleaf Herbicides on Three Annual Grasses. <i>Weed Technology</i> , 1996, 10, 914-922.	0.9	25
31	Giant foxtail (<i>Setaria faberi</i>) control in sethoxydim-resistant corn (<i>Zea mays</i>). <i>Weed Science</i> , 1997, 45, 771-776.	1.5	6
32	Mechanism of herbicidal activity of a new cyclohexane-1,3-dione, tepraloxym to <i>Poa annua</i> L.. <i>Weed Biology and Management</i> , 2002, 2, 84-91.	1.4	7
33	Rate response of select grass weeds to pinoxaden. <i>Weed Technology</i> , 2020, 34, 818-823.	0.9	7
34	Photosynthesis as an Index of Turfgrass Growth Following Application of Herbicides. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 1990, 25, 451-453.	1.0	4
35	Absorption, Translocation, and Metabolism of Sethoxydim in Centipedegrass and Goosegrass. <i>Journal of the American Society for Horticultural Science</i> , 1990, 115, 605-607.	1.0	4
36	Selectivity of Cyhalofop-butyl in Poaceae Species.. <i>Journal of Weed Science and Technology</i> , 1998, 43, 122-128.	0.1	7

#	ARTICLE	IF	CITATIONS
37	Assessment of Genetic Variations among Medicinal Plant Cassia tora from Different Geographic Regions of Central India Using RAPD Markers. , 2016, 05, .		7
38	Selectivity mechanisms of sethoxydim between red fescue (<i>Festuca rubra</i>) and tall fescue (<i>Festuca</i>) Tj ETQq1 1 0.784314 rgBT /Overlo	0.1	1