

Homogeneous Enzyme Immunoassay of Acetaminophe

American Journal of Clinical Pathology

81, 602-609

DOI: [10.1093/ajcp/81.5.602](https://doi.org/10.1093/ajcp/81.5.602)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Competitive enzyme immunoassay for anti-ulcer agent, (-)-cis-2,3-dihydro-3-(4-methylpiperazinylmethyl)-2-phenyl-1,5-benzothiazepin-4(5H)-one hydrochloride (BTM-1086).. Chemical and Pharmaceutical Bulletin, 1986, 34, 4238-4243.	1.3	12
2	Analytical Reviews in Clinical Chemistry: Methods for the Estimation of Salicylate and Paracetamol in Serum, Plasma and Urine. Annals of Clinical Biochemistry, 1987, 24, 552-565.	1.6	23
3	Design of the Immunogen and Label for Use in a Fluoroimmunoassay for Paracetamol. Annals of Clinical Biochemistry, 1988, 25, 42-48.	1.6	15
4	The estimation of paracetamol and its major metabolites in both plasma and urine by a single high-performance liquid chromatography assay. Journal of Pharmaceutical and Biomedical Analysis, 1994, 12, 1563-1572.	2.8	62
5	A Validated High-Performance Liquid Chromatographic Method for the Determination Of Paracetamol and Its Major Metabolites in Urine. Journal of Liquid Chromatography and Related Technologies, 1995, 18, 3257-3268.	1.0	30
6	Development and optimization of a reversed-phase high-performance liquid chromatographic method for the determination of acetaminophen and its major metabolites in rabbit plasma and urine after a toxic dose. Journal of Pharmaceutical and Biomedical Analysis, 2003, 32, 487-493.	2.8	43
7	Effect of hemolysis, icterus, and lipemia on three acetaminophen assays: Potential medical consequences of false positive results. Clinica Chimica Acta, 2018, 487, 287-292.	1.1	9
8	Application of Immunoassay Techniques in Psychopharmacology. , 1988, , 241-302.		0