

Nephrospris knutheierisp.n., an extant trissocyclid rad

Sarsia

70, 103-118

DOI: 10.1080/00364827.1985.10420623

Citation Report

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Distribution of coccoliths in surface sediments in the Norwegian-Greenland Sea. <i>Marine Micropaleontology</i> , 1990, 16, 65-75. | 1.2 | 31 |
| 2 | Holocene palaeo-oceanography of the northern Iceland Sea, indicated by Radiolaria and sponge spicules. <i>Journal of Quaternary Science</i> , 1991, 6, 303-312. | 2.1 | 20 |
| 3 | Tintinnids and Other Microplankton from the Greenland Sea: Abundance and Distribution in the Marginal Ice Zone (MayJune 1989). <i>Marine Ecology</i> , 1995, 16, 117-131. | 1.1 | 5 |
| 4 | Plankton in the Norwegian-Greenland Sea: from living communities to sediment assemblages ?an actualistic approach. <i>Geologische Rundschau: Zeitschrift Fur Allgemeine Geologie</i> , 1995, 84, 108. | 1.3 | 119 |
| 5 | Distribution of radiolaria in surface sediments and its relation to the oceanography of the Iceland and Greenland seas. <i>Sarsia</i> , 1996, 81, 315-328. | 0.5 | 16 |
| 6 | Radiolarian faunal provinces in surface sediments of the Greenland, Iceland and Norwegian (GIN) Seas. <i>Marine Micropaleontology</i> , 1998, 35, 105-140. | 1.2 | 62 |
| 7 | Polycystine radiolarians in surface sediments in the Arctic Ocean basins and marginal seas. <i>Marine Micropaleontology</i> , 2003, 49, 231-273. | 1.2 | 55 |
| 8 | Joergensenium rotatile n. gen., n. sp. (Entactinaria, Radiolaria): its distribution in west Norwegian fjords. <i>Micropaleontology</i> , 2007, 53, 457-468. | 1.0 | 10 |
| 9 | Locomotion in <i>Nebalia bipes</i> : a possible model for Palaeozoic phyllocarid crustaceans. <i>Lethaia</i> , 1997, 30, 89-104. | 1.4 | 43 |
| 10 | The Cambrian origin of the circulatory system of crustaceans. <i>Lethaia</i> , 1997, 30, 169-184. | 1.4 | 36 |
| 11 | Anatomy and lifestyle of <i>Kunmingella</i> (Arthropoda, Bradoriida) from the Chengjiang fossil Lagerstätte (lower Cambrian; Southwest China). <i>Lethaia</i> , 1999, 32, 279-298. | 1.4 | 81 |
| 12 | A new integrated morpho- and molecular systematic classification of Cenozoic radiolarians (Class) Tj ETQq1 1 0.784314 rgBT _{0.8} 14 Overlock | | |
| 13 | Distribution of Calcareous, Siliceous and Organic-Walled Planktic Microfossils in Surface Sediments of the Nordic Seas and their Relation to Surface-Water Masses. , 2001, , 105-127. | | 15 |
| 14 | Norwegian â€” Iceland Seas: Transfer Functions between Marine Planktic Diatoms and Surface Water Temperature. , 1990, , 337-361. | | 11 |
| 15 | Catalogue of Cenozoic radiolarian genera (Class Polycystinea). <i>Geodiversitas</i> , 2021, 43, . | 0.8 | 11 |
| 16 | Inventory of Cenozoic radiolarian species (Class Polycystinea) â€“ 1834-2020. <i>Geodiversitas</i> , 2022, 44, . | 0.8 | 2 |