

CITATION REPORT

List of articles citing

Risk stratification after acute myocardial infarction by means of exercise two-dimensional echocardiography

DOI: 10.1016/0002-8703(87)90530-8
American Heart Journal, 1987, 114, 1305-16.

Source: <https://exaly.com/paper-pdf/19126808/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
168	Exercise echocardiography: detection of coronary artery disease in patients with normal left ventricular wall motion at rest. <i>Journal of the American College of Cardiology</i> , 1988 , 11, 993-9	15.1	202
167	Stress testing. Directions for the future. 1988 , 6, 11-22		
166	Exercise echocardiography: ready, willing and able. <i>Journal of the American College of Cardiology</i> , 1988 , 11, 1359-61	15.1	37
165	Stress Echocardiography and Myocardial Contrast Echocardiography. <i>Cardiology Clinics</i> , 1989 , 7, 493-509	2.5	9
164	High dose dipyridamole echocardiography early after uncomplicated acute myocardial infarction: correlation with exercise testing and coronary angiography. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 357-63	15.1	103
163	Reproducibility of two-dimensional exercise echocardiography. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 923-8	15.1	59
162	Bibliography of biomedical ultrasound. 1989 , 15, 775-788		1
161	Role of echocardiography in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1990 , 66, 17H-22H		18
160	Echocardiography in coronary artery disease. 1990 , 15, 233-98		18
159	ACC/AHA guidelines for the clinical application of echocardiography. A report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Subcommittee to Develop Guidelines for the Clinical Application of Echocardiography). <i>Circulation</i> , 1990 , 82, 2323-45	16.7	34
158	Echocardiography (2). 1990 , 323, 165-72		31
157	ACC/AHA guidelines for the early management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (subcommittee to develop guidelines for the early management of patients with acute myocardial infarction). <i>Circulation</i> , 1990 , 82, 664-707	16.7	200
156	Bibliography of biomedical ultrasound. 1987. 1990 , 16, 851-1018		1
155	Prediction of cardiac events after uncomplicated myocardial infarction by cross-sectional echocardiography during transesophageal atrial pacing. 1990 , 28, 95-103		25
154	ACC/AHA guidelines for the clinical application of echocardiography. <i>Journal of the American College of Cardiology</i> , 1990 , 16, 1505-1528	15.1	21
153	Simultaneous transesophageal atrial pacing and transesophageal two-dimensional echocardiography: a new method of stress echocardiography. <i>Journal of the American College of Cardiology</i> , 1990 , 16, 1143-53	15.1	95
152	Guidelines for the early management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Subcommittee to Develop Guidelines for the Early Management of Patients with Acute Myocardial Infarction). <i>Journal of the American College of Cardiology</i> , 1990 , 16, 249-92	15.1	302

151	Improvement in rest and exercise-induced wall motion abnormalities after coronary angioplasty: an exercise echocardiographic study. <i>Journal of the American College of Cardiology</i> , 1990 , 15, 591-9	15.1	69
150	Exercise echocardiography after coronary angioplasty: expanding applications. <i>Journal of the American College of Cardiology</i> , 1990 , 15, 600-1	15.1	2
149	Diagnostic accuracy of peak exercise echocardiography in coronary artery disease: comparison with thallium-201 myocardial scintigraphy. <i>American Heart Journal</i> , 1991 , 122, 1609-16	4.9	60
148	Risk stratification in survivors of acute myocardial infarction: routine cardiac catheterization and angiography is a reasonable approach in most patients. <i>American Heart Journal</i> , 1991 , 121, 641-56	4.9	50
147	Uncomplicated first myocardial infarction: strategy for comprehensive prognostic studies. <i>Journal of the American College of Cardiology</i> , 1991 , 18, 1207-19	15.1	45
146	The comparative value of exercise echocardiography and 99m Tc MIBI single photon emission computed tomography in the diagnosis and localization of myocardial ischaemia. <i>European Heart Journal</i> , 1991 , 12, 1293-9	9.5	26
145	Role of Doppler echocardiography in coronary artery disease. 1991 , 6, 238-56		1
144	Noninvasive and invasive testing after myocardial infarction. 1991 , 16, 721-63		1
143	Trends in use of coronary angiography in subacute phase of myocardial infarction. <i>Circulation</i> , 1991 , 84, 1004-15	16.7	33
142	Exercise echocardiography versus 201Tl single-photon emission computed tomography in evaluation of coronary artery disease. Analysis of 292 patients. <i>Circulation</i> , 1992 , 85, 1026-31	16.7	266
141	Medical treatment after myocardial infarction. Results of studies using various methods. 1992 , 92, 84-90		
140	Cardiac imaging after acute myocardial infarction. Identification of patients at continued risk. 1992 , 92, 93-6, 99-100		
139	Stress echocardiography versus radionuclide stress techniques. <i>Echocardiography</i> , 1992 , 9, 199-209	1.5	4
138	Utility of stress echocardiography for postinfarct prognosis. <i>Echocardiography</i> , 1992 , 9, 211-8	1.5	4
137	Risk of cardiac events (in patients with established ischaemic heart disease). <i>European Heart Journal</i> , 1992 , 13 Suppl C, 14-9	9.5	8
136	Exercise Doppler: functional evaluation of right heart dynamics. <i>Echocardiography</i> , 1992 , 9, 225-33	1.5	7
135	Quantitative stress echocardiography. <i>Echocardiography</i> , 1992 , 9, 85-96	1.5	8
134	Exercise echocardiography: phase II, convincing the skeptics. <i>Journal of the American College of Cardiology</i> , 1992 , 19, 82-3	15.1	8

133	Silent versus symptomatic dipyridamole-induced ischemia after myocardial infarction: clinical and prognostic significance. <i>Journal of the American College of Cardiology</i> , 1992 , 19, 953-9	15.1	57
132	Exercise echocardiography: a methodological study comparing peak-exercise and post-exercise image information. 1992 , 12, 553-65		4
131	Relationship between exercise echocardiography and perfusion single-photon emission computed tomography in patients with single-vessel coronary artery disease. <i>American Heart Journal</i> , 1992 , 124, 75-83	4.9	50
130	Usefulness of dipyridamole-echocardiographic test to identify jeopardized myocardium after thrombolysis. Limited clinical predictivity of dipyridamole-echocardiographic test in convalescing acute myocardial infarction: correlation with coronary angiography. <i>European Heart Journal</i> , 1992 , 13, 1348-55	9.5	23
129	Exercise echocardiography. <i>American Journal of Cardiology</i> , 1992 , 69, 82H-89H	3	42
128	Predischarge exercise echocardiography in patients with unstable angina who respond to medical treatment. <i>Clinical Cardiology</i> , 1992 , 15, 417-23	3.3	17
127	Comparison of approaches in the assessment of myocardial viability and follow-up of PTCA/CABG. The role of echocardiography. 1993 , 9 Suppl 1, 11-7		5
126	Prognostic value of dobutamine echocardiography in patients with high pretest likelihood of coronary artery disease. <i>American Journal of Cardiology</i> , 1993 , 71, 33-9	3	97
125	Prognostic usefulness of positive or negative exercise stress echocardiography for predicting coronary events in ensuing twelve months. <i>American Journal of Cardiology</i> , 1993 , 71, 646-51	3	102
124	Prognostic significance of exercise thallium-201 myocardial perfusion imaging compared to stress echocardiography and clinical variables in patients with unstable angina who respond to medical treatment. 1993 , 39, 71-8		12
123	Stress echocardiography: time for critical reappraisal. <i>Heart</i> , 1993 , 70, 208-10	5.1	1
122	Clinical Utility of Echocardiography in Neurologic Disorders. 1993 , 11, 353-374		3
121	Usefulness of Exercise Echocardiography in Early Detection of Multivessel Coronary Artery Disease after Uncomplicated Myocardial Infarction. 1994 , 8, 215-220		
120	Reperfusion in acute myocardial infarction. International Society and Federation of Cardiology and World Health Organization Task Force on Myocardial Reperfusion. <i>Circulation</i> , 1994 , 90, 2091-102	16.7	34
119	A consideration of current clinical options for stress imaging in the diagnosis and evaluation of coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 1994 , 1, S147-70	2.1	5
118	Usefulness of transesophageal atrial pacing combined with two-dimensional echocardiography (echo-pacing) in predicting the presence and site of residual jeopardized myocardium after uncomplicated acute myocardial infarction. <i>American Journal of Cardiology</i> , 1994 , 73, 534-8	3	12
117	Risk stratification after myocardial infarction. 1994 , 17, 401-16		12
116	Comparison of modalities to diagnose coronary artery disease. 1994 , 24, 286-310		27

115	Dipyridamole echocardiographic test performed 3 days after an acute myocardial infarction: feasibility, tolerability, safety and in-hospital prognostic value. <i>European Heart Journal</i> , 1994 , 15, 842-50 ^{9.5}	29
114	Diagnostic and prognostic value of stress testing in older persons. 1995 , 43, 190-4	7
113	Stress imaging. Current clinical options for the diagnosis, localization, and evaluation of coronary artery disease. 1995 , 79, 1025-61	6
112	Usefulness of combined two-dimensional echocardiography and transesophageal atrial stimulation early after acute myocardial infarction. <i>American Journal of Cardiology</i> , 1995 , 76, 1112-4	3 6
111	Prognostic value of predischage exercise stress echocardiography after acute myocardial infarction. <i>American Journal of Cardiology</i> , 1995 , 76, 1115-21	3 36
110	Myocardial contrast echocardiography in coronary artery disease: potential applications using venous injections of contrast. <i>American Journal of Cardiology</i> , 1995 , 75, 61D-68D	3 21
109	Long-term prognosis of patients with a normal exercise echocardiogram and clinical suspicion of myocardial ischemia. <i>American Journal of Cardiology</i> , 1995 , 75, 934-6	3 21
108	Long-term prognostic importance of exercise echocardiography after an episode of unstable angina. 1995 , 86, 426-31	10
107	Prediction of Myocardium at Risk. <i>Cardiology Clinics</i> , 1995 , 13, 355-378	2.5 6
106	Stress echocardiography. Part I. Exercise echocardiography: techniques, implementation, clinical applications, and correlations. 1995 , 70, 5-15	88
105	Risk stratification after myocardial infarction: targets and tools. <i>Echocardiography</i> , 1995 , 12, 311-6	1.5
104	Strategies for prognostic assessment of uncomplicated first myocardial infarction: 5-year follow-up study. <i>Journal of the American College of Cardiology</i> , 1995 , 25, 815-22	15.1 29
103	Stress echocardiographic results predict risk of reinfarction early after uncomplicated acute myocardial infarction: large-scale multicenter study. Echo Persantine International Cooperative (EPIC) Study Group. <i>Journal of the American College of Cardiology</i> , 1995 , 26, 908-13	15.1 44
102	Stress Echocardiography. Part I. Exercise Echocardiography: Techniques, Implementation, Clinical Applications, and Correlations. 1995 , 70, 5-15	82
101	ACC/AHA guidelines for the management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction). <i>Journal of the American College of Cardiology</i> , 1996 , 28, 1328-428	15.1 871
100	Stress Testing for Coronary Artery Disease in the Elderly. 1996 , 12, 101-119	6
99	Value of dobutamine stress echocardiography in determining the prognosis of patients with known or suspected coronary artery disease. <i>American Journal of Cardiology</i> , 1996 , 78, 404-8	3 63
98	A metaanalysis of predischage risk stratification after acute myocardial infarction with stress electrocardiographic, myocardial perfusion, and ventricular function imaging. <i>American Journal of Cardiology</i> , 1996 , 78, 1327-37	3 97

97	Stress echocardiography for the assessment of myocardial ischemia and viability. 1996 , 21, 445-520		15
96	The role of stress echocardiography versus stress perfusion: a view from the other side. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, S66-74	2.1	11
95	Supine Bicycle Exercise Echocardiography: A Potent Immediately Available Tool for Detection and Localization of Myocardial Ischemia for the Initial Cardiologist. <i>Echocardiography</i> , 1996 , 13, 281-286	1.5	2
94	Angina pectoris in patients with a history of myocardial infarction. <i>European Heart Journal</i> , 1996 , 17 Suppl G, 25-9	9.5	35
93	Stress echocardiography for assessing myocardial ischaemia and viable myocardium. <i>Heart</i> , 1997 , 78 Suppl 1, 12-8	5.1	9
92	Stress echocardiography: new techniques. <i>European Heart Journal</i> , 1997 , 18 Suppl D, D49-56	9.5	14
91	Treadmill exercise echocardiography: methodology and clinical role. <i>European Heart Journal</i> , 1997 , 18 Suppl D, D2-8	9.5	38
90	Amlodipine Improves the Anti-Ischaemic Effect of Atenolol in Postinfarction Patients with Effort-Induced Ischaemia. 1997 , 13, 22-28		
89	Prognostic value of dobutamine echocardiography early after uncomplicated acute myocardial infarction: a comparison with exercise electrocardiography. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 261-7	15.1	49
88	Prognostic value of dobutamine-atropine stress echocardiography early after acute myocardial infarction. Echo Dobutamine International Cooperative (EDIC) Study. <i>Journal of the American College of Cardiology</i> , 1997 , 29, 254-60	15.1	148
87	Use of exercise echocardiography for prognostic evaluation of patients with known or suspected coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1997 , 30, 83-90	15.1	120
86	Prognostic implications of exercise echocardiography in women with known or suspected coronary artery disease. <i>Journal of the American College of Cardiology</i> , 1997 , 30, 414-20	15.1	67
85	Valor prognóstico da ecocardiografia com estresse pela dobutamina associada à atropina. 1997 , 69, 95-99		
84	Echocardiography in anesthesia and intensive care medicine I. 1997 , 41, 267-278		
83	Prediction of life-threatening arrhythmias: Multifactorial risk stratification following acute myocardial infarction. 1997 , 6, 241-253		1
82	Detection of Viability of Dysfunctional Myocardium in Coronary Heart Disease. II. Echocardiography. 1998 , 2, 207-233		4
81	Stress Echocardiography: A Review of the Principles and Practice. <i>Echocardiography</i> , 1998 , 15, 669-692	1.5	4
80	Editorial. <i>American Journal of Cardiology</i> , 1998 , 81, 1050-1053	3	14

79	Usefulness of exercise echocardiography in predicting cardiac events in an outpatient population. <i>American Journal of Cardiology</i> , 1998 , 82, 569-73	3	11
78	Current status of ECG stress testing. 1998 , 23, 353-423		12
77	Outcome after normal exercise echocardiography and predictors of subsequent cardiac events: follow-up of 1,325 patients. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 144-9	15.1	163
76	Prognostic value of myocardial viability and ischemia detected by dobutamine stress echocardiography early after acute myocardial infarction treated with thrombolysis. <i>Journal of the American College of Cardiology</i> , 1998 , 32, 380-6	15.1	46
75	Stress echocardiography: recommendations for performance and interpretation of stress echocardiography. Stress Echocardiography Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. <i>Journal of the American Society of Echocardiography</i> , 1998 , 11, 97-104	5.8	307
74	[The value of exercise echocardiography in the diagnosis of coronary disease in patients with left heart branch block in the bundle of His]. <i>Revista Espanola De Cardiologia</i> , 1998 , 51, 211-7	1.5	3
73	[Prognostic value of pre-discharge exertion echocardiography after acute myocardial infarct]. <i>Revista Espanola De Cardiologia</i> , 1998 , 51, 21-6	1.5	2
72	Use of Exercise Echocardiography to Evaluate Patients With Chest Pain. <i>American Journal of the Medical Sciences</i> , 1998 , 316, 345-350	2.2	
71	Advances in Exercise Echocardiography Can This Technique Still Thrive in the Era of Pharmacologic Stress Testing?. <i>Echocardiography</i> , 1999 , 16, 841-856	1.5	4
70	Diagnostic and Prognostic Use of Stress Echocardiography and Radionuclide Scintigraphy. <i>Echocardiography</i> , 1999 , 16, 857-877	1.5	1
69	Exercise echocardiography. Principles, methods, and clinical use. <i>Cardiology Clinics</i> , 1999 , 17, 447-60, vii	2.5	10
68	Use of stress echocardiography for risk assessment of patients after myocardial infarction. <i>Cardiology Clinics</i> , 1999 , 17, 521-38, ix	2.5	3
67	Valor pronóstico de la ecocardiografía con dobutamina después de un infarto agudo de miocardio no complicado. <i>Revista Espanola De Cardiologia</i> , 1999 , 52, 237-244	1.5	1
66	ACC/AHA guidelines for coronary angiography. A report of the American College of Cardiology/American Heart Association Task Force on practice guidelines (Committee on Coronary Angiography). Developed in collaboration with the Society for Cardiac Angiography and Intervention. <i>Circulation</i> , 1999 , 100, 1016-1044	15.1	794
65	Enhanced detection of ischemic myocardium by transesophageal dobutamine stress echocardiography: comparison with simultaneous transthoracic echocardiography. <i>Echocardiography</i> , 2000 , 17, 241-53	1.5	17
64	Diagnostic and prognostic use of stress echocardiography in stable patients. <i>Echocardiography</i> , 2000 , 17, 465-77	1.5	3
63	Stress Echocardiography. <i>Echocardiography</i> , 2000 , 17, 491-493	1.5	
62	Risk stratification after acute myocardial infarction in the reperfusion era. <i>Progress in Cardiovascular Diseases</i> , 2000 , 42, 273-309	8.5	37

61	Stress nuclear myocardial perfusion imaging versus stress echocardiography: prognostic comparisons. <i>Progress in Cardiovascular Diseases</i> , 2000 , 43, 231-44	8.5	5
60	The effects of acute coronary occlusion on noninvasive echocardiographically derived systolic and diastolic myocardial strain rates. <i>Journal of Surgical Education</i> , 2000 , 57, 466-472		13
59	Current status of stress echocardiography. <i>Clinical Cardiology</i> , 2000 , 23, 242-6	3.3	5
58	[Stress echocardiography]. <i>Clinical Research in Cardiology</i> , 2000 , 89 Suppl 1, 78-85		1
57	Prediction of cardiac mortality after myocardial infarction: the role of maximal treadmill stress echocardiography. <i>Journal of the American Society of Echocardiography</i> , 2001 , 14, 38-43	5.8	3
56	Post-myocardial infarction risk stratification with stress nuclear myocardial perfusion imaging versus echocardiography: separate but not equal. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 215-8	2.1	1
55	Incremental prognostic value of stress echocardiography as an adjunct to exercise electrocardiography after uncomplicated myocardial infarction. <i>British Heart Journal</i> , 2001 , 85, 417-23		13
54	Utility and limitation of treadmill exercise echocardiography for detecting significant coronary stenosis in infarct-related arteries in patients with healed myocardial infarction. <i>American Journal of Cardiology</i> , 2002 , 89, 159-63	3	3
53	Effect of infarcted myocardium on diagnostic accuracy of exercise echocardiography for detecting noninfarct-related coronary artery lesions. <i>American Heart Journal</i> , 2003 , 145, 162-8	4.9	2
52	[Risk stratification after acute myocardial infarction]. <i>Revista Espanola De Cardiologia</i> , 2003 , 56, 303-13	1.5	4
51	Stress echocardiography versus myocardial SPECT for risk stratification of patients with coronary artery disease. <i>Current Opinion in Cardiology</i> , 2003 , 18, 486-93	2.1	8
50	Stress radionuclide myocardial perfusion imaging detects more residual ischemia than stress echocardiography following acute myocardial infarction. <i>International Journal of Cardiovascular Imaging</i> , 2004 , 20, 145-54	2.5	
49	[Prognostic value of stress echocardiography]. <i>Herz</i> , 2005 , 30, 743-53	2.6	2
48	Stress echocardiography for the diagnosis and risk stratification of patients with suspected or known coronary artery disease: a critical appraisal. Supported by the British Society of Echocardiography. <i>Heart</i> , 2005 , 91, 427-36	5.1	60
47	Patency of the right coronary artery following implantation of metallic stent demonstrated by multislice computed tomography. <i>Heart</i> , 2005 , 91, 436	5.1	1
46	Long-term prognostic importance of transient left ventricular dilation during pharmacologic stress echocardiography. <i>Journal of the American Society of Echocardiography</i> , 2005 , 18, 57-62	5.8	8
45	Noninvasive assessment of coronary artery disease in diabetic patients: the role of stress echocardiography. <i>Canadian Journal of Cardiology</i> , 2006 , 22 Suppl A, 26A-33A	3.8	3
44	American Society of Echocardiography recommendations for performance, interpretation, and application of stress echocardiography. <i>Journal of the American Society of Echocardiography</i> , 2007 , 20, 1021-41	5.8	558

43	Harvey Feigenbaum: a retrospective. <i>Journal of the American Society of Echocardiography</i> , 2008 , 21, 3-6	5.8	3
42	Stress Echocardiography Expert Consensus Statement--Executive Summary: European Association of Echocardiography (EAE) (a registered branch of the ESC). <i>European Heart Journal</i> , 2009 , 30, 278-89	9.5	211
41	Echocardiography in Acute Coronary Syndrome. 2009 ,		1
40	Exercise Echocardiography. 2009 , 159-174		
39	Risk Stratification Following Acute Myocardial Infarction: Role of Dobutamine Stress Echocardiography. 2009 , 341-349		
38	The clinical use of stress echocardiography in ischemic heart disease. <i>Cardiovascular Ultrasound</i> , 2017 , 15, 7	2.4	33
37	Harvey Feigenbaum, MD, and the Creation of Clinical Echocardiography: A Conversation With Barry J. Maron, MD. <i>American Journal of Cardiology</i> , 2017 , 120, 2085-2099	3	2
36	Practical guidance for the implementation of stress echocardiography. <i>Journal of Echocardiography</i> , 2018 , 16, 105-129	1.6	12
35	Risk Stratification after Acute Myocardial Infarction: Theory and Practice. <i>Developments in Cardiovascular Medicine</i> , 1989 , 213-258		1
34	Prognosis. 1997 , 193-206		1
33	Risk stratification after myocardial infarction. Clinical science versus practice behavior. <i>Circulation</i> , 1997 , 95, 1352-4	16.7	7
32	Dobutamine stress echocardiography for risk stratification after myocardial infarction. <i>Circulation</i> , 1997 , 95, 1402-10	16.7	126
31	ACC/AHA Guidelines for the Clinical Application of Echocardiography. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Clinical Application of Echocardiography). Developed in collaboration with the American Society of Echocardiography. <i>Circulation</i> , 1997 , 95, 1686-744	16.7	431
30	[The Clinical use of Stress Echocardiography in Ischemic Heart Disease <i>Cardiovascular Ultrasound</i> (2017)15:7. Translation authors: Arystan A.Zh., Fettser D.V.]. <i>Kardiologiya</i> , 2019 , 59, 78-96	1.5	1
29	Echocardiography in Coronary Artery Disease. 2000 , 177-195		
28	Belastungsechokardiographie. 2001 , 119-145		
27	Interpreting Noninvasive Cardiac Tests. 2001 , 25-36		
26	Stress Echocardiography. <i>Developments in Cardiovascular Medicine</i> , 2002 , 109-125		

- 25 Application of stress echocardiography to the prediction of outcomes. *Developments in Cardiovascular Medicine*, **2003**, 167-205
- 24 Exercise Echocardiography. **2003**, 103-114
- 23 Echokardiographie. **2004**, 219-250 o
- 22 Introduction to Echocardiography. **2007**, 93-136
- 21 Exercise Echocardiography. **2007**, 326-352
- 20 Echocardiography in Acute Coronary Syndromes. **2011**, 129-147
- 19 Exercise echocardiography. *Developments in Cardiovascular Medicine*, **1989**, 483-494
- 18 Exercise 2D echocardiography. *Developments in Cardiovascular Medicine*, **1991**, 31-36
- 17 Stress echocardiography for identifying patients at risk after myocardial infarction. *Developments in Cardiovascular Medicine*, **1991**, 231-239
- 16 Prognostic Value of Stress Echocardiography. *Developments in Cardiovascular Medicine*, **1994**, 147-162
- 15 Stress Echocardiography and Prognosis. **1994**, 149-163
- 14 Stellenwert streßchokardiographischer Techniken im Spektrum möglicher Anwendungsgebiete. **1994**, 133-158
- 13 Echokardiographie. **1995**, 142-164
- 12 Viability of the Revascularized Myocardium. **1995**, 149-154
- 11 Echokardiographie. **1996**, 302-333
- 10 Exercise Echocardiography. **1997**, 85-94
- 9 Acute Myocardial Infarction. *Developments in Cardiovascular Medicine*, **1998**, 143-197
- 8 Stellenwert streßchokardiographischer Techniken im Spektrum möglicher Anwendungsgebiete. **1998**, 253-305

7	Use of exercise echocardiography to evaluate patients with chest pain. <i>American Journal of the Medical Sciences</i> , 1998 , 316, 345-50	2.2
6	Risk Stratification: Exercise Testing, Imaging, and Cardiac Catheterization. <i>Contemporary Cardiology</i> , 1999 , 383-405	0.1
5	Echocardiography in the Evaluation of Coronary Artery Disease. <i>Developments in Cardiovascular Medicine</i> , 1999 , 397-420	
4	Exercise Echocardiography. 2015 , 179-195	
3	Stress Echocardiography. 2009 , 143-166	
2	Dobutamine stress echocardiography in clinical practice with a review of the recent literature. <i>Texas Heart Institute Journal</i> , 1994 , 21, 202-10	0.8 4
1	STRESS ECHOCARDIOGRAPHY. 1997 , 51, 41-46	0