## Depositional environments within Middle Jurassic oyst integrated litho-, bio- and palynofacies study of the Dur

Transactions of the Royal Society of Edinburgh: Earth Sciences 81, 1-22 DOI: 10.1017/s0263593300005095

**Citation Report** 

#	Article	IF	CITATIONS
1	Toarcian to Bathonian (Jurassic) palynology of the Inner Hebrides, Northwest Scotland. Palynology, 1991, 15, 115-179.	1.5	82
2	Unusual nonmarine burrows from the Middle Jurassic of Scotland. Ichnos, 1991, 1, 247-253.	0.5	4
3	Mid-Jurassic lagoonal delta systems in the Hebridean basins: thickness and facies distribution patterns of potential reservoir sandbodies. Geological Society Special Publication, 1992, 62, 111-144.	1.3	11
4	Middle Jurassic clay-minerals from the Minch Basin: isotopic tracing of provenance and post-depositional alteration. Geological Society Special Publication, 1992, 62, 155-158.	1.3	2
5	Palaeoecology and evolution of Mesozoic salinity ontrolled benthic macroinvertebrate associations. Lethaia, 1993, 26, 327-346.	1.4	99
6	Ostracod biostratinomy at lagoonal shorelines: examples from the Great Estuarine Group, Middle Jurassic, Scotland. Proceedings of the Geologists Association, 1995, 106, 211-218.	1.1	14
7	Jurassic Paleosalinities and Brackish-Water Communities: A Case Study. Palaios, 1995, 10, 392.	1.3	32
8	Palaeoecology of benthic associations in salinity-controlled marginal marine environments: Examples from the Lower Bathonian (Jurassic) of the Causses (southern France). Palaeogeography, Palaeoclimatology, Palaeoecology, 1995, 113, 135-172.	2.3	41
9	Are oxygen and carbon isotopes of mollusc shells reliable palaeosalinity indicators in marginal marine environments? A case study from the Middle Jurassic of England. Journal of the Geological Society, 1997, 154, 321-333.	2.1	43
10	Evidence and cause of small size in Bathonian (Middle Jurassic) marine bivalves of north-western Europe. Palaeontology, 1999, 42, 605-624.	2.2	10
11	87Sr/86Sr and Sr/Ca Investigation of Jurassic mollusks from Scotland: Implications for paleosalinities and the Sr/Ca ratio of seawater. Bulletin of the Geological Society of America, 2003, 115, 1249.	3.3	36
12	Key Events in the Ecological Radiation of the Ostracoda. The Paleontological Society Papers, 2003, 9, 181-202.	0.6	37
13	Theropod dinosaur and other footprints from the Valtos Sandstone Formation (Bathonian, Middle) Tj ETQq0 0 0	rgBT /Ovei 0.1	rlock 10 Tf 50
14	Dinosaur Tracks from the Kilmaluag Formation (Bathonian, Middle Jurassic) of Score Bay, Isle of Skye, Scotland, UK. Ichnos, 2005, 12, 93-104.	0.5	24
15	Early Jurassic shell beds from marginal marine environments in southern Mendoza, Argentina. Palaeogeography, Palaeoclimatology, Palaeoecology, 2007, 250, 68-88.	2.3	9
16	A comparison between dinosaur footprints from the Middle Jurassic of the Isle of Skye, Scotland, UK, and Shell, Wyoming, USA. Scottish Journal of Geology, 2008, 44, 139-150.	0.1	11
17	Reconstruction of Jurassic (Bathonian) palaeosalinity using stable isotopes and faunal associations. Journal of the Geological Society, 2010, 167, 1199-1208.	2.1	23
18	An unusual small-bodied crocodyliform from the Middle Jurassic of Scotland, UK, and potential evidence for an early diversification of advanced neosuchians. Earth and Environmental Science Transactions of the Royal Society of Edinburgh, 2016, 107, 1-12.	0.3	11

**CITATION REPORT** ARTICLE IF CITATIONS Sauropod dinosaur trackways in a Middle Jurassic lagoon on the Isle of Skye, Scotland. Scottish 0.1 19 Journal of Geology, 2016, 52, 1-9. TAPHONOMY AND PALEOECOLOGY OF LOWER CRETACEOUS OYSTER MASS OCCURRENCES FROM WEST-CENTRAL ARGENTINA AND EVOLUTIONARY PALEOECOLOGY OF GREGARIOUSNESS IN OYSTERS. 1.3 Palaios, 2018, 33, 237-255. Role of pressure solution in the formation of bedding-parallel calcite veins in an immature shale 1.5 15 (Cretaceous, southern UK). Geological Magazine, 2019, 156, 918-934. Taxonomic revision and palaeobiogeographic affinities of Berriasian–Valanginian oysters from the Vaca Muerta and Mulichinco formations, southern Mendoza, Neuquà ©n Basin, Argentina. Cretaceous Research, 2020, 109, 104358. Bay-head deltas as hydrocarbon reservoirs: The Middle Jurassic Hugin Formation in Block 15/3 of the 3.3 2 South Viking Graben, Norway. Marine and Petroleum Geology, 2021, 126, 104841. Geology and petroleum prospectivity of the Sea of Hebrides Basin and Minch Basin, offshore NW Scotland. Petroleum Geoscience, 2021, 27, . 1.5 Back-reacted saponite in Jurassic mudstones and limestones intruded by a Tertiary sill, Isle of Skye. 0.6 6 Clay Minerals, 2005, 40, 263-282. Environmental controls on biotic distribution., 1998, , 24-65. Fossils as environmental indicators., 1998, , 179-217. 1

28The Lonfearn Member, Lealt Shale Formation, (Middle Jurassic) of the Inner Hebrides, Scotland.0.1329A global synthesis of ecosystem services provided and disrupted by freshwater bivalve molluscs.<br/>Biological Reviews, 2022, 97, 1967-1998.10.42830A Comparison of the Dinosaur Communities from the Middle Jurassic of the Cleveland (Yorkshire) and<br/>Hebrides (Skye) Basins, Based on Their Ichnites. Geosciences (Switzerland), 2018, 8, 327.2.24

 $_{31}$  Sedimentary evolution and distribution of benthic fauna of an Aptian protected bay (Oliete Subbasin,) Tj ETQq0 0 0 rg BT /Overlock 10 T

19

21

23

25

26

27