

Hydrophilic-interaction chromatography for the separation of other polar compounds

Journal of Chromatography A

499, 177-196

DOI: [10.1016/S0021-9673\(00\)96972-3](https://doi.org/10.1016/S0021-9673(00)96972-3)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Rapid metal-interaction chromatography of proteins and peptides on micropellicular sorbents. <i>Chromatographia</i> , 1990, 30, 484-488.	0.7	14
2	Assay of tyrosine protein kinase activity from HL-60 by high-performance liquid chromatography for specificity studies. <i>Analytical Biochemistry</i> , 1990, 190, 32-38.	1.1	17
3	The role of lactic acid in autocrine B-cell growth stimulation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 11081-11085.	3.3	28
4	Hydrophilic-interaction chromatography of peptides on hydrophilic and strong cation-exchange columns. <i>Journal of Chromatography A</i> , 1991, 548, 13-24.	1.8	102
5	Poly(methylglutamate)-coated surfaces in HPLC and CE. <i>Chromatographia</i> , 1991, 32, 171-178.	0.7	44
6	Chapter 13 Amino acids and peptides. <i>Journal of Chromatography Library</i> , 1992, 51, B75-B150.	0.1	4
7	Characterization of recombinant antistasin secreted by <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 1992, 3, 185-195.	0.6	15
8	Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase liquid chromatography for the separation of peptides. <i>Journal of Chromatography A</i> , 1992, 594, 75-86.	1.8	88
9	Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity. <i>Biomedical Applications</i> , 1992, 583, 137-143.	1.7	35
10	Separation of glucose polymers by hydrophilic interaction chromatography on aqueous size-exclusion columns using gradient elution with pulsed amperometric detection. <i>Journal of Chromatography A</i> , 1992, 607, 7-18.	1.8	11
11	Macromolecular prodrugs. II. Esters of l-dopa and \pm -methyldopa. <i>International Journal of Pharmaceutics</i> , 1993, 99, 135-143.	2.6	19
12	Studies of the potency of protein kinase inhibitors on ATPase activities. <i>Chemico-Biological Interactions</i> , 1993, 86, 17-27.	1.7	14
13	Separation of glucooligosaccharides and polysaccharide hydrolysates by gradient elution hydrophilic interaction chromatography with pulsed amperometric detection. <i>Journal of Chromatography A</i> , 1993, 630, 129-139.	1.8	34
14	Pharmacologic effects in vivo in brain by vector-mediated peptide drug delivery.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 2618-2622.	3.3	150
15	Protein import into yeast mitochondria: the inner membrane import site protein ISP45 is the MPI1 gene product.. <i>EMBO Journal</i> , 1993, 12, 3035-3041.	3.5	78
16	Preparation of amino-zirconia bonded phases and their evaluation in hydrophilic interaction chromatography of carbohydrates with pulsed amperometric detection. <i>Journal of High Resolution Chromatography</i> , 1994, 17, 773-778.	2.0	21
17	High-performance liquid chromatography-based assays of enzyme activities. <i>Biomedical Applications</i> , 1994, 656, 143-157.	1.7	12
18	Hydrophilic-interaction chromatography of complex carbohydrates. <i>Journal of Chromatography A</i> , 1994, 676, 191-202.	1.8	230

#	ARTICLE	IF	CITATIONS
19	Analysis of Protein Hydrolyzates. 1. Use of Poly(2-hydroxyethylaspartamide)-Silica Column in Size Exclusion Chromatography for the Fractionation of Casein Hydrolyzates. <i>Journal of Agricultural and Food Chemistry</i> , 1994, 42, 2778-2782.	2.4	59
21	[16] Isolation of protein import components from <i>Saccharomyces cerevisiae</i> mitochondria. <i>Methods in Enzymology</i> , 1995, 260, 232-241.	0.4	5
22	Strategies for the removal of ionic and non-ionic detergents from protein and peptide mixtures for on- and off-line liquid chromatography mass spectrometry (LCMS). <i>Techniques in Protein Chemistry</i> , 1995, 6, 267-275.	0.3	11
23	Retention and selectivity of flavanones on homopolypeptide-bonded stationary phases in both normal- and reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1995, 704, 289-305.	1.8	29
24	Normal-phase high-performance liquid chromatography with relay gradient elution I. Description of the method. <i>Journal of Chromatography A</i> , 1995, 696, 193-199.	1.8	16
25	Chromatographic Techniques for the Characterization of Proteins. <i>Pharmaceutical Biotechnology</i> , 1995, 7, 243-299.	0.3	3
26	Chapter 3 High Performance Hydrophilic Interaction Chromatography of Carbohydrates with Polar Sorbents. <i>Journal of Chromatography Library</i> , 1995, 58, 103-146.	0.1	12
27	[1] Analysis of peptides by high-performance liquid chromatography. <i>Methods in Enzymology</i> , 1996, 271, 3-50.	0.4	53
28	Separation of acetylated core histones by hydrophilic-interaction liquid chromatography. <i>Journal of Chromatography A</i> , 1996, 743, 137-144.	1.8	109
29	A Rapid High-Resolution High-Performance Liquid Chromatographic Method for Separating Glycan Mixtures and Analyzing Oligosaccharide Profiles. <i>Analytical Biochemistry</i> , 1996, 240, 210-226.	1.1	524
30	Improved protocol for the formation of N-(p-nitrobenzyloxy)aminoalditol derivatives of oligosaccharides. <i>Carbohydrate Research</i> , 1996, 282, 1-12.	1.1	24
31	Recent progress in carbohydrate separation by high-performance liquid chromatography based on hydrophilic interaction. <i>Journal of Chromatography A</i> , 1996, 720, 75-91.	1.8	173
32	Hydrophilic interaction chromatography on amino-silica phases complements reversed-phase high-performance liquid chromatography and capillary electrophoresis for peptide analysis. <i>Journal of Chromatography A</i> , 1996, 724, 378-383.	1.8	119
33	Enantioseparation in the synthesis of myo-inositol phosphates by high-performance liquid chromatography using a β -cyclodextrin-bonded column. <i>Journal of Chromatography A</i> , 1996, 749, 25-32.	1.8	5
34	Plant Cell Wall Analysis. <i>Modern Methods of Plant Analysis</i> , 1996, , .	0.1	11
35	Separation of High Molecular Weight Saponins of <i>Archidendron Ellipticum</i> by Hydrophilic Interaction Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1997, 20, 2415-2426.	0.5	13
36	Optimal Subsite Occupancy and Design of a Selective Inhibitor of Urokinase. <i>Journal of Biological Chemistry</i> , 1997, 272, 20456-20462.	1.6	85
37	Distinguishing the Specificities of Closely Related Proteases. <i>Journal of Biological Chemistry</i> , 1997, 272, 16603-16609.	1.6	50

#	ARTICLE	IF	CITATIONS
38	Analytic and Immunologic Characterization of Chickpea(<i>Cicer arietinum</i>)Protein Hydrolysates Obtained by Bromelain and $\hat{1}\pm$ -Chymotrypsin. <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 4758-4762.	2.4	7
39	[19] Analysis of synthetic peptides by high-performance liquid chromatography. <i>Methods in Enzymology</i> , 1997, 289, 426-469.	0.4	40
40	Peptide Separation in Normal Phase Liquid Chromatography. <i>Analytical Chemistry</i> , 1997, 69, 3038-3043.	3.2	139
41	Application of hydrophilic-interaction liquid chromatography to the separation of phosphorylated H1 histones. <i>Journal of Chromatography A</i> , 1997, 782, 55-62.	1.8	91
42	Strong cation-exchange high-performance liquid chromatography as a versatile tool for the characterization and purification of peptides. <i>Analytica Chimica Acta</i> , 1997, 352, 21-30.	2.6	20
43	Synthesis and purification of hydrophobic peptides for use in biomimetic ion channels. <i>Journal of Chromatography A</i> , 1998, 816, 11-20.	1.8	7
44	Hydrophilic interaction/cation-exchange chromatography for separation of cyclic peptides. <i>Journal of Chromatography A</i> , 1998, 816, 79-88.	1.8	59
45	Ion-exchange chromatography of proteins: modulation of selectivity by addition of organic solvents to mobile phase. <i>Journal of Chromatography A</i> , 1998, 825, 17-27.	1.8	3
46	Effect of polystyreneâ€“divinylbenzene resin sulfonation on solute retention in high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 1998, 797, 139-147.	1.8	48
47	Separation of a fructo-oligosaccharide mixture by hydrophilic interaction chromatography using silica-based micropellicular sorbents. <i>Journal of Chromatography A</i> , 1998, 803, 302-306.	1.8	30
48	Calculation of peptide retention coefficients in normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1998, 808, 105-112.	1.8	55
49	Prediction of peptide retention time in normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1998, 811, 61-67.	1.8	33
50	Hydrophilic interaction/cation-exchange chromatography for separation of amphipathic $\hat{1}\pm$ -helical peptides. <i>Journal of Chromatography A</i> , 1998, 816, 65-78.	1.8	66
51	Revised primary structures of rat pituitary $\hat{1}^3$ -lipotrophin and $\hat{1}^2$ -endorphin. <i>Neuropeptides</i> , 1998, 32, 339-349.	0.9	6
52	Quantitative analysis and process monitoring of site-specific glycosylation microheterogeneity in recombinant human interferon- $\hat{1}^3$ from Chinese hamster ovary cell culture by hydrophilic interaction chromatography. <i>Biomedical Applications</i> , 1998, 712, 73-82.	1.7	47
53	Practice and mechanism of HPLC oligosaccharide separation with a cyclodextrin bonded phase. <i>Talanta</i> , 1998, 47, 1001-1012.	2.9	72
54	Hydrophilic Interaction Chromatographyâ€“Electrospray Mass Spectrometry Analysis of Polar Compounds for Natural Product Drug Discovery. <i>Analytical Chemistry</i> , 1998, 70, 2439-2445.	3.2	208
55	Carbohydrates, Polymeric Sugars and their Constituents. <i>Journal of Chromatography Library</i> , 1998, , 257-314.	0.1	1

#	ARTICLE	IF	CITATIONS
56	Size Exclusion High-Performance Liquid Chromatography of Small Solutes. , 1999, , 249-266.		12
57	Hydrophilic interaction/cation-exchange chromatography for the purification of synthetic peptides from closely related impurities: serine side-chain acetylated peptides. <i>Chemical Biology and Drug Design</i> , 1999, 54, 1-11.	1.2	32
58	High-performance liquid chromatographicâ€“electrospray ionization mass spectrometric analyses for the integration of natural products with modern high-throughput screening. <i>Biomedical Applications</i> , 1999, 725, 67-78.	1.7	89
59	Peptide separation in normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 1999, 840, 1-9.	1.8	48
60	Prediction of peptide retention times in normal-phase liquid chromatography with only a single gradient run. <i>Journal of Chromatography A</i> , 1999, 841, 19-32.	1.8	13
61	Chromatographic Comparison of the UVA Sensitizers Present in Brunescant Cataracts and in Calf Lens Proteins Ascorbylated in vitro. <i>Experimental Eye Research</i> , 1999, 69, 375-384.	1.2	17
63	Determination of urea, allantoin and lysine pyroglutamate in cosmetic samples by hydrophilic interaction chromatography. <i>Biomedical Applications</i> , 2000, 742, 447-452.	1.7	59
64	Resolution of allelic and non-allelic variants of histone H1 by cation-exchange-hydrophilic-interaction chromatography. <i>Biomedical Applications</i> , 2000, 744, 33-46.	1.7	30
65	Calculation of amino acid hydrophilicity indices for retention of peptides on amide, diol and silica columns in normal-phase liquid chromatography. <i>Chromatographia</i> , 2000, 52, 418-424.	0.7	15
66	Chapter 1 Comparison of various modes and phase systems for analytical HPLC. <i>Handbook of Analytical Separations</i> , 2000, , 1-71.	0.8	6
67	Physicochemical factors in polypeptide and protein purification and analysis by high-performance liquid chromatographic techniques: current status and challenges for the future. <i>Separation Science and Technology</i> , 2000, 2, 71-235.	0.0	17
69	Mixed-Mode Anionâˆ“Cation Exchange/Hydrophilic Interaction Liquid Chromatographyâˆ“Electrospray Mass Spectrometry as an Alternative to Reversed Phase for Small Molecule Drug Discovery. <i>Analytical Chemistry</i> , 2000, 72, 4629-4633.	3.2	148
70	Chiral Separations of Polar Compounds by Hydrophilic Interaction Chromatography with Evaporative Light Scattering Detection. <i>Analytical Chemistry</i> , 2000, 72, 1736-1739.	3.2	91
71	CARBOHYDRATES Liquid Chromatography. , 2000, , 2224-2235.		1
72	HPLC of Peptides and Proteins. <i>Current Protocols in Protein Science</i> , 2001, 23, Unit8.7.	2.8	10
73	HPLC of Peptides and Proteins: Standard Operating Conditions. , 2001, Chapter 10, Unit 10.13.		1
74	Î²-Cyclodextrin dimers as potential tumor pretargeting agents. <i>Chemical Communications</i> , 2001, , 1312-1313.	2.2	15
75	Determination of Foliates in Human Plasma Using Hydrophilic Interaction Chromatographyâˆ“Tandem Mass Spectrometry. <i>Analytical Chemistry</i> , 2001, 73, 5358-5364.	3.2	108

#	ARTICLE	IF	CITATIONS
76	Silica and other materials as supports in liquid chromatography. Chromatographic tests and their importance for evaluating these supports. Part I. <i>Chromatographia</i> , 2001, 53, S113-S131.	0.7	74
77	Liquid chromatography ion trap mass spectrometric analysis of oligosaccharides using permethylated derivatives. <i>Rapid Communications in Mass Spectrometry</i> , 2001, 15, 325-334.	0.7	51
78	The use of silica for liquid chromatographic/mass spectrometric analysis of basic analytes. <i>Journal of Separation Science</i> , 2001, 24, 835-842.	1.3	30
79	Analysis of glucosinolates from broccoli and other cruciferous vegetables by hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2001, 919, 299-304.	1.8	63
80	Hydrophilic interaction chromatography using amino and silica columns for the determination of polar pharmaceuticals and impurities. <i>Journal of Chromatography A</i> , 2001, 913, 113-122.	1.8	251
81	Age-dependent deamidation of asparagine residues in proteins. <i>Experimental Gerontology</i> , 2001, 36, 1551-1563.	1.2	109
82	Effects of -aminopropionitrile on equine tendon metabolism in vitro and on effects of insulin-like growth factor-I on matrix production by equine tenocytes. <i>American Journal of Veterinary Research</i> , 2001, 62, 1557-1562.	0.3	28
83	Structural Investigations of Glycoconjugates at High Sensitivity. <i>Chemical Reviews</i> , 2002, 102, 321-370.	23.0	320
84	Chapter 4 High performance hydrophilic interaction chromatography of carbohydrates with polar sorbents. <i>Journal of Chromatography Library</i> , 2002, 66, 121-163.	0.1	11
85	Emprego de cromatografia líquida de alta eficiência hidrofílica na determinação dos aminoácidos de hidrolisados de caseína. <i>Food Science and Technology</i> , 2002, 22, 229-232.	0.8	4
86	Modeling and optimization for separation of ionic solutes in pressurized flow capillary electrochromatography. <i>Journal of Separation Science</i> , 2002, 25, 416-426.	1.3	27
87	Graphite powder as an alternative or supplement to reversed-phase material for desalting and concentration of peptide mixtures prior to matrix-assisted laser desorption/ionization-mass spectrometry. <i>Proteomics</i> , 2002, 2, 1277-1287.	1.3	129
88	New sample preparation technique for the determination of avoparcin in pressurized hot water extracts from kidney samples. <i>Journal of Chromatography A</i> , 2002, 954, 41-49.	1.8	51
89	Peptide separation in hydrophilic interaction capillary electrochromatography. <i>Electrophoresis</i> , 2003, 24, 2084-2091.	1.3	24
90	Analysis of the performance of antibody capture methods using fluorescent peptides with capillary zone electrophoresis with laser-induced fluorescence. <i>Electrophoresis</i> , 2003, 24, 892-896.	1.3	25
91	Chromatographic classification and comparison of commercially available perfluorinated stationary phases for reversed-phase liquid chromatography using Principal Component Analysis. <i>Journal of Separation Science</i> , 2003, 26, 295-306.	1.3	81
92	Development and validation of a liquid chromatography-tandem mass spectrometry method for the determination of pyridostigmine bromide from guinea pig plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2003, 796, 347-354.	1.2	10
93	Bioanalytical liquid chromatography tandem mass spectrometry methods on underivatized silica columns with aqueous/organic mobile phases. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2003, 796, 209-224.	1.2	243

#	ARTICLE	IF	CITATIONS
94	Characterization of a novel diol column for high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2003, 987, 119-125.	1.8	74
95	Comparison of reversed-phase liquid chromatography and hydrophilic interaction/cation-exchange chromatography for the separation of amphipathic L±-helical peptides with l- and d-amino acid substitutions in the hydrophilic face. <i>Journal of Chromatography A</i> , 2003, 1009, 61-71.	1.8	35
96	Determination of a small, highly polar aminopropylphosphinic acid as racemate in plasma and urine and as separated enantiomers in plasma by liquid chromatography and tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2003, 492, 261-269.	2.6	7
97	A HILIC method for the analysis of tromethamine as the counter ion in an investigational pharmaceutical salt. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2003, 31, 1191-1201.	1.4	66
98	Fungal metabolite screening: database of 474 mycotoxins and fungal metabolites for dereplication by standardised liquid chromatography-UV-mass spectrometry methodology. <i>Journal of Chromatography A</i> , 2003, 1002, 111-136.	1.8	394
99	Novel Method for the Determination of Piperazine in Pharmaceutical Drug Substances Using Hydrophilic Interaction Chromatography and Evaporative Light Scattering Detection. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2003, 26, 3093-3104.	0.5	36
100	Hydrophilic Interaction Liquid Chromatography Coupled to Electrospray Mass Spectrometry of Small Polar Compounds in Food Analysis. <i>Analytical Chemistry</i> , 2003, 75, 2349-2354.	3.2	142
101	Purification and Analyses of Histone H1 Variants and H1 Posttranslational Modifications. <i>Methods in Enzymology</i> , 2003, 375, 278-293.	0.4	8
102	Polar Polymeric Stationary Phases for Normal-Phase HPLC Based on Monodisperse Macroporous Poly(2,3-dihydroxypropyl methacrylate-co-ethylene dimethacrylate) Beads. <i>Analytical Chemistry</i> , 2003, 75, 1011-1021.	3.2	42
103	Isolation and Identification of Indigestible Pyroglutamyl Peptides in an Enzymatic Hydrolysate of Wheat Gluten Prepared on an Industrial Scale. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 8-13.	2.4	45
106	Hydrophilic Interaction Chromatography. , 2004, 251, 75-88.		3
107	Determination of the Stabilizer Sucrose in a Plasma-Derived Antithrombin Process Solution by Hydrophilic Interaction Chromatography with Evaporative Light-Scattering Detection. <i>Journal of Chromatographic Science</i> , 2004, 42, 361-365.	0.7	10
108	Histone H4 Hyperacetylation Precludes Histone H4 Lysine 20 Trimethylation. <i>Journal of Biological Chemistry</i> , 2004, 279, 53458-53464.	1.6	61
109	Chapter 18 Carbohydrates. <i>Journal of Chromatography Library</i> , 2004, , 839-904.	0.1	1
110	Analysis of Five Pharmacologically Active Compounds from <i>Rhodiola</i> for Natural Product Drug Discovery with Capillary Electrophoresis. <i>Chromatographia</i> , 2004, 60, 589-595.	0.7	14
111	Capillary electrochromatography with monolithic silica columns. <i>Journal of Chromatography A</i> , 2004, 1029, 239-247.	1.8	62
112	Determination of basic pharmaceuticals in human serum by hydrophilic interaction capillary electrochromatography. <i>Electrophoresis</i> , 2004, 25, 600-606.	1.3	20
113	Polar stationary phases for capillary electrochromatography. <i>Electrophoresis</i> , 2004, 25, 4095-4109.	1.3	28

#	ARTICLE	IF	CITATIONS
114	Determination of neomycin by LC-tandem mass spectrometry using hydrophilic interaction chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2004, 35, 633-638.	1.4	67
115	Chromatographic behavior of epirubicin and its analogues on high-purity silica in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2004, 1041, 163-169.	1.8	90
116	Quantitative determination of dityrosine in milk powders by liquid chromatography coupled to tandem mass spectrometry using isotope dilution. <i>Journal of Chromatography A</i> , 2004, 1052, 77-84.	1.8	35
117	Implications of column peak capacity on the separation of complex peptide mixtures in single- and two-dimensional high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2004, 1061, 183-192.	1.8	157
118	Determination of 1-Deoxynojirimycin in Mulberry Leaves Using Hydrophilic Interaction Chromatography with Evaporative Light Scattering Detection. <i>Journal of Agricultural and Food Chemistry</i> , 2004, 52, 1415-1418.	2.4	84
119	A New Strategy for Identification of N-Glycosylated Proteins and Unambiguous Assignment of Their Glycosylation Sites Using HILIC Enrichment and Partial Deglycosylation. <i>Journal of Proteome Research</i> , 2004, 3, 556-566.	1.8	452
120	Preparation and characterization of perhydroxyl-cucurbit[6]uril bonded silica stationary phase for hydrophilic-interaction chromatography. <i>Talanta</i> , 2004, 64, 929-934.	2.9	54
121	Peptide separation by Hydrophilic-Interaction Chromatography: a review. <i>Journal of Proteomics</i> , 2004, 60, 265-280.	2.4	224
122	Isotope-Coded Affinity Tag (ICAT) Approach to Redox Proteomics: Identification and Quantitation of Oxidant-Sensitive Cysteine Thiols in Complex Protein Mixtures. <i>Journal of Proteome Research</i> , 2004, 3, 1228-1233.	1.8	168
123	Hydrophilic Interaction Chromatography/Electrospray Ionization Mass Spectrometry for Determination of Allantoin in Human Biological Samples. <i>Bunseki Kagaku</i> , 2005, 54, 135-142.	0.1	8
124	Metabolic fingerprinting of rat urine by LC/MS Part 1. Analysis by hydrophilic interaction liquid chromatography-electrospray ionization mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2005, 828, 9-13.	1.2	137
125	Solute attributes and molecular interactions contributing to U-shape-retention on a fluorinated high-performance liquid chromatography stationary phase. <i>Journal of Chromatography A</i> , 2005, 1073, 99-109.	1.8	93
126	Evaluation of a monolithic silica column operated in the hydrophilic interaction chromatography mode with evaporative light scattering detection for the separation and detection of counter-ions. <i>Journal of Chromatography A</i> , 2005, 1073, 269-275.	1.8	68
127	Retention behavior of small polar compounds on polar stationary phases in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2005, 1074, 71-80.	1.8	436
128	Development and validation of a hydrophilic interaction chromatography-mass spectrometry assay for taurine and methionine in matrices rich in carbohydrates. <i>Journal of Chromatography A</i> , 2005, 1081, 174-181.	1.8	36
129	Hydrophilic interaction liquid chromatography-mass spectrometry for the analysis of paralytic shellfish poisoning (PSP) toxins. <i>Journal of Chromatography A</i> , 2005, 1081, 190-201.	1.8	246
130	Orthogonal method development using hydrophilic interaction chromatography and reversed-phase high-performance liquid chromatography for the determination of pharmaceuticals and impurities. <i>Journal of Chromatography A</i> , 2005, 1083, 58-62.	1.8	99
131	Alternative high-performance liquid chromatographic peptide separation and purification concept using a new mixed-mode reversed-phase/weak anion-exchange type stationary phase. <i>Journal of Chromatography A</i> , 2005, 1089, 158-169.	1.8	108

#	ARTICLE	IF	CITATIONS
132	Separation of monosaccharides by hydrophilic interaction chromatography with evaporative light scattering detection. <i>Journal of Chromatography A</i> , 2005, 1092, 246-249.	1.8	121
133	Influence of mobile phase composition and thermodynamics on the normal phase chromatography of echinocandins. <i>Journal of Chromatography A</i> , 2005, 1098, 55-65.	1.8	10
134	Potential Use of an Aminopropyl Stationary Phase in Hydrophilic Interaction Capillary Electrochromatography. Application to Tetracycline Antibiotics. <i>Chromatographia</i> , 2005, 62, 393-399.	0.7	13
135	Separation of proteins with a molecular mass difference of 2 kDa utilizing preparative double-inverted gradient polyacrylamide gel electrophoresis under nonreducing conditions: Application to the isolation of 24 kDa human growth hormone. <i>Electrophoresis</i> , 2005, 26, 4389-4395.	1.3	7
136	New hyphenated methodologies in high-sensitivity glycoprotein analysis. <i>Journal of Separation Science</i> , 2005, 28, 1956-1968.	1.3	100
137	Hydrophilic interaction liquid chromatography with tandem mass spectrometry for the determination of underivatized dencichine (β^2 -N-oxalyl-L- β - β^2 -diaminopropionic acid) in Panax medicinal plant species. <i>Rapid Communications in Mass Spectrometry</i> , 2005, 19, 1237-1244.	0.7	49
138	Hydrophilic interaction liquid chromatography/mass spectrometry for determination of domoic acid in Adriatic shellfish. <i>Rapid Communications in Mass Spectrometry</i> , 2005, 19, 2030-2038.	0.7	62
139	Analysis of acetylcholine and choline in microdialysis samples by liquid chromatography/tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2005, 19, 2950-2956.	0.7	100
140	O-Linked glycosylation in maize-expressed human IgA1. <i>Glycobiology</i> , 2005, 15, 965-981.	1.3	81
141	10 Free-flow isoelectric focusing. <i>Separation Science and Technology</i> , 2005, 7, 211-245.	0.0	3
142	6 HPLC method development. <i>Separation Science and Technology</i> , 2005, 6, 145-190.	0.0	5
143	Orthogonality of Separation in Two-Dimensional Liquid Chromatography. <i>Analytical Chemistry</i> , 2005, 77, 6426-6434.	3.2	726
144	Analysis of Quaternary Amine Compounds by Hydrophilic Interaction Chromatography/Mass Spectrometry (HILIC/MS). <i>Journal of Liquid Chromatography and Related Technologies</i> , 2005, 28, 497-512.	0.5	25
145	Single DNA Molecules as Probes of Chromatographic Surfaces. <i>Analytical Chemistry</i> , 2005, 77, 3256-3260.	3.2	21
146	A hydrophilic interaction chromatography coupled to a mass spectrometry for the determination of glutathione in plant somatic embryos. <i>Analyst, The</i> , 2006, 131, 1167-1174.	1.7	83
147	Speciation of non-covalent nickel species in plant tissue extracts by electrospray Q-TOFMS/MS after their isolation by 2D size exclusion-hydrophilic interaction LC (SEC-HILIC) monitored by ICP-MS. <i>Journal of Analytical Atomic Spectrometry</i> , 2006, 21, 676-683.	1.6	72
148	Isolation and Characterization of Glycosylphosphatidylinositol-Anchored Peptides by Hydrophilic Interaction Chromatography and MALDI Tandem Mass Spectrometry. <i>Analytical Chemistry</i> , 2006, 78, 3335-3341.	3.2	57
149	Hydrophilic Interaction Liquid Chromatographic (HILIC)/Ion Exchange Separation of Picolinic and Nicotinic Acids. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2006, 29, 2545-2558.	0.5	14

#	ARTICLE	IF	CITATIONS
150	Evaluation of a Preclinical Blood Test for Scrapie in Sheep Using Immunocapillary Electrophoresis. <i>Journal of AOAC INTERNATIONAL</i> , 2006, 89, 720-727.	0.7	13
151	Chapter 3.1 Liquid chromatographic methods used for microdialysis: an overview. <i>Handbook of Behavioral Neuroscience</i> , 2006, , 233-250.	0.7	1
153	Two-dimensional liquid chromatography normal-phase and reversed-phase separation of (co)oligomers. <i>Journal of Chromatography A</i> , 2006, 1119, 3-10.	1.8	69
154	Development of a hydrophilic interaction chromatography-mass spectrometry assay for spectinomycin and lincomycin in liquid hog manure supernatant and run-off from cropland. <i>Journal of Chromatography A</i> , 2006, 1107, 152-158.	1.8	61
155	Separation of isomeric 2-aminopyridine derivatized N-glycans and N-glycopeptides of human serum immunoglobulin G by using a zwitterionic type of hydrophilic-interaction chromatography. <i>Journal of Chromatography A</i> , 2006, 1113, 177-181.	1.8	120
156	Can the theory of gradient liquid chromatography be useful in solving practical problems?. <i>Journal of Chromatography A</i> , 2006, 1126, 195-218.	1.8	85
157	Zwitterionic stationary phase with covalently bonded phosphorylcholine type polymer grafts and its applicability to separation of peptides in the hydrophilic interaction liquid chromatography mode. <i>Journal of Chromatography A</i> , 2006, 1127, 82-91.	1.8	152
158	Separation and identification of phytosiderophores and their metal complexes in plants by zwitterionic hydrophilic interaction liquid chromatography coupled to electrospray ionization mass spectrometry. <i>Journal of Chromatography A</i> , 2006, 1136, 73-81.	1.8	70
159	Analysis of glutathione and glutathione disulfide in human saliva using hydrophilic interaction chromatography with mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2006, 839, 74-79.	1.2	72
160	Miniaturized separation techniques in glycomic investigations. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2006, 841, 65-78.	1.2	68
161	Chromatographic and electrophoretic characterization of protein variants. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2006, 841, 110-122.	1.2	52
162	Determination of Atenolol in Human Plasma by Pseudo Reversed Phase Liquid Chromatography-Tandem Mass Spectrometry. <i>Chromatographia</i> , 2006, 64, 583-586.	0.7	8
163	Simple and Rapid Determination of Denaturants in Alcohol Formulations by Hydrophilic Interaction Chromatography. <i>Chromatographia</i> , 2006, 63, 373-377.	0.7	16
164	Amino acid analysis by hydrophilic interaction chromatography coupled on-line to electrospray ionization mass spectrometry. <i>Amino Acids</i> , 2006, 30, 291-297.	1.2	108
165	HILIC mode separation of polar compounds by monolithic silica capillary columns coated with polyacrylamide. <i>Analytical and Bioanalytical Chemistry</i> , 2006, 386, 578-585.	1.9	82
166	Monolithic silica columns with mixed mode of hydrophilic interaction and weak anion-exchange stationary phase for pressurized capillary electrochromatography. <i>Electrophoresis</i> , 2006, 27, 3373-3380.	1.3	44
167	Analysis of mannitol in pharmaceutical formulations using hydrophilic interaction liquid chromatography with evaporative light-scattering detection. <i>Journal of Separation Science</i> , 2006, 29, 256-264.	1.3	37
168	Simple separation of isomeric sialylated N-glycopeptides by a zwitterionic type of hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2006, 29, 2533-2540.	1.3	134

#	ARTICLE	IF	CITATIONS
169	Hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2006, 29, 1784-1821.	1.3	1,053
170	ICAT (Isotope-Coded Affinity Tag) Approach to Redox Proteomics: Identification and Quantification of Oxidant-Sensitive Protein Thiols. , 2006, , 267-285.		2
174	Definition of the mitochondrial proteome by measurement of molecular masses of membrane proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 16170-16175.	3.3	83
175	Hydrophilic Interaction Chromatographic Determination of Epirubicin in Human Plasma using Solid Phase Extraction for Sample Cleanâ€šUp. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 2409-2418.	0.5	7
176	12 HPLC method development in early phase pharmaceutical development. <i>Separation Science and Technology</i> , 2007, , 353-371.	0.0	3
177	A novel chromatography system to isolate active ribosomes from pathogenic bacteria. <i>Rna</i> , 2008, 14, 188-195.	1.6	28
178	Separation of Polar Mushroom Toxins by Mixed-Mode Hydrophilic and Ionic Interaction Liquid Chromatography–Electrospray Ionization–Mass Spectrometry. <i>Journal of Chromatographic Science</i> , 2007, 45, 104-111.	0.7	40
179	Optimization of Enzyme Assisted Processes for Extracting and Hydrolysing Corn Proteins Aiming Phenylalanine Removal. <i>International Journal of Food Engineering</i> , 2007, 3, .	0.7	18
180	Rapid and Individual-specific Glycoprofiling of the Low Abundance N-Glycosylated Protein Tissue Inhibitor of Metalloproteinases-1. <i>Molecular and Cellular Proteomics</i> , 2007, 6, 638-647.	2.5	52
181	Determination of Diquat in Water Samples by Hydrophilic Interaction Chromatography/Electrospray Ionization/Mass Spectrometry Coupled with Solid-Phase Extraction Using a Stable Isotope Labeled Compound. <i>Bunseki Kagaku</i> , 2007, 56, 579-585.	0.1	6
182	The effects of arginine on protein binding and elution in hydrophobic interaction and ion-exchange chromatography. <i>Protein Expression and Purification</i> , 2007, 54, 110-116.	0.6	94
183	HPLC Analysis and Purification of Peptides. <i>Methods in Molecular Biology</i> , 2007, 386, 3-55.	0.4	36
184	Peptide Characterization and Application Protocols. <i>Methods in Molecular Biology</i> , 2007, , .	0.4	0
186	Application of Humic Acid Bondedâ€šSilica as a Hydrophilicâ€šInteraction Chromatographic Stationary Phase in Separation of Polar Compounds. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 31, 64-78.	0.5	14
187	Hydrophilic Interaction Chromatography Using Methacrylate-Based Monolithic Capillary Column for the Separation of Polar Analytes. <i>Analytical Chemistry</i> , 2007, 79, 1243-1250.	3.2	177
188	Analysis of Moniliformin in Maize Plants Using Hydrophilic Interaction Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 9764-9768.	2.4	33
189	Capillary LC Coupled with High-Mass Measurement Accuracy Mass Spectrometry for Metabolic Profiling. <i>Analytical Chemistry</i> , 2007, 79, 6081-6093.	3.2	47
190	Screening of Underivatized Oligosaccharides Extracted from the Stems of <i>Triticum aestivum</i> Using Porous Graphitized Carbon Liquid Chromatographyâ€šMass Spectrometry. <i>Analytical Chemistry</i> , 2007, 79, 2437-2445.	3.2	45

#	ARTICLE	IF	CITATIONS
191	Hydrophilic Chromatographic Determination of Carnosine, Anserine, Balenine, Creatine, and Creatinine. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 4664-4669.	2.4	107
192	Occurrence of Orally Administered Mulberry 1-Deoxynojirimycin in Rat Plasma. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 8928-8933.	2.4	63
193	â€œClick saccharidesâ€ novel separation materials for hydrophilic interaction liquid chromatography. <i>Chemical Communications</i> , 2007, , 2491-2493.	2.2	179
194	Modification-Specific Proteomic Analysis of Glycoproteins in Human Body Fluids by Mass Spectrometry. , 2007, , 107-128.		3
195	Direct detection method of oligosaccharides by high-performance liquid chromatography with charged aerosol detection. <i>Biomedical Chromatography</i> , 2007, 21, 338-342.	0.8	39
196	Ion-interaction CZE: The presence of high concentrations of ion-pairing reagents demonstrates the complex mechanisms involved in peptide separations. <i>Electrophoresis</i> , 2007, 28, 2181-2190.	1.3	5
197	Hydrophilic interaction chromatography coupled to nuclear magnetic resonance spectroscopy and mass spectroscopyâ€”A new approach for the separation and identification of extremely polar analytes in bodyfluids. <i>Journal of Chromatography A</i> , 2007, 1156, 87-93.	1.8	48
198	Retention behavior of oligomeric proanthocyanidins in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2007, 1143, 153-161.	1.8	82
199	Arginine improves protein elution in hydrophobic interaction chromatography. <i>Journal of Chromatography A</i> , 2007, 1154, 81-86.	1.8	49
200	Preparation and evaluation of hydroxylated poly(glycidyl methacrylate-co-ethylene dimethacrylate) monolithic capillary for in-tube solid-phase microextraction coupled to high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2007, 1160, 90-98.	1.8	48
201	Hydrophilic interaction liquid chromatography for separation and quantification of selected room-temperature ionic liquids. <i>Journal of Chromatography A</i> , 2007, 1164, 139-144.	1.8	40
202	Highly efficient monolithic silica capillary columns modified with poly(acrylic acid) for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2007, 1164, 198-205.	1.8	78
203	Is hydrophilic interaction chromatography with silica columns a viable alternative to reversed-phase liquid chromatography for the analysis of ionisable compounds?. <i>Journal of Chromatography A</i> , 2007, 1171, 46-55.	1.8	242
204	Metabolic profiling of rat hair and screening biomarkers using ultra performance liquid chromatography with electrospray ionization time-of-flight mass spectrometry. <i>Journal of Chromatography A</i> , 2007, 1176, 94-99.	1.8	32
205	Retention prediction of adrenoreceptor agonists and antagonists on a diol column in hydrophilic interaction chromatography. <i>Analytica Chimica Acta</i> , 2007, 598, 41-50.	2.6	48
206	Development of a liquid chromatography/tandem mass spectrometry method for the quantitation of acetylcholine and related neurotransmitters in brain microdialysis samples. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2007, 44, 586-593.	1.4	69
207	Evaluation of the intermediates generated during the degradation of Diuron and Linuron herbicides by the photo-Fenton reaction. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2007, 189, 364-373.	2.0	53
208	The determination of organophosphonate nerve agent metabolites in human urine by hydrophilic interaction liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007, 852, 235-243.	1.2	86

#	ARTICLE	IF	CITATIONS
209	Determination of zanamivir in rat and monkey plasma by positive ion hydrophilic interaction chromatography (HILIC)/tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007, 852, 505-511.	1.2	32
210	Determination of ascorbic acid and its related compounds in foods and beverages by hydrophilic interaction liquid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007, 853, 214-220.	1.2	69
211	Development and validation of a selective and sensitive bioanalytical procedure for the quantitative determination of gaboxadol in human plasma employing mixed mode solid phase extraction and hydrophilic interaction liquid chromatography with tandem mass spectroscopic detection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007, 858, 168-176.	1.2	23
212	Capillary hydrophilic interaction chromatography/mass spectrometry for simultaneous determination of multiple neurotransmitters in primate cerebral cortex. <i>Rapid Communications in Mass Spectrometry</i> , 2007, 21, 3621-3628.	0.7	79
213	Hydroxymethyl methacrylate-based monolithic columns designed for separation of oligonucleotides in hydrophilic-interaction capillary liquid chromatography. <i>Journal of Proteomics</i> , 2007, 70, 23-29.	2.4	77
214	Evaluation and Optimization of ZIC-HILIC-RP as an Alternative MudPIT Strategy. <i>Journal of Proteome Research</i> , 2007, 6, 937-946.	1.8	182
215	New Approaches for Analysis of Metabolism Compounds in Hydrophilic Interaction Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 2117-2126.	0.5	24
216	Investigating the Effect of Chromatographic Conditions on Retention of Organic Acids in Hydrophilic Interaction Chromatography Using a Design of Experiment. <i>Chromatographia</i> , 2007, 66, 223-229.	0.7	101
217	Simultaneous determination of the advanced glycation end product N ^ε -carboxymethyllysine and its precursor, lysine, in exhaled breath condensate using isotope-dilution ¹³ C-hydrophilic-interaction liquid chromatography coupled to tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 387, 2783-2791.	1.9	30
218	Retention prediction of adrenoreceptor agonists and antagonists on unmodified silica phase in hydrophilic interaction chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 388, 1693-1706.	1.9	35
219	A new hydrophilic interaction liquid chromatographic (HILIC) procedure for the simultaneous determination of pseudoephedrine hydrochloride (PSH), diphenhydramine hydrochloride (DPH) and dextromethorphan hydrobromide (DXH) in cough-cold formulations. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2007, 43, 158-167.	1.4	57
220	Application of hydrophilic interaction chromatography for simultaneous separation of six impurities of mildronate substance. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2008, 48, 649-656.	1.4	38
221	Molecular mass analysis of antibodies by on-line SEC-MS. <i>Journal of the American Society for Mass Spectrometry</i> , 2008, 19, 502-509.	1.2	61
222	Hydrophilic interaction liquid chromatography (HILIC) in proteomics. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 151-159.	1.9	234
223	Application of at-line two-dimensional liquid chromatography ¹³ C-mass spectrometry for identification of small hydrophilic angiotensin I-inhibiting peptides in milk hydrolysates. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 299-307.	1.9	60
224	Highly efficient analysis of underivatized carbohydrates using monolithic-silica-based capillary hydrophilic interaction (HILIC) HPLC. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 2533-2542.	1.9	96
225	Analysis of amino acids without derivatization in barley extracts by LC-MS-MS. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 2663-2672.	1.9	95
226	Probing genetic algorithms for feature selection in comprehensive metabolic profiling approach. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 1312-1324.	0.7	25

#	ARTICLE	IF	CITATIONS
227	Hydrophilic interaction chromatography/electrospray mass spectrometry analysis of carbohydrate-related metabolites from <i>Arabidopsis thaliana</i> leaf tissue. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 1399-1407.	0.7	95
228	Quantification of acetylcholine in microdialysate of subcutaneous tissue by hydrophilic interaction chromatography/tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 1497-1502.	0.7	13
229	Determination of polar organophosphorus pesticides in water samples by hydrophilic interaction liquid chromatography with tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 2203-2210.	0.7	40
230	Liquid chromatography/mass spectrometry determination of endogenous plasma acetyl and palmitoyl carnitines as potential biomarkers of β -oxidation in mice. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 3434-3442.	0.7	23
231	Alternative organic solvents for HILIC separation of cisplatin species with online ICP-MS detection. <i>Journal of Separation Science</i> , 2008, 31, 599-603.	1.3	34
232	Stability-indicating assay of sodium cromoglicate in ophthalmic solution using mixed-mode hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1645-1650.	1.3	18
233	Development of retention prediction models for adrenoreceptor agonists and antagonists on a polyvinyl alcohol-bonded stationary phase in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1537-1549.	1.3	32
234	Mixed-mode hydrophilic interaction/cation-exchange chromatography: Separation of complex mixtures of peptides of varying charge and hydrophobicity. <i>Journal of Separation Science</i> , 2008, 31, 1573-1584.	1.3	42
235	Impact of column temperature and mobile phase components on selectivity of hydrophilic interaction chromatography (HILIC). <i>Journal of Separation Science</i> , 2008, 31, 1449-1464.	1.3	238
236	The advantages of ESI-MS detection in conjunction with HILIC mode separations: Fundamentals and applications. <i>Journal of Separation Science</i> , 2008, 31, 1465-1480.	1.3	250
237	Hydrophilic interaction and reversed-phase ultra-performance liquid chromatography TOF-MS for metabonomic analysis of Zucker rat urine. <i>Journal of Separation Science</i> , 2008, 31, 1598-1608.	1.3	121
238	HILIC for separation of co-eluted flavonoids under RP-HPLC mode. <i>Journal of Separation Science</i> , 2008, 31, 1623-1627.	1.3	40
239	Chromatographic deuterium isotope effects of derivatized N-glycans and N-glycopeptides in a zwitterionic type of hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1594-1597.	1.3	12
240	The application of novel 1.7 μ m ethylene bridged hybrid particles for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1511-1518.	1.3	83
241	Method development for HILIC assays. <i>Journal of Separation Science</i> , 2008, 31, 1438-1448.	1.3	126
242	Pressurized CEC of neutral and charged solutes using silica monolithic stationary phases functionalized with 3-(2-aminoethylamino)propyl ligands. <i>Journal of Separation Science</i> , 2008, 31, 2244-2251.	1.3	13
243	Preparation of a sorbitol methacrylate grafted silica as stationary phase for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1504-1510.	1.3	20
244	Stationary phases for hydrophilic interaction chromatography, their characterization and implementation into multidimensional chromatography concepts. <i>Journal of Separation Science</i> , 2008, 31, 1421-1437.	1.3	219

#	ARTICLE	IF	CITATIONS
245	Hydrophilic interaction chromatography of small metal species in plants using sulfobetaine- and phosphorylcholine-type zwitterionic stationary phases. <i>Journal of Separation Science</i> , 2008, 31, 1615-1622.	1.3	41
246	Multiple linear regression and artificial neural network retention prediction models for ginsenosides on a polyamine-bonded stationary phase in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2008, 31, 1550-1563.	1.3	27
247	Multidimensional liquid phase separations for mass spectrometry. <i>Journal of Separation Science</i> , 2008, 31, 1964-1979.	1.3	27
248	Evaluation of mobile phase, ion pairing, and temperature influence on an HILIC-MS/MS method for arginine and its dimethylated derivatives detection. <i>Journal of Separation Science</i> , 2008, 31, 2424-2429.	1.3	27
249	Considerations on HILIC and polar organic solvent-based separations: Use of cyclodextrin and macrocyclic glycopeptide stationary phases. <i>Journal of Separation Science</i> , 2008, 31, 1980-1990.	1.3	66
250	Mixed-mode ion-exchangers and their comparative chromatographic characterization in reversed-phase and hydrophilic interaction chromatography elution modes. <i>Journal of Separation Science</i> , 2008, 31, 2572-2588.	1.3	148
251	Polymethacrylate monolithic columns for capillary liquid chromatography. <i>Journal of Separation Science</i> , 2008, 31, 2521-2540.	1.3	118
252	Mixed-mode hydrophilic interaction/cation-exchange chromatography (HILIC/CEX) of peptides and proteins. <i>Journal of Separation Science</i> , 2008, 31, 2754-2773.	1.3	74
253	Hydrophilic interaction chromatography in nonaqueous elution mode for separation of hydrophilic analytes on silica-based packings with noncharged polar bondings*. <i>Journal of Separation Science</i> , 2008, 31, 2971-2987.	1.3	79
254	Monolithic column with double mixed-modes of hydrophilic interaction/ cation-exchange and reverse-phase/ cation-exchange stationary phase for pressurized capillary electrochromatography. <i>Electrophoresis</i> , 2008, 29, 928-935.	1.3	26
255	Preparation and evaluation of a lysine-bonded silica monolith as polar stationary phase for hydrophilic interaction pressurized capillary electrochromatography. <i>Electrophoresis</i> , 2008, 29, 3896-3904.	1.3	16
256	Methacrylate-based monolithic column with mixed-mode hydrophilic interaction/strong cation-exchange stationary phase for capillary liquid chromatography and pressure-assisted CEC. <i>Electrophoresis</i> , 2008, 29, 4055-4065.	1.3	39
257	Analysis of histones, histone variants, and their post-translationally modified forms. <i>Electrophoresis</i> , 2008, 29, 2516-2532.	1.3	26
258	Highly efficient peptide separations in proteomics. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 866, 48-63.	1.2	87
259	Rapid determination of the applicability of hydrophilic interaction chromatography utilizing ACD Labs Log D Suite: A bioanalytical application. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 863, 1-8.	1.2	31
260	A novel method for the determination of 1,5-anhydroglucitol, a glycemic marker, in human urine utilizing hydrophilic interaction liquid chromatography/MS3. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 873, 144-150.	1.2	21
261	Quantitative metabolome analysis using liquid chromatography-high-resolution mass spectrometry. <i>Analytical Biochemistry</i> , 2008, 382, 94-100.	1.1	91
262	Monitoring of sulfonamide antibacterial residues in milk and egg by polymer monolith microextraction coupled to hydrophilic interaction chromatography/mass spectrometry. <i>Analytica Chimica Acta</i> , 2008, 625, 160-172.	2.6	86

#	ARTICLE	IF	CITATIONS
263	Developments in the use and fabrication of organic monolithic phases for use with high-performance liquid chromatography and capillary electrochromatography. <i>Journal of Chromatography A</i> , 2008, 1184, 416-440.	1.8	98
264	Synthesis and characterization of silica-based hyper-crosslinked sulfonate-modified reversed stationary phases. <i>Journal of Chromatography A</i> , 2008, 1182, 41-55.	1.8	21
265	New hydrophilic interaction/reversed-phase mixed-mode stationary phase and its application for analysis of nonionic ethoxylated surfactants. <i>Journal of Chromatography A</i> , 2008, 1191, 83-89.	1.8	74
266	Development of a comprehensive two-dimensional hydrophilic interaction chromatography/quadrupole time-of-flight mass spectrometry system and its application in separation and identification of saponins from <i>Quillaja saponaria</i> . <i>Journal of Chromatography A</i> , 2008, 1181, 51-59.	1.8	68
267	Separation efficiencies in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2008, 1184, 474-503.	1.8	395
268	Hydrophilic interaction chromatography of nucleotides and their pathway intermediates on titania. <i>Journal of Chromatography A</i> , 2008, 1187, 87-93.	1.8	77
269	Method development for identification of ketobemidone metabolites in microdialysate samples by coupled-column capillary liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2008, 1189, 503-513.	1.8	11
270	Stability-indicating hydrophilic interaction liquid chromatography method for highly polar and basic compounds. <i>Journal of Chromatography A</i> , 2008, 1188, 255-263.	1.8	82
271	Electrochromatographic characterization of methacrylate-based monolith with mixed mode of hydrophilic and weak electrostatic interactions by pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , 2008, 1190, 365-371.	1.8	43
272	Evaluation of the properties of a superficially porous silica stationary phase in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2008, 1193, 85-91.	1.8	108
273	Application of silica-based hyper-crosslinked sulfonate-modified reversed stationary phases for separating highly hydrophilic basic compounds. <i>Journal of Chromatography A</i> , 2008, 1202, 8-18.	1.8	30
274	Novel two-dimensional reversed-phase liquid chromatography/hydrophilic interaction chromatography, an excellent orthogonal system for practical analysis. <i>Journal of Chromatography A</i> , 2008, 1208, 133-140.	1.8	107
275	Matrix solid-phase dispersion extraction of carbadox and olaquinox in feed followed by hydrophilic interaction ultra-high-pressure liquid chromatographic analysis. <i>Journal of Chromatography A</i> , 2008, 1209, 83-87.	1.8	41
276	Development and validation of a fast quantitative method for plasma dimethylarginines analysis using liquid chromatography-tandem mass spectrometry. <i>Clinical Biochemistry</i> , 2008, 41, 1391-1395.	0.8	34
277	Two-dimensional hydrophilic interaction chromatography coupling anion-exchange and hydrophilic interaction columns for separation of 2-pyridylamino derivatives of neutral and sialylated N-glycans. <i>Journal of Chromatography A</i> , 2008, 1189, 169-174.	1.8	48
278	Fast sample preparation and liquid chromatography-tandem mass spectrometry method for assaying cell lysate acetylcholine. <i>Journal of Chromatography A</i> , 2008, 1183, 100-107.	1.8	24
279	Retention of opioids and their glucuronides on a combined zwitterion and hydrophilic interaction stationary phase. <i>Journal of Chromatography A</i> , 2008, 1187, 46-52.	1.8	28
280	Estimation of the extent of the water-rich layer associated with the silica surface in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2008, 1192, 225-229.	1.8	162

#	ARTICLE	IF	CITATIONS
281	Serial coupling of reversed-phase and hydrophilic interaction liquid chromatography to broaden the elution window for the analysis of pharmaceutical compounds. <i>Journal of Chromatography A</i> , 2008, 1208, 90-94.	1.8	72
282	Separation of Organoarsenicals by Means of Zwitterionic Hydrophilic Interaction Chromatography (ZIC [®] -HILIC) and Parallel ICP-MS/ESI-MS Detection. <i>Engineering in Life Sciences</i> , 2008, 8, 582-588.	2.0	19
283	Simultaneous Determination of Metformin Hydrochloride, Cyanoguanidine and Melamine in Tablets by Mixed-Mode HILIC. <i>Chromatographia</i> , 2008, 67, 517-525.	0.7	41
284	Chromatographic Behavior of Uric Acid and Methyl Uric Acids on a Diol Column in HILIC. <i>Chromatographia</i> , 2008, 67, 507-515.	0.7	37
285	Hydrophilic di- and tripeptides are not a precondition for savoury flavour in mature Cheddar cheese. <i>Dairy Science and Technology</i> , 2008, 88, 467-475.	2.2	14
286	Speciation Analysis of Gadolinium-Based MRI Contrast Agents in Blood Plasma by Hydrophilic Interaction Chromatography/Electrospray Mass Spectrometry. <i>Analytical Chemistry</i> , 2008, 80, 8163-8170.	3.2	50
287	A New HPLC Method for the Simultaneous Determination of Acetaminophen, Phenylephrine, Dextromethorphan and Chlorpheniramine in Pharmaceutical Formulations. <i>Analytical Letters</i> , 2008, 41, 965-976.	1.0	38
288	2D PAGE: Sample Preparation and Fractionation. <i>Methods in Molecular Biology</i> , 2008, , .	0.4	1
289	Multidimensional LC Separations in Shotgun Proteomics. <i>Analytical Chemistry</i> , 2008, 80, 7187-7193.	3.2	202
290	Electrostatic Repulsion Hydrophilic Interaction Chromatography for Isocratic Separation of Charged Solutes and Selective Isolation of Phosphopeptides. <i>Analytical Chemistry</i> , 2008, 80, 62-76.	3.2	479
291	Study on the retention equation in hydrophilic interaction liquid chromatography. <i>Talanta</i> , 2008, 76, 522-527.	2.9	168
292	Speciation Analysis of Selenium Metabolites in Yeast-Based Food Supplements by ICPMS-Assisted Hydrophilic Interaction HPLC-Hybrid Linear Ion Trap/Orbitrap MS. <i>Analytical Chemistry</i> , 2008, 80, 3975-3984.	3.2	65
293	Solvent structures of mixed water/acetonitrile mixtures at chromatographic interfaces from computer simulations. <i>Physical Chemistry Chemical Physics</i> , 2008, 10, 4765.	1.3	40
294	Hydrophilic interaction liquid chromatography (HILIC) coupled to inductively coupled plasma mass spectrometry (ICPMS) utilizing a mobile phase with a low-volatile organic modifier for the determination of cisplatin, and its monohydrolyzed metabolite. <i>Journal of Analytical Atomic Spectrometry</i> , 2008, 23, 948.	1.6	29
295	Development of an On-Line SPE-LC-ESI-MS Method for Urinary Nucleosides: Hyphenation of Aprotic Boronic Acid Chromatography with Hydrophilic Interaction LC-ESI-MS. <i>Analytical Chemistry</i> , 2008, 80, 1263-1271.	3.2	75
296	Rapid and Sensitive HILIC-ESI-MS/MS Quantitation of Polar Metabolites of Acrylamide in Human Urine Using Column Switching with an Online Trap Column. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 9828-9834.	2.4	35
297	Mass spectrometry-based neurochemical analysis: perspectives for primate research. <i>Expert Review of Proteomics</i> , 2008, 5, 641-652.	1.3	7
298	Combination of Two Hydrophilic Interaction Chromatography Methods That Facilitates Identification of 2-Aminobenzamide-Labeled Oligosaccharides. <i>Journal of Chromatographic Science</i> , 2008, 46, 68-73.	0.7	16

#	ARTICLE	IF	CITATIONS
299	Hydrophilic Interaction Chromatography Reduces the Complexity of the Phosphoproteome and Improves Global Phosphopeptide Isolation and Detection. <i>Molecular and Cellular Proteomics</i> , 2008, 7, 971-980.	2.5	329
300	Evaluation of BEH C18, BEH HILIC, and HSS T3 (C18) Column Chemistries for the UPLC-MS-MS Analysis of Glutathione, Glutathione Disulfide, and Ophthalmic Acid in Mouse Liver and Human Plasma. <i>Journal of Chromatographic Science</i> , 2008, 46, 209-214.	0.7	88
301	Identification and Characterization of Toxicity of Contaminants in Pet Food Leading to an Outbreak of Renal Toxicity in Cats and Dogs. <i>Toxicological Sciences</i> , 2008, 106, 251-262.	1.4	408
302	Retention Prediction Modeling of Ginsenosides on a Polyvinyl Alcohol-bonded Stationary Phase at Subambient Temperatures Using Multiple Linear Regression and Artificial Neural Network. <i>Analytical Sciences</i> , 2008, 24, 139-148.	0.8	16
304	Some Insights on Retention and Selectivity for Hydrophilic Interaction Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 628-646.	0.5	6
305	A Bidentate Amino Stationary Phase for Hydrophilic Interaction Chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 2164-2175.	0.5	7
306	On-line nano-HPLC/ESI QTOF MS monitoring of β and α sialylation in granulocyte glycosphingolipidome. <i>Biological Chemistry</i> , 2009, 390, 657-72.	1.2	32
307	Determination of γ -Aminobutyric Acid in Food Matrices by Isotope Dilution Hydrophilic Interaction Chromatography Coupled to Mass Spectrometry. <i>Journal of Chromatographic Science</i> , 2009, 47, 564-568.	0.7	11
308	Substrate phosphorylation capacities of the major tyrosine protein kinase from the human promyelocytic cell line, HL60. <i>International Journal of Peptide and Protein Research</i> , 1994, 43, 496-504.	0.1	8
309	Biochemical and Synthetic Polymer Separations. , 0, , 569-663.		0
310	Normal-Phase Chromatography. , 0, , 361-402.		0
311	HPLC methods for the purification of [11C]-labelled radiopharmaceuticals: reversal of the retention order of products and precursors. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 2009, 52, 177-181.	0.5	12
312	Structural glycomics using hydrophilic interaction chromatography (HILIC) with mass spectrometry. <i>Mass Spectrometry Reviews</i> , 2009, 28, 192-206.	2.8	230
313	Online separations combined with MS for analysis of glycosaminoglycans. <i>Mass Spectrometry Reviews</i> , 2009, 28, 254-272.	2.8	112
314	Simultaneous determination of three anticonvulsants using hydrophilic interaction LC-MS. <i>Journal of Separation Science</i> , 2009, 32, 238-243.	1.3	42
315	HILIC separation and quantitation of water-soluble vitamins using diol column. <i>Journal of Separation Science</i> , 2009, 32, 909-917.	1.3	60
316	Purification of flavonoids and triterpene saponins from the licorice extract using preparative HPLC under RP and HILIC mode. <i>Journal of Separation Science</i> , 2009, 32, 526-535.	1.3	28
317	Grafting of silica with sulfobetaine polymers via aqueous reversible addition fragmentation chain transfer polymerization and its use as a stationary phase in HILIC. <i>Journal of Separation Science</i> , 2009, 32, 2008-2016.	1.3	37

#	ARTICLE	IF	CITATIONS
318	Polymer monolith microextraction online coupled to hydrophilic interaction chromatography/mass spectrometry for analysis of β_2 -agonist in human urine. <i>Journal of Separation Science</i> , 2009, 32, 1965-1974.	1.3	32
319	Development of orthogonal two-dimensional hydrophilic interaction chromatography systems with the introduction of novel stationary phases. <i>Journal of Separation Science</i> , 2009, 32, 2871-2876.	1.3	18
320	Glycin α -bonded silica monolithic column as zwitterionic stationary phase for hydrophilic interaction pressurized capillary electrochromatography. <i>Journal of Separation Science</i> , 2009, 32, 2767-2775.	1.3	13
321	Novel highly hydrophilic zwitterionic monolithic column for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2009, 32, 2544-2555.	1.3	45
322	Profiling of polar metabolites in biological extracts using diamond hydride α -based aqueous normal phase chromatography. <i>Journal of Separation Science</i> , 2009, 32, 2273-2280.	1.3	63
323	Preparation and characterization of polymethacrylate monolithic capillary columns with dual hydrophilic interaction reversed α -phase retention mechanism for polar compounds. <i>Journal of Separation Science</i> , 2009, 32, 2530-2543.	1.3	65
324	The acetonitrile shortage: Is reversed HILIC with water an alternative for the analysis of highly polar ionizable solutes?. <i>Journal of Separation Science</i> , 2009, 32, 2001-2007.	1.3	101
325	Utilization of dual retention mechanism on columns with bonded PEG and diol stationary phases for adjusting the separation selectivity of phenolic and flavone natural antioxidants. <i>Journal of Separation Science</i> , 2009, 32, 3603-3619.	1.3	93
326	Evaluation of sulfobetaine α -type polymer resin as an SPE adsorbent in the analysis of trace tetracycline antibiotics in honey. <i>Journal of Separation Science</i> , 2009, 32, 3591-3595.	1.3	18
327	Application of HILIC stationary phase to determination of dimethindene maleate in topical gel. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 50, 23-26.	1.4	9
328	Investigation of an on-line two-dimensional chromatographic approach for peptide analysis in plasma by LC α -MS α -MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 1873-1881.	1.2	39
329	Simultaneous determination of OZ277, a synthetic 1,2,4-trioxolane antimalarial, and its polar metabolites in rat plasma using hydrophilic interaction chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 2989-2995.	1.2	9
330	Modeling of retention of adrenoceptor agonists and antagonists on polar stationary phases in hydrophilic interaction chromatography: a review. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 393, 137-153.	1.9	26
331	Analytical strategies for phosphoproteomics. <i>Proteomics</i> , 2009, 9, 1451-1468.	1.3	434
332	An integrated metabonomic method for profiling of metabolic changes in carbon tetrachloride induced rat urine. <i>Toxicology</i> , 2009, 256, 191-200.	2.0	31
333	Remediation of undesirable secondary interactions encountered in hydrophilic interaction chromatography during development of a quantitative LC α -MS/MS assay for a dipeptidyl peptidase IV (DPP-IV) inhibitor in monkey serum. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 471-476.	1.2	12
334	Quantification of peramivir (a novel anti-influenza drug) in human plasma by hydrophilic interaction chromatography/tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 933-938.	1.2	16
335	Highly efficient peptide separations in proteomicsPart 2: Bi- and multidimensional liquid-based separation techniques. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 1019-1039.	1.2	94

#	ARTICLE	IF	CITATIONS
336	Chromatographic and mass spectrometric analysis of glutathione in biological samples. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 3309-3317.	1.2	95
337	Determination of metformin in human plasma using hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 3695-3700.	1.2	61
338	Comparison of the sensitivity of evaporative universal detectors and LC/MS in the HILIC and the reversed-phase HPLC modes. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 4133-4139.	1.2	73
339	Determination of metformin and its prodrugs in human and rat blood by hydrophilic interaction liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 50, 469-474.	1.4	40
340	Fingerprint analysis of <i>Ligusticum chuanxiong</i> using hydrophilic interaction chromatography and reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 2136-2141.	1.8	62
341	Qualitative and quantitative analysis in quality control of traditional Chinese medicines. <i>Journal of Chromatography A</i> , 2009, 1216, 2033-2044.	1.8	199
342	Hydrophilic interaction 10 μ m I.D. porous layer open tubular columns for ultratrace glycan analysis by liquid chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2009, 1216, 1223-1231.	1.8	65
343	Synthesis, chromatographic evaluation and hydrophilic interaction/reversed-phase mixed-mode behavior of a β -cyclodextrin-stationary phase. <i>Journal of Chromatography A</i> , 2009, 1216, 257-263.	1.8	160
344	Preparation of a mixed-mode hydrophilic interaction/anion-exchange polymeric monolithic stationary phase for capillary liquid chromatography of polar analytes. <i>Journal of Chromatography A</i> , 2009, 1216, 801-806.	1.8	38
345	Challenge of high polarity and low concentrations in analysis of cytostatics and metabolites in wastewater by hydrophilic interaction chromatography/tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2009, 1216, 1100-1108.	1.8	106
346	Separation, detection and quantitation of peptides by liquid chromatography and capillary electrochromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 1825-1837.	1.8	56
347	Hydrophilic interaction liquid chromatography with alcohol as a weak eluent. <i>Journal of Chromatography A</i> , 2009, 1216, 2362-2370.	1.8	74
348	Novel zwitterionic polyphosphorylcholine monolithic column for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 2439-2448.	1.8	75
349	Multidimensional system enabling deglycosylation of proteins using a capillary reactor with peptide-N-glycosidase F immobilized on a porous polymer monolith and hydrophilic interaction liquid chromatography-mass spectrometry of glycans. <i>Journal of Chromatography A</i> , 2009, 1216, 3252-3259.	1.8	61
350	Rapid and efficient purification of positron emission tomography probes by hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 3933-3940.	1.8	4
351	Analysis of chlormequat and mepiquat by hydrophilic interaction chromatography coupled to tandem mass spectrometry in food samples. <i>Journal of Chromatography A</i> , 2009, 1216, 4402-4406.	1.8	58
352	Preparation and evaluation of a neutral methacrylate-based monolithic column for hydrophilic interaction stationary phase by pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 4611-4617.	1.8	53
353	Hydrophilic interaction chromatography coupled to electrospray mass spectrometry for the separation of peptides and protein digests. <i>Journal of Chromatography A</i> , 2009, 1216, 5518-5524.	1.8	49

#	ARTICLE	IF	CITATIONS
354	Comparison of the adsorption mechanisms of pyridine in hydrophilic interaction chromatography and in reversed-phase aqueous liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 8496-8504.	1.8	62
355	Quantitative site-specific analysis of protein glycosylation by LC-MS using different glycopeptide-enrichment strategies. <i>Analytical Biochemistry</i> , 2009, 395, 178-188.	1.1	109
356	Analysis of highly polar metabolites in human plasma by ultra-performance hydrophilic interaction liquid chromatography coupled with quadrupole-time of flight mass spectrometry. <i>Analytica Chimica Acta</i> , 2009, 650, 10-15.	2.6	54
357	Synthesis of Novel Sulfobetaine-Type Adsorbents and Characteristics of Their Adsorption of Polar Solutes in Hydrophilic SPE. <i>Chromatographia</i> , 2009, 70, 1525-1530.	0.7	27
358	A Stability-Indicating Assay of Brimonidine Tartrate Ophthalmic Solution and Stress Testing Using HILIC. <i>Chromatographia</i> , 2009, 70, 539-544.	0.7	18
359	Validated HPLC Method for the Quantitative Analysis of a 4-Methanesulfonyl-Piperidine Hydrochloride Salt. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 1000-1009.	0.5	5
360	Lipophilicity of Basic Drugs Measured by Hydrophilic Interaction Chromatography. <i>Journal of Medicinal Chemistry</i> , 2009, 52, 3416-3419.	2.9	28
361	Development of a Robust and High Throughput Method for Profiling N-Linked Glycans Derived from Plasma Glycoproteins by NanoLC-FTICR Mass Spectrometry. <i>Journal of Proteome Research</i> , 2009, 8, 3764-3770.	1.8	42
362	Development of a nanoLC LTQ Orbitrap Mass Spectrometric Method for Profiling Glycans Derived from Plasma from Healthy, Benign Tumor Control, and Epithelial Ovarian Cancer Patients. <i>Analytical Chemistry</i> , 2009, 81, 1130-1136.	3.2	74
363	Mass spectrometry-based technologies for high-throughput metabolomics. <i>Bioanalysis</i> , 2009, 1, 1665-1684.	0.6	60
364	Phosphoproteomics—finally fulfilling the promise?. <i>Molecular BioSystems</i> , 2009, 5, 1122.	2.9	39
365	Quantitative Profiling of Polar Cationic Metabolites in Human Cerebrospinal Fluid by Reversed-Phase Nanoliquid Chromatography/Mass Spectrometry. <i>Analytical Chemistry</i> , 2009, 81, 1121-1129.	3.2	71
366	Bioanalytical hydrophilic interaction chromatography: recent challenges, solutions and applications. <i>Bioanalysis</i> , 2009, 1, 239-253.	0.6	18
367	Metallomics: the concept and methodology. <i>Chemical Society Reviews</i> , 2009, 38, 1119.	18.7	309
368	Quantification of cisplatin, carboplatin and oxaliplatin in spiked human plasma samples by ICP-SFMS and hydrophilic interaction liquid chromatography (HILIC) combined with ICP-MS detection. <i>Journal of Analytical Atomic Spectrometry</i> , 2009, 24, 1336.	1.6	66
369	Hydrophilic-interaction liquid chromatography (HILIC) with dad and mass spectroscopic detection for direct analysis of glyphosate and glufosinate residues and for product quality control. <i>Acta Chromatographica</i> , 2009, 21, 559-576.	0.7	30
371	Selection of HILIC columns to handle matrix effect due to phospholipids. <i>Bioanalysis</i> , 2009, 1, 57-62.	0.6	32
372	Comprehensive Analytical Strategy for Biomarker Identification based on Liquid Chromatography Coupled to Mass Spectrometry and New Candidate Confirmation Tools. <i>Analytical Chemistry</i> , 2009, 81, 7677-7694.	3.2	31

#	ARTICLE	IF	CITATIONS
373	Pattern Recognition and Pathway Analysis with Genetic Algorithms in Mass Spectrometry Based Metabolomics. <i>Algorithms</i> , 2009, 2, 638-666.	1.2	16
374	Hydrophilic-interaction planar chromatography of some water-soluble Co(III) complexes on different adsorbents. <i>Journal of Planar Chromatography - Modern TLC</i> , 2009, 22, 249-253.	0.6	12
375	Recent Applications of Phosphoproteomics. <i>Current Proteomics</i> , 2010, 7, 168-176.	0.1	2
376	Emerging Analytical Separation Techniques with High Throughput Potential for Pharmaceutical Analysis, Part II: Novel Chromatographic Modes. <i>Combinatorial Chemistry and High Throughput Screening</i> , 2010, 13, 530-547.	0.6	8
377	Development of Polymer-Coated Packing Materials for High-Performance Liquid Chromatography. <i>Bunseki Kagaku</i> , 2010, 59, 175-192.	0.1	0
378	éŸä@â%â...”ăăăăă,LC-MS/MS†æž• <i>Kagaku To Seibutsu</i> , 2010, 48, 201-209.	0.0	1
379	Regularities of chromatographic retention of phenols on microdispersed sintered detonation nanodiamond in aqueousâ€”organic solvents. <i>Russian Journal of Physical Chemistry A</i> , 2010, 84, 476-480.	0.1	26
380	Study of the Behavior of Water-Soluble Vitamins in HILIC on a Diol Column. <i>Chromatographia</i> , 2010, 71, 751-759.	0.7	55
381	Analysis of Pterins in Urine by HILIC. <i>Chromatographia</i> , 2010, 71, 761-768.	0.7	13
382	Determination of Propylenethiourea, the Main Metabolite of Propineb, in Tomato by HILIC-MS. <i>Chromatographia</i> , 2010, 71, 75-80.	0.7	12
383	Glycoproteomic Profile in Wine: A â€”Sweetâ€”™ Molecular Renaissance. <i>Journal of Proteome Research</i> , 2010, 9, 6148-6159.	1.8	45
384	Separation and identification of oligonucleotides by hydrophilic interaction liquid chromatography (HILIC)â€”inductively coupled plasma mass spectrometry (ICPMS). <i>Analyst, The</i> , 2010, 135, 2560.	1.7	57
385	Crossâ€”linked agarose for separation of low molecular weight natural products in hydrophilic interaction liquid chromatography. <i>Biotechnology Journal</i> , 2010, 5, 505-510.	1.8	19
386	Separations of Cycloinulooligosaccharides via Hydrophilic Interaction Chromatography (HILIC) and Ligand-Exchange Chromatography. <i>Separation Science and Technology</i> , 2010, 45, 447-452.	1.3	8
387	The significance, development and progress of high-throughput combinatorial histone code analysis. <i>Cellular and Molecular Life Sciences</i> , 2010, 67, 3983-4000.	2.4	90
388	Separation of peptides by HPLC using a surface-confined ionic liquid stationary phase. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 396, 775-781.	1.9	43
389	Multitarget quantitative metabolic profiling of hydrophilic metabolites in fermentation broths of Î²-lactam antibiotics production by HILICâ€”ESIâ€”MS/MS. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 396, 1655-1679.	1.9	57
390	Hydrophilic interaction liquid chromatography/positive ion electrospray ionization mass spectrometry method for the quantification of perindopril and its main metabolite in human plasma. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 397, 2161-2170.	1.9	8

#	ARTICLE	IF	CITATIONS
391	A hydrophilic interaction ultraperformance liquid chromatography (HILIC-UPLC) method for genomic DNA methylation assessment by UV detection. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 396, 2937-2941.	1.9	17
392	Determination of isoascorbic acid in fish tissue by hydrophilic interaction liquid chromatography-ultraviolet detection. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 397, 2199-2210.	1.9	12
393	Structure elucidation of the thermal degradation products of the nucleotide cofactors NADH and NADPH by nano-ESI-FTICR-MS and HPLC-MS. <i>Analytical and Bioanalytical Chemistry</i> , 2010, 398, 2803-2811.	1.9	31
394	LC-MS/MS method using unbonded silica column and aqueous/methanol mobile phase for the simultaneous quantification of a drug candidate and co-administered metformin in rat plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010, 878, 1583-1589.	1.2	17
395	Hydrophilic interaction liquid chromatography and accurate mass measurement for quantification and confirmation of morphine, codeine and their glucuronide conjugates in human urine. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010, 878, 2959-2966.	1.2	41
396	Quantitative determination of un-derivatised amino acids in artistic mural paintings using high-performance liquid chromatography/electrospray ionization triple quadrupole mass spectrometry. <i>Analytica Chimica Acta</i> , 2010, 675, 1-7.	2.6	26
397	Efficient hydrophilic interaction liquid chromatography-tandem mass spectrometry for the multiclass analysis of veterinary drugs in chicken muscle. <i>Analytica Chimica Acta</i> , 2010, 682, 117-129.	2.6	89
398	Simultaneous determination of positive and negative pharmaceutical counterions using mixed-mode chromatography coupled with charged aerosol detector. <i>Journal of Chromatography A</i> , 2010, 1217, 5776-5784.	1.8	86
399	Determination of deoxynivalenol-sulfonate (DONS) in cereals by hydrophilic interaction chromatography coupled to tandem mass spectrometry. <i>Mycotoxin Research</i> , 2010, 26, 109-117.	1.3	23
400	Preparation and evaluation of an enzymatic microreactor based on HILIC matrix for digestion and identification of proteins. <i>Science Bulletin</i> , 2010, 55, 910-914.	1.7	6
401	In vivo nicotine exposure in the zebra finch: A promising innovative animal model to use in neurodegenerative disorders related research. <i>Pharmacology Biochemistry and Behavior</i> , 2010, 96, 152-159.	1.3	14
402	Predicting human developmental toxicity of pharmaceuticals using human embryonic stem cells and metabolomics. <i>Toxicology and Applied Pharmacology</i> , 2010, 247, 18-27.	1.3	176
403	Sulfoalkylbetaine-based monolithic column with mixed-mode of hydrophilic interaction and strong anion-exchange stationary phase for capillary electrochromatography. <i>Electrophoresis</i> , 2010, 31, 2997-3005.	1.3	31
405	Synthesis of Sugar-Based Silica Gels by Copper-Catalysed Azide-Alkyne Cycloaddition via a Single-Step Azido-Activated Silica Intermediate and the Use of the Gels in Hydrophilic Interaction Chromatography. <i>Chemistry - A European Journal</i> , 2010, 16, 5712-5722.	1.7	63
406	Quantification of acetylcholine, an essential neurotransmitter, in brain microdialysis samples by liquid chromatography mass spectrometry. <i>Biomedical Chromatography</i> , 2010, 24, 39-48.	0.8	43
407	Hydrophilic interaction LC of peptides: Columns comparison and clustering. <i>Journal of Separation Science</i> , 2010, 33, 728-739.	1.3	51
408	Application of hydrophilic interaction chromatography retention coefficients for predicting peptide elution with TFA and methanesulfonic acid ion-pairing reagents. <i>Journal of Separation Science</i> , 2010, 33, 826-833.	1.3	11
409	HILIC behavior of a reversed-phase/cation-exchange/anion-exchange trimode column. <i>Journal of Separation Science</i> , 2010, 33, 779-786.	1.3	85

#	ARTICLE	IF	CITATIONS
410	Influence of stationary phase chemistry and mobile phase composition on retention, selectivity, and MS response in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2010, 33, 740-751.	1.3	77
411	Separation of purine and pyrimidine bases and nucleosides by hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2010, 33, 803-816.	1.3	81
412	Dual hydrophilic interaction-RP retention mechanism on polar columns: Structural correlations and implementation for 2D separations on a single column. <i>Journal of Separation Science</i> , 2010, 33, 841-852.	1.3	88
413	Retention behavior of estrogen metabolites on hydrophilic interaction chromatography stationary phases. <i>Journal of Separation Science</i> , 2010, 33, 793-802.	1.3	28
414	Zwitterionic polymeric monoliths for HILIC/RP mixed mode for CEC separation applications. <i>Journal of Separation Science</i> , 2010, 33, 787-792.	1.3	40
415	Recent advances in application of hydrophilic interaction chromatography for quantitative bioanalysis. <i>Journal of Separation Science</i> , 2010, 33, 681-697.	1.3	139
416	Evaluation of hybrid hydrophilic interaction chromatography stationary phases for ultra-HPLC in analysis of polar pteridines. <i>Journal of Separation Science</i> , 2010, 33, 765-772.	1.3	34
417	Retention of arsenic species on zwitterionic stationary phase in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2010, 33, 817-825.	1.3	15
418	HILIC methods in pharmaceutical analysis. <i>Journal of Separation Science</i> , 2010, 33, 698-715.	1.3	167
419	Evaluation of various HILIC materials for the fast separation of polar compounds. <i>Journal of Separation Science</i> , 2010, 33, 752-764.	1.3	107
420	Orotic acid quantification in dried blood spots and biological fluids by hydrophilic interaction liquid chromatography tandem mass spectrometry. <i>Journal of Separation Science</i> , 2010, 33, 966-973.	1.3	21
421	Silica-based zwitterionic monolithic stationary phase for separation of neutral and ionized solutes using pressurized CEC. <i>Journal of Separation Science</i> , 2010, 33, 1625-1632.	1.3	4
422	Enhanced glyco-profiling by specific glycopeptide enrichment and complementary monolithic nano-LC (ZIC-HILIC/RP18e)/ESI-MS analysis. <i>Journal of Separation Science</i> , 2010, 33, 880-890.	1.3	59
423	Green hydrophilic interaction chromatography using ethanol-water-carbon dioxide mixtures. <i>Journal of Separation Science</i> , 2010, 33, 834-837.	1.3	52
424	Hydrophilic interaction chromatography coupled to MS for metabonomic/metabolomic studies. <i>Journal of Separation Science</i> , 2010, 33, 716-727.	1.3	180
425	Determination of melamine in milk powder using zwitterionic HILIC stationary phase with UV detection. <i>Journal of Separation Science</i> , 2010, 33, 988-995.	1.3	39
426	Purification of active bufadienolides from toad skin by preparative reversed-phase liquid chromatography coupled with hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2010, 33, 1487-1494.	1.3	31
427	Tris(hydroxymethyl)aminomethane-functionalized silica particles and their application for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2010, 33, 2965-2976.	1.3	31

#	ARTICLE	IF	CITATIONS
428	Chromatographic behavior of a new hybrid type RP material containing silica bonded 1,3-bis(2-cyano-propyloxy)propane. Journal of Separation Science, 2010, 33, 2956-2964.	2.8	151
429	Metabolomic applications of HILIC-LC-MS. Mass Spectrometry Reviews, 2010, 29, 671-684.	2.8	151
430	Separation of 2-aminobenzamide labeled glycans using hydrophilic interaction chromatography columns packed with 1.7µm sorbent. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2010, 878, 403-408.	1.2	182
431	Quantitation of polar analytes using column-switching: Application to oxycodone and three metabolites in human plasma. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2010, 878, 637-644.	1.2	17
432	Quantitative analysis of acetylcholine in rat brain microdialysates by liquid chromatography coupled with electrospray ionization tandem mass spectrometry. Journal of Neuroscience Methods, 2010, 194, 87-93.	1.3	26
433	Investigation of the adsorption behavior of glycine peptides on 12% cross-linked agarose gel media. Journal of Chromatography A, 2010, 1217, 1916-1925.	1.8	16
434	Approach to hydrophilic interaction chromatography column selection: Application to neurotransmitters analysis. Journal of Chromatography A, 2010, 1217, 3091-3104.	1.8	110
435	Determination of iminosugars in mulberry leaves and silkworms using hydrophilic interaction chromatography-tandem mass spectrometry. Analytical Biochemistry, 2010, 404, 217-222.	1.1	54
436	Multi-dimensional liquid chromatography in proteomics—A review. Analytica Chimica Acta, 2010, 664, 101-113.	2.6	158
437	Determination of aromatic amines in aqueous extracts of polyurethane foam using hydrophilic interaction liquid chromatography and mass spectrometry. Analytica Chimica Acta, 2010, 678, 117-123.	2.6	21
438	Separation of carboxylates by hydrophilic interaction liquid chromatography on titania. Journal of Chromatography A, 2010, 1217, 82-88.	1.8	50
439	Further investigations of the effect of pressure on retention in ultra-high-pressure liquid chromatography. Journal of Chromatography A, 2010, 1217, 276-284.	1.8	59
440	The challenges of the analysis of basic compounds by high performance liquid chromatography: Some possible approaches for improved separations. Journal of Chromatography A, 2010, 1217, 858-880.	1.8	210
441	High-efficiency hydrophilic interaction chromatography by coupling 25cm ² —4.6mm ID—5µm silica columns and operation at 80°C. Journal of Chromatography A, 2010, 1217, 514-521.	1.8	20
442	Efficiency of the same neat silica column in hydrophilic interaction chromatography and per aqueous liquid chromatography. Journal of Chromatography A, 2010, 1217, 683-688.	1.8	71
443	Characterization of adsorption processes in analytical liquid-solid chromatography. Journal of Chromatography A, 2010, 1217, 792-812.	1.8	92
444	Zirconia based monoliths used in hydrophilic-interaction chromatography for original selectivity of xanthenes. Journal of Chromatography A, 2010, 1217, 1496-1500.	1.8	51
445	Development and validation of a hydrophilic interaction liquid chromatographic method for determination of aromatic amines in environmental water. Journal of Chromatography A, 2010, 1217, 1799-1805.	1.8	39

#	ARTICLE	IF	CITATIONS
466	Challenges in Mass Spectrometry Based Targeted Metabolomics. <i>Current Molecular Medicine</i> , 2010, 10, 216-226.	0.6	78
468	Chemically bonded maltose via click chemistry as stationary phase for HILIC. <i>Analytical Methods</i> , 2010, 2, 217-224.	1.3	59
469	Novel Application of Electrostatic Repulsion-Hydrophilic Interaction Chromatography (ERLIC) in Shotgun Proteomics: Comprehensive Profiling of Rat Kidney Proteome. <i>Journal of Proteome Research</i> , 2010, 9, 3520-3526.	1.8	84
470	Liquid Chromatography Electrospray Ionization Fourier Transform Ion Cyclotron Resonance Mass Spectrometric Characterization of N-Linked Glycans and Glycopeptides. <i>Analytical Chemistry</i> , 2010, 82, 6542-6548.	3.2	28
471	Utilizing Ion-Pairing Hydrophilic Interaction Chromatography Solid Phase Extraction for Efficient Glycopeptide Enrichment in Glycoproteomics. <i>Analytical Chemistry</i> , 2010, 82, 5598-5609.	3.2	264
472	Chip-based Reversed-phase Liquid Chromatography-Mass Spectrometry of Permethylated N-Linked Glycans: A Potential Methodology for Cancer-biomarker Discovery. <i>Analytical Chemistry</i> , 2010, 82, 5095-5106.	3.2	153
473	Elimination of LC-MS/MS matrix effect due to phospholipids using specific solid-phase extraction elution conditions. <i>Bioanalysis</i> , 2010, 2, 1011-1021.	0.6	45
474	Practical metabolomics in drug discovery. <i>Expert Opinion on Drug Discovery</i> , 2010, 5, 249-263.	2.5	24
475	Interest of a chemometric approach in understanding the retention behaviour of three columns in hydrophilic interaction liquid chromatography: Application to the separation of glycerol carbonate, glycerol and urea. <i>Talanta</i> , 2010, 81, 1281-1287.	2.9	15
476	Orthogonal separation on one beta-cyclodextrin column by switching reversed-phase liquid chromatography and hydrophilic interaction chromatography. <i>Talanta</i> , 2010, 81, 1870-1876.	2.9	30
477	Ultra Performance Liquid Chromatographic Profiling of Serum N-Glycans for Fast and Efficient Identification of Cancer Associated Alterations in Glycosylation. <i>Analytical Chemistry</i> , 2010, 82, 10208-10215.	3.2	162
478	Selective enrichment of N-linked glycopeptides by using a highly hydrophilic matrix synthesized via click chemistry. <i>Analytical Methods</i> , 2010, 2, 1667.	1.3	17
479	Determination of cytokinins in plant samples by polymer monolith microextraction coupled with hydrophilic interaction chromatography-tandem mass spectrometry. <i>Analytical Methods</i> , 2010, 2, 1676.	1.3	55
480	Separation of Multiphosphorylated Peptide Isomers by Hydrophilic Interaction Chromatography on an Aminopropyl Phase. <i>Analytical Chemistry</i> , 2010, 82, 6409-6414.	3.2	38
481	Application of click chemistry on preparation of separation materials for liquid chromatography. <i>Chemical Society Reviews</i> , 2011, 40, 2177.	18.7	195
482	HYDROPHILIC INTERACTION CHROMATOGRAPHIC DETERMINATION OF OXYTETRACYCLINE IN THE ENVIRONMENTAL WATER USING SILICA COLUMN. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2011, 34, 511-520.	0.5	19
484	A Novel Multidimensional Protein Identification Technology Approach Combining Protein Size Exclusion Prefractionation, Peptide Zwitterion-Ion Hydrophilic Interaction Chromatography, and Nano-Ultrapformance RP Chromatography/nESI-MS ² for the in-Depth Analysis of the Serum Proteome and Phosphoproteome: Application to Clinical Sera Derived from Humans with Benign Prostate Hyperplasia. <i>Analytical Chemistry</i> , 2011, 83, 708-718.	3.2	50
485	Comparison of ERLIC-TiO ₂ , HILIC-TiO ₂ , and SCX-TiO ₂ for Global Phosphoproteomics Approaches. <i>Journal of Proteome Research</i> , 2011, 10, 3474-3483.	1.8	83

#	ARTICLE	IF	CITATIONS
486	A novel zwitterionic HILIC stationary phase based on α -thiol-ene click chemistry between cysteine and vinyl silica. <i>Chemical Communications</i> , 2011, 47, 4550.	2.2	162
487	Zwitterionic Hydrophilic Interaction Liquid Chromatography (ZIC-HILIC and ZIC-cHILIC) Provide High Resolution Separation and Increase Sensitivity in Proteome Analysis. <i>Analytical Chemistry</i> , 2011, 83, 3440-3447.	3.2	128
489	RECENT APPLICATIONS OF HYDROPHILIC INTERACTION CHROMATOGRAPHY IN ENVIRONMENTAL ANALYSIS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2011, 34, 1112-1132.	0.5	6
490	Identification of coexisting pentose, hexose, and disaccharides with preliminary separation through hydrophilic interaction on silica HPTLC plate using aqueous sodium deoxycholate-acetonitrile mobile phase system. <i>Journal of Planar Chromatography - Modern TLC</i> , 2011, 24, 491-496.	0.6	3
491	Multidimensional Strategy for Sensitive Phosphoproteomics Incorporating Protein Prefractionation Combined with SIMAC, HILIC, and TiO ₂ Chromatography Applied to Proximal EGF Signaling. <i>Journal of Proteome Research</i> , 2011, 10, 5383-5397.	1.8	63
492	Ultrasensitive Identification of Localization Variants of Modified Peptides Using Ion Mobility Spectrometry. <i>Analytical Chemistry</i> , 2011, 83, 5617-5623.	3.2	35
493	Quantifying tetrapeptide SS-20 in rat plasma using hydrophilic interaction liquid chromatography coupled with electrospray ionization tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2011, 879, 3353-3359.	1.2	2
494	Analysis of vitamin B1 in dry-cured sausages by hydrophilic interaction liquid chromatography (HILIC) and diode array detection. <i>Meat Science</i> , 2011, 87, 234-238.	2.7	29
495	Titanium dioxide as chemo-affinity chromatographic sorbent of biomolecular compounds α Applications in acidic modification-specific proteomics. <i>Journal of Proteomics</i> , 2011, 75, 317-328.	1.2	61
496	Effect of chromatographic parameters and detector settings on the response of HILIC-evaporative light-scattering detection system using experimental design approach and multicriteria optimization methodology. <i>Talanta</i> , 2011, 83, 1126-1133.	2.9	14
497	Determination of uric acid and creatinine in human urine using hydrophilic interaction chromatography. <i>Talanta</i> , 2011, 83, 1707-1710.	2.9	79
498	Towards multimodal HPLC separations on humic acid-bonded aminopropyl silica: RPLC and HILIC behavior. <i>Talanta</i> , 2011, 85, 1472-1482.	2.9	30
499	Identification and quantification of nucleosides and nucleobases in Geosaurus and Leech by hydrophilic-interaction chromatography. <i>Talanta</i> , 2011, 85, 1634-1641.	2.9	44
501	Quantification of Total and Free Carnitine in Human Plasma by Hydrophilic Interaction Liquid Chromatography Tandem Mass Spectrometry. <i>Journal of Chromatographic Science</i> , 2011, 49, 463-468.	0.7	26
502	Evaluation of the Deuterium Isotope Effect in Zwitterionic Hydrophilic Interaction Liquid Chromatography Separations for Implementation in a Quantitative Proteomic Approach. <i>Analytical Chemistry</i> , 2011, 83, 8352-8356.	3.2	28
503	Recent Advancements on Greening Analytical Separation. <i>Critical Reviews in Analytical Chemistry</i> , 2011, 41, 2-20.	1.8	58
504	Separation of inorganic anions using a series of sulfobetaine exchangers. <i>Journal of Chromatography A</i> , 2011, 1218, 1185-1194.	1.8	15
505	Retention Behavior of Polar Pesticides in Hydrophilic Interaction Chromatography. <i>Journal of Environmental Chemistry</i> , 2011, 21, 293-301.	0.1	0

#	ARTICLE	IF	CITATIONS
506	Development of Mulberry Leaf Extract for Suppressing Postprandial Blood Glucose Elevation. , 0, , .		6
508	Analysis of Glycosaminoglycans Using Mass Spectrometry. <i>Current Proteomics</i> , 2011, 8, 325-336.	0.1	46
509	Method Development and Analysis of Mono- and Diphosphorylated Nucleotides by HILIC HPLC-ESI-MS. <i>Chromatographic Science</i> , 2011, , 445-468.	0.1	0
510	Retention and Selectivity of Polar Stationary Phases for Hydrophilic Interaction Chromatography. <i>Chromatographic Science</i> , 2011, , 401-426.	0.1	3
511	Counter-ionic Effect on the Separation of Water-soluble Compounds Applying a Hydrophilic Stationary Phase Bonded with a Zwitter-ionic Polymer. <i>Analytical Sciences</i> , 2011, 27, 885.	0.8	10
512	Quantification of Non-Volatile Amines in Beer Using Hydrophilic Interaction Liquid Chromatograph-Tandem Mass Spectrometer. <i>Bunseki Kagaku</i> , 2011, 60, 157-162.	0.1	8
513	How Should Ionic Liquids be Analyzed?. <i>Current Organic Chemistry</i> , 2011, 15, 1873-1887.	0.9	27
514	Ultrasonic-assisted liquid-liquid extraction and HILIC-ELSD analysis of ginsenoside Rb1, astragaloside IV and dulcitol in sugar-free Fufangfufangteng Heji. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011, 56, 836-840.	1.4	12
515	One-year plasma N-linked glycome intra-individual and inter-individual variability in the chicken model of spontaneous ovarian adenocarcinoma. <i>International Journal of Mass Spectrometry</i> , 2011, 305, 79-86.	0.7	8
516	Retention behavior of peptides in hydrophilic-interaction chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 8890-8896.	1.8	54
517	Hydrophilic interaction liquid chromatography in food analysis. <i>Journal of Chromatography A</i> , 2011, 1218, 7438-7452.	1.8	107
518	Intact protein separation by chromatographic and/or electrophoretic techniques for top-down proteomics. <i>Journal of Chromatography A</i> , 2011, 1218, 8760-8776.	1.8	76
519	Protein UTLC-MALDI-MS using thin films of submicrometer silica particles. <i>Journal of Chromatography A</i> , 2011, 1218, 7196-7202.	1.8	23
520	The retention behaviour of polar compounds on zirconia based stationary phases under hydrophilic interaction liquid chromatography conditions. <i>Journal of Chromatography A</i> , 2011, 1218, 6981-6986.	1.8	29
521	Determination of organic acid impurities in lactic acid obtained by fermentation of sugarcane juice. <i>Journal of Chromatography A</i> , 2011, 1218, 7147-7157.	1.8	17
522	Retention behaviour of imidazolium ionic liquid cations on 1.7 µm ethylene bridged hybrid silica column using acetonitrile-rich and water-rich mobile phases. <i>Journal of Chromatography A</i> , 2011, 1218, 6884-6891.	1.8	13
523	Rapid and simultaneous determination of hexapeptides (Ac-EEMQRR-amide and H2N-EEMQRR-amide) in anti-wrinkle cosmetics by hydrophilic interaction liquid chromatography-solid phase extraction preparation and hydrophilic interaction liquid chromatography with tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2011, 1218, 7956-7963.	1.8	33
524	Development and evaluation of new zwitterionic Hydrophilic interaction liquid chromatography stationary phases based on 3-P,P-diphenylphosphonium-propylsulfonate. <i>Journal of Chromatography A</i> , 2011, 1218, 8075-8082.	1.8	49

#	ARTICLE	IF	CITATIONS
525	Screening and confirmation analysis of stimulants, narcotics and beta-adrenergic agents in human urine by hydrophilic interaction liquid chromatography coupled to mass spectrometry. <i>Journal of Chromatography A</i> , 2011, 1218, 8156-8167.	1.8	42
526	Characterization of enhanced-fluidity liquid hydrophilic interaction chromatography for the separation of nucleosides and nucleotides. <i>Journal of Chromatography A</i> , 2011, 1218, 8222-8230.	1.8	38
527	Mixed-mode chromatography and its applications to biopolymers. <i>Journal of Chromatography A</i> , 2011, 1218, 8813-8825.	1.8	167
528	Development and validation of a hydrophilic interaction chromatography-tandem mass spectrometry method with on-line polar extraction for the analysis of urinary nucleosides. Potential application in clinical diagnosis. <i>Journal of Chromatography A</i> , 2011, 1218, 9055-9063.	1.8	54
529	Hydrophilic interaction chromatography of seized drugs and related compounds with sub 2 μ m particle columns. <i>Journal of Chromatography A</i> , 2011, 1218, 9336-9344.	1.8	19
530	Characterizing plant cell wall derived oligosaccharides using hydrophilic interaction chromatography with mass spectrometry detection. <i>Journal of Chromatography A</i> , 2011, 1218, 9227-9235.	1.8	76
531	Green Analytical Separation Methods. <i>RSC Green Chemistry</i> , 2011, , 168-198.	0.0	6
532	HILIC-UPLC-MS for Exploratory Urinary Metabolic Profiling in Toxicological Studies. <i>Analytical Chemistry</i> , 2011, 83, 382-390.	3.2	135
534	Characterization of glycoprotein digests with hydrophilic interaction chromatography and mass spectrometry. <i>Analytical Biochemistry</i> , 2011, 417, 80-88.	1.1	65
535	Extraction and Separation Properties of Hydrophilic Compounds Using Novel Water-Holding Adsorbents Bonded with a Zwitter-Ionic Polymer. <i>Chromatographia</i> , 2011, 73, 849-855.	0.7	21
536	Tetrazole-Functionalized Silica for Hydrophilic Interaction Chromatography of Polar Solutes. <i>Chromatographia</i> , 2011, 73, 865-870.	0.7	14
537	Metabonomic Study of Lung Cancer and the Effects of Radiotherapy on Lung Cancer Patients: Analysis of Highly Polar Metabolites by Ultraperformance HILIC Coupled with Q-TOF MS. <i>Chromatographia</i> , 2011, 74, 391-398.	0.7	11
538	Preparation and Evaluation of Poly-L-Lysine Stationary Phase for Hydrophilic Interaction/Reversed-Phase Mixed-Mode Chromatography. <i>Chromatographia</i> , 2011, 74, 523-530.	0.7	41
539	Development of silica-based stationary phases for high-performance liquid chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 399, 3307-3322.	1.9	126
540	Chocolate HILIC phases: development and characterization of novel saccharide-based stationary phases by applying non-enzymatic browning (Maillard reaction) on amino-modified silica surfaces. <i>Analytical and Bioanalytical Chemistry</i> , 2011, 400, 2539-2554.	1.9	20
541	Increasing phosphoproteome coverage and identification of phosphorylation motifs through combination of different HPLC fractionation methods. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2011, 879, 25-34.	1.2	33
542	Chromatographic separation techniques and data handling methods for herbal fingerprints: A review. <i>Analytica Chimica Acta</i> , 2011, 690, 148-161.	2.6	270
543	Liquid chromatography/mass spectrometry compatible approaches for the quantitation of folic acid in fortified juices and cereals using aqueous normal phase conditions. <i>Journal of Chromatography A</i> , 2011, 1218, 2121-2126.	1.8	37

#	ARTICLE	IF	CITATIONS
544	Cyclofructan 6 based stationary phases for hydrophilic interaction liquid chromatography. Journal of Chromatography A, 2011, 1218, 270-279.	1.8	73
545	Enhanced fluidity liquid chromatography for hydrophilic interaction separation of nucleosides. Journal of Chromatography A, 2011, 1218, 5897-5902.	1.8	30
546	Hydrophilic interaction chromatography of nucleoside triphosphates with temperature as a separation parameter. Journal of Chromatography A, 2011, 1218, 5981-5986.	1.8	58
547	Evaluation of different column types for the hydrophilic interaction chromatographic separation of iron-citrate and copper-histidine species from plants. Journal of Chromatography A, 2011, 1218, 4934-4943.	1.8	16
548	Imidazoline type stationary phase for hydrophilic interaction chromatography and reversed-phase liquid chromatography. Journal of Chromatography A, 2011, 1218, 5987-5994.	1.8	38
549	Characterization of carboxylic acids in atmospheric aerosols using hydrophilic interaction liquid chromatography tandem mass spectrometry. Journal of Chromatography A, 2011, 1218, 4417-4425.	1.8	48
550	Hydrophilic Interaction Chromatography (HILIC) in the Analysis of Relevant Quality and Safety Biochemical Compounds in Meat, Poultry and Processed Meats. Food Analytical Methods, 2011, 4, 121-129.	1.3	19
551	Hydrophobic Derivatization of <i>N</i> -linked Glycans for Increased Ion Abundance in Electrospray Ionization Mass Spectrometry. Journal of the American Society for Mass Spectrometry, 2011, 22, 1309-1317.	1.2	71
552	Catch me if you can: Mass spectrometry-based phosphoproteomics and quantification strategies. Proteomics, 2011, 11, 554-570.	1.3	90
553	Current methodologies for proteomics of bacterial surface-exposed and cell envelope proteins. Proteomics, 2011, 11, 3169-3189.	1.3	69
554	Systematic evaluation of acetone and acetonitrile for use in hydrophilic interaction liquid chromatography coupled with electrospray ionization mass spectrometry of basic small molecules. Rapid Communications in Mass Spectrometry, 2011, 25, 3666-3674.	0.7	12
555	Sulfonated cyclofructan 6 based stationary phase for hydrophilic interaction chromatography. Journal of Separation Science, 2011, 34, 1636-1647.	1.3	43
556	Hydrophilic interaction chromatography separation mechanisms of tetracyclines on amino-bonded silica column. Journal of Separation Science, 2011, 34, 1508-1516.	1.3	50
557	Preparation of zwitterionic polymeric monolithic columns for hydrophilic interaction capillary liquid chromatography. Journal of Separation Science, 2011, 34, 2088-2096.	1.3	16
558	Retention of \hat{I}^2 blockers on native titania stationary phase. Journal of Separation Science, 2011, 34, 1805-1810.	1.3	5
559	Simultaneous determination nucleosides in marine organisms using ultrasound-assisted extraction followed by hydrophilic interaction liquid chromatography-electrospray ionization time-of-flight mass spectrometry. Journal of Separation Science, 2011, 34, 2594-2601.	1.3	20
560	Preparation of hydrophilic polymer-grafted polystyrene beads for hydrophilic interaction chromatography via surface-initiated atom transfer radical polymerization. Journal of Separation Science, 2011, 34, 3115-3122.	1.3	18
561	Separation strategies for untargeted metabolomics. Journal of Separation Science, 2011, 34, 3460-3469.	1.3	109

#	ARTICLE	IF	CITATIONS
562	HILIC-MS/MS method development for targeted quantitation of metabolites: Practical considerations from a clinical diagnostic perspective. <i>Journal of Separation Science</i> , 2011, 34, 3527-3537.	1.3	15
563	Grafting of silica with a hydrophilic triol acrylamide polymer via surface-initiated "grafting from" method for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2011, 34, 3123-3130.	1.3	20
565	Recent advances in hydrophilic interaction liquid chromatography (HILIC) for structural glycomics. <i>Electrophoresis</i> , 2011, 32, 3456-3466.	1.3	169
566	Development and validation of a hydrophilic interaction chromatography-tandem mass spectrometry for quantification of nicotine and its metabolites in human maternal and cord sera. <i>Biomedical Chromatography</i> , 2011, 25, 503-510.	0.8	16
567	Quantitative liquid chromatography tandem mass spectrometry analysis of macromolecules using signature peptides in biological fluids. <i>Biomedical Chromatography</i> , 2011, 25, 47-58.	0.8	21
568	LC-MS in analytical toxicology: some practical considerations. <i>Biomedical Chromatography</i> , 2011, 25, 100-123.	0.8	58
569	A novel hydrophilic polystyrene-based beads for hydrophilic interaction chromatography by surface-initiated atom transfer radical polymerization. <i>Chinese Chemical Letters</i> , 2011, 22, 245-248.	4.8	8
570	Hydrophilic interaction liquid chromatography in the separation of a moderately lipophilic drug from its highly polar metabolites—the cardioprotectant dexrazoxane as a model case. <i>Journal of Chromatography A</i> , 2011, 1218, 416-426.	1.8	18
571	The hydrophilic interaction like properties of some reversed phase high performance liquid chromatography columns in the analysis of basic compounds. <i>Journal of Chromatography A</i> , 2011, 1218, 1486-1491.	1.8	18
572	Covalently bonded polysaccharide-modified stationary phase for per aqueous liquid chromatography and hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 1503-1508.	1.8	21
573	Preparation and application of hydrophilic monolithic columns. <i>Journal of Chromatography A</i> , 2011, 1218, 2350-2361.	1.8	89
574	A revisit to the retention mechanism of hydrophilic interaction liquid chromatography using model organic compounds. <i>Journal of Chromatography A</i> , 2011, 1218, 2871-2879.	1.8	117
575	Chromatographic evaluation of reversed-phase/anion-exchange/cation-exchange trimodal stationary phases prepared by electrostatically driven self-assembly process. <i>Journal of Chromatography A</i> , 2011, 1218, 3407-3412.	1.8	51
576	Immobilized metallacarborane as a new type of stationary phase for high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 3029-3036.	1.8	11
577	Investigations on the chromatographic behaviour of zwitterionic stationary phases used in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 5939-5963.	1.8	160
578	Study of retention behaviour and mass spectrometry compatibility in zwitterionic hydrophilic interaction chromatography for the separation of modified nucleosides and nucleobases. <i>Journal of Chromatography A</i> , 2011, 1218, 3994-4001.	1.8	29
579	Novel highly hydrophilic methacrylate-based monolithic column with mixed-mode of hydrophilic and strong cation-exchange interactions for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 4671-4677.	1.8	46
580	Method development and validation for optimised separation of salicylic, acetyl salicylic and ascorbic acid in pharmaceutical formulations by hydrophilic interaction chromatography and response surface methodology. <i>Journal of Chromatography A</i> , 2011, 1218, 5995-6003.	1.8	60

#	ARTICLE	IF	CITATIONS
581	Analysis of oligonucleotides by hydrophilic interaction liquid chromatography coupled to negative ion electrospray ionization mass spectrometry. <i>Journal of Chromatography A</i> , 2011, 1218, 5480-5486.	1.8	60
582	Chromatographic characterization of hydrophilic interaction liquid chromatography stationary phases: Hydrophilicity, charge effects, structural selectivity, and separation efficiency. <i>Journal of Chromatography A</i> , 2011, 1218, 5903-5919.	1.8	168
583	Retention and selectivity of stationary phases for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 5920-5938.	1.8	277
584	Analysis of global DNA methylation by hydrophilic interaction ultra high-pressure liquid chromatography tandem mass spectrometry. <i>Analytical Biochemistry</i> , 2011, 413, 164-170.	1.1	44
585	Investigation of polar stationary phases for the separation of sympathomimetic drugs with nano-liquid chromatography in hydrophilic interaction liquid chromatography mode. <i>Analytica Chimica Acta</i> , 2011, 685, 103-110.	2.6	38
586	Stationary and mobile phases in hydrophilic interaction chromatography: a review. <i>Analytica Chimica Acta</i> , 2011, 692, 1-25.	2.6	610
587	Hydrophilic interaction chromatography of intact, soluble proteins. <i>Journal of Chromatography A</i> , 2011, 1218, 5892-5896.	1.8	46
588	Chromatographic behavior of xanthenes in aqueous normal phase chromatography using titania stationary phase. <i>Journal of Chromatography A</i> , 2011, 1218, 721-725.	1.8	24
589	On-line solid-phase extraction coupled to hydrophilic interaction chromatography- ¹³ C mass spectrometry for the determination of polar drugs. <i>Journal of Chromatography A</i> , 2011, 1218, 5975-5980.	1.8	42
590	HILIC at 21: Reflections and perspective. <i>Journal of Chromatography A</i> , 2011, 1218, 5879.	1.8	17
591	Probing the interaction mode in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2011, 1218, 5880-5891.	1.8	171
592	Spherical mesoporous silica particles by spray drying: Doubling the retention factor of HPLC columns. <i>Microporous and Mesoporous Materials</i> , 2011, 142, 282-291.	2.2	24
593	Analysis of cephalosporins by hydrophilic interaction chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011, 54, 623-628.	1.4	28
594	Rapid hydrophilic interaction chromatography determination of lysine in pharmaceutical preparations with fluorescence detection after postcolumn derivatization with o-phthalaldehyde. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011, 54, 972-978.	1.4	38
595	Chemical profiling of the deacetylase activity of acetyl xylan esterase A (AxeA) variants on chitooligosaccharides using hydrophilic interaction chromatography- ¹³ C mass spectrometry. <i>Journal of Biotechnology</i> , 2011, 155, 257-265.	1.9	19
596	Analysis of polar metabolites by hydrophilic interaction chromatography- ¹³ C MS/MS. <i>Bioanalysis</i> , 2011, 3, 899-912.	0.6	27
597	Determination of Cotinine in Urine and Saliva Using Hydrophilic Interaction Liquid Chromatography and Tandem Mass Spectrometry. <i>Analytical Chemistry Letters</i> , 2011, 1, 25-33.	0.4	0
598	Application of hydrophilic interaction TLC systems for separation of highly polar glycosidic compounds from the flowers of selected <i>Verbascum</i> species. <i>Journal of Planar Chromatography - Modern TLC</i> , 2011, 24, 295-300.	0.6	21

#	ARTICLE	IF	CITATIONS
599	Effect-Directed Analysis of Complex Environmental Contamination. Handbook of Environmental Chemistry, 2011, , .	0.2	16
600	Bioanalytical Methods for the Analysis of Hypoglycemic Drugs and their Main Metabolites. Current Pharmaceutical Analysis, 2011, 7, 141-159.	0.3	0
601	Lipidomics of Glycosphingolipids. Metabolites, 2012, 2, 134-164.	1.3	46
602	Hydrophilic Interaction Liquid Chromatography and Per Aqueous Liquid Chromatography in Fungicides Analysis. Journal of AOAC INTERNATIONAL, 2012, 95, 1362-1370.	0.7	12
603	Increased serum clearance of oligomannose species present on a human IgG1 molecule. MAbs, 2012, 4, 509-520.	2.6	93
604	Evaluation of the Electrostatic Contribution to the Retention of Modified Nucleosides and Nucleobases by Zwitterionic Hydrophilic Interaction Chromatography. , 2012, 2012, 1-5.		1
605	Enrichment strategies for phosphoproteomics: state-of-the-art. Reviews in Analytical Chemistry, 2012, 31, .	1.5	9
606	Utilization of hydrophilic-interaction LC to minimize matrix effects caused by phospholipids. Bioanalysis, 2012, 4, 2049-2058.	0.6	22
607	The Use of Proteomics for Systematic Analysis of Normal and Transformed Hematopoietic Stem Cells. Current Pharmaceutical Design, 2012, 18, 1730-1750.	0.9	9
608	Control of the Electrostatic Effect on DAM-adsorbent for the Water-soluble Compounds by HPLC. Analytical Sciences, 2012, 28, 1109-1111.	0.8	3
609	Preparation of a Hydrophilic Stationary Phase Immobilizing an Acetylated Diallylamineâ€“Acrylamide Copolymer and Its Retention Characteristics for Water-soluble Compounds. Bunseki Kagaku, 2012, 61, 613-619.	0.1	2
610	Recent Developments in High-Performance Liquid Chromatography. , 2012, , 18-49.		1
611	Recent Developments in High-Performance Liquidâ€“Chromatography. , 2012, , 1-32.		0
612	Determination of Histamine in Seafood by Hydrophilic Interaction Chromatography/Tandem Mass Spectrometry. Analytical Sciences, 2012, 28, 179.	0.8	14
613	Simultaneous Analysis of Heterocyclic Amines in Beer and Beer-like Drinks by Hydrophilic Interaction Liquid Chromatography-Tandem Mass Spectrometry. Bunseki Kagaku, 2012, 61, 391-396.	0.1	3
614	Aqueous normal phase liquid chromatography coupled with tandem time-of-flight quadrupole mass spectrometry for determination of zanamivir in human serum. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2012, 906, 58-62.	1.2	15
615	8.12 Chromatographic Separations and Analysis: Macrocyclic Glycopeptide Chiral Stationary Phases. , 2012, , 227-262.		5
616	Analytical strategies for shotgun phosphoproteomics: Status and prospects. Seminars in Cell and Developmental Biology, 2012, 23, 836-842.	2.3	20

#	ARTICLE	IF	CITATIONS
617	Synthesis of poly(<i>N</i> -[tris(hydroxymethyl)methyl]acrylamide) functionalized porous silica for application in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2012, 35, 3257-3269.	1.3	9
618	Toward reading the sequence of short oligonucleotides from their retention factors obtained by means of hydrophilic interaction chromatography and ion-pair interaction reversed-phase liquid chromatography. <i>Journal of Separation Science</i> , 2012, 35, 3227-3234.	1.3	6
619	Evaluation of peptide fractionation strategies used in proteome analysis. <i>Journal of Separation Science</i> , 2012, 35, 3293-3301.	1.3	15
620	Development of Mucoadhesive Drug Delivery System Using Phenylboronic Acid Functionalized Poly(D,L-lactide) β -Dextran Nanoparticles. <i>Macromolecular Bioscience</i> , 2012, 12, 1622-1626.	2.1	26
621	The effect of temperature and mobile phase composition on separation mechanism of flavonoid compounds on hydrosilated silica-based columns. <i>Journal of Chromatography A</i> , 2012, 1245, 98-108.	1.8	43
622	Phosphoproteomics and molecular cardiology: Techniques, applications and challenges. <i>Journal of Molecular and Cellular Cardiology</i> , 2012, 53, 354-368.	0.9	17
623	A comparison of the chromatographic properties of silica gel and silicon hydride modified silica gels. <i>Journal of Chromatography A</i> , 2012, 1263, 61-67.	1.8	23
624	New Silica Gel-Based Monolithic Column for Nano-Liquid Chromatography, used in the HILIC Mode. <i>Journal of Chromatographic Science</i> , 2012, 50, 649-657.	0.7	17
625	Maximizing the speed of separations for industrial problems. <i>Journal of Chromatography A</i> , 2012, 1261, 69-77.	1.8	3
626	Comprehensive hydrophilic interaction and ion-pair reversed-phase liquid chromatography for analysis of di- to deca-oligonucleotides. <i>Journal of Chromatography A</i> , 2012, 1255, 237-243.	1.8	58
627	Determination of alkane/water partition coefficients of polar compounds using hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2012, 1260, 164-168.	1.8	9
628	Effect of silica gel modification with cyclofructans on properties of hydrophilic interaction liquid chromatography stationary phases. <i>Journal of Chromatography A</i> , 2012, 1257, 58-65.	1.8	35
629	HYDROPHILIC INTERACTION PLANAR CHROMATOGRAPHY OF GEOMETRICAL ISOMERS OF SELECTED Co(III) COMPLEXES. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2012, 35, 1289-1297.	0.5	2
630	Identification of mercury and other metals complexes with metallothioneins in dolphin liver by hydrophilic interaction liquid chromatography with the parallel detection by ICP MS and electrospray hybrid linear/orbital trap MS/MS. <i>Metallomics</i> , 2012, 4, 473.	1.0	31
631	Systematic Comparison of Reverse Phase and Hydrophilic Interaction Liquid Chromatography Platforms for the Analysis of N-Linked Glycans. <i>Analytical Chemistry</i> , 2012, 84, 8198-8206.	3.2	23
632	Analytical techniques for the detection of β -amino- β -methylaminopropionic acid. <i>Analyst, The</i> , 2012, 137, 1991.	1.7	59
633	Expression, purification, and reconstitution of the transmembrane domain of the human amyloid precursor protein for NMR studies. <i>Protein Expression and Purification</i> , 2012, 81, 11-17.	0.6	9
634	Hydrophilic interaction chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2012, 37, 48-60.	5.8	135

#	ARTICLE	IF	CITATIONS
635	A new peptide-silica bio-inspired stationary phase with an improved approach for hydrophilic interaction liquid chromatography. <i>Analyst</i> , The, 2012, 137, 4907.	1.7	18
636	LC-MS metabolomics of polar compounds. <i>Bioanalysis</i> , 2012, 4, 1235-1243.	0.6	47
637	Hydrophilic interaction chromatography (HILIC) for LC-MS/MS analysis of monoamine neurotransmitters. <i>Bioanalysis</i> , 2012, 4, 783-794.	0.6	33
638	Determination of caprolactam and 6-aminocaproic acid in human urine using hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 885-886, 61-65.	1.2	12
639	Rapid and reliable quantitation of amino acids and myo-inositol in mouse brain by high performance liquid chromatography and tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 893-894, 15-20.	1.2	27
640	Hydrophilic interaction liquid chromatography/positive ion electrospray mass spectrometry for the quantification of deferasirox, an oral iron chelator, in human plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 893-894, 114-120.	1.2	14
641	Analytical aspects of biosimilarity issues of protein drugs. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 69, 185-195.	1.4	54
642	Recent advances in peptide separation by multidimensional liquid chromatography for proteome analysis. <i>Journal of Proteomics</i> , 2012, 75, 3791-3813.	1.2	141
643	Matrine- and oxymatrine-imprinted monodisperse polymers prepared by precipitation polymerization and their applications for the selective extraction of matrine-type alkaloids from <i>Sophora flavescens</i> Aiton. <i>Journal of Chromatography A</i> , 2012, 1248, 18-23.	1.8	45
644	Evaluation of hydrophilic interaction chromatography (HILIC) versus C18 reversed-phase chromatography for targeted quantification of peptides by mass spectrometry. <i>Journal of Chromatography A</i> , 2012, 1264, 31-39.	1.8	34
645	Hypercrosslinked Large Surface Area Porous Polymer Monoliths for Hydrophilic Interaction Liquid Chromatography of Small Molecules Featuring Zwitterionic Functionalities Attached to Gold Nanoparticles Held in Layered Structure. <i>Analytical Chemistry</i> , 2012, 84, 8457-8460.	3.2	101
646	Retention characteristics of water-soluble compounds on water-holding adsorbents with immobilized zwitterionic copolymers having different ionicities. <i>Journal of Separation Science</i> , 2012, 35, 2834-2839.	1.3	6
647	Glycopeptide enrichment and separation for protein glycosylation analysis. <i>Journal of Separation Science</i> , 2012, 35, 2341-2372.	1.3	138
648	Rapid and effective removal of perfluorooctanoic acid from proteomics samples. <i>Proteomics</i> , 2012, 12, 2271-2275.	1.3	0
649	Temporal profiling of the heat-stable proteome during late maturation of <i>Medicago truncatula</i> seeds identifies a restricted subset of late embryogenesis abundant proteins associated with longevity. <i>Plant, Cell and Environment</i> , 2012, 35, 1440-1455.	2.8	122
651	Preparation of imidazole-functionalized silica by surface-initiated atom transfer radical polymerization and its application for hydrophilic interaction chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 404, 1477-1484.	1.9	13
653	ZIC-CHILIC as a fractionation method for sensitive and powerful shotgun proteomics. <i>Nature Protocols</i> , 2012, 7, 2041-2055.	5.5	31
654	Methods to Isolate and Identify New Plant Signaling Peptides. <i>Signaling and Communication in Plants</i> , 2012, , 217-239.	0.5	3

#	ARTICLE	IF	CITATIONS
656	Modern approach for determination of lactulose, mannitol and sucrose in human urine using HPLC-MS/MS for the studies of intestinal and upper digestive tract permeability. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 907, 34-40.	1.2	40
657	Quantitation of cotinine and its metabolites in rat plasma and brain tissue by hydrophilic interaction chromatography tandem mass spectrometry (HILIC-MS/MS). <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 907, 117-125.	1.2	26
658	Hydrophilic interaction/weak cation-exchange mixed-mode chromatography for chitooligosaccharides separation. <i>Carbohydrate Research</i> , 2012, 361, 195-199.	1.1	22
659	Temperature Pulsing for Controlling Chromatographic Resolution in Capillary Liquid Chromatography. <i>Analytical Chemistry</i> , 2012, 84, 3362-3368.	3.2	17
660	Reaching for the deep proteome: Recent nano liquid chromatography coupled with tandem mass spectrometry-based studies on the deep proteome. <i>Archives of Pharmacal Research</i> , 2012, 35, 1861-1870.	2.7	11
661	A simple graphical representation of selectivity in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1260, 126-131.	1.8	51
662	Mass spectrometry-based phosphoproteomics in cancer research. <i>Frontiers in Biology</i> , 2012, 7, 566-586.	0.7	1
663	New surface-confined ionic liquid stationary phases with enhanced chromatographic selectivity and stability by co-immobilization of polymerizable anion and cation pairs. <i>Chemical Communications</i> , 2012, 48, 1299-1301.	2.2	71
664	Enrichment techniques employed in phosphoproteomics. <i>Amino Acids</i> , 2012, 43, 1025-1047.	1.2	170
665	Mixed mode HILIC/anion exchange separations on latex coated silica monoliths. <i>Talanta</i> , 2012, 100, 313-319.	2.9	34
667	Plant Signaling Peptides. <i>Signaling and Communication in Plants</i> , 2012, , .	0.5	2
668	Ultra-high performance liquid chromatography-tandem mass spectrometry for the analysis of phenicol drugs and florfenicol-amine in foods. <i>Analyst, The</i> , 2012, 137, 2486.	1.7	41
669	Facile Preparation of Zwitterionic Organic-Silica Hybrid Monolithic Capillary Column with an Improved "One-Pot" Approach for Hydrophilic-Interaction Liquid Chromatography (HILIC). <i>Analytical Chemistry</i> , 2012, 84, 2721-2728.	3.2	111
670	New poly(ionic liquid)-grafted silica multi-mode stationary phase for anion-exchange/reversed-phase/hydrophilic interaction liquid chromatography. <i>Analyst, The</i> , 2012, 137, 2553.	1.7	108
671	Zwitterionic silica-based monolithic capillary columns for isocratic and gradient hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1270, 178-185.	1.8	33
672	Advances in Quantitative Phosphoproteomics. <i>Analytical Chemistry</i> , 2012, 84, 735-746.	3.2	87
673	Direct Analysis of Underivatized Amino Acids in Plant Extracts by LC-MS-MS. <i>Methods in Molecular Biology</i> , 2012, 828, 317-328.	0.4	24
674	Investigation on synthesis of spheres-on-sphere silica particles and their assessment for high performance liquid chromatography applications. <i>Journal of Chromatography A</i> , 2012, 1270, 194-203.	1.8	30

#	ARTICLE	IF	CITATIONS
675	Separation Technique for the Determination of Highly Polar Metabolites in Biological Samples. <i>Metabolites</i> , 2012, 2, 496-515.	1.3	43
676	HPLC-MS/MS of Highly Polar Compounds. , 0, , .		1
677	Clinical Implementation of Metabolomics. , 0, , .		8
679	Vol 2, No 1 (2012). <i>Journal of Integrated OMICS</i> , 2012, 2, .	0.5	0
680	High Performance Liquid Chromatographic Approaches to Mass Spectrometry Based Metabolomics. <i>Current Metabolomics</i> , 2012, 1, 58-83.	0.5	3
681	Development of a liquid chromatographic method based on ultraviolet-visible and electrospray ionization mass spectrometric detection for the identification of nitrocatechols and related tracers in biomass burning atmospheric organic aerosol. <i>Rapid Communications in Mass Spectrometry</i> , 2012, 26, 793-804.	0.7	61
682	Metabolite Profiling Identifies a Key Role for Glycine in Rapid Cancer Cell Proliferation. <i>Science</i> , 2012, 336, 1040-1044.	6.0	1,201
683	Comprehensive HILIC – RPLC with mass spectrometry detection for the analysis of saponins in <i>Panax notoginseng</i> . <i>Analyst</i> , The, 2012, 137, 2239.	1.7	46
684	Limitation of predictive 2D liquid chromatography in reducing the database search space in shotgun proteomics: <i>in silico</i> studies. <i>Journal of Separation Science</i> , 2012, 35, 1771-1778.	1.3	4
685	Comprehensive characterization of <i>Stevia Rebaudiana</i> using two-dimensional reversed-phase liquid chromatography/hydrophilic interaction liquid chromatography. <i>Journal of Separation Science</i> , 2012, 35, 1821-1827.	1.3	61
686	Assessment of Î²-lactams retention in hydrophilic interaction chromatography applying <i>in silico</i> design. <i>Journal of Separation Science</i> , 2012, 35, 1424-1431.	1.3	11
687	Profiling Phlorotannins in Brown Macroalgae by Liquid Chromatography-High Resolution Mass Spectrometry. <i>Phytochemical Analysis</i> , 2012, 23, 547-553.	1.2	103
688	Quantitation of Galactosides in <i>Rehmannia glutinosa</i> by Hydrophilic Interaction Chromatography-Evaporative Light Scattering Detector. <i>Phytochemical Analysis</i> , 2012, 23, 607-612.	1.2	4
689	Detection of <i>myo</i> -inositol trispyrophosphate in equine urine and plasma by hydrophilic interaction chromatography-tandem mass spectrometry. <i>Drug Testing and Analysis</i> , 2012, 4, 355-361.	1.6	7
691	A Molecular Dynamics Study on the Partitioning Mechanism in Hydrophilic Interaction Chromatography. <i>Angewandte Chemie - International Edition</i> , 2012, 51, 6251-6254.	7.2	102
692	Superficially porous particles columns for super fast HPLC separations. <i>Biomedical Chromatography</i> , 2012, 26, 1001-1008.	0.8	47
693	Mass spectrometry-based proteomics and peptidomics for systems biology and biomarker discovery. <i>Frontiers in Biology</i> , 2012, 7, 313-335.	0.7	31
694	Determination of trehalose-6-phosphate in <i>Arabidopsis thaliana</i> seedlings by hydrophilic-interaction liquid chromatography-mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 403, 1353-1360.	1.9	21

#	ARTICLE	IF	CITATIONS
695	Strategies for the Analysis of Pharmaceutical Cocrystals Using HPLC with Charged Aerosol Detection. <i>Chromatographia</i> , 2012, 75, 321-328.	0.7	2
696	Behavior and Retention Models of Melamine and Its Hydrolysis Products. <i>Chromatographia</i> , 2012, 75, 457-467.	0.7	12
697	Hydrosilated silica-based columns: The effects of mobile phase and temperature on dual hydrophilic-reversed-phase separation mechanism of phenolic acids. <i>Journal of Chromatography A</i> , 2012, 1228, 125-134.	1.8	35
698	Preparation of methacrylate-based monolith for capillary hydrophilic interaction chromatography and its application in determination of nucleosides in urine. <i>Journal of Chromatography A</i> , 2012, 1228, 183-192.	1.8	47
699	Preparation and chromatographic evaluation of a cysteine-bonded zwitterionic hydrophilic interaction liquid chromatography stationary phase. <i>Journal of Chromatography A</i> , 2012, 1228, 175-182.	1.8	100
700	New trends in fast liquid chromatography for food and environmental analysis. <i>Journal of Chromatography A</i> , 2012, 1228, 298-323.	1.8	211
701	An efficient hydrophilic interaction liquid chromatography separation of 7 phospholipid classes based on a diol column. <i>Journal of Chromatography A</i> , 2012, 1220, 26-34.	1.8	75
702	A molecular simulation study of the effects of stationary phase and solute chain length in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1223, 24-34.	1.8	32
703	Hydrophilic interaction liquid chromatography coupled to mass spectrometry for the characterization of prebiotic galactooligosaccharides. <i>Journal of Chromatography A</i> , 2012, 1220, 57-67.	1.8	53
704	Comparison of reversed-phase/cation-exchange/anion-exchange trimodal stationary phases and their use in active pharmaceutical ingredient and counterion determinations. <i>Journal of Chromatography A</i> , 2012, 1232, 190-195.	1.8	52
705	Design and evaluation of hydrolytically stable bidentate urea-type stationary phases for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2012, 1232, 196-211.	1.8	31
706	A novel click lysine zwitterionic stationary phase for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2012, 1223, 47-52.	1.8	42
707	Combination of off-line two-dimensional hydrophilic interaction liquid chromatography for polar fraction and two-dimensional hydrophilic interaction liquid chromatography—reversed-phase liquid chromatography for medium-polar fraction in a traditional Chinese medicine. <i>Journal of Chromatography A</i> , 2012, 1224, 61-69.	1.8	53
708	Magnetic bead-based hydrophilic interaction liquid chromatography for glycopeptide enrichments. <i>Journal of Chromatography A</i> , 2012, 1224, 70-78.	1.8	51
709	The influence of a carbon layer deposited on a zirconia surface on the retention of polar analytes in an organic rich mobile phase. <i>Journal of Chromatography A</i> , 2012, 1232, 242-247.	1.8	6
710	The utility of porous graphitic carbon as a stationary phase in proteomics workflows: Two-dimensional chromatography of complex peptide samples. <i>Journal of Chromatography A</i> , 2012, 1232, 276-280.	1.8	13
711	Development of a hydrophilic interaction liquid chromatography—mass spectrometry method for detection and quantification of urea thermal decomposition by-products in emission from diesel engine employing selective catalytic reduction technology. <i>Journal of Chromatography A</i> , 2012, 1229, 208-215.	1.8	15
712	Preparation and characterization of methacrylate-based monolith for capillary hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2012, 1230, 54-60.	1.8	63

#	ARTICLE	IF	CITATIONS
713	Study of the retention behavior in zwitterionic hydrophilic interaction chromatography of isomeric hydroxy- and aminobenzoic acids. <i>Journal of Chromatography A</i> , 2012, 1235, 60-67.	1.8	76
714	Preparation and characterization of hybrid-silica monolithic column with mixed-mode of hydrophilic and strong anion-exchange interactions for pressurized capillary electrochromatography. <i>Journal of Chromatography A</i> , 2012, 1239, 56-63.	1.8	31
715	Screening for low molecular weight compounds in fish meal solubles by hydrophilic interaction liquid chromatography coupled to mass spectrometry. <i>Food Chemistry</i> , 2012, 130, 739-745.	4.2	13
716	Optimization of a solid phase extraction and hydrophilic interaction liquid chromatography-tandem mass spectrometry method for the determination of metformin in dietary supplements and herbal medicines. <i>Food Chemistry</i> , 2012, 133, 482-488.	4.2	38
717	Comparison of the quantitative performances and measurement uncertainty estimates obtained during method validation versus routine applications of a novel hydrophilic interaction chromatography method for the determination of cidofovir in human plasma. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 57, 153-165.	1.4	18
718	Method development for pharmaceuticals: Some solutions for tuning selectivity in reversed phase and hydrophilic interaction liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 63, 95-105.	1.4	33
719	Determination of swainsonine in the endophytic <i>Undifilum</i> fungi by high-performance liquid chromatography with evaporative light-scattering detector. <i>Toxicon</i> , 2012, 60, 44-49.	0.8	13
720	Hydrophilic monolith with ethylene glycol-based grafts prepared via surface confined thiol-ene click photoaddition. <i>Polymer</i> , 2012, 53, 29-36.	1.8	52
721	Hydrophilic interaction chromatography in drug analysis. <i>Open Chemistry</i> , 2012, 10, 534-553.	1.0	11
722	Organic monoliths for hydrophilic interaction electrochromatography/chromatography and immunoaffinity chromatography. <i>Electrophoresis</i> , 2012, 33, 251-261.	1.3	37
723	Hydrophilic interaction chromatography for the analysis of aminoglycosides. <i>Journal of Separation Science</i> , 2012, 35, 498-504.	1.3	49
724	Size-tunable nanoparticles composed of dextran-b-poly(D,L-lactide) for drug delivery applications. <i>Nano Research</i> , 2012, 5, 49-61.	5.8	64
725	Simultaneous quantitation of histamine and its major metabolite 1-methylhistamine in brain dialysates by using precolumn derivatization prior to HILIC-MS/MS analysis. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 449-459.	1.9	25
726	Hydrophilic interaction liquid chromatography (HILIC) - a powerful separation technique. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 231-247.	1.9	950
727	Measuring multiple neurochemicals and related metabolites in blood and brain of the rhesus monkey by using dual microdialysis sampling and capillary hydrophilic interaction chromatography-mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 2545-2554.	1.9	14
728	The herbalome - an attempt to globalize Chinese herbal medicine. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 402, 573-581.	1.9	25
729	The application of quantification techniques in proteomics for biomedical research. <i>Mass Spectrometry Reviews</i> , 2013, 32, 1-26.	2.8	60
730	Preparation, characterization, and application of a new stationary phase containing different kinds of amine groups. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 8311-8318.	1.9	13

#	ARTICLE	IF	CITATIONS
731	Rapid and sensitive HILIC-MS/MS analysis of carnitine and acetylcarnitine in biological fluids. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 7397-7404.	1.9	14
732	Thorough investigation of the retention mechanisms and retention behavior of amides and sulfonamides on amino column in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1301, 27-37.	1.8	15
733	Additional investigations into the retention mechanism of hydrophilic interaction liquid chromatography by linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2013, 1301, 98-110.	1.8	41
734	Dehydration of Ethanol by Facile Synthesized Glucose-Based Silica. <i>Applied Biochemistry and Biotechnology</i> , 2013, 169, 1056-1068.	1.4	7
735	Green chromatography. <i>Journal of Chromatography A</i> , 2013, 1307, 1-20.	1.8	217
736	Selectivity differences of water-soluble vitamins separated on hydrophilic interaction stationary phases. <i>Journal of Separation Science</i> , 2013, 36, 1897-1903.	1.3	5
737	Chromatographic properties of silica modified with 6,10-ionene and sodium lignosulfonate. <i>Journal of Analytical Chemistry</i> , 2013, 68, 450-457.	0.4	2
738	Effects of the operation parameters on HILIC separation of flavonoids on zwitterionic column. <i>Talanta</i> , 2013, 115, 284-290.	2.9	37
740	Retention Mechanism Studies of Selected Amino Acids and Vitamin B6 on HILIC Columns with Evaporative Light Scattering Detection. <i>Chromatographia</i> , 2013, 76, 929-937.	0.7	21
741	Identification of Sialylated Oligosaccharides Derived from Ovine and Caprine Caseinomacropeptide by Graphitized Carbon Liquid Chromatography-Electrospray Ionization Ion Trap Tandem Mass Spectrometry. <i>Food Analytical Methods</i> , 2013, 6, 814-825.	1.3	3
743	Liquid chromatography-mass spectrometry for metabolic footprinting of co-cultures of lactic and propionic acid bacteria. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 8151-8170.	1.9	13
744	Retention behavior of proton pump inhibitors using immobilized polysaccharide-derived chiral stationary phases with organic-aqueous mobile phases. <i>Journal of Chromatography A</i> , 2013, 1304, 147-153.	1.8	28
745	Liquid chromatographic-mass spectrometric method for simultaneous determination of small organic acids potentially contributing to acidosis in severe malaria. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 941, 116-122.	1.2	33
746	Determination of naloxone-3-glucuronide in human plasma and urine by HILIC-MS/MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 942-943, 83-87.	1.2	11
747	A Sulfonic-Azobenzene-Grafted Silica Amphiphilic Material: A Versatile Stationary Phase for Mixed-Mode Chromatography. <i>Chemistry - A European Journal</i> , 2013, 19, 18004-18010.	1.7	44
748	An improved approach to hydrophilic interaction chromatography of peptides: Salt gradients in the presence of high isocratic acetonitrile concentrations. <i>Journal of Chromatography A</i> , 2013, 1277, 15-25.	1.8	16
749	Separation of carbohydrates using hydrophilic interaction liquid chromatography. <i>Carbohydrate Research</i> , 2013, 379, 13-17.	1.1	58
750	Milestones in the Development of Liquid Chromatography. , 2013, , 1-17.		3

#	ARTICLE	IF	CITATIONS
751	How Ternary Mobile Phases Allow Tuning of Analyte Retention in Hydrophilic Interaction Liquid Chromatography. <i>Analytical Chemistry</i> , 2013, 85, 8850-8856.	3.2	27
752	Advances in the development of organic polymer monolithic columns and their applications in food analysisâ€”A review. <i>Journal of Chromatography A</i> , 2013, 1313, 37-53.	1.8	105
753	Ion-exchange and hydrophobic interactions affecting selectivity for neutral and charged solutes on three structurally similar agglomerated ion-exchange and mixed-mode stationary phases. <i>Analytica Chimica Acta</i> , 2013, 803, 143-153.	2.6	37
754	Thermodynamic studies of a zwitterionic stationary phase in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1272, 81-89.	1.8	32
755	Water uptake on polar stationary phases under conditions for hydrophilic interaction chromatography and its relation to solute retention. <i>Journal of Chromatography A</i> , 2013, 1320, 33-47.	1.8	100
756	Preparation of a Novel Amino-Phosphate Zwitterionic Stationary Phase for Hydrophilic Interaction Chromatography. <i>Chromatographia</i> , 2013, 76, 1569-1576.	0.7	12
757	Comparative characterization of hydrophilic interaction liquid chromatography columns by linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2013, 1273, 73-94.	1.8	84
758	Hydrophilic interaction chromatographic analysis of anthocyanins. <i>Journal of Chromatography A</i> , 2013, 1319, 127-140.	1.8	50
759	Main Interactions and Influences of the Chromatographic Parameters in HILIC Separations. <i>Journal of Chromatographic Science</i> , 2013, 51, 684-693.	0.7	132
760	Properties of two amideâ€”based hydrophilic interaction liquid chromatography columns. <i>Journal of Separation Science</i> , 2013, 36, 2421-2429.	1.3	27
761	Zwitterionic hydrophilic interaction chromatography coupled with post-column derivatization for the analysis of glutathione in wine samples. <i>Analytica Chimica Acta</i> , 2013, 795, 75-81.	2.6	18
762	Tandem LC columns for the simultaneous retention of polar and nonpolar molecules in comprehensive metabolomics analysis. <i>Journal of Separation Science</i> , 2013, 36, 3478-3485.	1.3	45
763	A facile and efficient strategy to enhance hydrophilicity of zwitterionic sulfoalkylbetaine type monoliths. <i>Journal of Chromatography A</i> , 2013, 1301, 88-97.	1.8	33
764	Polyacrylamide brush layer for hydrophilic interaction liquid chromatography of intact glycoproteins. <i>Journal of Chromatography A</i> , 2013, 1301, 156-161.	1.8	36
765	Domoic acid at trace levels in lagoon waters: assessment of a method using internal standard quantification. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 9113-9123.	1.9	17
766	Preparation and Characterization of the Neomycin-Bonded Silica Stationary Phase for Hydrophilic-Interaction Chromatography. <i>Chromatographia</i> , 2013, 76, 459-465.	0.7	7
767	Structural analysis of glycoprotein sialylation â€” Part I: pre-LC-MS analytical strategies. <i>RSC Advances</i> , 2013, 3, 22683.	1.7	46
768	Glutathione-based zwitterionic stationary phase for hydrophilic interaction/cation-exchange mixed-mode chromatography. <i>Journal of Chromatography A</i> , 2013, 1314, 63-69.	1.8	76

#	ARTICLE	IF	CITATIONS
769	Quantitative proteomics analysis of varicose veins: Identification of a set of differentially expressed proteins related to ATP generation and utilization. <i>Kaohsiung Journal of Medical Sciences</i> , 2013, 29, 594-605.	0.8	12
770	Comparison of Resins for Metal Oxide Affinity Chromatography with Mass Spectrometry Detection for the Determination of Phosphopeptides. <i>Analytical Letters</i> , 2013, 46, 1505-1524.	1.0	7
771	Determination of eight artificial sweeteners in wastewater by hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Analytical Methods</i> , 2013, 5, 3825.	1.3	38
772	An eco-friendly hydrophilic interaction HPLC method for the determination of renal function biomarkers, creatinine and uric acid, in human fluids. <i>Analytical Methods</i> , 2013, 5, 1307.	1.3	46
773	Equipment and metabolite identification (ID) strategies for mass-based metabolomic analysis. , 2013, , 3-28.		1
774	Hydrophilic Strong Anion Exchange (hSAX) Chromatography for Highly Orthogonal Peptide Separation of Complex Proteomes. <i>Journal of Proteome Research</i> , 2013, 12, 2449-2457.	1.8	60
775	Efficient separation of oxidized cello-oligosaccharides generated by cellulose degrading lytic polysaccharide monooxygenases. <i>Journal of Chromatography A</i> , 2013, 1271, 144-152.	1.8	151
776	The role of stearic acid in ascorbic acid protection from degradation: a heterogeneous system for homogeneous thermodynamic data. <i>New Journal of Chemistry</i> , 2013, 37, 761.	1.4	21
777	HPLC Analysis. , 2013, , 465-519.		2
778	Poly(triallyl isocyanurate-co-ethylene dimethacrylate-co-alkyl methacrylate) stationary phases in the chromatographic separation of hydrophilic solutes. <i>Journal of Chromatography A</i> , 2013, 1272, 65-72.	1.8	11
779	Recent Developments in High-Performance Liquid Chromatography Stationary Phases. <i>Analytical Chemistry</i> , 2013, 85, 579-589.	3.2	122
780	Analytical glycobiology at high sensitivity: current approaches and directions. <i>Glycoconjugate Journal</i> , 2013, 30, 89-117.	1.4	57
781	Preparation and application of novel zwitterionic monolithic column for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2013, 36, 262-269.	1.3	30
782	Large volume injection of aqueous peptide samples on a monolithic silica based zwitterionic-hydrophilic interaction liquid chromatography system for characterization of posttranslational modifications. <i>Journal of Chromatography A</i> , 2013, 1317, 129-137.	1.8	16
783	Gradient elution in aqueous normal-phase liquid chromatography on hydrosilated silica-based stationary phases. <i>Journal of Chromatography A</i> , 2013, 1286, 111-118.	1.8	44
784	Practical investigation of the factors that affect the selectivity in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2013, 1276, 33-46.	1.8	93
785	Neutral, charged and stratified polar monoliths for hydrophilic interaction capillary electrochromatography. <i>Journal of Chromatography A</i> , 2013, 1317, 77-84.	1.8	20
786	Application of two new LC-ESI-MS methods for improved detection of intact polar lipids (IPLs) in environmental samples. <i>Organic Geochemistry</i> , 2013, 59, 10-21.	0.9	106

#	ARTICLE	IF	CITATIONS
787	Hydrophilic interaction liquid chromatography with tandem mass spectrometric detection applied for analysis of pteridines in two Graphosoma species (Insecta: Heteroptera). <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013, 930, 82-89.	1.2	16
788	A novel amide stationary phase for hydrophilic interaction liquid chromatography and ion chromatography. <i>Talanta</i> , 2013, 115, 129-132.	2.9	40
789	Epoxy-based monoliths for capillary liquid chromatography of small and large molecules. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 2233-2244.	1.9	18
790	Nanoscale reversed-phase liquid chromatography–mass spectrometry of permethylated N-glycans. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 2469-2480.	1.9	25
791	Analysis of magnesium from magnesium stearate in pharmaceutical tablet formulations using hydrophilic interaction liquid chromatography with nano quantity analyte detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2013, 78-79, 112-117.	1.4	8
792	Discovery of the Antibiotic Phosacetamycin via a New Mass Spectrometry-Based Method for Phosphonic Acid Detection. <i>ACS Chemical Biology</i> , 2013, 8, 908-913.	1.6	30
793	New Advances in Separation Science for Metabolomics: Resolving Chemical Diversity in a Post-Genomic Era. <i>Chemical Reviews</i> , 2013, 113, 2437-2468.	23.0	298
794	Cross-linker effects on the separation efficiency on (poly)methacrylate capillary monolithic columns. Part II. Aqueous normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2013, 1289, 47-57.	1.8	42
795	Protein Analysis by Shotgun/Bottom-up Proteomics. <i>Chemical Reviews</i> , 2013, 113, 2343-2394.	23.0	1,171
796	Facile preparation of organic-silica hybrid monolith for capillary hydrophilic liquid chromatography based on “thiol-ene”-click chemistry. <i>Journal of Chromatography A</i> , 2013, 1284, 118-125.	1.8	53
797	Glycocapture–based proteomics for secretome analysis. <i>Proteomics</i> , 2013, 13, 512-525.	1.3	24
798	Natural product isolation – how to get from biological material to pure compounds. <i>Natural Product Reports</i> , 2013, 30, 525.	5.2	290
799	A rapid determination method for ethylenethiourea in potato and cucumber by modified QuEChERS – High performance liquid chromatography–tandem mass spectrometry. <i>Food Chemistry</i> , 2013, 138, 1355-1359.	4.2	21
800	Improved sulfoalkylbetaine–based organic–silica hybrid monolith for high efficient hydrophilic interaction liquid chromatography of polar compounds. <i>Electrophoresis</i> , 2013, 34, 1877-1885.	1.3	12
801	Preparation of a dextran sulfate functionalized monolithic silica column for hydrophilic interaction/cation exchange chromatography. <i>Analytical Methods</i> , 2013, 5, 3074.	1.3	7
802	A novel ionic-bonded cellulose stationary phase for saccharide separation. <i>Journal of Chromatography A</i> , 2013, 1291, 56-63.	1.8	15
803	Hydrophilic interaction chromatography with aerosol-based detectors (ELSD, CAD, NQAD) for polar compounds lacking a UV chromophore in an intravenous formulation. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2013, 78-79, 14-18.	1.4	14
804	Comparison of reversed-phase and hydrophilic interaction liquid chromatography for the quantification of ephedrines using medium-resolution accurate mass spectrometry. <i>Journal of Chromatography A</i> , 2013, 1289, 37-46.	1.8	43

#	ARTICLE	IF	CITATIONS
805	Disaccharide Analysis of Glycosaminoglycans Using Hydrophilic Interaction Chromatography and Mass Spectrometry. <i>Analytical Chemistry</i> , 2013, 85, 1138-1145.	3.2	64
806	Hydrophilic interaction liquid chromatography columns classification by effect of solvation and chemometric methods. <i>Journal of Chromatography A</i> , 2013, 1278, 89-97.	1.8	64
807	Polyanionic and polyzwitterionic azobenzene ionic liquid-functionalized silica materials and their chromatographic applications. <i>Chemical Communications</i> , 2013, 49, 2454.	2.2	40
808	Optimization of hydrophilic interaction LC by univariate and multivariate methods and its combination with salting-out liquid-liquid extraction for the determination of antihypertensive drugs in the environmental waters. <i>Journal of Separation Science</i> , 2013, 36, 1007-1014.	1.3	14
809	Application of hydrophilic interaction chromatography in phytochemical analysis. <i>Acta Chromatographica</i> , 2013, 25, 1-25.	0.7	18
810	Determination of saccharides in aerosols by hydrophilic interaction liquid chromatography-evaporative light scattering detection after preconcentration/clean-up on magnetic titania. <i>Analytical Methods</i> , 2013, 5, 2802.	1.3	2
811	Simultaneous separation of hydrophobic and polar bases using a silica hydride stationary phase. <i>Journal of Separation Science</i> , 2013, 36, 1209-1216.	1.3	16
812	Screening of the most relevant parameters for method development in ultra-high performance hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2013, 1282, 72-83.	1.8	68
813	Development of a fast robust derivatization method of an extremely polar PET radiopharmaceutical: a critical aspect for starting a clinical trial. <i>Tetrahedron Letters</i> , 2013, 54, 2583-2586.	0.7	2
814	High-sensitivity Analytical Approaches for the Structural Characterization of Glycoproteins. <i>Chemical Reviews</i> , 2013, 113, 2668-2732.	23.0	276
815	Clinical Chemistry, Including Therapeutic Drug Monitoring and Biomarkers for Diagnosis. , 2013, , 611-631.		2
816	Comparison between the intra-particle diffusivity in the hydrophilic interaction chromatography and reversed phase liquid chromatography modes. Impact on the column efficiency. <i>Journal of Chromatography A</i> , 2013, 1297, 85-95.	1.8	56
817	Development and evaluation of new imidazolium-based zwitterionic stationary phases for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2013, 1286, 137-145.	1.8	83
819	Hydrophilic Interaction Liquid Chromatography. , 2013, , 105-119.		3
820	Bioanalytical LC separation techniques for quantitative analysis of free amino acids in human plasma. <i>Bioanalysis</i> , 2013, 5, 495-512.	0.6	19
821	Fast and simultaneous analysis of biothiols by high-performance liquid chromatography with fluorescence detection under hydrophilic interaction chromatography conditions. <i>Analyst</i> , The, 2013, 138, 3802.	1.7	64
822	Mass transfer mechanism in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2013, 1302, 55-64.	1.8	33
823	Structural Glycomic Analyses at High Sensitivity: A Decade of Progress. <i>Annual Review of Analytical Chemistry</i> , 2013, 6, 237-265.	2.8	57

#	ARTICLE	IF	CITATIONS
824	Automated Assignments of N- and O-Site Specific Glycosylation with Extensive Glycan Heterogeneity of Glycoprotein Mixtures. <i>Analytical Chemistry</i> , 2013, 85, 5666-5675.	3.2	69
825	Reversed-phase fused-core HPLC modeling of peptides. <i>Journal of Pharmaceutical Analysis</i> , 2013, 3, 93-101.	2.4	15
826	Preparation, chromatographic evaluation and comparison between linear peptide- and cyclopeptide-bonded stationary phases. <i>Talanta</i> , 2013, 109, 152-159.	2.9	16
830	Five different columns in the analysis of basic drugs in hydrophilic interaction liquid chromatography. <i>Open Chemistry</i> , 2013, 11, 1150-1162.	1.0	5
831	Carboxylate Modified Porous Graphitic Carbon: A New Class of Hydrophilic Interaction Liquid Chromatography Phases. <i>Analytical Chemistry</i> , 2013, 85, 5684-5691.	3.2	38
832	Determination of nucleosides and nucleobases in <i>Isatidis Radix</i> by HILIC-UPLC-MS/MS. <i>Analytical Methods</i> , 2013, 5, 6395.	1.3	10
833	Adsorption of Water/Acetonitrile Mixtures to Model Silica Surfaces. <i>Journal of Physical Chemistry C</i> , 2013, 117, 6620-6631.	1.5	49
834	Multidimensional separation of tryptic peptides from human serum proteins using reversed-phase, strong cation exchange, weak anion exchange, and fused-core fluorinated stationary phases. <i>Journal of Separation Science</i> , 2013, 36, 3463-3470.	1.3	26
835	Molecular Dynamics Simulation of Water/Acetonitrile Mixtures in a Silica Slit. <i>Journal of Physical Chemistry C</i> , 2013, 117, 3923-3929.	1.5	41
836	Separation of nucleotides by hydrophilic interaction chromatography using the FRULIC-N column. <i>Analytical and Bioanalytical Chemistry</i> , 2013, 405, 8837-8848.	1.9	32
837	Dissipation and residue behavior of mepiquat on wheat and potato field application. <i>Environmental Monitoring and Assessment</i> , 2013, 185, 9101-9109.	1.3	7
838	Glycomic Analysis by Glycoprotein Immobilization for Glycan Extraction and Liquid Chromatography on Microfluidic Chip. <i>Analytical Chemistry</i> , 2013, 85, 10117-10125.	3.2	31
839	Determination of glutathione and cysteine in yeasts by hydrophilic interaction liquid chromatography followed by online postcolumn derivatization. <i>Journal of Separation Science</i> , 2013, 36, 1877-1882.	1.3	13
840	Preparation of a novel carboxyl stationary phase by click chemistry for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2013, 36, 2571-2577.	1.3	31
841	Underivatized amylose and cellulose as new stationary phases for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2013, 36, 3345-3350.	1.3	1
842	A facile and efficient one-step strategy for the preparation of cyclodextrin monoliths. <i>Journal of Separation Science</i> , 2013, 36, 2441-2449.	1.3	28
843	New zwitterionic polymethacrylate monolithic columns for one- and two-dimensional microliquid chromatography. <i>Journal of Separation Science</i> , 2013, 36, 2430-2440.	1.3	36
844	Improved 2D-HPLC of red wine by incorporating pre-process signal-smoothing algorithms. <i>Journal of Separation Science</i> , 2013, 36, 3503-3510.	1.3	6

#	ARTICLE	IF	CITATIONS
845	Quantification of L-ergothioneine in whole blood by hydrophilic interaction ultra-performance liquid chromatography and UV-detection. <i>Journal of Separation Science</i> , 2013, 36, 1002-1006.	1.3	15
846	Comparison of nonaqueous hydrophilic interaction chromatography with aqueous normal-phase chromatography on hydrosilated silica-based stationary phases. <i>Journal of Separation Science</i> , 2013, 36, 2753-2759.	1.3	19
847	Analysis of polar peptides using a silica hydride column and high aqueous content mobile phases. <i>Journal of Separation Science</i> , 2013, 36, 3019-3025.	1.3	20
848	Performance of zwitterionic hydrophilic interaction LC for the determination of iodinated X-ray contrast agents. <i>Journal of Separation Science</i> , 2013, 36, 3688-3695.	1.3	8
849	The retention behaviour of amino acids in hydrophilic interaction liquid chromatography on zwitterionic stationary phases. <i>Journal of Separation Science</i> , 2013, 36, 2217-2222.	1.3	12
850	Analysis of Aqueous Alkanolamines Using Hydrophilic Interaction Liquid Chromatography and Mass Spectrometry. <i>Analytical Chemistry Letters</i> , 2013, 3, 298-313.	0.4	8
851	Methacrylate-based Diol Monolithic Stationary Phase for the Separation of Polar and Non-Polar Compounds in Capillary Liquid Chromatography. <i>Analytical Sciences</i> , 2013, 29, 631-635.	0.8	4
853	^#. ^t22: TLC of Antiepileptics. , 2013, , 466-477.		0
854	A Comprehensive Workflow of Mass Spectrometry-Based Untargeted Metabolomics in Cancer Metabolic Biomarker Discovery Using Human Plasma and Urine. <i>Metabolites</i> , 2013, 3, 787-819.	1.3	31
855	Determination of nitenpyram and its metabolites in agricultural products by using hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Journal of Pesticide Sciences</i> , 2013, 38, 27-32.	0.8	19
856	Plasma L-Ergothioneine Measurement by High-Performance Liquid Chromatography and Capillary Electrophoresis after a Pre-Column Derivatization with 5-Iodoacetamidofluorescein (5-IAF) and Fluorescence Detection. <i>PLoS ONE</i> , 2013, 8, e70374.	1.1	22
857	Characterization of Nucleosides and Nucleobases in Natural Cordyceps by HILIC-ESI/TOF/MS and HILIC-ESI/MS. <i>Molecules</i> , 2013, 18, 9755-9769.	1.7	38
858	Stationary Phases for the Separation of Reducing Sugars by Normal-Phase Partition Chromatography. <i>Chromatography</i> , 2014, 35, 63-72.	0.8	6
859	Evolution and Current Trends in Liquid and Supercritical Fluid Chromatography. <i>Current Chromatography</i> , 2014, 1, 15-40.	0.1	18
864	Data Analytics for Traditional Chinese Medicine Research. , 2014, , .		9
865	NATIVE/DERIVATIZED CYCLOFRUCTAN 6 BOUND TO RESINS VIA "CLICK"-CHEMISTRY AS STATIONARY PHASES FOR ACHIRAL/CHIRAL SEPARATIONS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 2302-2326.	0.5	5
866	An Overview of Quantitative Proteomic Approaches. <i>Comprehensive Analytical Chemistry</i> , 2014, 63, 111-135.	0.7	0
867	Recent developments in liquid-phase separation techniques for metabolomics. <i>Bioanalysis</i> , 2014, 6, 1011-1026.	0.6	47

#	ARTICLE	IF	CITATIONS
868	Hypoxia-induced Changes to Integrin $\alpha 3$ Glycosylation Facilitate Invasion in Epidermoid Carcinoma Cell Line A431. <i>Molecular and Cellular Proteomics</i> , 2014, 13, 3126-3137.	2.5	38
869	Targeted proteomics: a bridge between discovery and validation. <i>Expert Review of Proteomics</i> , 2014, 11, 657-661.	1.3	32
870	Recent advances in hydrophilic interaction chromatography for quantitative analysis of endogenous and pharmaceutical compounds in plasma samples. <i>Bioanalysis</i> , 2014, 6, 2421-2439.	0.6	35
871	The science of laboratory and project management in regulated bioanalysis. <i>Bioanalysis</i> , 2014, 6, 1357-1372.	0.6	5
872	Supercritical Fluid Chromatography: A Green Approach for Separation and Purification of Organic and Inorganic Analytes. , 2014, , 55-80.		1
873	Screening and confirmation of myo α -inositol trispyrophosphate (ITPP) in human urine by hydrophilic interaction liquid chromatography high resolution / high accuracy mass spectrometry for doping control purposes. <i>Drug Testing and Analysis</i> , 2014, 6, 1102-1107.	1.6	12
874	Separation of nucleobases, nucleosides, and nucleotides using two zwitterionic silica-based monolithic capillary columns coupled with tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1373, 90-96.	1.8	33
875	Contribution of Cytosine α -Containing Cyclobutane Dimers to DNA Damage Produced by Photosensitized Triplet α -Triplet Energy Transfer. <i>Chemistry - A European Journal</i> , 2014, 20, 5787-5794.	1.7	20
876	The impact of pressure and frictional heating on retention, selectivity and efficiency in ultra-high-pressure liquid chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2014, 63, 31-43.	5.8	36
877	Hydrophilic interaction electrokinetic chromatography using bio α -based nanofillers. <i>Electrophoresis</i> , 2014, 35, 2229-2236.	1.3	3
878	Evaluation of different hydrophilic stationary phases for the simultaneous determination of iminosugars and other low molecular weight carbohydrates in vegetable extracts by liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2014, 1372, 81-90.	1.8	9
879	Zirconia-A stationary phase capable of the separation of polar markers of myocardial metabolism in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2014, 37, 1089-1093.	1.3	9
880	Post-translational Modifications in the Human Proteome. <i>Translational Bioinformatics</i> , 2014, , 101-136.	0.0	2
881	Preconcentration based on paramagnetic microparticles for the separation of sarcosine using hydrophilic interaction liquid chromatography coupled with coulometric detection. <i>Journal of Separation Science</i> , 2014, 37, 465-575.	1.3	12
882	Adsorption of water from aqueous acetonitrile on silica-based stationary phases in aqueous normal-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1374, 102-111.	1.8	70
883	Influence of the crosslinker type on the chromatographic properties of hydrophilic sulfoalkylbetaine-type monolithic columns. <i>Journal of Chromatography A</i> , 2014, 1373, 73-80.	1.8	25
884	Analysis of <i>Dioscorea japonica</i> by Hydrophilic Interaction Liquid Chromatography. <i>Analytical Letters</i> , 2014, 47, 901-910.	1.0	1
886	Investigation on performance of zirconia and magnesia α -zirconia stationary phases in hydrophilic interaction chromatography. <i>Talanta</i> , 2014, 129, 438-447.	2.9	18

#	ARTICLE	IF	CITATIONS
887	Separation of complex sugar mixtures on a hydrolytically stable bidentate urea-type stationary phase for hydrophilic interaction near ultra high performance liquid chromatography. <i>Journal of Separation Science</i> , 2014, 37, 527-535.	1.3	19
888	Determination of Strong Acidic Drugs in Biological Matrices: A Review of Separation Methods. <i>Chromatography Research International</i> , 2014, 2014, 1-10.	0.4	3
889	Fluorous affinity-based separation techniques for the analysis of biogenic and related molecules. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 101, 151-160.	1.4	24
890	DETERMINATION OF CETYLPYRIDINIUM CHLORIDE RESIDUE IN CHICKEN MEAT BY HYDROPHILIC INTERACTION CHROMATOGRAPHY. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 538-547.	0.5	4
891	Validation of a LC-MS Method for the Determination of Urea Contamination in Market Teas. <i>Food Analytical Methods</i> , 2014, 7, 13-20.	1.3	5
892	SPIN TIP SOLID-PHASE EXTRACTION AND HILIC-MS-MS FOR QUANTITATIVE DETERMINATION OF METHAMPHETAMINE AND AMPHETAMINE IN HUMAN PLASMA. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 420-432.	0.5	5
893	A new type of monodisperse porous, hydrophilic microspheres with reactive chloroalkyl functionality: synthesis and derivatization properties. <i>Colloid and Polymer Science</i> , 2014, 292, 219-228.	1.0	21
894	Recent advances in enrichment and separation strategies for mass spectrometry-based phosphoproteomics. <i>Electrophoresis</i> , 2014, 35, 3418-3429.	1.3	41
895	Direct separation of the enantiomers of oxaliplatin on a cellulose-based chiral stationary phase in hydrophilic interaction liquid chromatography mode. <i>Journal of Chromatography A</i> , 2014, 1339, 210-213.	1.8	33
896	Retention behavior of a homologous series and positional isomers of aliphatic amino acids in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2014, 37, 739-747.	1.3	20
897	Improving peak shapes with counter gradients in two-dimensional high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1337, 147-154.	1.8	28
898	Chemical and technical challenges in the analysis of central carbon metabolites by liquid-chromatography mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 966, 21-33.	1.2	49
899	New reversed-phase/anion-exchange/hydrophilic interaction mixed-mode stationary phase based on dendritic polymer-modified porous silica. <i>Journal of Chromatography A</i> , 2014, 1337, 133-139.	1.8	62
900	Anti-diabetic and antihypertensive activities of two flaxseed protein hydrolysate fractions revealed following their simultaneous separation by electrodialysis with ultrafiltration membranes. <i>Food Chemistry</i> , 2014, 145, 66-76.	4.2	101
901	Practical observations on the performance of bare silica in hydrophilic interaction compared with C18 reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1328, 7-15.	1.8	43
902	Retention modeling and method development in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2014, 1337, 116-127.	1.8	63
903	Study of the retention behavior of small polar molecules on different types of stationary phases used in hydrophilic interaction liquid chromatography. <i>Journal of Separation Science</i> , 2014, 37, 1297-1307.	1.3	24
904	Retention mechanism assessment and method development for the analysis of iohexol and its related compounds in hydrophilic interaction liquid chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4217-4232.	1.9	7

#	ARTICLE	IF	CITATIONS
905	The next level of complexity: Crosstalk of posttranslational modifications. <i>Proteomics</i> , 2014, 14, 513-524.	1.3	246
906	Protein-Centric N-Glycoproteomics Analysis of Membrane and Plasma Membrane Proteins. <i>Journal of Proteome Research</i> , 2014, 13, 2705-2714.	1.8	24
907	Efficient purification of active bufadienolides by a class separation method based on hydrophilic solid-phase extraction and reversed-phase high performance liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 97, 54-64.	1.4	25
908	HPLC-fluorescence determination of thiol compounds in the serum of human male and female subjects using HILIC-mode column. <i>Biomedical Chromatography</i> , 2014, 28, 589-593.	0.8	17
910	Core-shell particles: Preparation, fundamentals and applications in high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1357, 36-52.	1.8	375
911	Sensitive hydrophilic interaction liquid chromatography/tandem mass spectrometry method for rapid detection, quantification and confirmation of cathinone-derived designer drugs for doping control in equine plasma. <i>Rapid Communications in Mass Spectrometry</i> , 2014, 28, 217-229.	0.7	13
912	High-throughput determination of nonsteroidal anti-inflammatory drugs in human plasma by HILIC-MS/MS. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 88, 71-80.	1.4	31
913	Determination of Deoxynivalenol-3-Glucoside in Cereals by Hydrophilic Interaction Chromatography with Ultraviolet Detection. <i>Food Analytical Methods</i> , 2014, 7, 1139-1146.	1.3	11
914	Metabolic profiling of amino acids in cellular samples via zwitterionic sub-2 μ m particle size HILIC-MS/MS and a uniformly ^{13}C labeled internal standard. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 915-922.	1.9	21
915	Analysis of plant nucleotide sugars by hydrophilic interaction liquid chromatography and tandem mass spectrometry. <i>Analytical Biochemistry</i> , 2014, 448, 14-22.	1.1	49
916	Study of surface-bonded dicationic ionic liquids as stationary phases for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2014, 1330, 40-50.	1.8	41
919	Separation of xanthines in hydro-organic and polar-organic elution modes on a titania stationary phase. <i>Journal of Separation Science</i> , 2014, 37, 536-542.	1.3	4
920	Preparation and characterization of surfactin-modified silica stationary phase for reversed-phase and hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1371, 257-260.	1.8	27
921	Comparison of the kinetic performance and retentivity of sub-2 μ m core-shell, hybrid and conventional bare silica phases in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2014, 1371, 106-116.	1.8	33
922	Determination of acysole in whole blood by high-performance liquid chromatography. <i>Journal of Analytical Chemistry</i> , 2014, 69, 800-804.	0.4	0
923	THEORETICAL AND EMPIRICAL MODELS IN HYDROPHILIC INTERACTION LIQUID CHROMATOGRAPHY. <i>Instrumentation Science and Technology</i> , 2014, 42, 230-266.	0.9	16
924	Ultrasensitive proteome analysis using paramagnetic bead technology. <i>Molecular Systems Biology</i> , 2014, 10, 757.	3.2	835
925	Micellar per aqueous liquid chromatography: an alternative to micellar liquid chromatography for determination of polar compounds?. <i>Analytical Methods</i> , 2014, 6, 7426.	1.3	4

#	ARTICLE	IF	CITATIONS
926	Facile preparation of polyvinyl alcohol coated SiO ₂ stationary phases for high performance liquid chromatography. <i>Analyst, The</i> , 2014, 139, 5594-5599.	1.7	18
927	Amino-terminated ionic liquid modified graphene oxide coated silica composite stationary phase for hydrophilic interaction chromatography. <i>RSC Advances</i> , 2014, 4, 37381-37388.	1.7	23
928	A novel fullerene oxide functionalized silica composite as stationary phase for high performance liquid chromatography. <i>RSC Advances</i> , 2014, 4, 17541-17548.	1.7	20
929	Preparation, chromatographic evaluation and comparison of cystine- and cysteine-bonded stationary phases. <i>Analytical Methods</i> , 2014, 6, 2205-2214.	1.3	7
930	Interaction modes and approaches to glycopeptide and glycoprotein enrichment. <i>Analyst, The</i> , 2014, 139, 688-704.	1.7	111
931	Development and Application of a Robust N-glycan Profiling Method for Heightened Characterization of Monoclonal Antibodies and Related Glycoproteins. <i>Journal of Pharmaceutical Sciences</i> , 2014, 103, 1967-1978.	1.6	36
932	Evaluation and comparison of the kinetic performance of ultra-high performance liquid chromatography and high-performance liquid chromatography columns in hydrophilic interaction and reversed-phase liquid chromatography conditions. <i>Journal of Chromatography A</i> , 2014, 1369, 83-91.	1.8	28
933	Expression, purification, and micelle reconstitution of antimicrobial piscidin 1 and piscidin 3 for NMR studies. <i>Protein Expression and Purification</i> , 2014, 102, 63-68.	0.6	13
934	Simultaneous determination of the repertoire of classical neurotransmitters released from embryonal carcinoma stem cells using online microdialysis coupled with hydrophilic interaction chromatography-tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2014, 849, 70-79.	2.6	24
935	Simultaneous determination of metformin and vildagliptin in human plasma by a HILIC-MS/MS method. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 965, 133-141.	1.2	46
936	A new HILIC-MS/MS method for the simultaneous analysis of carbidopa, levodopa, and its metabolites in human plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 967, 41-49.	1.2	33
937	High-performance liquid chromatography enantioseparation of polyhalogenated 4,4'-bipyridines on polysaccharide-based chiral stationary phases under multimodal elution. <i>Journal of Separation Science</i> , 2014, 37, 2481-2489.	1.3	13
938	Highly hydrophilic and nonionic poly(2-vinylloxazoline)-grafted silica: a novel organic phase for high-selectivity hydrophilic interaction chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4585-4593.	1.9	6
939	Mixed-mode liquid chromatography coupled to tandem mass spectrometry for the analysis of aminoglycosides in meat. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 4941-4953.	1.9	21
940	Polyhydroxy glucose functionalized silica for the dehydration of bio-ethanol distillate. <i>Bioprocess and Biosystems Engineering</i> , 2014, 37, 1417-1425.	1.7	2
941	A Comparison of Silica C and Silica Gel in HILIC Mode: The Effect of Stationary Phase Surface Area. <i>Chromatographia</i> , 2014, 77, 873-881.	0.7	11
942	Comprehensive Separation and Structural Analyses of Polyphenols and Related Compounds from Bracts of Hops (<i>Humulus lupulus</i> L.). <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 2198-2206.	2.4	39
943	Recent progress in polar metabolite quantification in plants using liquid chromatography-mass spectrometry. <i>Journal of Integrative Plant Biology</i> , 2014, 56, 816-825.	4.1	20

#	ARTICLE	IF	CITATIONS
944	Study of hydration process on silica hydride surfaces by microcalorimetry and water adsorption. <i>Journal of Colloid and Interface Science</i> , 2014, 416, 161-166.	5.0	17
945	Hybrid carbon nanoparticles modified core-shell silica: A high efficiency carbon-based phase for hydrophilic interaction liquid chromatography. <i>Analytica Chimica Acta</i> , 2014, 820, 187-194.	2.6	13
946	A new stationary phase for high performance liquid chromatography: Calix[4]arene derivatized chitosan bonded silica gel. <i>Journal of Chromatography A</i> , 2014, 1350, 61-67.	1.8	34
947	An integrated sample pretreatment platform for quantitative N-glycoproteome analysis with combination of on-line glycopeptide enrichment, deglycosylation and dimethyl labeling. <i>Analytica Chimica Acta</i> , 2014, 833, 1-8.	2.6	24
948	Hydrophilic interaction liquid chromatography-solid phase extraction directly combined with protein precipitation for the determination of triptorelin in plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014, 960, 214-221.	1.2	14
949	Characterization of intact N- and O-linked glycopeptides using higher energy collisional dissociation. <i>Analytical Biochemistry</i> , 2014, 452, 96-102.	1.1	56
950	Polyethylenimine attached-poly(3-chloro-2-hydroxypropyl methacrylate-co-ethylene dimethacrylate) monosized-porous microspheres as a new separation medium for polar compounds. <i>Colloids and Surfaces A: Physicochemical and Engineering Aspects</i> , 2014, 441, 629-637.	2.3	8
951	Hydrophilic interaction chromatography-mass spectrometry for anionic metabolic profiling of urine from antibiotic-treated rats. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 92, 98-104.	1.4	11
952	Hydrophilic interaction chromatography for selective separation of isomeric saponins. <i>Journal of Chromatography A</i> , 2014, 1325, 121-128.	1.8	41
953	Investigation and identification of potential biomarkers in human saliva for the early diagnosis of oral squamous cell carcinoma. <i>Clinica Chimica Acta</i> , 2014, 427, 79-85.	0.5	81
955	Novel orthogonal liquid chromatography methods to dose neurotransmitters involved in Parkinson's disease. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 98, 253-259.	1.4	10
956	Synthetic oligonucleotide separations by mixed-mode reversed-phase/weak anion-exchange liquid chromatography. <i>Journal of Chromatography A</i> , 2014, 1354, 43-55.	1.8	56
957	Monolithic capillary columns based on silica and zirconium oxides for use in hydrophilic interaction liquid chromatography. <i>Microchemical Journal</i> , 2014, 116, 249-254.	2.3	9
958	Hydrophilic interaction chromatography (HILIC) in the analysis of antibiotics. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 87, 142-154.	1.4	83
959	Preparative separation of steviol glycosides from <i>Stevia rebaudiana</i> by macroporous resin and preparative HPLC. <i>Acta Chromatographica</i> , 2014, 26, 123-135.	0.7	4
962	(1 \rightarrow 3)- β -D-Glucan oligosaccharides monomers purification and its H ₂ O ₂ induction effect study. <i>International Journal of Biological Macromolecules</i> , 2015, 81, 1069-1073.	3.6	7
963	Genetically Modified Organisms in Food. , 2015, , 1197-1212.		1
967	The Effects of the Column Length on the Efficiency of Capillary Zwitterionic Organic Polymer Monolithic Columns in HILIC Chromatography. <i>Chromatographia</i> , 2015, 78, 853-859.	0.7	6

#	ARTICLE	IF	CITATIONS
968	Hydrophilic interaction liquid chromatography for the separation, purification, and quantification of raffinose family oligosaccharides from <i>Lycopus lucidus</i> Turcz. Journal of Separation Science, 2015, 38, 2607-2613.	1.3	11
969	Enhanced fluidity liquid chromatography using mixed-mode hydrophilic interaction liquid chromatography/strong cation-exchange retention mechanisms. Journal of Separation Science, 2015, 38, 3119-3129.	1.3	18
970	Comprehensive analysis of fatty alcohol ethoxylates by ultra high pressure hydrophilic interaction chromatography coupled with ion mobility spectrometry mass spectrometry using a custom-designed sub-2 μ m column. Journal of Separation Science, 2015, 38, 2182-2191.	1.3	5
971	Silica modified with a thiourea derivative as a new stationary phase for hydrophilic interaction liquid chromatography. Journal of Separation Science, 2015, 38, 3852-3861.	1.3	2
972	Hydrophilic Interaction Chromatographic Analysis of Quercetin and its Glycosides. Current Analytical Chemistry, 2015, 12, 60-64.	0.6	3
973	Affinity-Based Methods in Drug-Target Discovery. Current Drug Targets, 2015, 16, 60-76.	1.0	11
974	Metabonomic Evaluation of Chronic Unpredictable Mild Stress-Induced Changes in Rats by Intervention of Fluoxetine by HILIC-UHPLC/MS. PLoS ONE, 2015, 10, e0129146.	1.1	18
975	Amino acid analysis using chromatography-mass spectrometry: An inter platform comparison study. Journal of Pharmaceutical and Biomedical Analysis, 2015, 114, 398-407.	1.4	60
976	Nucleoside Analysis by Hydrophilic Interaction Liquid Chromatography Coupled with Mass Spectrometry. Methods in Enzymology, 2015, 560, 19-28.	0.4	61
977	Synthesis and characterization of phosphodiester stationary bonded phases for liquid chromatography. Talanta, 2015, 143, 35-41.	2.9	14
978	Rapid Quantification of Four Anthocyanins in Red Grape Wine by Hydrophilic Interaction Liquid Chromatography/ Triple Quadrupole Linear Ion Trap Mass Spectrometry. Journal of AOAC INTERNATIONAL, 2015, 98, 1628-1631.	0.7	5
979	Simultaneous Determination of Five Alkaloid Compounds in a Drug Based on a Hydrophilic Monolithic Column by Capillary Electrochromatography. Journal of Chromatographic Science, 2015, 54, bmv096.	0.7	5
980	Recent advances in the application of hydrophilic interaction chromatography for the analysis of biological matrices. Bioanalysis, 2015, 7, 2927-2945.	0.6	20
981	Evaluating the use of HILIC in large-scale, multi dimensional proteomics: Horses for courses?. International Journal of Mass Spectrometry, 2015, 391, 105-114.	0.7	10
982	Evaluation of the Kinetic Performance Differences between Hydrophilic-Interaction Liquid Chromatography and Reversed-Phase Liquid Chromatography under Conditions of Identical Packing Structure. Analytical Chemistry, 2015, 87, 12331-12339.	3.2	24
983	Comprehensive Analysis of Microbial Lipids in Environmental Samples Through HPLC-MS Protocols. Springer Protocols, 2015, , 289-317.	0.1	19
984	Chemically Bonded Polyacrylamide via Thiol-Ene Click Chemistry as Separation Materials for Hydrophilic Interaction Liquid Chromatography. Chinese Journal of Analytical Chemistry, 2015, 43, 1439-1444.	0.9	8
985	Preparation and chromatographic performance evaluation of cucurbit[7]uril immobilized silica. Journal of Chromatography A, 2015, 1376, 64-73.	1.8	11

#	ARTICLE	IF	CITATIONS
986	The relative importance of the adsorption and partitioning mechanisms in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1376, 112-125.	1.8	51
987	Development of analytical method for catechol compounds in mouse urine using hydrophilic interaction liquid chromatography with fluorescence detection. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 985, 142-148.	1.2	39
988	Molecular Speciated Isotope Dilution Mass Spectrometric Methods for Accurate, Reproducible and Direct Quantification of Reduced, Oxidized and Total Glutathione in Biological Samples. <i>Analytical Chemistry</i> , 2015, 87, 1232-1240.	3.2	36
989	Evaluation of Aqueous and Nonaqueous Binary Solvent Mixtures as Mobile Phase Alternatives to Water/Acetonitrile Mixtures for Hydrophilic Interaction Liquid Chromatography by Molecular Dynamics Simulations. <i>Journal of Physical Chemistry C</i> , 2015, 119, 512-523.	1.5	33
990	Preparation and chromatographic evaluation of a newly designed steviol glycoside modified-silica stationary phase in hydrophilic interaction liquid chromatography and reversed phase liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1388, 110-118.	1.8	48
991	Separation of therapeutic peptides with cyclofructan and glycopeptide based columns in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1390, 50-61.	1.8	15
992	Quality by Design approach in the development of hydrophilic interaction liquid chromatographic method for the analysis of iohexol and its impurities. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015, 110, 42-48.	1.4	21
993	An efficient hydrophilic interaction liquid chromatographic method for the simultaneous determination of metformin and pioglitazone using high-purity silica column. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 997, 16-22.	1.2	21
994	An L-lysine derived organogelator-based stationary phase for mixed-mode liquid chromatography. <i>Analytical Methods</i> , 2015, 7, 3320-3323.	1.3	2
995	Hydrophilic-subtraction model for the characterization and comparison of hydrophilic interaction liquid chromatography columns. <i>Journal of Chromatography A</i> , 2015, 1398, 29-46.	1.8	38
996	A Platform for Discovery and Quantification of Modified Ribonucleosides in RNA. <i>Methods in Enzymology</i> , 2015, 560, 29-71.	0.4	69
997	Ionic liquid-based zwitterionic organic polymer monolithic column for capillary hydrophilic interaction chromatography. <i>Analyst</i> , 2015, 140, 5585-5592.	1.7	15
998	Quantification of N-hydroxysuccinimide and N-hydroxysulfosuccinimide by hydrophilic interaction chromatography (HILIC). <i>Analytical Methods</i> , 2015, 7, 6443-6448.	1.3	34
999	Determination of free quinic acid in food matrices by Hydrophilic Interaction Liquid Chromatography with UV detection. <i>Journal of Food Composition and Analysis</i> , 2015, 44, 80-85.	1.9	11
1000	The relevance of pharmacognosy in pharmacological research on herbal medicinal products. <i>Epilepsy and Behavior</i> , 2015, 52, 344-362.	0.9	76
1001	Design of C ₁₈ Organic Phases with Multiple Embedded Polar Groups for Ultraversatile Applications with Ultrahigh Selectivity. <i>Analytical Chemistry</i> , 2015, 87, 6614-6621.	3.2	47
1002	Preparation and chromatographic evaluation of zwitterionic stationary phases with controllable ratio of positively and negatively charged groups. <i>Talanta</i> , 2015, 141, 8-14.	2.9	16
1003	Aminoglycoside analysis in food of animal origin with a zwitterionic stationary phase and liquid chromatography-tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2015, 882, 127-139.	2.6	64

#	ARTICLE	IF	CITATIONS
1004	Versatile ligands for high-performance liquid chromatography: An overview of ionic liquid-functionalized stationary phases. <i>Analytica Chimica Acta</i> , 2015, 887, 1-16.	2.6	73
1005	Grafting zwitterionic polymer onto cryogel surface enhances protein retention in steric exclusion chromatography on cryogel monolith. <i>Journal of Chromatography A</i> , 2015, 1389, 104-111.	1.8	26
1006	Hydrophilic Interaction Liquid Chromatography (HILIC) and Perfluorinated Stationary Phases. , 2015, , 149-184.		0
1007	Bioassay-Guided Separation and Identification of a New Anti-Lung Cancer Compound from <i>Cordyceps militaris</i> by Means of Off-Line Two-Dimensional Preparative Chromatography, Real-Time Cell Analysis, and X-ray Single-Crystal Diffraction. <i>Chromatographia</i> , 2015, 78, 495-506.	0.7	12
1008	“Dilute-and-inject” multi-target screening assay for highly polar doping agents using hydrophilic interaction liquid chromatography high resolution/high accuracy mass spectrometry for sports drug testing. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 5365-5379.	1.9	48
1009	Synthesis of a reactive polymethacrylate capillary monolith and its use as a starting material for the preparation of a stationary phase for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2015, 1396, 86-97.	1.8	17
1010	Rapid quantification of imidazolium-based ionic liquids by hydrophilic interaction liquid chromatography: Methodology and an investigation of the retention mechanisms. <i>Journal of Chromatography A</i> , 2015, 1400, 54-64.	1.8	25
1011	An arginine-functionalized stationary phase for hydrophilic interaction liquid chromatography. <i>Analyst, The</i> , 2015, 140, 3921-3924.	1.7	33
1012	Evaluation of a silicon oxynitride hydrophilic interaction liquid chromatography column in saccharide and glycoside separations. <i>Journal of Separation Science</i> , 2015, 38, 1271-1276.	1.3	11
1013	A versatile reversed phase-strong cation exchange-reversed phase (RP-SCX-RP) multidimensional liquid chromatography platform for qualitative and quantitative shotgun proteomics. <i>Analyst, The</i> , 2015, 140, 1237-1252.	1.7	20
1014	Onco-proteogenomics identifies urinary S100A9 and GRN as potential combinatorial biomarkers for early diagnosis of hepatocellular carcinoma. <i>BBA Clinical</i> , 2015, 3, 205-213.	4.1	33
1015	Determination of creatinine, uric and ascorbic acid in bovine milk and orange juice by hydrophilic interaction HPLC. <i>Food Chemistry</i> , 2015, 182, 242-245.	4.2	75
1016	Multidimensional proteomics for cell biology. <i>Nature Reviews Molecular Cell Biology</i> , 2015, 16, 269-280.	16.1	375
1017	Development of liquid chromatography methods coupled to mass spectrometry for the analysis of substances with a wide variety of polarity in meconium. <i>Talanta</i> , 2015, 138, 231-239.	2.9	6
1018	Diazonium modification of porous graphitic carbon with catechol and amide groups for hydrophilic interaction and attenuated reversed phase liquid chromatography. <i>Journal of Chromatography A</i> , 2015, 1422, 186-193.	1.8	2
1019	Hydrophilic interaction liquid chromatography with indirect ultraviolet detection for the separation and quantification of pyrrolidinium ionic liquid cations. <i>Chinese Chemical Letters</i> , 2015, 26, 1371-1375.	4.8	10
1020	Liquid chromatography-mass spectrometry platform for both small neurotransmitters and neuropeptides in blood, with automatic and robust solid phase extraction. <i>Scientific Reports</i> , 2015, 5, 9308.	1.6	54
1021	Recent progress in the fundamental understanding of hydrophilic interaction chromatography (HILIC). <i>Analyst, The</i> , 2015, 140, 6452-6466.	1.7	114

#	ARTICLE	IF	CITATIONS
1022	Direct injection human plasma analysis for the quantification of antihypertensive drugs for therapeutic drug monitoring using hydrophilic interaction liquid chromatography/electrospray ionization mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 1004, 1-9.	1.2	14
1023	Preparation and Analysis of Cello- and Xylooligosaccharides. <i>Advances in Polymer Science</i> , 2015, , 53-92.	0.4	4
1024	A remarkable enhancement of selectivity towards versatile analytes by a strategically integrated H-bonding site containing phase. <i>Chemical Communications</i> , 2015, 51, 14243-14246.	2.2	9
1025	Preparative separation of a challenging anthocyanin from <i>Lycium ruthenicum</i> Murr. by two-dimensional reversed-phase liquid chromatography/hydrophilic interaction chromatography. <i>RSC Advances</i> , 2015, 5, 62134-62141.	1.7	35
1026	Chromatographic methods for the isolation, separation and characterisation of dissolved organic matter. <i>Environmental Sciences: Processes and Impacts</i> , 2015, 17, 1531-1567.	1.7	52
1027	A rapid and simple method for the simultaneous determination of four endogenous monoamine neurotransmitters in rat brain using hydrophilic interaction liquid chromatography coupled with atmospheric-pressure chemical ionization tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 1002, 379-386.	1.2	16
1028	Hydrophilic interaction chromatography coupled to tandem mass spectrometry in the presence of hydrophilic ion-pairing reagents for the separation of nucleosides and nucleotide mono-, di- and triphosphates. <i>Journal of Chromatography A</i> , 2015, 1414, 129-137.	1.8	34
1029	LC-MS-based quantification of intact proteins: perspective for clinical and bioanalytical applications. <i>Bioanalysis</i> , 2015, 7, 1943-1958.	0.6	52
1030	Optimization of Chromatographic Separation of Acetylsalicylic Acid, Amlodipine, Impurity A of Amlodipine, and Atenolol in Hydrophilic Interaction Liquid Chromatography Employing DoE Methodology. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 1555-1560.	0.5	4
1031	Uncovering dynamic phosphorylation signaling using mass spectrometry. <i>International Journal of Mass Spectrometry</i> , 2015, 391, 123-138.	0.7	5
1032	Opiorphin analysis in equine plasma and urine using hydrophilic interaction LC-MS. <i>Bioanalysis</i> , 2015, 7, 593-603.	0.6	2
1033	Reduction of urinary uric acid excretion in patients with proteinuria. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 1006, 59-64.	1.2	10
1034	Rapid Determination of Ethephon in Grapes by Hydrophilic Interaction Chromatography Tandem Mass Spectrometry. <i>Food Analytical Methods</i> , 2015, 8, 524-530.	1.3	13
1035	A novel method for the simultaneous analysis of seven biothiols in rice (<i>Oryza sativa</i> L.) using hydrophilic interaction chromatography coupled with electrospray tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015, 976-977, 19-26.	1.2	18
1036	Applications of hydrophilic interaction chromatography to amino acids, peptides, and proteins. <i>Journal of Separation Science</i> , 2015, 38, 357-367.	1.3	61
1037	Improved short peptide identification using HILIC-MS/MS: Retention time prediction model based on the impact of amino acid position in the peptide sequence. <i>Food Chemistry</i> , 2015, 173, 847-854.	4.2	64
1038	A simple LC-MS/MS method using HILIC chromatography for the determination of fosfomycin in plasma and urine: Application to a pilot pharmacokinetic study in humans. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015, 105, 39-45.	1.4	28
1039	Design of Experiments in Optimization and Validation of a Hydrophilic Interaction Liquid Chromatography Method for Determination of Amlodipine Besylate and Bisoprolol Fumarate. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 919-928.	0.5	13

#	ARTICLE	IF	CITATIONS
1040	Recent advances in nonpolar and polar organic monoliths for HPLC and CEC. <i>Electrophoresis</i> , 2015, 36, 76-100.	1.3	41
1041	Preparation, characterization and application of N-methylene phosphonic acid chitosan grafted magnesia/zirconia stationary phase. <i>Analytica Chimica Acta</i> , 2015, 854, 191-201.	2.6	9
1042	Direct and Indirect Applications of Sub- and Supercritical Water in Food-Related Analysis. <i>Food Engineering Series</i> , 2015, , 269-302.	0.3	0
1043	Separation properties of saccharides on a hydrophilic stationary phase having hydration layer formed zwitterionic copolymer. <i>Analytica Chimica Acta</i> , 2015, 853, 602-607.	2.6	9
1044	Relative quantification of amine-containing metabolites using isobaric N,N-dimethyl leucine (DiLeu) reagents via LC-ESI-MS/MS and CE-ESI-MS/MS. <i>Analyst</i> , The, 2015, 140, 467-475.	1.7	52
1045	Quantitative mass spectrometric analysis of glycoproteins combined with enrichment methods. <i>Mass Spectrometry Reviews</i> , 2015, 34, 148-165.	2.8	69
1046	The application of targeted mass spectrometry-based strategies to the detection and localization of post-translational modifications. <i>Mass Spectrometry Reviews</i> , 2015, 34, 595-626.	2.8	27
1047	Direct injection HILIC-MS/MS analysis of darunavir in rat plasma applying supported liquid extraction. <i>Journal of Pharmaceutical Analysis</i> , 2015, 5, 43-50.	2.4	10
1048	High Pressure Fluid Technology for Green Food Processing. <i>Food Engineering Series</i> , 2015, , .	0.3	13
1049	Hydrophilic solid-phase extraction of melamine with ampholine-modified hybrid organic-inorganic silica material. <i>Journal of Separation Science</i> , 2015, 38, 87-92.	1.3	23
1050	The early diagnosis and monitoring of squamous cell carcinoma via saliva metabolomics. <i>Scientific Reports</i> , 2014, 4, 6802.	1.6	112
1051	Comparison of capillary electrophoresis-mass spectrometry and hydrophilic interaction chromatography-mass spectrometry for anionic metabolic profiling of urine. <i>Talanta</i> , 2015, 132, 1-7.	2.9	33
1052	Development of a simple method for quantitation of methanesulfonic acid at low ppm level using hydrophilic interaction chromatography coupled with ESI-MS. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2015, 102, 17-24.	1.4	18
1053	Hydrophilic interaction liquid chromatography coupled with tandem mass spectrometry for acidic herbicides and metabolites analysis in fresh water. <i>Environmental Science and Pollution Research</i> , 2015, 22, 3988-3996.	2.7	18
1054	HILIC-MS for metabolomics: An attractive and complementary approach to RPLC-MS. <i>Mass Spectrometry Reviews</i> , 2016, 35, 574-600.	2.8	191
1055	HILIC Chromatography - An Insight on the Retention Mechanism. <i>Journal of Chromatography & Separation Techniques</i> , 2016, 7, .	0.2	1
1056	Thin-layer chromatography and pressurized planar electrochromatography of amino acids in systems with silica gel and water mobile phase. <i>Journal of Planar Chromatography - Modern TLC</i> , 2016, 29, 30-37.	0.6	5
1057	Application of Hydrophilic Interaction Liquid Chromatography for the Quantification of Flavonoids in <i>Genista tinctoria</i> Extract. <i>Journal of Analytical Methods in Chemistry</i> , 2016, 2016, 1-9.	0.7	13

#	ARTICLE	IF	CITATIONS
1058	Analytical Approaches for the Quantitation of Redox-active Pyridine Dinucleotides in Biological Matrices. <i>Periodica Polytechnica: Chemical Engineering</i> , 2016, 60, 218-230.	0.5	4
1059	High-Throughput Metabolic Screening. , 2016, , 111-136.		0
1060	Quantitative Proteomic Analysis of Differentially Expressed Protein Profiles Involved in Pancreatic Ductal Adenocarcinoma. <i>Pancreas</i> , 2016, 45, 71-83.	0.5	17
1061	The hydrophilicity vs. ion interaction selectivity plot revisited: The effect of mobile phase pH and buffer concentration on hydrophilic interaction liquid chromatography selectivity behavior. <i>Journal of Chromatography A</i> , 2016, 1458, 82-89.	1.8	13
1062	Quantification of structurally related aliphatic amino alcohols in <sc>valinol by hydrophilic interaction liquid chromatography separation combined with postcolumn derivatization and fluorescence detection. <i>Journal of Separation Science</i> , 2016, 39, 851-856.	1.3	9
1063	Thiol-Maleimide Michael Addition Click Reaction: A New Route to Surface Modification of Porous Polymeric Monolith. <i>Macromolecular Chemistry and Physics</i> , 2016, 217, 997-1006.	1.1	23
1064	Evaluation of coverage, retention patterns, and selectivity of seven liquid chromatographic methods for metabolomics. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 6079-6091.	1.9	72
1065	Automated SDS Depletion for Mass Spectrometry of Intact Membrane Proteins through Transmembrane Electrophoresis. <i>Journal of Proteome Research</i> , 2016, 15, 2634-2642.	1.8	23
1066	Monoliths in capillary electrochromatography and capillary liquid chromatography in conjunction with mass spectrometry. <i>Electrophoresis</i> , 2016, 37, 880-912.	1.3	23
1067	Analysis of low abundant trehalose-6-phosphate and related metabolites in <i>Medicago truncatula</i> by hydrophilic interaction liquid chromatography-triple quadrupole mass spectrometry. <i>Journal of Chromatography A</i> , 2016, 1477, 30-38.	1.8	7
1068	Hydrophilic interaction chromatography with Fourier transform mass spectrometry of monosaccharide anhydrides. <i>Journal of Separation Science</i> , 2016, 39, 4748-4755.	1.3	4
1069	Improving Proteome Coverage by Reducing Sample Complexity via Chromatography. <i>Advances in Experimental Medicine and Biology</i> , 2016, 919, 83-143.	0.8	5
1072	Exometabolomics for Linking Soil Carbon Dynamics to Microbial Communities. , 2016, , 119-145.		0
1073	Modern Proteomics – Sample Preparation, Analysis and Practical Applications. <i>Advances in Experimental Medicine and Biology</i> , 2016, , .	0.8	13
1074	Dual Retention Mechanism in Two-Dimensional LC Separations of Barbiturates, Sulfonamides, Nucleic Bases and Nucleosides on Polymethacrylate Zwitterionic Monolithic Micro-Columns. <i>Chromatographia</i> , 2016, 79, 657-666.	0.7	14
1075	Amino acids, peptides, and proteins as chemically bonded stationary phases – A review. <i>Journal of Separation Science</i> , 2016, 39, 83-92.	1.3	40
1076	Recent advances in development and characterization of stationary phases for hydrophilic interaction chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2016, 81, 23-33.	5.8	96
1077	Development of a low level detection method for 2-chloromalonaldehyde in active pharmaceutical ingredients by HILIC separation. <i>Analytical Methods</i> , 2016, 8, 4659-4663.	1.3	2

#	ARTICLE	IF	CITATIONS
1078	A sensitive non-derivatization method for apramycin and impurities analysis using hydrophilic interaction liquid chromatography and charged aerosol detection. <i>Talanta</i> , 2016, 146, 423-429.	2.9	20
1079	Development, validation and application of a hydrophilic interaction liquid chromatography-evaporative light scattering detection based method for process control of hydrolysis of xylans obtained from different agricultural wastes. <i>Food Chemistry</i> , 2016, 212, 155-161.	4.2	13
1080	Poly(vinyl alcohol) Modified Porous Graphitic Carbon Stationary Phase for Hydrophilic Interaction Liquid Chromatography. <i>Analytical Chemistry</i> , 2016, 88, 4676-4681.	3.2	47
1081	Determination of catecholamines and related compounds in mouse urine using column-switching HPLC. <i>The Analyst</i> , 2016, 141, 2568-2573.	1.7	20
1082	Chromatographic methods enabling the characterization of stationary phases and retention prediction in high-performance liquid chromatography and supercritical fluid chromatography. <i>Journal of Separation Science</i> , 2016, 39, 115-131.	1.3	21
1083	Monolithic stationary phases with incorporated fumed silica nanoparticles. Part I. Polymethacrylate-based monolithic column with incorporated bare fumed silica nanoparticles for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2016, 1445, 55-61.	1.8	34
1084	Hydrophilic interaction chromatography as an alternative to reversed-phase HPLC in determining anthocyanins and betacyanins. <i>Journal of Analytical Chemistry</i> , 2016, 71, 297-301.	0.4	7
1085	Recent advances in materials for stationary phases of mixed-mode high-performance liquid chromatography. <i>TrAC - Trends in Analytical Chemistry</i> , 2016, 80, 495-506.	5.8	88
1086	Hydroxypropyl beta cyclodextrin bonded superficially porous particle-based HILIC stationary phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 459-464.	0.5	15
1087	Quantifying the Metabolites of the Methylerythritol 4-Phosphate (MEP) Pathway in Plants and Bacteria by Liquid Chromatography-Triple Quadrupole Mass Spectrometry. <i>Methods in Enzymology</i> , 2016, 576, 225-249.	0.4	18
1088	Multiple, simultaneous, independent gradients for a versatile multidimensional liquid chromatography. Part II: Application 1 - Large increases in isoform resolution of human transferrin by use of dual simultaneous independent gradients of pH & acetonitrile on a mixed bed (anion exchange) Tj ETQq0 0 0 r gBT /Overlock 10 T	1.8	5
1089	Dipeptide-bonded stationary phases for hydrophilic interaction liquid chromatography. <i>RSC Advances</i> , 2016, 6, 96389-96397.	1.7	17
1090	Determination of 3'-phosphoadenosine-5'-phosphosulfate in cells and Golgi fractions using hydrophilic interaction liquid chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2016, 1470, 70-75.	1.8	5
1091	Chromatographic behaviour of synthetic high pressure high temperature diamond in aqueous normal phase chromatography. <i>Journal of Chromatography A</i> , 2016, 1470, 59-69.	1.8	10
1092	Postpolymerization modification of a hydroxy monolith precursor. Part III. Activation of poly(hydroxyethyl methacrylate-co-pentaerythritol triacrylate) monolith with epoxy functionalities followed by bonding of glycerol, polyamines, and hydroxypropyl-beta-cyclodextrin for hydrophilic interaction and chiral capillary electrochromatography. <i>Electrophoresis</i> , 2016, 37, 3178-3185.	1.3	18
1093	Preparation and evaluation of 3 m open tubular capillary columns with a zwitterionic polymeric porous layer for liquid chromatography. <i>Journal of Separation Science</i> , 2016, 39, 3736-3744.	1.3	22
1094	The effect of charged groups on hydrophilic monolithic stationary phases on their chromatographic properties. <i>Journal of Chromatography A</i> , 2016, 1469, 77-87.	1.8	15
1095	Carboxylated cyclofructan 6 as a hydrolytically stable high efficiency stationary phase for hydrophilic interaction liquid chromatography and mixed mode separations. <i>Analytical Methods</i> , 2016, 8, 6038-6045.	1.3	34

#	ARTICLE	IF	CITATIONS
1096	Deconvoluting the effects of buffer salt concentration in hydrophilic interaction chromatography on a zwitterionic stationary phase. <i>Journal of Chromatography A</i> , 2016, 1461, 92-97.	1.8	43
1097	RP-HPLC—HILIC chromatography for quantifying ertapenem sodium with a look at green chemistry. <i>Talanta</i> , 2016, 160, 745-753.	2.9	29
1099	Coupling normal phase liquid chromatography with electrospray ionization mass spectrometry: strategies and applications. <i>Analytical Methods</i> , 2016, 8, 6478-6488.	1.3	17
1100	Imidazolium-embedded iodoacetamide-functionalized silica-based stationary phase for hydrophilic interaction/reversed-phase mixed-mode chromatography. <i>Journal of Separation Science</i> , 2016, 39, 3498-3504.	1.3	5
1101	Detailed insights into the retention mechanism of caffeine metabolites on the amide stationary phase in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2016, 1463, 121-127.	1.8	18
1102	Facile synthesis of carboxymethyl- α -cyclodextrin conjugated magnetic nanoparticles for selective enrichment of glycopeptides. <i>Rapid Communications in Mass Spectrometry</i> , 2016, 30, 190-195.	0.7	11
1103	Green analytical chemistry: Opportunities for pharmaceutical quality control. <i>Journal of Analytical Chemistry</i> , 2016, 71, 861-871.	0.4	32
1104	Manual of Cardiovascular Proteomics. , 2016, , .		4
1105	Post-translational Modifications in the Cardiovascular Proteome. , 2016, , 293-320.		0
1106	Current landscape of protein glycosylation analysis and recent progress toward a novel paradigm of glycoscience research. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 130, 273-300.	1.4	34
1107	Synthesis and evaluation of a maltose-bonded silica gel stationary phase for hydrophilic interaction chromatography and its application in Ginkgo Biloba extract separation in two-dimensional systems. <i>Journal of Separation Science</i> , 2016, 39, 3339-3347.	1.3	16
1108	The effects of temperature and mobile phase on the retention of aliphatic carboxylic acids in hydrophilic interaction chromatography on zwitterionic stationary phases. <i>Journal of Separation Science</i> , 2016, 39, 4732-4739.	1.3	10
1110	Sorption of nitrogen-containing aromatic compounds on ultradispersed diamonds. <i>Russian Chemical Bulletin</i> , 2016, 65, 110-119.	0.4	10
1111	Preparation of phosphorylcholine-based hydrophilic monolithic column and application for analysis of drug-related impurities with capillary electrochromatography. <i>Electrophoresis</i> , 2016, 37, 1725-1732.	1.3	21
1112	Metal reduction at point-of-use filtration. , 2016, , .		1
1113	A Molecular Dynamics View on Hydrophilic Interaction Chromatography with Polar-Bonded Phases: Properties of the Water-Rich Layer at a Silica Surface Modified with Diol-Functionalized Alkyl Chains. <i>Journal of Physical Chemistry C</i> , 2016, 120, 13126-13138.	1.5	19
1114	Halogenated methanesulfonic acids: A new class of organic micropollutants in the water cycle. <i>Water Research</i> , 2016, 101, 292-299.	5.3	67
1115	Thiol-ene click chemistry for the design of diol porous monoliths with hydrophilic surface interaction ability: a capillary electrochromatography study. <i>New Journal of Chemistry</i> , 2016, 40, 6916-6923.	1.4	10

#	ARTICLE	IF	CITATIONS
1116	Inflammatory glycoproteins in cardiometabolic disorders, autoimmune diseases and cancer. <i>Clinica Chimica Acta</i> , 2016, 459, 177-186.	0.5	66
1117	Recent development in liquid chromatography stationary phases for separation of Traditional Chinese Medicine components. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 130, 336-346.	1.4	28
1118	An aqueous normal-phase chromatography coupled with tandem mass spectrometry method for determining unbound brain-to-plasma concentration ratio of AZD1775, a Wee1 kinase inhibitor, in patients with glioblastoma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016, 1028, 25-32.	1.2	11
1119	Determination of polar pesticides in olive oil and olives by hydrophilic interaction liquid chromatography coupled to tandem mass spectrometry and high resolution mass spectrometry. <i>Talanta</i> , 2016, 158, 222-228.	2.9	46
1120	Automated dual two-dimensional liquid chromatography approach for fast acquisition of three-dimensional data using combinations of zwitterionic polymethacrylate and silica-based monolithic columns. <i>Journal of Chromatography A</i> , 2016, 1446, 91-102.	1.8	26
1121	Facile one-pot-synthesis of poly(methacrylic acid)-based hybrid monolith via thiol-ene click reaction for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2016, 1454, 49-57.	1.8	21
1122	Identification of Low Abundant Isomeric N-Glycan Structures in Biological Therapeutics by LC/MS. <i>Analytical Chemistry</i> , 2016, 88, 7049-7059.	3.2	38
1123	Advances in the analysis of iminocyclitols: Methods, sources and bioavailability. <i>Talanta</i> , 2016, 151, 157-171.	2.9	3
1124	Separation of substrates and closely related glucuronide metabolites using various chromatographic modes. <i>Journal of Chromatography A</i> , 2016, 1435, 54-65.	1.8	18
1125	Bioassay-guided separation and identification of anticancer compounds in <i>Tagetes erecta</i> L. flowers. <i>Analytical Methods</i> , 2016, 8, 3255-3262.	1.3	11
1126	A hyperbranched polyethylenimine functionalized stationary phase for hydrophilic interaction liquid chromatography. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 3633-3638.	1.9	20
1127	Direct determination of amino acids by hydrophilic interaction liquid chromatography with charged aerosol detection. <i>Journal of Chromatography A</i> , 2016, 1446, 41-49.	1.8	20
1128	Quantification of Lactulose and Epilactose in the Presence of Lactose in Milk using a dual HPLC analysis. <i>Food Analytical Methods</i> , 2016, 9, 2210-2222.	1.3	15
1129	A comparative study of phosphopeptide-selective techniques for a sub-proteome of a complex biological sample. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 2347-2356.	1.9	4
1130	Gradient enhanced-fluidity liquid hydrophilic interaction chromatography of ribonucleic acid nucleosides and nucleotides: A "green" technique. <i>Journal of Chromatography A</i> , 2016, 1436, 84-90.	1.8	37
1131	Enhancement of glucose uptake in muscular cell by peptide fractions separated by electro dialysis with filtration membrane from salmon frame protein hydrolysate. <i>Journal of Functional Foods</i> , 2016, 22, 337-346.	1.6	49
1132	Determination of d -glucaric acid and/or d -glucaro-1,4-lacton in different apple varieties through hydrophilic interaction chromatography. <i>Food Chemistry</i> , 2016, 203, 1-7.	4.2	6
1133	Streamlined discovery of cross-linked chromatin complexes and associated histone modifications by mass spectrometry. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 1784-1789.	3.3	17

#	ARTICLE	IF	CITATIONS
1134	Advances in preparation, analysis and biological activities of single chitooligosaccharides. <i>Carbohydrate Polymers</i> , 2016, 139, 178-190.	5.1	112
1135	Target-based metabolomics for the quantitative measurement of 37 pathway metabolites in rat brain and serum using hydrophilic interaction ultra-high-performance liquid chromatography-tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 2527-2542.	1.9	33
1136	Separation of urea, hexamethylenetetramine, and their reaction products in sol-gel feed solution by zwitterionic HILIC. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2016, 39, 119-127.	0.5	0
1137	Analysis of phospholipids in <i>Schizochytrium</i> sp. S31 by using UPLC-Q-TOF-MS. <i>Analytical Methods</i> , 2016, 8, 763-770.	1.3	17
1138	Hydrophilic interaction chromatography combined with dispersive liquid-liquid microextraction as a preconcentration tool for the simultaneous determination of the panel of underivatized neurotransmitters in human urine samples. <i>Journal of Chromatography A</i> , 2016, 1431, 111-121.	1.8	49
1139	Hydrophilic interaction liquid chromatography for separation and determination of pyrrolidinium ionic liquid cations. <i>Analytical Methods</i> , 2016, 8, 840-845.	1.3	9
1140	Chromatographic behavior of 12 polar pteridines in hydrophilic interaction chromatography using five different HILIC columns coupled with tandem mass spectrometry. <i>Talanta</i> , 2016, 150, 493-502.	2.9	28
1141	Recent advances and trends in the liquid-chromatography-mass spectrometry analysis of flavonoids. <i>Journal of Chromatography A</i> , 2016, 1430, 16-78.	1.8	155
1142	Development and validation of a hydrophilic interaction chromatography method coupled with a charged aerosol detector for quantitative analysis of nonchromophoric hydroxylamines, organic impurities of metoprolol. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 118, 242-250.	1.4	16
1143	Preparation of a novel ionic hybrid stationary phase by non-covalent functionalization of single-walled carbon nanotubes with amino-derivatized silica gel for fast HPLC separation of aromatic compounds. <i>Talanta</i> , 2016, 149, 21-29.	2.9	17
1144	Characterization of post-translationally modified peptides by hydrophilic interaction and reverse phase liquid chromatography coupled to quadrupole-time-of-flight mass spectrometry. <i>Journal of Chromatography A</i> , 2016, 1428, 202-211.	1.8	15
1145	Possibilities and Limitations of Computer-Assisted Method Development in HILIC: A Case Study. <i>Chromatographia</i> , 2017, 80, 771-781.	0.7	9
1146	A positively charged porous graphitic carbon stationary phase for hydrophilic interaction liquid chromatography. <i>Talanta</i> , 2017, 164, 159-163.	2.9	20
1147	Stability-Indicating HILIC Method for the Determination of Citicoline and Characterization of its Degradation Products by LC-MS/TOF, 1H and 13C NMR. <i>Chromatographia</i> , 2017, 80, 265-274.	0.7	2
1148	Preparation and application of covalently bonded polysaccharide-modified stationary phase for per aqueous liquid chromatography. <i>Analytica Chimica Acta</i> , 2017, 964, 195-202.	2.6	26
1149	Evaluation of a hydrophilic interaction liquid chromatography design space for sugars and sugar alcohols. <i>Journal of Chromatography A</i> , 2017, 1489, 65-74.	1.8	31
1150	Hydrophilic interaction liquid chromatography for dalargin separation from its structural analogues and side products. <i>Journal of Chromatography A</i> , 2017, 1498, 155-162.	1.8	8
1151	Hydrophilic interaction chromatography and evaporative light scattering detection for the determination of polar analytes in Belgian endive. <i>Food Chemistry</i> , 2017, 229, 296-303.	4.2	6

#	ARTICLE	IF	CITATIONS
1152	Using the fundamentals of adsorption to understand peak distortion due to strong solvent effect in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2017, 1489, 95-106.	1.8	15
1153	Gradient separation of oligosaccharides and suppressing anomeric mutarotation with enhanced-fluidity liquid hydrophilic interaction chromatography. <i>Analytica Chimica Acta</i> , 2017, 960, 151-159.	2.6	28
1154	Recent advances in stationary phases and understanding of retention in hydrophilic interaction chromatography. A review. <i>Analytica Chimica Acta</i> , 2017, 967, 12-32.	2.6	226
1155	Hydrophilic Strong Anion Exchange (hSAX) Chromatography Enables Deep Fractionation of Tissue Proteomes. <i>Methods in Molecular Biology</i> , 2017, 1550, 69-82.	0.4	13
1156	Investigation of the temperature dependence of water adsorption on silica-based stationary phases in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1489, 143-148.	1.8	12
1157	Retention of glycopeptides analyzed using hydrophilic interaction chromatography is influenced by charge and carbon chain length of ion-pairing reagent for mobile phase. <i>Biomedical Chromatography</i> , 2017, 31, e3988.	0.8	14
1158	Glycoprotein Enrichment Analytical Techniques. <i>Methods in Enzymology</i> , 2017, 585, 397-429.	0.4	52
1159	Silica grafted with silanized carbon dots as a nano-on-micro packing material with enhanced hydrophilic selectivity. <i>Mikrochimica Acta</i> , 2017, 184, 2629-2636.	2.5	38
1160	HILIC Stationary Phase Based on Monodisperse-Porous Polymethacrylate Beads Functionalized with Zwitterionic Molecular Brushes. <i>Chromatographia</i> , 2017, 80, 565-575.	0.7	11
1161	Peptide Retention Time Prediction in Hydrophilic Interaction Liquid Chromatography: Data Collection Methods and Features of Additive and Sequence-Specific Models. <i>Analytical Chemistry</i> , 2017, 89, 5526-5533.	3.2	24
1162	Understanding and manipulating the separation in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1523, 49-71.	1.8	179
1163	Analysis of recombinant monoclonal antibodies in hydrophilic interaction chromatography: A generic method development approach. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 24-32.	1.4	32
1164	Modeling of HILIC retention behavior with theoretical models and new spline interpolation technique. <i>Journal of Chemometrics</i> , 2017, 31, e2910.	0.7	5
1165	Structural characterization of pyoverdines produced by <i>Pseudomonas putida</i> KT2440 and <i>Pseudomonas taiwanensis</i> VLB120. <i>BioMetals</i> , 2017, 30, 589-597.	1.8	14
1166	Poly(vinyl alcohol)-cationic cellulose copolymer encapsulated SiO ₂ stationary phase for hydrophilic interaction liquid chromatography. <i>RSC Advances</i> , 2017, 7, 21336-21341.	1.7	9
1167	Recent liquid chromatographic approaches and developments for the separation and purification of carbohydrates. <i>Analytical Methods</i> , 2017, 9, 3579-3593.	1.3	47
1169	Novel contribution of chromatography in the development and analyses of metformin hydrochloride in biological and environmental samples. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2017, 40, 599-615.	0.5	3
1170	Simultaneous determination of nine β -lactam antibiotics in human plasma by an ultrafast hydrophilic-interaction chromatography-tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1060, 138-143.	1.2	60

#	ARTICLE	IF	CITATIONS
1171	Analysis of illegal peptide drugs via HILIC-DAD-MS. <i>Talanta</i> , 2017, 174, 562-571.	2.9	21
1172	Characterization of isomeric glycan structures by LC-MS/MS. <i>Electrophoresis</i> , 2017, 38, 2100-2114.	1.3	123
1173	Controllable preparation of a hydrophilic/ion-exchange mixed-mode stationary phase by surface-initiated atom transfer radical polymerization using a mixture of two functional monomers. <i>Journal of Separation Science</i> , 2017, 40, 1861-1868.	1.3	12
1174	Separation of ten phosphorylated mono- and disaccharides using HILIC and ion-pairing interactions. <i>Analytica Chimica Acta</i> , 2017, 972, 102-110.	2.6	25
1175	Hydrophilic interaction chromatography with a focus on the drug-phosphate interaction in drug screening to determine the phospholipidosis induction risk. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1051, 33-40.	1.2	2
1176	The why and how of amino acid analytics in cancer diagnostics and therapy. <i>Journal of Biotechnology</i> , 2017, 242, 30-54.	1.9	48
1177	The Separation and Quantitation of Peptides with and without Oxidation of Methionine and Deamidation of Asparagine Using Hydrophilic Interaction Liquid Chromatography with Mass Spectrometry (HILIC-MS). <i>Journal of the American Society for Mass Spectrometry</i> , 2017, 28, 818-826.	1.2	19
1178	Preparation and characterization of hydrophilic hybrid monoliths via thiol-ene click polymerization and their applications in chromatographic analysis and glycopeptides enrichment. <i>Journal of Chromatography A</i> , 2017, 1498, 37-45.	1.8	26
1179	It is all about the solvent: on the importance of the mobile phase for ZIC-HILIC glycopeptide enrichment. <i>Analytical and Bioanalytical Chemistry</i> , 2017, 409, 529-538.	1.9	63
1180	Prediction of retention in hydrophilic interaction liquid chromatography using solute molecular descriptors based on chemical structures. <i>Journal of Chromatography A</i> , 2017, 1486, 59-67.	1.8	47
1181	Preparation and evaluation of surface-grafted block copolymers and random copolymers via surface-initiated atom transfer radical polymerization for hydrophilic/ion-exchange stationary phases. <i>RSC Advances</i> , 2017, 7, 46812-46822.	1.7	5
1182	Analysis of oligonucleotides by ion-pairing hydrophilic interaction liquid chromatography/electrospray ionization mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2017, 31, 2125-2134.	0.7	18
1183	A polyacrylamide-based silica stationary phase for the separation of carbohydrates using alcohols as the weak eluent in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1524, 153-159.	1.8	26
1184	The preparation of a poly (pentaerythritol tetraglycidyl ether-co-poly ethylene imine) organic monolithic capillary column and its application in hydrophilic interaction chromatography for polar molecules. <i>Analytica Chimica Acta</i> , 2017, 988, 104-113.	2.6	12
1185	Accurate prediction of retention in hydrophilic interaction chromatography by back calculation of high pressure liquid chromatography gradient profiles. <i>Journal of Chromatography A</i> , 2017, 1520, 75-82.	1.8	6
1186	The effect of solvation processes on amino acid- and peptide-silica stationary phases. <i>Journal of Separation Science</i> , 2017, 40, 4152-4159.	1.3	9
1187	Advances in Hydrophilic Interaction Liquid Chromatography. , 2017, , 39-87.		3
1189	Hydrophilic interaction liquid chromatography tandem mass spectrometry analysis of malonyl-coenzyme A in breast cancer cell cultures applying online solid-phase extraction. <i>Journal of Separation Science</i> , 2017, 40, 4303-4310.	1.3	4

#	ARTICLE	IF	CITATIONS
1191	Protein separations using enhanced-fluidity liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1523, 257-264.	1.8	27
1192	Alternative method for lactulose quantification in the presence of lactose in milk using HILIC with refractive index detection. <i>Analytical Methods</i> , 2017, 9, 4657-4662.	1.3	6
1193	Determination of Glyphosate, Maleic Hydrazide, Fosetyl Aluminum, and Ethephon in Grapes by Liquid Chromatography/Tandem Mass Spectrometry. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 7535-7541.	2.4	29
1194	Trimodal Mixed Mode Chromatography That Enables Efficient Offline Two-Dimensional Peptide Fractionation for Proteome Analysis. <i>Analytical Chemistry</i> , 2017, 89, 8884-8891.	3.2	22
1195	A fast and sensitive LC-MS/MS method for the quantification of fosfomycin in human urine and plasma using one sample preparation method and HILIC chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1061-1062, 263-269.	1.2	33
1196	Hydrophilic interaction liquid chromatography coupled to mass spectrometry-based detection to determine emerging organic contaminants in environmental samples. <i>TrAC - Trends in Analytical Chemistry</i> , 2017, 94, 141-149.	5.8	38
1197	Comparison of newly developed hydroxyl-functionalized monodisperse HILIC columns new HILIC column. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2017, 40, 649-655.	0.5	1
1198	Applicability of retention modelling in hydrophilic-interaction liquid chromatography for algorithmic optimization programs with gradient-scanning techniques. <i>Journal of Chromatography A</i> , 2017, 1530, 104-111.	1.8	36
1199	LIQUID CHROMATOGRAPHY-MASS SPECTROMETRY ANALYSIS OF DIALYSATE AND APPLICATIONS WITH SELECTIVE NEURONAL STIMULATION. , 2017, , 171-192.		0
1200	Milestones in the development of liquid chromatography. , 2017, , 1-15.		3
1201	Hydrophilic interaction liquid chromatography. , 2017, , 147-169.		1
1202	Hydrophilic interaction liquid chromatography of hydroxy aromatic carboxylic acid positional isomers. <i>Analytica Chimica Acta</i> , 2017, 996, 98-105.	2.6	5
1203	Polar silica-based stationary phases. Part II- Neutral silica stationary phases with surface bound maltose and sorbitol for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1508, 24-32.	1.8	11
1204	Comparative genetic, proteomic and phosphoproteomic analysis of <i>C. elegans</i> embryos with a focus on ham-1/STOX and pig-1/MELK in dopaminergic neuron development. <i>Scientific Reports</i> , 2017, 7, 4314.	1.6	11
1205	Polar and nonpolar organic polymer-based monolithic columns for capillary electrochromatography and high-performance liquid chromatography. <i>Electrophoresis</i> , 2017, 38, 60-79.	1.3	29
1206	Application of hydrophilic interaction liquid chromatography coupled with mass spectrometry in the analysis of phosphorothioate oligonucleotides in serum. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1040, 282-288.	1.2	30
1207	Hydrophilic materials in sample pretreatment. <i>TrAC - Trends in Analytical Chemistry</i> , 2017, 86, 172-184.	5.8	64
1208	Characterization of lectin binding affinities via direct LC-MS profiling: implications for glycopeptide enrichment and separation strategies. <i>Analyst</i> , The, 2017, 142, 65-74.	1.7	19

#	ARTICLE	IF	CITATIONS
1209	Study of Separation and Identification of the Active Ingredients in <i>Gardenia jasminoides</i> Ellis Based on a Two-Dimensional Liquid Chromatography by Coupling Reversed Phase Liquid Chromatography and Hydrophilic Interaction Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 2017, 55, 75-81.	0.7	8
1210	Studies on Behaviors of Interactions Between New Polymer-based ZIC-HILIC Stationary Phases and Carboxylic Acids. <i>Journal of Chromatographic Science</i> , 2017, 55, 52-59.	0.7	22
1211	Reversed-phase separation methods for glycan analysis. <i>Analytical and Bioanalytical Chemistry</i> , 2017, 409, 359-378.	1.9	94
1212	Purification of high-purity glycyrrhizin from licorice using hydrophilic interaction solid phase extraction coupled with preparative reversed-phase liquid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1040, 47-52.	1.2	27
1213	New Materials for Stationary Phases in Liquid Chromatography/Mass Spectrometry. , 2017, , 179-225.		2
1214	Separation of structurally related primary aliphatic amines using hydrophilic interaction chromatography with fluorescence detection after postcolumn derivatization with <i>o</i> -phthaldialdehyde/mercaptoethanol. <i>Journal of Separation Science</i> , 2017, 40, 4689-4699.	1.3	6
1215	Fast and Simple Determination of 3-Aminopiperidine without Derivatization Using High Performance Liquid Chromatography–Charged Aerosol Detector with an Ion-Exchange/Reversed-Phase Mixed-mode Column. <i>Analytical Sciences</i> , 2017, 33, 293-298.	0.8	4
1216	Separation of bioactive chamazulene from chamomile extract using metal-organic framework. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 146, 126-134.	1.4	43
1217	HILIC-MS rat brain analysis, a new approach for the study of ischemic attack. <i>Translational Neuroscience</i> , 2017, 8, 70-75.	0.7	6
1218	Predictive Liquid Chromatography of Peptides Based on Hydrophilic Interactions for Mass Spectrometry-Based Proteomics. <i>Journal of Analytical Chemistry</i> , 2017, 72, 1375-1382.	0.4	3
1219	Capillary Electrophoresis as a Green Alternative Separation Technique. , 2017, , 517-532.		8
1220	Modelling of Hydrophilic Interaction Liquid Chromatography Stationary Phases Using Chemometric Approaches. <i>Metabolites</i> , 2017, 7, 54.	1.3	6
1221	Diverse Mechanisms of Sulfur Decoration in Bacterial tRNA and Their Cellular Functions. <i>Biomolecules</i> , 2017, 7, 33.	1.8	25
1222	LC/MS Analysis of Marine Toxins. <i>Comprehensive Analytical Chemistry</i> , 2017, 78, 137-192.	0.7	3
1223	Nanosized Drug Carriers for Oral Delivery of Anticancer Compounds and the Importance of the Chromatographic Techniques. , 2017, , 165-195.		9
1224	Applications of liquid chromatography in the quality control of traditional Chinese medicines: An overview. , 2017, , 617-663.		0
1225	Retention Behavior of Inorganic Anions in Hydrophilic Interaction Chromatography. <i>Analytical Sciences</i> , 2017, 33, 619-623.	0.8	11
1226	A quantitative metabolomics study of bacterial metabolites in different domains. <i>Analytica Chimica Acta</i> , 2018, 1037, 237-244.	2.6	9

#	ARTICLE	IF	CITATIONS
1227	A sulfonated chito-oligosaccharide modified silica material for hydrophilic interaction liquid chromatography and its chromatographic evaluation. <i>Analytical Methods</i> , 2018, 10, 1258-1265.	1.3	8
1228	Cell-Free DNA Blood Collection Tubes Are Appropriate for Clinical Proteomics: A Demonstration in Colorectal Cancer. <i>Proteomics - Clinical Applications</i> , 2018, 12, e1700121.	0.8	11
1229	The versatility of <i>N</i> , <i>O</i> -dialkylphosphoramidate stationary phase-separations in HILIC, highly aqueous RP LC conditions and purely aqueous mobile phase. <i>Analyst</i> , The, 2018, 143, 1217-1223.	1.7	9
1230	High speed hydrophilic interaction liquid chromatographic method for simultaneous determination of selected pharmaceuticals in wastewater using a cyano-bonded silica column. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018, 41, 180-187.	0.5	17
1231	Mobile phase effects in reversed-phase and hydrophilic interaction liquid chromatography revisited. <i>Journal of Chromatography A</i> , 2018, 1543, 48-57.	1.8	19
1232	The determination of pharmaceutically active thiols using hydrophilic interaction chromatography followed postcolumn derivatization with <i>o</i> -phthalaldehyde and fluorescence detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 156, 1-7.	1.4	15
1233	Simultaneous determination of twelve polar pteridines including dihydro- and tetrahydropteridine in human urine by hydrophilic interaction liquid chromatography with tandem mass spectrometry. <i>Biomedical Chromatography</i> , 2018, 32, e4244.	0.8	4
1234	New stationary phase for hydrophilic interaction chromatography to separate chito-oligosaccharides with degree of polymerization 2-6. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2018, 1081-1082, 33-40.	1.2	5
1235	Free silanols and ionic liquids as their suppressors in liquid chromatography. <i>Journal of Chromatography A</i> , 2018, 1559, 17-43.	1.8	29
1236	Characterization of Retention Mechanisms in Mixed-Mode HPLC with a Bimodal Reversed-Phase/Cation-Exchange Stationary Phase. <i>Chromatographia</i> , 2018, 81, 387-399.	0.7	13
1238	Effect of salts on retention in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2018, 1538, 45-53.	1.8	61
1239	Chemical, computational and functional insights into the chemical stability of the Hedgehog pathway inhibitor GANT61. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2018, 33, 349-358.	2.5	45
1240	Peptide retention time prediction in hydrophilic interaction liquid chromatography. Comparison of separation selectivity between bare silica and bonded stationary phases. <i>Journal of Chromatography A</i> , 2018, 1534, 75-84.	1.8	20
1241	Terminology of separation methods (IUPAC Recommendations 2017). <i>Pure and Applied Chemistry</i> , 2018, 90, 181-231.	0.9	32
1242	Comprehensive characterization of neurochemicals in three zebrafish chemical models of human acute organophosphorus poisoning using liquid chromatography-tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 1735-1748.	1.9	28
1243	Analysis of Morpholinium Ionic Liquid Cations by Hydrophilic Interaction Columns Coupled with Indirect UV Detection. <i>Journal of the Chinese Chemical Society</i> , 2018, 65, 726-734.	0.8	2
1244	Hyperbranched Polyglycerol Functionalized Silica Stationary Phase for Hydrophilic Interaction Liquid Chromatography. <i>Analytical Sciences</i> , 2018, 34, 433-438.	0.8	6
1245	Determination of alkyl ammonium ionic liquid cations by hydrophilic interaction liquid chromatography and its application in analysis of environmental water. <i>Analytical Methods</i> , 2018, 10, 2812-2820.	1.3	13

#	ARTICLE	IF	CITATIONS
1246	Development of a HILIC method for the determination of 5-fluorouracil from nano drug delivery systems and rat skin extracts. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 154, 285-293.	1.4	17
1247	Extending the Compatibility of the SP3 Paramagnetic Bead Processing Approach for Proteomics. <i>Journal of Proteome Research</i> , 2018, 17, 1730-1740.	1.8	186
1248	Simultaneous determination of ribonucleoside and deoxyribonucleoside triphosphates in biological samples by hydrophilic interaction liquid chromatography coupled with tandem mass spectrometry. <i>Nucleic Acids Research</i> , 2018, 46, e66-e66.	6.5	40
1249	Plasma phospholipid profiling of a mouse model of anxiety disorder by hydrophilic interaction liquid chromatography coupled to high-resolution mass spectrometry. <i>Biomedical Chromatography</i> , 2018, 32, e4202.	0.8	2
1250	Simultaneous determination of 14 oral antihyperglycaemic drugs in human urine by liquid chromatography-tandem mass spectrometry. <i>Archives of Pharmacal Research</i> , 2018, 41, 530-543.	2.7	8
1251	Determination of piperidinium ionic liquid cations in environmental water samples by solid phase extraction and hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2018, 1559, 136-140.	1.8	21
1252	The benefits (and misfortunes) of SDS in top-down proteomics. <i>Journal of Proteomics</i> , 2018, 175, 75-86.	1.2	25
1253	Characterization of the Retention of Artificial Sweeteners by Hydrophilic Interaction Liquid Chromatography. <i>Analytical Letters</i> , 2018, 51, 49-72.	1.0	5
1254	Surface-bonded amide-functionalized imidazolium ionic liquid as stationary phase for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2018, 1559, 141-148.	1.8	37
1255	Mobile phase effects on the retention on polar columns with special attention to the dual hydrophilic interaction-reversed-phase liquid chromatography mechanism, a review. <i>Journal of Separation Science</i> , 2018, 41, 145-162.	1.3	45
1256	Thiol-ene-grafting of silica particles with three-dimensional branched copolymer for HILIC/cation-exchange chromatographic separation and N-glycopeptide enrichment. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 1019-1027.	1.9	24
1257	Optimizing separations in online comprehensive two-dimensional liquid chromatography. <i>Journal of Separation Science</i> , 2018, 41, 68-98.	1.3	176
1258	Chemometric-assisted method development in hydrophilic interaction liquid chromatography: A review. <i>Analytica Chimica Acta</i> , 2018, 1000, 20-40.	2.6	81
1259	Studies on the retention mechanism of solutes in hydrophilic interaction chromatography using stoichiometric displacement theory I. The linear relationship of $\lg k'$ vs. $\lg [H_2O]$. <i>Talanta</i> , 2018, 176, 499-508.	2.9	10
1260	Utilization of large volume zwitterionic hydrophilic interaction liquid chromatography for the analysis of polar pharmaceuticals in aqueous environmental samples: Benefits and limitations. <i>Journal of Chromatography A</i> , 2018, 1535, 27-43.	1.8	49
1261	Mass spectrometry for protein sialoglycosylation. <i>Mass Spectrometry Reviews</i> , 2018, 37, 652-680.	2.8	35
1262	Mass Spectrometry Analyses of Multicellular Tumor Spheroids. <i>Proteomics - Clinical Applications</i> , 2018, 12, e1700124.	0.8	20
1263	High-performance liquid chromatography mass spectrometry of gold and alloy clusters protected by hydrophilic thiolates. <i>Nanoscale</i> , 2018, 10, 1641-1649.	2.8	42

#	ARTICLE	IF	CITATIONS
1264	1H-Tetrazole-5-amine Immobilized on Substituted Polymer Gel/Silica as a New Stationary Phase for Hydrophilic Interaction Chromatography. <i>Chromatographia</i> , 2018, 81, 349-357.	0.7	6
1265	Comprehensive phospholipid and sphingomyelin profiling of different brain regions in mouse model of anxiety disorder using online two-dimensional (HILIC/RP)-LC/MS method. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 149, 308-317.	1.4	23
1266	Stability and assessment of amino acids in parenteral nutrition solutions. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 147, 125-139.	1.4	26
1268	Multidimensional Liquid Chromatography. , 2018, , .		0
1269	Solid-phase Extraction Properties of Novel Sorbents Modified with Adenine for Nucleic Acid Related Compounds. <i>Bunseki Kagaku</i> , 2018, 67, 445-451.	0.1	4
1270	Fast and sensitive method for the determination of trace 1,2,4-triazole and 4-amino-1,2,4-triazole by hydrophilic interaction liquid chromatography (HILIC) in antifungal drug substance. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018, 41, 770-779.	0.5	1
1271	Column Technology for Modern High Performance Liquid Chromatography (HPLC). , 2018, , .		1
1272	Liquid Chromatographic Strategies for Separation of Bioactive Compounds in Food Matrices. <i>Molecules</i> , 2018, 23, 3091.	1.7	18
1273	Development of a hydrophilic interaction liquid chromatography (HILIC) method for the chemical characterization of water-soluble isoprene epoxydiol (IEPOX)-derived secondary organic aerosol. <i>Environmental Sciences: Processes and Impacts</i> , 2018, 20, 1524-1536.	1.7	66
1274	Novel zwitterionic HILIC stationary phase for the determination of ethyl glucuronide in human hair by LC-MS/MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2018, 1100-1101, 33-38.	1.2	17
1275	Simultaneous quantification of urea and allantoin in cosmetic products by nano-HPLC using a highly hydrophilic monolith. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018, 41, 780-785.	0.5	2
1276	Poly(2-carboxyethyl acrylate-co-ethylene glycol dimethacrylate) monolithic precursor. Part II. Carbodiimide assisted post-polymerization modification with tris and d-Glucamine for use in hydrophilic interaction capillary liquid chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018, 41, 684-691.	0.5	4
1277	Clinical application of quantitative glycomics. <i>Expert Review of Proteomics</i> , 2018, 15, 1007-1031.	1.3	40
1278	Regularities of adsorption of water-soluble vitamins on the surface of microdispersed sintered detonation nanodiamond. <i>Adsorption</i> , 2018, 24, 637-645.	1.4	2
1279	Trace Phosphate Improves ZIC-pHILIC Peak Shape, Sensitivity, and Coverage for Untargeted Metabolomics. <i>Journal of Proteome Research</i> , 2018, 17, 3537-3546.	1.8	32
1280	Post-polymerization modification of a new reactive monolith for reversed phase and hydrophilic interaction capillary electrochromatography of neutral, polar, and biologically active compounds. <i>Polymers for Advanced Technologies</i> , 2018, 29, 2110-2120.	1.6	6
1281	Method optimization of hydrophilic interaction chromatography separation of nucleotides using design of experiment approaches I: Comparison of several zwitterionic columns. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 158, 307-316.	1.4	13
1282	Quantification of Low-Abundant Phosphorylated Carbohydrates Using HILIC-QqQ-MS/MS. <i>Methods in Molecular Biology</i> , 2018, 1778, 71-86.	0.4	2

#	ARTICLE	IF	CITATIONS
1283	Analysis of sulfates on low molecular weight heparin using mass spectrometry: structural characterization of enoxaparin. <i>Expert Review of Proteomics</i> , 2018, 15, 503-513.	1.3	12
1284	High-performance liquid chromatographic separation of 8-aminopyrene-1,3,6-trisulfonic acid labeled N-glycans using a functional tetrazole hydrophilic interaction liquid chromatography column. <i>Journal of Chromatography A</i> , 2018, 1566, 44-50.	1.8	9
1285	Mass spectrometric characterization of siderophores produced by <i>Pseudomonas taiwanensis</i> VLB120 assisted by stable isotope labeling of nitrogen source. <i>BioMetals</i> , 2018, 31, 785-795.	1.8	3
1286	Peptide-Based Derivative-Grafted Silica for Molecular Recognition System. , 2018, , 235-294.		0
1287	Preparation of attapulgite nanoparticles-based hybrid monolithic column with covalent bond for hydrophilic interaction liquid chromatography. <i>Talanta</i> , 2018, 189, 397-403.	2.9	9
1288	Separation of highly charged compounds using competing ions with hydrophilic interaction liquid chromatography " Application to assay of cellular nucleotides. <i>Journal of Chromatography A</i> , 2018, 1567, 233-238.	1.8	3
1289	Quantification of ondansetron, granisetron and tropisetron in goat plasma using hydrophilic interaction liquid chromatography-solid phase extraction coupled with hydrophilic interaction liquid chromatography-triple quadrupole tandem mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2018, 1095, 50-58.	1.2	8
1290	Preparation and evaluation of surface-bonded phenylglycine zwitterionic stationary phase. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 5941-5950.	1.9	7
1291	Determination of glucosamine and monitoring of its mutarotation by hydrophilic interaction liquid chromatography with evaporative light scattering detector. <i>Biomedical Chromatography</i> , 2018, 32, e4368.	0.8	8
1292	Hydrophilic Interaction Chromatography: Current Trends and Applications. , 2018, , 100-100.		0
1293	Investigating hydrophilic and electrostatic properties of surfactants using retention on two mixed-mode liquid chromatographic columns. <i>Journal of Chromatography A</i> , 2018, 1571, 185-192.	1.8	19
1294	A rapid UHPLC-HILIC method for algal guanosine 5'-diphosphate 3'-diphosphate (ppGpp) and the potential separation mechanism. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2018, 1096, 143-153.	1.2	14
1295	Hydrophilic interaction liquid chromatography of aminoglycoside antibiotics with a diol-type stationary phase. <i>Analytica Chimica Acta</i> , 2018, 1044, 174-180.	2.6	15
1296	Definition of HILIC System and Quantitative Analysis of Retention Mechanisms. <i>Current Chromatography</i> , 2018, 5, 43-52.	0.1	9
1297	Class Separation of the Trace Polar Species Present in Diesel Using Hydrophilic Interaction Chromatography and High Resolution Mass Spectrometry. <i>Energy & Fuels</i> , 2018, 32, 8944-8954.	2.5	5
1298	Molecular Structure Characterization of Riverine and Coastal Dissolved Organic Matter with Ion Mobility Quadrupole Time-of-Flight LCMS (IM Q-TOF LCMS). <i>Environmental Science & Technology</i> , 2018, 52, 7182-7191.	4.6	59
1299	Hydrophilic interaction liquid chromatography coupled with quadrupole-orbitrap ultra high resolution mass spectrometry to quantitate nucleobases, nucleosides, and nucleotides during white tea withering process. <i>Food Chemistry</i> , 2018, 266, 343-349.	4.2	29
1300	Geopolymers as a New Class of High pH Stable Supports with Different Chromatographic Selectivity. <i>Analytical Chemistry</i> , 2018, 90, 8139-8146.	3.2	18

#	ARTICLE	IF	CITATIONS
1301	Online Removal of Sodium Dodecyl Sulfate via Weak Cation Exchange in Liquid Chromatographyâ€“Mass Spectrometry Based Proteomics. <i>Journal of Proteome Research</i> , 2018, 17, 2390-2400.	1.8	9
1302	Application of deep eutectic solvents in chromatography: A review. <i>TrAC - Trends in Analytical Chemistry</i> , 2019, 120, 115623.	5.8	91
1303	Direct Analysis of Underivatized Amino Acids in Plant Extracts by LC-MS/MS (Improved Method). <i>Methods in Molecular Biology</i> , 2019, 2030, 403-414.	0.4	7
1304	Hydrophilic Interaction Liquid Chromatography Coupled to Mass Spectrometry and Multivariate Analysis of the De Novo Pyrimidine Pathway Metabolites. <i>Biomolecules</i> , 2019, 9, 328.	1.8	2
1305	High efficiency and simple preparation of polyacrylamide coated silica stationary phase for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2019, 1605, 360357.	1.8	17
1306	Finding the Sweet Spot in ERLIC Mobile Phase for Simultaneous Enrichment of N-Glyco and Phosphopeptides. <i>Journal of the American Society for Mass Spectrometry</i> , 2019, 30, 2491-2501.	1.2	23
1307	Analytical Methods for Mass Spectrometry-Based Metabolomics Studies. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1140, 635-647.	0.8	28
1308	Advances in hydrophilic nanomaterials for glycoproteomics. <i>Chemical Communications</i> , 2019, 55, 10359-10375.	2.2	62
1309	A comprehensive approach to detecting multitudinous bioactive peptides in equine plasma and urine using hydrophilic interaction liquid chromatography coupled to high resolution mass spectrometry. <i>Drug Testing and Analysis</i> , 2019, 11, 1308-1325.	1.6	9
1310	Evaluating Relative Retention of Polar Stationary Phases in Hydrophilic Interaction Chromatography. <i>Separations</i> , 2019, 6, 42.	1.1	4
1312	An optimized mixedâ€“mode stationary phase based on silica monolith particles for the separation of peptides and proteins in highâ€“performance liquid chromatography. <i>Journal of Separation Science</i> , 2019, 42, 3621-3630.	1.3	9
1313	Isolation and characterization of glycosylated neuropeptides. <i>Methods in Enzymology</i> , 2019, 626, 147-202.	0.4	7
1314	Analytical strategies for the determination of amino acids: Past, present and future trends. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1132, 121819.	1.2	63
1316	Characterization of hydrophilic interaction liquid chromatography retention by a linear free energy relationship. Comparison to reversed- and normal-phase retentions. <i>Analytica Chimica Acta</i> , 2019, 1092, 132-143.	2.6	26
1317	The role of fluoroalcohols as counter anions for ionâ€“pairing reversedâ€“phase liquid chromatography/highâ€“resolution electrospray ionization mass spectrometry analysis of oligonucleotides. <i>Rapid Communications in Mass Spectrometry</i> , 2019, 33, 697-709.	0.7	21
1319	Protocol to compare column performance applied to hydrophilic interaction liquid chromatography. <i>Microchemical Journal</i> , 2019, 149, 103973.	2.3	9
1320	Facile High-Performance Liquid Chromatography Mass Spectrometry Method for Analysis of Cyclocreatine and Phosphocyclocreatine in Complex Mixtures of Amino Acids. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 7190-7196.	2.4	4
1321	Hydrophilic Interaction Chromatography. <i>Journal of Analytical Chemistry</i> , 2019, 74, 415-424.	0.4	24

#	ARTICLE	IF	CITATIONS
1322	The non-destructive separation of diverse astrobiologically relevant organic molecules by customizable capillary zone electrophoresis and monolithic capillary electrochromatography. <i>International Journal of Astrobiology</i> , 2019, 18, 562-574.	0.9	5
1323	Recent trends in analytical methods for water-soluble vitamins. <i>Journal of Chromatography A</i> , 2019, 1606, 360245.	1.8	42
1324	HPLC Enantioseparations with Polysaccharide-Based Chiral Stationary Phases in HILIC Conditions. <i>Methods in Molecular Biology</i> , 2019, 1985, 127-146.	0.4	6
1325	Accurate and efficient amino acid analysis for protein quantification using hydrophilic interaction chromatography coupled tandem mass spectrometry. <i>Plant Methods</i> , 2019, 15, 46.	1.9	67
1326	Development and validation of a novel UPLC-ELSD method for the assessment of lipid composition of nanomedicine formulation. <i>International Journal of Pharmaceutics</i> , 2019, 566, 11-23.	2.6	16
1327	Application of a Sensitive and Reproducible Label-Free Proteomic Approach to Explore the Proteome of Individual Meiotic-Phase Barley Anthers. <i>Frontiers in Plant Science</i> , 2019, 10, 393.	1.7	12
1328	Comparison of reversed-phase, anion-exchange, and hydrophilic interaction HPLC for the analysis of nucleotides involved in biological enzymatic pathways. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2019, 42, 184-193.	0.5	7
1329	Simultaneous determination of cartap and its metabolite in tea using hydrophilic interaction chromatography tandem mass spectrometry and the combination of dispersive solid phase extraction and solid phase extraction. <i>Journal of Chromatography A</i> , 2019, 1600, 148-157.	1.8	22
1330	Comparison of different fractionation strategies for in-depth phosphoproteomics by liquid chromatography tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 3417-3424.	1.9	10
1331	Nanogold hybrid silica gel and its 1-octadecanethiol self-assembled modified composite as a stationary phase for liquid chromatography. <i>Analyst</i> , 2019, 144, 3072-3079.	1.7	3
1332	Protein Aggregation Capture on Microparticles Enables Multipurpose Proteomics Sample Preparation*. <i>Molecular and Cellular Proteomics</i> , 2019, 18, 1027a-1035.	2.5	189
1333	L-lysine-derived highly selective stationary phases for hydrophilic interaction chromatography: Effect of chain length on selectivity, efficiency, resolution, and asymmetry. <i>Separation Science Plus</i> , 2019, 2, 42-50.	0.3	5
1334	In Vitro Antiatherothrombotic Effects of Extracts from <i>Berberis Vulgaris</i> L., <i>Teucrium Polium</i> L., and <i>Orthosiphon Stamineus</i> Benth. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019, 2019, 1-10.	0.5	8
1335	Polyethyleneimine-functionalized carbon dots and their precursor co-immobilized on silica for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2019, 1597, 142-148.	1.8	55
1336	Quantitative determination of beta-cyclodextrin in a powder insulin formulation for nasal delivery using hydrophilic interaction chromatography with evaporative light scattering detection. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2019, 42, 74-78.	0.5	5
1337	Determination of Piperidinium Cations by Hydrophilic Interaction Chromatography with Imidazolium Ionic Liquids as Mobile Phase Additives. <i>Journal of Analytical Chemistry</i> , 2019, 74, 126-133.	0.4	1
1338	Analysis of Non-Ionic Surfactant Triton X-100 Using Hydrophilic Interaction Liquid Chromatography and Mass Spectrometry. <i>Molecules</i> , 2019, 24, 1223.	1.7	14
1339	Evaluating the Adsorbed Water Layer on Polar Stationary Phases for Hydrophilic Interaction Chromatography (HILIC). <i>Separations</i> , 2019, 6, 19.	1.1	15

#	ARTICLE	IF	CITATIONS
1340	SP2: Rapid and Automatable Contaminant Removal from Peptide Samples for Proteomic Analyses. <i>Journal of Proteome Research</i> , 2019, 18, 1644-1656.	1.8	36
1341	Oligonucleotide analysis by hydrophilic interaction liquid chromatography-mass spectrometry in the absence of ion-pair reagents. <i>Journal of Chromatography A</i> , 2019, 1595, 39-48.	1.8	70
1342	Development and application of a HILIC UHPLC-MS method for polar fecal metabolome profiling. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1109, 142-148.	1.2	26
1343	Further Insights into Structural Diversity of Phosphorus-Based Decomposition Products in Lithium Ion Battery Electrolytes via Liquid Chromatographic Techniques Hyphenated to Ion Trap-Time-of-Flight Mass Spectrometry. <i>Analytical Chemistry</i> , 2019, 91, 3980-3988.	3.2	27
1344	Enhanced fluidity liquid chromatography: A guide to scaling up from analytical to preparative separations. <i>Journal of Chromatography A</i> , 2019, 1595, 190-198.	1.8	31
1345	Towards a high peak capacity of 130 using nanoflow hydrophilic interaction liquid chromatography. <i>Analytica Chimica Acta</i> , 2019, 1062, 147-155.	2.6	16
1346	Fabrication of hydrophilic multilayer magnetic probe for salivary glycopeptidome analysis. <i>Journal of Chromatography A</i> , 2019, 1587, 24-33.	1.8	19
1347	Zwitterionic HILIC stationary phase as a valuable alternative in separative techniques: Application to the analysis of gamma-hydroxybutyric acid and its metabolite in hair. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1134-1135, 121876.	1.2	8
1349	Ammonium Fluoride as Suitable Additive for HILIC-Based LC-HRMS Metabolomics. <i>Metabolites</i> , 2019, 9, 292.	1.3	19
1351	Poly(itaconic acid)-grafted silica stationary phase prepared in deep eutectic solvents and its unique performance in hydrophilic interaction chromatography. <i>Talanta</i> , 2019, 191, 265-271.	2.9	32
1352	Glycopolymer Grafted Silica Gel as Chromatographic Packing Materials. <i>International Journal of Molecular Sciences</i> , 2019, 20, 10.	1.8	22
1353	The fabrication of poly (polyethylene glycol diacrylate) monolithic porous layer open tubular (mono-PLOT) columns and applications in hydrophilic interaction chromatography and capillary gas chromatography for small molecules. <i>Electrophoresis</i> , 2019, 40, 521-529.	1.3	7
1354	The History of the Core-Shell Particles and Applications in Active Pharmaceutical Ingredients Via Liquid Chromatography. <i>Chromatographia</i> , 2019, 82, 17-48.	0.7	26
1355	Interaction of toluene with polar stationary phases under conditions typical of hydrophilic interaction chromatography probed by saturation transfer difference nuclear magnetic resonance spectroscopy. <i>Journal of Chromatography A</i> , 2019, 1588, 58-67.	1.8	6
1356	Universal retention standard for peptide separations using various modes of high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2019, 1588, 163-168.	1.8	14
1357	Column Characterization and Selection Systems in Reversed-Phase High-Performance Liquid Chromatography. <i>Chemical Reviews</i> , 2019, 119, 3674-3729.	23.0	191
1358	Effect of nature of electrolytes on retention and selectivity in hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2019, 1584, 80-86.	1.8	9
1359	Preparation of a monolithic column with a mixed-mode stationary phase of reversed-phase/hydrophilic interaction for capillary liquid chromatography. <i>Journal of Separation Science</i> , 2019, 42, 662-669.	1.3	12

#	ARTICLE	IF	CITATIONS
1360	Investigation and prediction of retention characteristics of imidazoline and serotonin receptor ligands and their related compounds on mixed-mode stationary phase. <i>Journal of Chromatography A</i> , 2019, 1585, 92-104.	1.8	15
1361	Facile one-pot synthesized hydrothermal carbon from cyclodextrin: A stationary phase for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2019, 1585, 144-151.	1.8	10
1362	Single-pot, solid-phase-enhanced sample preparation for proteomics experiments. <i>Nature Protocols</i> , 2019, 14, 68-85.	5.5	802
1363	Sensitive analysis of trehalose-6-phosphate and related sugar phosphates in plant tissues by chemical derivatization combined with hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2019, 1592, 82-90.	1.8	13
1364	Facile fabrication of zwitterionic magnetic composites by one-step distillation-precipitation polymerization for highly specific enrichment of glycopeptides. <i>Analytica Chimica Acta</i> , 2019, 1053, 43-53.	2.6	27
1365	Deepening the Understanding of Thiolate-Protected Metal Clusters Using High-Performance Liquid Chromatography. <i>Bulletin of the Chemical Society of Japan</i> , 2019, 92, 664-695.	2.0	29
1366	Hydrophilic-interaction planar chromatography in ultra-sensitive determination of β -aminocephalosporin antibiotics. Application to analysis of cefalexin in goat milk samples using modified QuEChERS extraction technique. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019, 166, 421-434.	1.4	6
1367	Fundamental Properties of Packing Materials for Liquid Chromatography. <i>Separations</i> , 2019, 6, 2.	1.1	9
1368	HILIC Chromatography: Powerful Technique in the Analysis of Polyphenols. , 2019, , 341-351.		1
1369	Application of lipidomics in bivalve aquaculture, a review. <i>Reviews in Aquaculture</i> , 2020, 12, 678-702.	4.6	21
1370	Retention behavior of ginsenosides in a sulfo-based high performance liquid chromatography column. <i>Journal of Chromatography A</i> , 2020, 1610, 460542.	1.8	5
1371	Retention characteristics of poly(N-(1H-tetrazole-5-yl)-methacrylamide)-bonded stationary phase in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2020, 1609, 460500.	1.8	8
1372	Hydrophobic and hydrophilic interactions in countercurrent chromatography. <i>Journal of Chromatography A</i> , 2020, 1611, 460576.	1.8	13
1373	Enhancing the selectivity of polar hydrophilic analytes with a low concentration of barium ions in the mobile phase using geopolymers and silica supports. <i>Talanta</i> , 2020, 207, 120339.	2.9	1
1374	Analysis of the behaviour of confined molecules using ^2H nuclear magnetic relaxation dispersion. <i>Molecular Physics</i> , 2020, 118, .	0.8	2
1375	Recent advances in hydrophilic interaction liquid chromatography materials for glycopeptide enrichment and glycan separation. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 124, 115570.	5.8	103
1376	Quantitative structure-retention relationships of amino acids on the amino acid- and peptide-silica stationary phases for liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1609, 460514.	1.8	8
1377	Accurate modelling of the retention behaviour of peptides in gradient-elution hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1614, 460650.	1.8	19

#	ARTICLE	IF	CITATIONS
1378	Sustainable development and environmental analysis. , 2020, , 343-379.		0
1379	Application of two-dimensional liquid chromatography in the separation of traditional Chinese medicine. <i>Journal of Separation Science</i> , 2020, 43, 87-104.	1.3	37
1380	Dual-mode hydrophilic interaction normal phase and reversed phase liquid chromatography of polar compounds on a single column. <i>Journal of Separation Science</i> , 2020, 43, 70-86.	1.3	14
1381	Citrullinome of <i>Porphyromonas gingivalis</i> Outer Membrane Vesicles: Confident Identification of Citrullinated Peptides. <i>Molecular and Cellular Proteomics</i> , 2020, 19, 167-180.	2.5	18
1382	Chemical separation and characterization of complex samples with herbal medicine. <i>TrAC - Trends in Analytical Chemistry</i> , 2020, 124, 115775.	5.8	11
1383	Investigation of Selectivity of Amino Functionalised Phases for Pharmaceutical Applications. <i>Chromatographia</i> , 2020, 83, 159-171.	0.7	2
1384	Imidazolium ionic liquids-derived carbon dots-modified silica stationary phase for hydrophilic interaction chromatography. <i>Talanta</i> , 2020, 209, 120518.	2.9	43
1385	Profiling of post-translational modifications by chemical and computational proteomics. <i>Chemical Communications</i> , 2020, 56, 13506-13519.	2.2	15
1386	One-pot hydrothermal cross-linking preparation of poly(vinylpyrrolidone) immobilized silica stationary phase for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2020, 1633, 461656.	1.8	5
1387	An ultrafast and highly efficient enrichment method for both N-Glycopeptides and N-Glycans by bacterial cellulose. <i>Analytica Chimica Acta</i> , 2020, 1140, 60-68.	2.6	10
1388	Application of High-Performance Liquid Chromatography for Simultaneous Determination of Tenofovir and Creatinine in Human Urine and Plasma Samples. <i>Pharmaceuticals</i> , 2020, 13, 367.	1.7	4
1389	Mass Spectrometry for Natural Product Discovery. , 2020, , 263-306.		13
1390	HILIC characterization: Estimation of phase volumes and composition for a zwitterionic column. <i>Analytica Chimica Acta</i> , 2020, 1130, 39-48.	2.6	15
1391	Liquid chromatography-high resolution mass spectrometry for the analysis of bioactive natural products. <i>Studies in Natural Products Chemistry</i> , 2020, , 331-353.	0.8	4
1392	RNA abasic sites in yeast and human cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 20689-20695.	3.3	27
1395	Application of hydrophilic interaction liquid chromatography coupled with tandem mass spectrometry for the retention and sensitivity studies of antisense oligonucleotides. <i>Journal of Chromatography A</i> , 2020, 1622, 461100.	1.8	19
1396	Probing the retention mechanism of small hydrophilic molecules in hydrophilic interaction chromatography using saturation transfer difference nuclear magnetic resonance spectroscopy. <i>Journal of Chromatography A</i> , 2020, 1623, 461130.	1.8	12
1397	Nitrogen-doping to enhance the separation selectivity of glucose-based carbon dots-modified silica stationary phase for hydrophilic interaction chromatography. <i>Talanta</i> , 2020, 218, 121140.	2.9	29

#	ARTICLE	IF	CITATIONS
1398	A novel process for the preparation of Cys-Si-NIPAM as a stationary phase of hydrophilic interaction liquid chromatography (HILIC). <i>Talanta</i> , 2020, 218, 121154.	2.9	9
1399	LC-HRMS based on mixed-mode chromatography for the separation of teicoplanin and the unravelment of its composition. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 186, 113308.	1.4	2
1400	Optimized hidden target screening for very polar molecules in surface waters including a compound database inquiry. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 4953-4966.	1.9	6
1401	Method for evaluation of ionic interactions in liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1625, 461301.	1.8	18
1402	Comparison of various modes and phase systems for analytical HPLC. <i>Handbook of Analytical Separations</i> , 2020, 8, 1-91.	0.8	1
1403	A novel imidazolium bonding stationary phase derived from N-(3-aminopropyl)-imidazole for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1625, 461331.	1.8	16
1404	Multiple heart-cutting mixed-mode chromatography-reversed-phase 2D-liquid chromatography method for separation and mass spectrometric characterization of synthetic oligonucleotides. <i>Journal of Chromatography A</i> , 2020, 1625, 461338.	1.8	30
1405	A click tyrosine zwitterionic stationary phases for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1621, 461045.	1.8	11
1406	HILIC-MS/MS for Linkage-Specific Separation of Sialylated Glycopeptides to Quantify Prostate-Specific Antigen Proteoforms. <i>Journal of Proteome Research</i> , 2020, 19, 2708-2716.	1.8	27
1407	The combination of partition, size exclusion, and hydrodynamic models in chromatography, and application to bonded phases on porous supports. <i>Journal of Chromatography A</i> , 2020, 1620, 461011.	1.8	5
1408	Hyphenations of 2D capillary-based LC with mass spectrometry. , 2020, , 369-412.		1
1409	Enhancing Host-Cell Protein Detection in Protein Therapeutics Using HILIC Enrichment and Proteomic Analysis. <i>Analytical Chemistry</i> , 2020, 92, 10327-10335.	3.2	20
1410	Review of Three-Dimensional Liquid Chromatography Platforms for Bottom-Up Proteomics. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1524.	1.8	50
1411	Progress in Monolithic Column-based Separation and Enrichment of Glycoproteins. <i>Chinese Journal of Analytical Chemistry</i> , 2020, 48, 13-21.	0.9	6
1412	Retention of stevioside polar compounds on a sulfonic acid-functionalized stationary phase. <i>Journal of Chromatography A</i> , 2020, 1620, 460978.	1.8	9
1413	A simple, versatile, and automated pulse-diffusion-focusing strategy for sensitive milliliter-level-injection HILIC-MS/MS analysis of hydrophilic toxins. <i>Journal of Hazardous Materials</i> , 2020, 392, 122318.	6.5	5
1414	Analysis of mobile chemicals in the aquatic environment—current capabilities, limitations and future perspectives. <i>Analytical and Bioanalytical Chemistry</i> , 2020, 412, 4763-4784.	1.9	35
1415	Novel computational approaches to retention modeling in dual hydrophilic interactions/reversed phase chromatography. <i>Journal of Chromatography A</i> , 2020, 1619, 460951.	1.8	11

#	ARTICLE	IF	CITATIONS
1416	Rapid Quantification of Peptide Oxidation Isomers From Complex Mixtures. <i>Analytical Chemistry</i> , 2020, 92, 3834-3843.	3.2	3
1417	Quality by design applied to olanzapine and quetiapine LC-MS/MS bioanalysis. <i>Journal of Chromatographic Science</i> , 2020, 58, 117-126.	0.7	2
1418	Characterization of therapeutic oligonucleotides by liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 182, 113105.	1.4	66
1419	The relationship between polymer structures on silica particles and the separation characteristics of the corresponding columns for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2020, 1618, 460837.	1.8	9
1420	Glucose-based carbon dots-modified silica stationary phase for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2020, 1619, 460930.	1.8	30
1421	Peptide retention time prediction in hydrophilic interaction liquid chromatography: Zwitter-ionic sulfoalkylbetaine and phosphorylcholine stationary phases. <i>Journal of Chromatography A</i> , 2020, 1619, 460909.	1.8	16
1422	Evaluating MISER chromatography as a tool for characterizing HILIC column equilibration. <i>Journal of Chromatography A</i> , 2020, 1619, 460931.	1.8	8
1423	Impurity profiling of l-aspartic acid and glycine using high-performance liquid chromatography coupled with charged aerosol and ultraviolet detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 183, 113149.	1.4	8
1424	Glycoprotein biomarkers and analysis in chronic obstructive pulmonary disease and lung cancer with special focus on serum immunoglobulin G. <i>Clinica Chimica Acta</i> , 2020, 506, 204-213.	0.5	18
1425	A polymer-based zwitterionic stationary phase for hydrophilic interaction chromatography. <i>Talanta</i> , 2020, 216, 120927.	2.9	21
1426	A Monolithic Column with a Sorbent Based on 1-Vinyl-1,2,4-Triazole for Hydrophilic HPLC. <i>Protection of Metals and Physical Chemistry of Surfaces</i> , 2020, 56, 49-53.	0.3	5
1427	Oligonucleotides: Current Trends and Innovative Applications in the Synthesis, Characterization, and Purification. <i>Biotechnology Journal</i> , 2020, 15, e1900226.	1.8	32
1428	Challenges in Analysis of Hydrophilic Metabolites Using Chromatography Coupled with Mass Spectrometry. <i>Journal of Analysis and Testing</i> , 2020, 4, 140-162.	2.5	20
1429	HILIC LC-MS/MS method for the quantification of cefepime, imipenem and meropenem. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 186, 113289.	1.4	26
1430	Strong hydrophilic monolithic column functionalized with amphiphilic benzyl quinine for capillary electrochromatography and application in pharmaceutical analysis. <i>Journal of Chromatography A</i> , 2020, 1621, 461031.	1.8	3
1431	BIOANALYSIS AND BIOTRANSFORMATION OF OLIGONUCLEOTIDE THERAPEUTICS BY LIQUID CHROMATOGRAPHY-MASS SPECTROMETRY. <i>Mass Spectrometry Reviews</i> , 2021, 40, 334-358.	2.8	23
1432	Comparison of determination of sugar-PMP derivatives by two different stationary phases and two HPLC detectors: C18 vs. amide columns and DAD vs. ELSD. <i>Journal of Food Composition and Analysis</i> , 2021, 96, 103715.	1.9	25
1433	Characterization of synthetic dyes for environmental and forensic assessments: A chromatography and mass spectrometry approach. <i>Journal of Separation Science</i> , 2021, 44, 387-402.	1.3	6

#	ARTICLE	IF	CITATIONS
1434	THERAPEUTIC OLIGONUCLEOTIDES, IMPURITIES, DEGRADANTS, AND THEIR CHARACTERIZATION BY MASS SPECTROMETRY. <i>Mass Spectrometry Reviews</i> , 2021, 40, 75-109.	2.8	35
1435	Hydrophilic interaction liquid chromatography-mass spectrometry for the characterization of glycoproteins at the glycan, peptide, subunit, and intact level. , 2021, , 209-278.		2
1436	Chapter 3. Ion Pair Liquid Chromatographyâ€“Mass Spectrometry for Probing the Polar Metabolome. <i>New Developments in Mass Spectrometry</i> , 2021, , 41-68.	0.2	0
1437	Application of Liquid Chromatography for the Analysis of Flavonoids in Food: An Overview. <i>Food Bioactive Ingredients</i> , 2021, , 11-29.	0.3	1
1438	Carboxylate-Modified Magnetic Bead (CMMB)-Based Isopropanol Gradient Peptide Fractionation (CIF) Enables Rapid and Robust Off-Line Peptide Mixture Fractionation in Bottom-Up Proteomics. <i>Molecular and Cellular Proteomics</i> , 2021, 20, 100039.	2.5	10
1439	Mass Spectrometry-Based Methods for Characterization of Hypomodifications in Transfer RNA. <i>RNA Technologies</i> , 2021, , 555-592.	0.2	1
1440	Application of magnetic solid phase extraction in separation and enrichment of glycoproteins and glycopeptides. <i>Chinese Chemical Letters</i> , 2021, 32, 2629-2636.	4.8	42
1441	Pro-apoptotic property of phytochemicals from <i>Naringi crenulata</i> in HER2+ breast cancer cells in vitro. <i>Biotechnology and Biotechnological Equipment</i> , 2021, 35, 311-322.	0.5	1
1442	Polymethacrylate-based monolithic column with incorporated carbamide-modified fumed silica nanoparticles for hydrophilic liquid interaction chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2021, 44, 255-264.	0.5	5
1443	Evaluation of a linear free energy relationship for the determination of the column void volume in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2021, 1638, 461849.	1.8	8
1444	Development of Fast and Selective Analytical Methods of Pharmaceuticals and Herbal Medicines by High-Performance Liquid Chromatography and Capillary Electrophoresis. <i>Chromatography</i> , 2021, 42, 1-16.	0.8	4
1445	Functionalization using polymer or silane? A practical test method to characterize hydrophilic interaction chromatography phases in terms of their functionalization method. <i>Journal of Chromatography A</i> , 2021, 1638, 461850.	1.8	13
1446	Performance evaluation of silica microspheres functionalized by different amine-ligands for hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2021, 1640, 461967.	1.8	8
1447	Facile Preparation of Peptides for Mass Spectrometry Analysis in Bottom-Up Proteomics Workflows. <i>Current Protocols</i> , 2021, 1, e85.	1.3	10
1448	A general new method for calculating the molecular nonpolar surface for analysis of LC-MS data. <i>International Journal of Mass Spectrometry</i> , 2021, 461, 116495.	0.7	3
1449	Fundamental to achieving fast separations with high efficiency: A review of chromatography with superficially porous particles. <i>Biomedical Chromatography</i> , 2021, 35, e5087.	0.8	2
1450	Enzymatic-fingerprinting workflow of polysaccharides in <i>Heridium erinaceus</i> fruiting bodies: From HILIC-ESI ⁺ -MS screening to targeted MIM profiling. <i>International Journal of Biological Macromolecules</i> , 2021, 173, 491-503.	3.6	7
1451	Separation and identification of commercial reactive dyes with hydrophilic interaction liquid chromatography and quadrupole time-of-flight mass spectrometry. <i>Coloration Technology</i> , 2021, 137, 407-417.	0.7	4

#	ARTICLE	IF	CITATIONS
1452	Multiplexed small molecule impurity monitoring in antibody-based therapeutics by mixed-mode chromatography paired with charged aerosol detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 197, 113952.	1.4	4
1453	Polyethyleneimine-functionalized Fe ₃ O ₄ /attapulgite particles for hydrophilic interaction-based magnetic dispersive solid-phase extraction of fluoroquinolones in chicken muscle. <i>Analytical and Bioanalytical Chemistry</i> , 2021, 413, 3529-3540.	1.9	11
1454	Determination of selected water-soluble vitamins (thiamine, riboflavin, nicotinamide and pyridoxine) from a food matrix using hydrophilic interaction liquid chromatography coupled with mass spectroscopy. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2021, 1171, 122541.	1.2	15
1455	Glycosylation Biomarkers Associated with Age-Related Diseases and Current Methods for Glycan Analysis. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5788.	1.8	24
1456	Characterization and Separation of Platinum-Based Antineoplastic Drugs by Zwitterionic Hydrophilic Interaction Liquid Chromatography (HILIC)-Tandem Mass Spectrometry, and Its Application in Surface Wipe Sampling. <i>Separations</i> , 2021, 8, 69.	1.1	3
1457	Counterion Optimization Dramatically Improves Selectivity for Phosphopeptides and Glycopeptides in Electrostatic Repulsion-Hydrophilic Interaction Chromatography. <i>Analytical Chemistry</i> , 2021, 93, 7908-7916.	3.2	18
1458	Influence of different hydrophilic interaction liquid chromatography stationary phases on method performance for the determination of highly polar anionic pesticides in complex feed matrices. <i>Journal of Separation Science</i> , 2021, 44, 2165-2176.	1.3	14
1459	Benzoic acid-modified monolithic column for separation of hydrophilic compounds by capillary electrochromatography with high content of water in mobile phase. <i>Journal of Chromatography A</i> , 2021, 1647, 462166.	1.8	7
1460	Intermolecular Interactions at the Silica-Liquid Interface Modulate the Fermi Resonance Coupling in Surface Methanol. <i>Journal of Physical Chemistry Letters</i> , 2021, 12, 5695-5702.	2.1	7
1461	Dual-Functional Ti(IV)-IMAC Material Enables Simultaneous Enrichment and Separation of Diverse Glycopeptides and Phosphopeptides. <i>Analytical Chemistry</i> , 2021, 93, 8568-8576.	3.2	32
1463	Ionizable Cyclofructan 6-Based Stationary Phases for Hydrophilic Interaction Liquid Chromatography Using Superficially Porous Particles. <i>Chromatographia</i> , 2021, 84, 821-832.	0.7	2
1465	Simple TLC-spectrodensitometric method for studying lipophilicity and quantitative analysis of hypoglycemic drugs in their binary mixture. <i>Biomedical Chromatography</i> , 2021, 35, e5154.	0.8	2
1466	Analytical characterization of DNA and RNA oligonucleotides by hydrophilic interaction liquid chromatography-tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2021, 1648, 462184.	1.8	30
1467	Separation of multiphosphorylated cyclopeptides and their positional isomers by hydrophilic interaction liquid chromatography (HILIC) coupled to electrospray ionization mass spectrometry (ESI-MS). <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2021, 1177, 122792.	1.2	2
1468	PrSM-Level Side-by-Side Comparison of Online LC-MS Methods with Intact Histone H3 and H4 Proteoforms. <i>Journal of Proteome Research</i> , 2021, 20, 4331-4345.	1.8	3
1469	Proteomics for Low Cell Numbers: How to Optimize the Sample Preparation Workflow for Mass Spectrometry Analysis. <i>Journal of Proteome Research</i> , 2021, 20, 4217-4230.	1.8	16
1470	Preparation and evaluation of a polymer-based sulfobetaine zwitterionic stationary phase. <i>Journal of Chromatography A</i> , 2021, 1649, 462229.	1.8	12
1472	A New Definition of the Stationary Phase Volume in Mixed-Mode Chromatographic Columns in Hydrophilic Liquid Chromatography. <i>Molecules</i> , 2021, 26, 4819.	1.7	2

#	ARTICLE	IF	CITATIONS
1473	Separation of labeled isomeric oligosaccharides by hydrophilic interaction liquid chromatography â€” the role of organic solvent in manipulating separation selectivity of the amide stationary phase. <i>Journal of Chromatography A</i> , 2021, 1651, 462303.	1.8	5
1474	Evaluation of multiple hydrophilic interaction chromatography columns and surrogate matrix for arginine quantification in saliva by high-resolution mass spectrometry. <i>Journal of Separation Science</i> , 2021, 44, 3580-3593.	1.3	3
1475	Downstream Processing of Therapeutic Peptides by Means of Preparative Liquid Chromatography. <i>Molecules</i> , 2021, 26, 4688.	1.7	28
1476	Analytical techniques currently used in the pharmaceutical industry for the quality control of RNA-based therapeutics and ongoing developments. <i>Journal of Chromatography A</i> , 2021, 1651, 462283.	1.8	12
1477	Quantitation of Acetyl Hexapeptide-8 in Cosmetics by Hydrophilic Interaction Liquid Chromatography Coupled to Photo Diode Array Detection. <i>Separations</i> , 2021, 8, 125.	1.1	2
1478	Comparison of the steric selectivity on hydrophilic interaction chromatography columns modified with poly(acrylamide) possessing different morphology. <i>Journal of Chromatography A</i> , 2021, 1650, 462207.	1.8	5
1480	What do we know about the ecotoxicological implications of the rare earth element gadolinium in aquatic ecosystems?. <i>Science of the Total Environment</i> , 2021, 781, 146273.	3.9	46
1481	A new high-throughput method based on hydrophilic interaction liquid chromatography-tandem mass spectrometry to determine 18 short-chain carboxylic acids in foods. <i>Food Control</i> , 2021, 126, 108042.	2.8	3
1482	Direct Target Site Identification of a Sulfonamide-Triazole Covalent Kinase Probe by LC-MS Chemical Proteomics. <i>Analytical Chemistry</i> , 2021, 93, 11946-11955.	3.2	10
1483	Advances in mass spectrometry-based glycomicsâ€”An update covering the period 2017â€”2021. <i>Electrophoresis</i> , 2022, 43, 119-142.	1.3	21
1484	Constraints on isomers of dissolved organic matter in aquatic environments: Insights from ion mobility mass spectrometry. <i>Geochimica Et Cosmochimica Acta</i> , 2021, 308, 353-372.	1.6	23
1485	Simultaneous quantitation of inorganic ions in oligonucleotides using mixed-mode liquid chromatography coupled with a charged aerosol detector. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 204, 114244.	1.4	5
1486	Peak shape enhancement using diethylamine in hydrophilic liquid interaction chromatography: Application in simultaneous determination of methionine and paracetamol. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 203, 114214.	1.4	2
1487	Genetically Encoded Green Fluorescent Biosensors for Monitoring UDP-GlcNAc in Live Cells. <i>ACS Central Science</i> , 2021, 7, 1763-1770.	5.3	7
1488	Titania-Based Adsorbents for Hydrophilic Interaction Liquid Chromatography. <i>Journal of Analytical Chemistry</i> , 2021, 76, 1153-1162.	0.4	0
1489	HILIC-MS-based metabolomics reveal that <i>Astragalus</i> polysaccharide alleviates doxorubicin-induced cardiomyopathy by regulating sphingolipid and glycerophospholipid homeostasis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2021, 203, 114177.	1.4	11
1490	Hydrophilic Interaction Liquid Chromatography Coupled with Fluorescence Detection (HILIC-FL) for the Quantitation of Octreotide in Injection Forms. <i>Analyticaâ€”A Journal of Analytical Chemistry and Chemical Analysis</i> , 2021, 2, 121-129.	0.8	3
1491	A systematic investigation of the effect of sample solvent on peak shape in nano- and microflow hydrophilic interaction liquid chromatography columns. <i>Journal of Chromatography A</i> , 2021, 1655, 462498.	1.8	6

#	ARTICLE	IF	CITATIONS
1492	Volume and composition of semi-adsorbed stationary phases in hydrophilic interaction liquid chromatography. Comparison of water adsorption in common stationary phases and eluents. <i>Journal of Chromatography A</i> , 2021, 1656, 462543.	1.8	8
1493	Relative significance of hydrophilic partitioning and surface adsorption to the retention of polar compounds in hydrophilic interaction chromatography. <i>Analytica Chimica Acta</i> , 2021, 1184, 339025.	2.6	3
1494	Preparation and evaluation of a bacitracin-bonded silica stationary phase for hydrophilic interaction liquid chromatography. <i>Microchemical Journal</i> , 2021, 170, 106661.	2.3	4
1495	Surface properties of stationary phases with embedded polar group based on secondary interaction, zeta potential measurement and linear solvation energy relationship studies. <i>Journal of Chromatography A</i> , 2021, 1637, 461853.	1.8	5
1496	Molecular dynamics simulations of amino acid adsorption and transport at the acetonitrile-water-silica interface: the role of side chains. <i>RSC Advances</i> , 2021, 11, 21666-21677.	1.7	0
1497	HPLC- and CE-based methods for the characterization of therapeutic glycoproteins. , 2021, , 761-814.		3
1498	LC-MS/MS in glycomics and glycoproteomics analyses. , 2021, , 391-441.		3
1499	Chapter 4. Hydrophilic Interaction Chromatography-Mass Spectrometry (HILIC-MS) Approaches for Probing the Polar Metabolome. <i>New Developments in Mass Spectrometry</i> , 2021, , 69-90.	0.2	0
1500	Analysis of pancreatic extracellular matrix protein post-translational modifications <i>via</i> electrostatic repulsion-hydrophilic interaction chromatography coupled with mass spectrometry. <i>Molecular Omics</i> , 2021, 17, 652-664.	1.4	7
1501	Free Amino Acid Analysis in Honey Samples by Hydrophilic Interaction Liquid Chromatography with UV Detection Using Precolumn Derivatization with Dansyl Chloride. <i>Chromatographia</i> , 2021, 84, 127-133.	0.7	2
1502	Bioactive peptides from foods: production, function, and application. <i>Food and Function</i> , 2021, 12, 7108-7125.	2.1	49
1503	Novel Adsorbents for Hydrophilic Chromatography Based on Silica Gel Covalently Modified with Poly(ethylene glycol). <i>Moscow University Chemistry Bulletin</i> , 2021, 76, 75-84.	0.2	4
1504	State-of-the-art Mass Spectrometry-based Approaches to Explore the Polar Metabolome: Alzheimer's Disease as a Case Study. <i>New Developments in Mass Spectrometry</i> , 2021, , 219-241.	0.2	1
1507	Liquid Chromatography Methods for Separation of Polar and Charged Intracellular Metabolites for ¹³ C Metabolic Flux Analysis. <i>Methods in Molecular Biology</i> , 2020, 2088, 33-50.	0.4	4
1508	Two Birds with One Stone: Parallel Quantification of Proteome and Phosphoproteome Using iTRAQ. <i>Methods in Molecular Biology</i> , 2016, 1394, 25-41.	0.4	24
1509	Analysis of Hydroxyproline in Collagen Hydrolysates. <i>Methods in Molecular Biology</i> , 2019, 2030, 47-56.	0.4	10
1510	Protein Concentration by Hydrophilic Interaction Chromatography Combined with Solid Phase Extraction. <i>Methods in Molecular Biology</i> , 2008, 424, 63-69.	0.4	3
1511	Electroelution of Proteins from Polyacrylamide Gels. <i>Springer Protocols</i> , 1996, , 207-214.	0.1	5

#	ARTICLE	IF	CITATIONS
1512	Hydrophilic Interaction Chromatography for Fractionation and Enrichment of the Phosphoproteome. <i>Methods in Molecular Biology</i> , 2009, 527, 93-105.	0.4	37
1513	Historical Overview of Glycoanalysis. <i>Methods in Molecular Biology</i> , 2010, 600, 9-30.	0.4	37
1514	Analysis of Hydroxyproline in Collagen Hydrolysates. <i>Methods in Molecular Biology</i> , 2012, 828, 271-280.	0.4	3
1515	Chromatographic Fingerprinting and Chemometric Techniques for Quality Control of Herb Medicines. , 2014, , 133-153.		4
1517	Analysis of Pectin Structure by HPAEC-PAD. <i>Modern Methods of Plant Analysis</i> , 1996, , 129-146.	0.1	6
1518	Reversed-Phase HPLC of Peptides – A Valuable Sample Preparation Tool in Bottom-Up Proteomics: Separation Selectivity in Single and Multi-dimensional Separation Modes. , 2011, , 1031-1050.		2
1519	High Performance Thin-Layer Chromatography. , 2014, , 81-101.		5
1520	Mixed-mode hydrophilic and ionic interaction chromatography rivals reversed-phase chromatography for the separation of peptides. , 1992, , 546-548.		2
1521	Monitoring the hydrophilicity/hydrophobicity of amino acid side-chains in the non-polar and polar faces of amphipathic α -helices by reversed-phase and hydrophilic interaction/cation-exchange chromatography. <i>Journal of Chromatography A</i> , 2004, 1053, 161-172.	1.8	14
1523	Fragment-based Design of Zwitterionic, Strong Cation- and Weak Anion-Exchange Type Mixed-mode Liquid Chromatography Ligands and their Chromatographic Exploration. <i>Journal of Chromatography A</i> , 2020, 1621, 461075.	1.8	16
1524	Recent advances on protein separation and purification methods. <i>Advances in Colloid and Interface Science</i> , 2020, 284, 102254.	7.0	98
1525	Insight into the hydrophilic interaction liquid chromatographic retention behaviors of hydrophilic compounds on different stationary phases. <i>Talanta</i> , 2020, 219, 121363.	2.9	5
1526	Application of Carbonaceous Materials in Separation Science. <i>RSC Green Chemistry</i> , 2015, , 103-126.	0.0	3
1527	The Application of Thiol-ene/Yne Radical Click Chemistry in Surface Modification and Functionalization. <i>RSC Polymer Chemistry Series</i> , 2013, , 286-308.	0.1	4
1528	Determination of Gemcitabine in Plasma of Bladder Cancer Patients by Hydrophilic Interaction Chromatography with Ultraviolet Detection. <i>Journal of Chromatographic Science</i> , 2020, 58, 587-590.	0.7	3
1530	Influence of nonideal mixing properties on viscous fingering in micropillar array columns. <i>Physical Review Fluids</i> , 2016, 1, .	1.0	5
1531	Separation of Hydrophilic Oligomers and Polymers Using Monodisperse Poly(2,3-dihydroxypropyl) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Collection of Czechoslovak Chemical Communications, 2001, 66, 1047-1061.	1.0	8
1532	Organic polymer monolithic columns with incorporated bare and cyano-modified fumed silica nanoparticles for use in hydrophilic interaction liquid chromatography. <i>Journal of Analytical Science and Technology</i> , 2020, 11, .	1.0	3

#	ARTICLE	IF	CITATIONS
1533	Reducing Protein Sample Complexity with Free Flow Electrophoresis (FFE). , 2005, , 187-204.		1
1536	HILIC Retention Behavior and Method Development for Highly Polar Basic Compounds Used in Pharmaceutical Synthesis. <i>Chromatographic Science</i> , 2011, , 345-372.	0.1	2
1537	Hydrophilic Interaction Liquid Chromatography-Based Enrichment Protocol Coupled to Mass Spectrometry for Glycoproteome Analysis. <i>Chromatographic Science</i> , 2011, , 469-490.	0.1	1
1538	Development and Application of Methods for Separation of Carbohydrates by Hydrophilic Interaction Liquid Chromatography. <i>Chromatographic Science</i> , 2011, , 491-522.	0.1	4
1539	Hydrophilic Interaction Liquid Chromatography in the Characterization of Glycoproteins. <i>Chromatographic Science</i> , 2011, , 523-550.	0.1	3
1540	Analysis of Protein Glycosylation and Phosphorylation Using HILIC-MS. <i>Chromatographic Science</i> , 2011, , 551-576.	0.1	5
1541	Hydrophilic Interaction Liquid Chromatography-Mass Spectrometry (HILIC-MS) of Paralytic Shellfish Poisoning Toxins, Domoic Acid, and Assorted Cyanobacterial Toxins. <i>Chromatographic Science</i> , 2011, , 105-132.	0.1	1
1542	Thin-layer chromatography in the analysis of biologically active ionic and ionizable compounds. <i>Journal of Planar Chromatography - Modern TLC</i> , 2015, 28, 413-425.	0.6	3
1543	Metals from cell to environment: Connecting Metallomics with other omics. <i>Open Journal of Plant Science</i> , 0, , 001-014.	0.2	3
1545	Investigation on Selectivity of Mercaptopropylsilica and Silver Nanoparticles- Functionalised Mercaptopropylsilica. <i>Current Chromatography</i> , 2015, 2, 122-135.	0.1	5
1550	An Investigation of Vanillin Imposed Oxidative Stress in Corn (<i>Zea mays</i> L.) and the Activities of Antioxidative Enzymes. <i>Pakistan Journal of Biological Sciences</i> , 2005, 9, 34-38.	0.2	2
1551	Hydrophilic Interaction Liquid Chromatography for LC-MS. <i>Mass Spectrometry & Purification Techniques</i> , 2018, 04, .	0.2	1
1552	Determination of Free Amino Acids in <i>Isatis Radix</i> By HILIC-UPLC-MS/MS. <i>Bulletin of the Korean Chemical Society</i> , 2014, 35, 197-203.	1.0	19
1553	Development of Isotope Dilution LC-MS/MS Method for Accurate Determination of Arsenobetaine in Oyster Certified Reference Material. <i>Bulletin of the Korean Chemical Society</i> , 2014, 35, 821-827.	1.0	6
1554	Hypoglycemia - Causes and Occurrences. , 2011, , .		2
1555	Retention dependence on temperature in reversed-phase and hydrophilic interaction liquid chromatography. <i>Biologija (Vilnius, Lithuania)</i> , 2011, 57, .	0.3	10
1556	The isolation of water-soluble natural products - challenges, strategies and perspectives. <i>Natural Product Reports</i> , 2022, 39, 596-669.	5.2	18
1557	Analytical Platforms for Mass Spectrometry-Based Metabolomics of Polar and Ionizable Metabolites. <i>Advances in Experimental Medicine and Biology</i> , 2021, 1336, 215-242.	0.8	4

#	ARTICLE	IF	CITATIONS
1558	Mass Spectrometry-Based Methods for Immunoglobulin G N-Glycosylation Analysis. <i>Experientia Supplementum</i> (2012), 2021, 112, 73-135.	0.5	0
1559	PDA-CAD method for the determination of magnesium, pyridoxine and thiamine in a dietary supplement supported by analytical quality by design methodology. <i>Arhiv Za Farmaciju</i> , 2021, 71, 378-392.	0.2	1
1560	Hydrophilic interaction-based magnetically assisted matrix solid-phase dispersion extraction of carbadox and olaquinox in feeds. <i>Journal of the Science of Food and Agriculture</i> , 2021, , .	1.7	1
1561	Empathy evaluation by the physical elements of the advertising. <i>Multimedia Tools and Applications</i> , 2022, 81, 2241-2257.	2.6	3
1562	State-of-the-art glycosaminoglycan characterization. <i>Mass Spectrometry Reviews</i> , 2022, 41, 1040-1071.	2.8	16
1563	Grafting copolymer brushes on polyhedral oligomeric silsesquioxanes silsesquioxane-decorated silica stationary phase for hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2021, 1659, 462627.	1.8	9
1564	Detection of the Prion Protein in Blood Using a Fluorescence Immunoassay. , 2003, , .		0
1565	Use of Reversed-Phase Liquid Chromatography in Biopharmaceutical Development. , 2005, , 27-65.		0
1566	Determination of Endogenous GA3 Levels in Different Parts of Artichoke (<i>Cynara scolymus</i> L. cv. Sakiz) Plants in Various Developmental Stages. <i>Pakistan Journal of Biological Sciences</i> , 2005, 9, 30-33.	0.2	0
1567	N-Linked Global Glycan Profiling by NanoLC Mass Spectrometry. <i>Methods in Molecular Biology</i> , 2011, 790, 87-97.	0.4	1
1568	A Multidimensional Chromatography Strategy Using HILIC and IMAC for Quantitative Phosphoproteome Analysis. , 2011, , 487-495.		0
1569	Retention of Polar Acidic and Basic Drugs by HILIC. <i>Chromatographic Science</i> , 2011, , 373-400.	0.1	0
1570	Hydrophilic Interaction Chromatography-Mass Spectrometry. <i>Chromatographic Science</i> , 2011, , 157-176.	0.1	0
1571	Dimethylarginine Analysis by HILIC-MS/MS and Its Applications in Clinical Chemistry and In Vivo Animal Models. <i>Chromatographic Science</i> , 2011, , 203-224.	0.1	0
1572	Fast In-Process Method for the Determination of loversol and Related Polar Compounds by Hydrophilic Interaction Chromatography (HILIC) and UPLC. <i>Chromatographic Science</i> , 2011, , 291-308.	0.1	0
1573	Retention Behavior of Hydrazine Derivatives in HILIC Mode. <i>Chromatographic Science</i> , 2011, , 309-344.	0.1	0
1574	Mechanisms Controlling the Separation and Retention of Proanthocyanidins and Other Plant Polyphenols in Hydrophilic Interaction Chromatography. <i>Chromatographic Science</i> , 2011, , 77-88.	0.1	0
1575	HILIC Behavior of Reversed-Phase/Ion-Exchange Mixed-Mode Stationary Phases and Their Applications. <i>Chromatographic Science</i> , 2011, , 47-76.	0.1	1

#	ARTICLE	IF	CITATIONS
1576	Application of the Calculated Physicochemical Property, Log D, to Hydrophilic Interaction Chromatography in the Bioanalytical Laboratory. <i>Chromatographic Science</i> , 2011, , 27-46.	0.1	0
1577	Clinical Applications of Hydrophilic Interaction Liquid Chromatography. <i>Chromatographic Science</i> , 2011, , 177-202.	0.1	0
1579	Recent Developments and Applications in Nonlinear Reversed Phase Liquid Chromatography. <i>Advances in Chromatography</i> , 2012, 50, 415-440.	1.0	0
1581	Separation Methods. <i>Chromatographic Science</i> , 2013, , 69-90.	0.1	0
1582	Paralytic Shellfish Poisoning (PSP) Analysis using Liquid Chromatography-Tandem Mass Spectrometry. <i>Han'guk Susan Hakhoe Chi = Bulletin of the Korean Fisheries Society</i> , 2013, 46, 154-159.	0.1	0
1583	New Developments in Targeted Analysis of Protein Posttranslational Modifications. <i>Current Proteomics</i> , 2013, 10, 98-119.	0.1	0
1584	Metabolomics. <i>Springer Protocols</i> , 2014, , 425-451.	0.1	0
1585	Next Generation Proteomics: PTMs in Space and Time. <i>New Developments in Mass Spectrometry</i> , 2014, , 233-256.	0.2	0
1586	Quantitative Proteomics Analysis of Differentially Expressed Proteins Involved in Renal Clear Cell Carcinoma by Shotgun Approach Coupled with Stable Isotope Dimethyl Labeling. <i>Journal of Proteomics and Bioinformatics</i> , 2014, s7, .	0.4	1
1587	Shotgun Proteomics Analysis of Differentially Expressed Urinary Proteins Involved in the Hepatocellular Carcinoma. <i>Journal of Proteomics and Bioinformatics</i> , 2014, 07, .	0.4	0
1589	The Use of Electrostatic Repulsion-Hydrophilic Interaction Chromatography (ERLIC) for Proteomics Research. <i>Mass Spectrometry Letters</i> , 2014, 5, 95-103.	0.5	1
1590	Optimization of Normal-Phase and Reversed-Phase Systems for Analysis of Pesticides: Choice of the Mode of Elution—Isocratic and Gradient Elution. <i>Chromatographic Science</i> , 2015, , 215-242.	0.1	0
1593	SÄubern und Putzen. , 2016, , 151-197.		0
1594	Comparative study of different column types for the separation of polar basic hallucinogenic alkaloids. <i>South African Journal of Chemistry</i> , 2016, 69, .	0.3	0
1595	Liquid Chromatography-Mass Spectrometry of Glycerophospholipids. , 2016, , 1-9.		0
1596	Metal reduction at bulk chemical filtration. , 2017, , .		0
1597	Application of Novel Materials in Sample Treatment and Separation. , 2017, , 199-220.		0
1598	Open Journal of Plant Science. <i>Open Journal of Plant Science</i> , 0, , .	0.2	0

#	ARTICLE	IF	CITATIONS
1599	Exploring the Effect of Buffer Strength on the Retention Time of Weak Acids, Neutral and Weak Bases in Hydrophilic Interaction Liquid Chromatography (HILIC) Mode. <i>Current Analytical Chemistry</i> , 2018, 15, 34-46.	0.6	0
1601	A HILIC stationary phase functionalized with glutathione by thiol-ene chemistry on monodisperse-porous polymer microparticles. <i>Hacettepe Journal of Biology and Chemistry</i> , 0, , .	0.3	0
1602	Mobile Phase Effects in Reversed-Phase and Hydrophilic Interaction Liquid Chromatography. , 2020, , 1-58.		0
1603	Rapid polymerization of polyhedral oligomeric siloxane-based zwitterionic sulfoalkylbetaine monolithic column in ionic liquid for hydrophilic interaction capillary electrochromatography. <i>Journal of Chromatography A</i> , 2021, 1659, 462651.	1.8	6
1604	Mass Spectrometry-Based Technology and Workflows for Studying the Chemistry of Fungal Endophyte Derived Bioactive Compounds. <i>ACS Chemical Biology</i> , 2021, 16, 2068-2086.	1.6	16
1605	Fundamentals of Mass Spectrometry-Based Metabolomics. <i>NATO Science for Peace and Security Series A: Chemistry and Biology</i> , 2020, , 61-81.	0.5	0
1606	Polar polymer-immobilized stationary phase for aqueous reversed-phase liquid chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2021, 44, 25-32.	0.5	1
1607	Mass spectrometry-based proteomics and peptidomics for systems biology and biomarker discovery. <i>Frontiers in Biology</i> , 0, , .	0.7	0
1609	Protein import into yeast mitochondria: the inner membrane import site protein ISP45 is the MPI1 gene product. <i>EMBO Journal</i> , 1993, 12, 3035-41.	3.5	28
1610	Technique development of high-throughput and high-sensitivity sample preparation and separation for proteomics. <i>Bioanalysis</i> , 2022, 14, 101-111.	0.6	1
1611	Hydrophilic interaction chromatography " mass spectrometry for metabolomics and proteomics: state-of-the-art and current trends. <i>Microchemical Journal</i> , 2022, 175, 106986.	2.3	16
1612	Liquid Chromatography in Bioanalysis. , 2022, , 377-403.		0
1613	Coupling Mixed Mode Chromatography/ESI Negative MS Detection with Message-Passing Neural Network Modeling for Enhanced Metabolome Coverage and Structural Identification. <i>Metabolites</i> , 2021, 11, 772.	1.3	2
1614	Paralytic shellfish toxins (PSTs) and tetrodotoxin (TTX) of Korean pufferfish. <i>Fisheries and Aquatic Sciences</i> , 2021, 24, 360-369.	0.3	5
1615	Graph Convolutional Networks for Improved Prediction and Interpretability of Chromatographic Retention Data. <i>Analytical Chemistry</i> , 2021, 93, 15633-15641.	3.2	18
1616	A survey of polar stationary phases for hydrophilic interaction chromatography and recent progress in understanding retention and selectivity. <i>Biomedical Chromatography</i> , 2022, 36, e5332.	0.8	13
1617	Investigate Retention Behavior of 2-deoxycytidine in Hydrophilic Interaction Liquid Chromatography. <i>International Journal of Drug Delivery Technology</i> , 2020, 10, 349-353.	0.0	3
1618	Expanding Antineoplastic Drugs Surface Monitoring Profiles: Enhancing of Zwitterionic Hydrophilic Interaction Methods. <i>Separations</i> , 2022, 9, 34.	1.1	1

#	ARTICLE	IF	CITATIONS
1619	Characterization of a highly stable zwitterionic hydrophilic interaction chromatography stationary phase based on hybrid organic–inorganic particles. <i>Journal of Separation Science</i> , 2022, 45, 1389-1399.	1.3	14
1620	High performance liquid chromatography: Separations. , 2022, , 283-336.		0
1621	Multicomponent Ugi reaction as a tool for fast and easy preparation of stationary phases for hydrophilic interaction liquid chromatography. Part I: The influence of attachment and spacing of the functional ligand obtained via the Ugi reaction. <i>Journal of Chromatography A</i> , 2022, 1666, 462804.	1.8	12
1622	A simple generic method for analyzing water sensitive pinacol boronate compounds by hydrophilic interaction liquid chromatography. <i>Journal of Chromatography Open</i> , 2022, 2, 100036.	0.8	1
1623	Recent Strategies for Using Monolithic Materials in Glycoprotein and Glycopeptide Analysis. <i>Separations</i> , 2022, 9, 44.	1.1	6
1624	Enhancement of protein crystallization with the application of Taylor vortex and Poly(ionic liquid)s. <i>Chemical Engineering Science</i> , 2022, 252, 117501.	1.9	10
1625	Preparation, Characterization and Chromatographic Evaluation of Nickel Oxide Deposited Silica Stationary Phase in Hydrophilic Interaction Liquid Chromatography. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
1627	Cell culture metabolomics and lipidomics. , 2022, , 415-456.		0
1628	Application of a hybrid zwitterionic hydrophilic interaction liquid chromatography column in metabolic profiling studies. <i>Journal of Chromatography A</i> , 2022, 1672, 463013.	1.8	8
1629	Robust co-immunoprecipitation with mass spectrometry for <i>Caenorhabditis elegans</i> using solid-phase enhanced sample preparation. <i>BioTechniques</i> , 2022, 72, 175-184.	0.8	2
1630	Contribution of ionic interactions to stationary phase selectivity in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2022, 45, 3264-3275.	1.3	11
1631	Preparation and evaluation of a double-hydrophilic interaction stationary phase based on bovine serum albumin and graphene quantum dots modified silica. <i>Journal of Chromatography A</i> , 2022, 1669, 462933.	1.8	7
1632	The multifaceted roles of mass spectrometric analysis in nucleic acids drug discovery and development. <i>Mass Spectrometry Reviews</i> , 2023, 42, 1332-1357.	2.8	4
1633	Metabolomics Work Flow and Analytics in Systems Biology. <i>Current Molecular Medicine</i> , 2022, 22, 870-881.	0.6	6
1634	Ion-pair Reversed-phase Liquid Chromatographic Separation of Oligonucleotides. <i>Analytical Sciences</i> , 2021, 37, 1811-1814.	0.8	1
1636	Characterizing the N- and O-linked glycans of the PGF-CTERM sorting domain-containing S-layer protein of <i>Methanoculleus marisnigri</i> . <i>Glycobiology</i> , 2022, 32, 629-644.	1.3	4
1642	Diet- and microbiota-related metabolite, 5-aminovaleric acid betaine (5-AVAB), in health and disease. <i>Trends in Endocrinology and Metabolism</i> , 2022, 33, 463-480.	3.1	19
1643	A Spin-Tip Enrichment Strategy for Simultaneous Analysis of N-Glycopeptides and Phosphopeptides from Human Pancreatic Tissues. <i>Journal of Visualized Experiments</i> , 2022, , .	0.2	1

#	ARTICLE	IF	CITATIONS
1644	Solute-Solvent Interactions in Hydrophilic Interaction Liquid Chromatography: Characterization of the Retention in a Silica Column by the Abraham Linear Free Energy Relationship Model. <i>Journal of Solution Chemistry</i> , 2022, 51, 1081-1100.	0.6	6
1645	Natural Products™ Extraction and Isolation-Between Conventional and Modern Techniques. , 2022, 1, .		10
1646	Functionalization of graphene composites using ionic liquids and applications. , 2022, , 445-461.		1
1647	Quality Markers of <i>Dendrobium officinale</i> by Oligosaccharide-Spectrum-Effect Relationships. <i>Frontiers in Nutrition</i> , 0, 9, .	1.6	3
1648	Structure and function of non-digestible carbohydrates in the gut microbiome. <i>Beneficial Microbes</i> , 2022, 13, 95-168.	1.0	26
1649	Evaluation of the Base Stability of Hydrophilic Interaction Chromatography Columns Packed with Silica or Ethylene-Bridged Hybrid Particles. <i>Separations</i> , 2022, 9, 146.	1.1	3
1650	Preparation, Characterization and Chromatographic Evaluation of Nickel Oxide Deposited Silica Stationary Phase in Hydrophilic Interaction Liquid Chromatography. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
1651	Medical Bioanalytics: Separation Techniques in Medical Diagnostics of Neurological Diseases and Disorders on Selected Examples. , 2022, , 49-66.		0
1652	Hydrophilic interaction liquid chromatography. , 2022, , 447-477.		0
1653	Advances in glycopeptide enrichment methods for the analysis of protein glycosylation over the past decade. <i>Journal of Separation Science</i> , 2022, 45, 3169-3186.	1.3	18
1654	Ultrasonic Processing of Food Waste to Generate Value-Added Products. <i>Foods</i> , 2022, 11, 2035.	1.9	13
1655	Studies on the Effects of Process Conditions on Separation of B1, B2 and B3 Vitamin Mixture Using HILIC and RPLC Chromatography. <i>AgriEngineering</i> , 2022, 4, 566-591.	1.7	1
1656	The impact of low adsorption surfaces for the analysis of DNA and RNA oligonucleotides. <i>Journal of Chromatography A</i> , 2022, 1677, 463324.	1.8	15
1658	New mode of liquid chromatography separation: A new mode of separation using multi-charged ions in the mobile phase for high-performance liquid chromatography. <i>Journal of Separation Science</i> , 2022, 45, 3520-3528.	1.3	2
1659	Insights on the Extraction and Analysis of Phenolic Compounds from Citrus Fruits: Green Perspectives and Current Status. <i>Critical Reviews in Analytical Chemistry</i> , 0, , 1-27.	1.8	6
1660	Separation of nicotinamide metabolites using a PBr column packed with pentabromobenzyl group modified silica gel. <i>Analytical Biochemistry</i> , 2022, 655, 114837.	1.1	3
1661	Quantification of Arbutin in Cosmetics, Drugs and Food Supplements by Hydrophilic-Interaction Chromatography. <i>Molecules</i> , 2022, 27, 5673.	1.7	4
1662	Evaluation and Correction of Injection Order Effects in Lc-MS/Ms Based Targeted Metabolomics. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0

#	ARTICLE	IF	CITATIONS
1663	Macrocyclic Glycopeptide Chiral Stationary Phases. , 2022, , .		1
1664	Adsorbents with a Covalently Bonded Polymer Layer for Hydrophilic Interaction Liquid Chromatography. <i>Journal of Analytical Chemistry</i> , 2022, 77, 1173-1183.	0.4	3
1665	Preparation of thioglycerol-modified silica through thiol-epoxy click reaction and its application in HILIC for detection of oligosaccharide in beverages. <i>Food Chemistry</i> , 2023, 402, 134486.	4.2	2
1666	On the occurrence of very low intra-particle diffusion rates in zwitterionic hydrophilic interaction liquid chromatography polymer columns. <i>Journal of Chromatography A</i> , 2022, 1683, 463531.	1.8	5
1667	Recent development in hydrophilic interaction liquid chromatography stationary materials for glycopeptide analysis. <i>Analytical Methods</i> , 2022, 14, 4437-4448.	1.3	8
1668	Comparison of solid-phase extraction methods for efficient purification of phosphopeptides with low sample amounts. <i>Journal of Chromatography A</i> , 2022, 1685, 463597.	1.8	5
1669	QuEChERS-based extraction and two-dimensional liquid chromatography-high resolution mass spectrometry for the determination of long chain chlorinated paraffins in sediments. <i>Journal of Chromatography A</i> , 2022, 1684, 463585.	1.8	2
1670	Evaluation and correction of injection order effects in LC-MS/MS based targeted metabolomics. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2022, 1212, 123513.	1.2	2
1671	Dynamic acylome reveals metabolite driven modifications in <i>Syntrophomonas wolfei</i> . <i>Frontiers in Microbiology</i> , 0, 13, .	1.5	1
1672	Effect of spacer alkyl chain length on retention among three imidazolium stationary phases under various modes in high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2022, 1685, 463646.	1.8	5
1673	Hydrophilic-Phase Extraction: A New Avenue of Solid-Phase Extraction for Oligonucleotide Bioanalysis. <i>Lc-gc Europe</i> , 2022, , 46-51.	0.2	0
1674	Multidimensional separation and analysis of alpha-1-acid glycoprotein N-glycopeptides using high-field asymmetric waveform ion mobility spectrometry (FAIMS) and nano-liquid chromatography tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2023, 415, 379-390.	1.9	3
1675	Covalent organic framework-based monolithic column with hydrophilic and π - π stacking interaction for efficient in-tube solid-phase microextraction of synthetic phenolic antioxidants. <i>Microchemical Journal</i> , 2023, 186, 108311.	2.3	4
1676	Pursuing the Polar Routes with Penguins and Polar Bears: Balancing the Choice of Chromatographic Mode. <i>LC-GC North America</i> , 2022, , 20-22.	0.1	0
1677	PEG-modified halloysite as a hydrophilic interaction and cation exchange mixed-mode sorbent for solid-phase extraction of biogenic amines in fish samples. <i>Analytical and Bioanalytical Chemistry</i> , 0, , .	1.9	1
1678	Unified-Hydrophilic-Interaction/Anion-Exchange Liquid Chromatography Mass Spectrometry (Unified-HILIC/AEX/MS): A Single-Run Method for Comprehensive and Simultaneous Analysis of Polar Metabolome. <i>Analytical Chemistry</i> , 2022, 94, 16877-16886.	3.2	9
1679	Stable isotope tracing reveals compartmentalized nitrogen assimilation in scleractinian corals. <i>Frontiers in Marine Science</i> , 0, 9, .	1.2	2
1681	Immobilized Artificial Membrane Chromatography Using Acetonitrile-Rich Mobile Phase for Comparison of Retention Properties Between Phospholipidosis-Inducing and Non-inducing Basic Drugs. <i>Chromatographia</i> , 0, , .	0.7	0

#	ARTICLE	IF	CITATIONS
1682	A polyethyleneimine-functionalized polymer substrate polar stationary phase. <i>Journal of Chromatography A</i> , 2023, 1689, 463711.	1.8	3
1683	Defining the altered glycoproteomic space of the early secretory pathway by class I mannosidase pharmacological inhibition. <i>Frontiers in Molecular Biosciences</i> , 0, 9, .	1.6	1
1684	Preparation, Characterization, and Chromatographic Evaluation of Nickel Oxide Deposited Silica Stationary Phase in Hydrophilic Interaction Liquid Chromatography. <i>Chromatographia</i> , 2023, 86, 125-134.	0.7	1
1685	A Compendium of the Principal Stationary Phases Used in Hydrophilic Interaction Chromatography: Where Have We Arrived?. <i>Separations</i> , 2023, 10, 22.	1.1	4
1686	Dimethyl Carbonate as a Mobile-Phase Modifier for Normal-Phase and Hydrophilic Interaction Liquid Chromatography. <i>Separations</i> , 2023, 10, 70.	1.1	5
1687	Separation in <i>Analytical Chemistry</i> , , 2023, , 154-189.		0
1688	Hydrophilic Interaction Chromatography. <i>Separations</i> , 2023, 10, 120.	1.1	0
1689	Comparison of chromatographic conditions for the targeted tandem mass spectrometric determination of 354 mammalian metabolites. <i>Journal of Chromatography A</i> , 2023, 1697, 463985.	1.8	1
1690	Separation of therapeutic oligonucleotides using ion-pair reversed-phase chromatography based on fundamental separation science. <i>Journal of Chromatography Open</i> , 2023, 3, 100079.	0.8	9
1691	Clinical Metabolomics: Expanding the Metabolome Coverage Using Advanced Analytical Techniques. <i>Lc-gc Europe</i> , 2019, , 465-480.	0.2	1
1692	Silica microparticles modified with ionic liquid bonded chitosan as hydrophilic moieties for preparation of high-performance liquid chromatographic stationary phases. <i>Mikrochimica Acta</i> , 2023, 190, .	2.5	7
1693	Microflow liquid chromatography “ multi-emitter nanoelectrospray mass spectrometry of oligonucleotides. <i>Journal of Chromatography A</i> , 2023, 1696, 463976.	1.8	2
1694	Retention and mass transfer properties of the series of unbonded, amide-bonded, and alkylsulfobetaine-bonded ethylene bridged hybrid hydrophilic interaction liquid chromatography columns. <i>Journal of Chromatography A</i> , 2023, 1692, 463828.	1.8	3
1695	Preparation of poly(2-vinylpyrrolidone-co-pentaerythritol triacrylate) monolithic column for hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , 2023, 46, .	1.3	1
1696	Managing nonspecific adsorption to liquid chromatography hardware: A review. <i>Analytica Chimica Acta</i> , 2023, 1250, 340994.	2.6	5
1697	Zwitterionic polymer-terminated porous silica stationary phases for highly selective separation in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2023, 1693, 463885.	1.8	2
1698	Determination of 5-methyldeoxycytosine and oxidized derivatives by nano-liquid chromatography with zwitterionic monolithic capillary column. <i>Journal of Chromatography A</i> , 2023, 1693, 463895.	1.8	1
1699	Clean and green analytical techniques. , 2023, , 3-45.		0

#	ARTICLE	IF	CITATIONS
1700	Further Evaluation of the Base Stability of Hydrophilic Interaction Chromatography Columns Packed with Silica or Ethylene-Bridged Hybrid Particles. <i>Separations</i> , 2023, 10, 175.	1.1	1
1701	ZIF-8 Assisted Polyacrylamide Functionalized Silica Core-Shell Stationary Phase for Hydrophilic Interaction Liquid Chromatography. <i>Separations</i> , 2023, 10, 219.	1.1	0
1702	In-Depth Performance Analysis and Comparison of Monolithic and Particulate Zwitterionic Hydrophilic Interaction Liquid Chromatography Polymer Columns. <i>Molecules</i> , 2023, 28, 2902.	1.7	2
1703	Design of β -Alanine-Derived Novel C18 Stationary Phase Containing Embedded Urea and Amide Groups for the Separation of Versatile Analytes. <i>Analytical Chemistry</i> , 2023, 95, 6172-6181.	3.2	5
1704	Simultaneous determination of five lignans from <i>Schisandra chinensis</i> by matrix solid-phase dispersion extraction-high performance liquid chromatography. <i>Chinese Journal of Chromatography (Se Pu)</i> , 2023, 41, 257-264.	0.1	0
1705	HPLC/HILIC determination of biogenic amines in wines produced by different winemaking technologies. <i>Acta Alimentaria</i> , 2023, 52, 209-219.	0.3	1
1706	Green and Scalable Fractionation of Gold Nanoclusters by Anion Exchange Chromatography: Proof of Principle and Scale-Up. <i>ACS Applied Nano Materials</i> , 2023, 6, 6953-6962.	2.4	3
1711	Flavor perception and health benefits of tea. <i>Advances in Food and Nutrition Research</i> , 2023, , .	1.5	0
1714	Analysis of oligonucleotides by liquid chromatography. , 2023, , 357-380.		1
1715	Milestones in the development of liquid chromatography. , 2023, , 1-15.		0
1716	Basic principles for the selection of liquid chromatographic modes for specific applications. , 2023, , 81-157.		1
1717	Hydrophilic interaction liquid chromatography. , 2023, , 227-249.		1
1740	Introduction of Metabolomics: An Overview. , 2023, , 1-37.		0
1741	Analysis of water pollutants. , 2024, , 131-165.		0
1755	Biological big-data sources, problems of storage, computational issues, and applications: a comprehensive review. <i>Knowledge and Information Systems</i> , 0, , .	2.1	0