

CITATION REPORT

List of articles citing

A geriatric-anesthesiologic program to reduce acute confusional states in elderly patients treated for femoral neck fractures

DOI: 10.1111/j.1532-5415.1991.tb03618.x

Journal of the American Geriatrics Society, 1991, 39, 655-62.

Source: <https://exaly.com/paper-pdf/22284620/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
187	Underdiagnosis and poor documentation of acute confusional states in elderly hip fracture patients. <i>Journal of the American Geriatrics Society</i> , 1991 , 39, 760-5	5.6	130
186	Delirium in older patients. <i>Journal of the American Geriatrics Society</i> , 1992 , 40, 829-38	5.6	129
185	Preventing postoperative medical complications. How primary care physicians can help. 1992 , 92, 189-96, 202		1
184	The ageing process. 1993 , 7, 17-37		1
183	Delirium in the older person with cancer. 1994 , 74, 2079-91		50
182	Management of trauma in the elderly patient. 1994 , 74, 163-86		53
181	Postoperative delirium in the elderly. <i>Anesthesia and Analgesia</i> , 1995 , 80, 1223-32	3.9	67
180	Postoperative Delirium in the Elderly. <i>Anesthesia and Analgesia</i> , 1995 , 80, 1223-1232	3.9	191
179	A simple model for predicting postoperative delirium in older patients undergoing elective orthopedic surgery. <i>Journal of the American Geriatrics Society</i> , 1995 , 43, 175-8	5.6	92
178	Psychosocial Aspects of Osteoporosis. 1995 , 6, 441-453		9
177	Delirium. 1997 , 917-922		
176	The appearance and disappearance of cognitive impairment in elderly patients during treatment for hip fracture. <i>Scandinavian Journal of Caring Sciences</i> , 1997 , 11, 167-75	2.3	49
175	Outcomes of delirium: can systems of care make a difference?. <i>Journal of the American Geriatrics Society</i> , 1997 , 45, 247-8	5.6	11
174	Cerebral dysfunction after anaesthesia. <i>Acta Anaesthesiologica Scandinavica</i> , 1997 , 110, 13-6	1.9	13
173	Management of acute delirium in hospitalized elderly: a process improvement project. 1997 , 18, 150-4		20
172	Enhanced isoflurane suppression of excitatory synaptic transmission in the aged rat hippocampus. 1998 , 124, 1075-82		11
171	Postoperative variation in neurocognitive and functional status in elderly hip fracture patients. 1998 , 27, 59-67		49

170	Frequently missed diagnosis in geriatric psychiatry. 1998 , 21, 941-71, viii		14
169	Delirium in the elderly. Optimal management. 1998 , 13, 119-30		32
168	Delirium: prevention, treatment, and outcome studies. 1998 , 11, 126-37; discussion 157-8		51
167	Prevalence of dementia, delirium and psychiatric symptoms in various care settings for the elderly. 1998 , 26, 56-62		56
166	Attention, frailty, and falls: the effect of a manual task on basic mobility. <i>Journal of the American Geriatrics Society</i> , 1998 , 46, 758-61	5.6	174
165	The Influence of an Aging Surgical Population on the Anesthesia Workload. <i>Anesthesia and Analgesia</i> , 1998 , 86, 1165-1170	3.9	
164	The influence of an aging surgical population on the anesthesia workload: a ten-year survey. <i>Anesthesia and Analgesia</i> , 1998 , 86, 1165-70	3.9	57
163	Patient's age is single best predictor for success after hemiarthroplasty of the hip joint. 1999 , 9, 9-11		
162	GERIATRICS AND ANESTHESIA. 1999 , 17, 453-465		10
161	Intensive and high-dependency care for the high-risk surgical patient. 1999 , 13, 427-436		1
160	Postoperative mental impairment in hip fracture patients. A randomized study of reorientation measures in 223 patients. 1999 , 70, 250-5		16
159	Cognitive decline after major noncardiac operations: a preliminary prospective study. 1999 , 68, 1786-91		34
158	A multicomponent intervention to prevent delirium in hospitalized older patients. 1999 , 340, 669-76		2010
157	Reorganization of Nursing and Medical Care to Reduce the Incidence of Postoperative Delirium and Improve Rehabilitation Outcome in Elderly Patients Treated for Femoral Neck Fractures. <i>Scandinavian Journal of Caring Sciences</i> , 1999 , 13, 193-200	2.3	32
156	Continuity, Assessment and Feedback in Orthopaedic Nursing Care Practice is Cost-effective. <i>Scandinavian Journal of Caring Sciences</i> , 1999 , 13, 177-185	2.3	2
155	Geriatric-based versus general wards for older acute medical patients: a randomized comparison of outcomes and use of resources. <i>Journal of the American Geriatrics Society</i> , 2000 , 48, 1381-8	5.6	110
154	Delirium is independently associated with poor functional recovery after hip fracture. <i>Journal of the American Geriatrics Society</i> , 2000 , 48, 618-24	5.6	460
153	Combined lumbar and sacral plexus block compared with plain bupivacaine spinal anesthesia for hip fractures in the elderly. 2000 , 25, 158-162		66

152	The effectiveness of old age psychiatry services. <i>International Journal of Geriatric Psychiatry</i> , 2000 , 15, 687-703	3.9	68
151	Central nervous system dysfunction after anesthesia in the geriatric patient. 2000 , 18, 59-70, vi		36
150	Perioperative temporal profile of cognitive function in elderly patients undergoing hip surgery. 2000 , 13, 206-9		6
149	Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: subgroup analysis of patients with dementia. <i>BMJ: British Medical Journal</i> , 2000 , 321, 1107-11		179
148	Stroke, a major and increasing risk factor for femoral neck fracture. <i>Stroke</i> , 2000 , 31, 1572-7	6.7	173
147	The cost of delirium in the surgical patient. 2001 , 42, 68-73		171
146	Alcohol withdrawal. 2001 , 85, 1191-212		32
145	Postoperative delirium. 2001 , 85, 1229-39		58
144	Delirium in the Hospitalised Elderly. 2001 , 31, 35-40		14
143	Sleep apnea, delirium, depressed mood, cognition, and ADL ability after stroke. <i>Journal of the American Geriatrics Society</i> , 2001 , 49, 391-7	5.6	120
142	Reducing delirium after hip fracture: a randomized trial. <i>Journal of the American Geriatrics Society</i> , 2001 , 49, 516-22	5.6	1006
141	Delirium before and after operation for femoral neck fracture. <i>Journal of the American Geriatrics Society</i> , 2001 , 49, 1335-40	5.6	114
140	Acute confusional state in the elderly following hip surgery: incidence, risk factors and complications. <i>International Journal of Geriatric Psychiatry</i> , 2001 , 16, 349-55	3.9	168
139	The recovery of cognitive function after general anesthesia in elderly patients: a comparison of desflurane and sevoflurane. <i>Anesthesia and Analgesia</i> , 2001 , 93, 1489-94, table of contents	3.9	116
138	Minimizing perioperative adverse events in the elderly. <i>British Journal of Anaesthesia</i> , 2001 , 87, 608-24	5.4	194
137	Preoperative factors associated with postoperative change in confusion assessment method score in hip fracture patients. <i>Anesthesia and Analgesia</i> , 2002 , 94, 1628-32, table of contents	3.9	30
136	Preoperative Factors Associated with Postoperative Change in Confusion Assessment Method Score in Hip Fracture Patients. <i>Anesthesia and Analgesia</i> , 2002 , 94, 1628-1632	3.9	73
135	Small-dose ketamine improves the postoperative state of depressed patients. <i>Anesthesia and Analgesia</i> , 2002 , 95, 114-8, table of contents	3.9	159

134	The incidence and consequences of falls in stroke patients during inpatient rehabilitation: factors associated with high risk. 2002 , 83, 329-33		207
133	Perioperative management of the hospitalized patient. 2002 , 86, 731-48		15
132	Titration of isoflurane using BIS index improves early recovery of elderly patients undergoing orthopedic surgeries. <i>Canadian Journal of Anaesthesia</i> , 2002 , 49, 13-8	3	75
131	Serum concentration of S-100 protein in assessment of cognitive dysfunction after general anesthesia in different types of surgery. <i>Acta Anaesthesiologica Scandinavica</i> , 2002 , 46, 384-9	1.9	61
130	Treatment of delirium in older medical inpatients: a challenge for geriatric specialists. <i>Journal of the American Geriatrics Society</i> , 2002 , 50, 2101-3	5.6	13
129	Dementia after delirium in patients with femoral neck fractures. <i>Journal of the American Geriatrics Society</i> , 2003 , 51, 1002-6	5.6	175
128	Delirium, depression, and anxiety. 2003 , 19, 771-87, viii		13
127	Medical Aspects of Hip Fracture Management. 2003 , 669-680		3
126	Patient safety in geriatrics: a call for action. 2003 , 58, M813-9		33
125	Predicting delirium after vascular surgery: a model based on pre- and intraoperative data. 2003 , 238, 149-56		122
124	A 3 Week Geriatric Education Program for 4 Year Medical Students at Dalhousie University. 2003 , 8, 4342		1
123	Postoperative delirium--a challenge for the orthopedic team. 2004 , 75, 375-7		9
122	Delirium: behavioural changes before and during the prodromal phase. 2004 , 13, 609-16		38
121	Pathogenesis of and management strategies for postoperative delirium after hip fracture: a review. 2004 , 75, 378-89		139
120	[Factors affecting quality of care for elderly subjects undergoing surgery for hip fracture: review of the literature]. 2004 , 90, 504-16		4
119	Delirium in Elderly Patients. 2004 , 12, 7-21		172
118	How do delirium and dementia increase length of stay of elderly general medical inpatients?. 2004 , 45, 235-42		82
117	Anaesthesia in elderly patients with neurodegenerative disorders: special considerations. 2004 , 21, 229-42		57

116	Management of dementia and acute confusional states in the perioperative period. 2004 , 22, vii-viii, 413-22		15
115	Effect of postoperative delirium on outcome after hip fracture. 2004 , 195-200		103
114	Nightly enteral nutrition support of elderly hip fracture patients: a pilot study. 2004 , 23, 683-91		41
113	Notable Names in Anaesthesia. <i>European Journal of Anaesthesiology</i> , 2005 , 22, 564	2.3	
112	Mediastinal haemorrhage mimicking tamponade during en-bloc oesophagectomy. <i>European Journal of Anaesthesiology</i> , 2005 , 22, 555-6	2.3	1
111	The impact of the type of anaesthesia on cognitive status and delirium during the first postoperative days in elderly patients. <i>European Journal of Anaesthesiology</i> , 2005 , 22, 492-9	2.3	76
110	Cardiovascular and Thoracic Anaesthesia. <i>European Journal of Anaesthesiology</i> , 2005 , 22, 565	2.3	
109	Delirium is associated with poor rehabilitation outcome in elderly patients treated for femoral neck fractures. <i>Scandinavian Journal of Caring Sciences</i> , 2005 , 19, 119-27	2.3	99
108	A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. <i>Journal of the American Geriatrics Society</i> , 2005 , 53, 622-8	5.6	253
107	Multicomponent intervention strategies for managing delirium in hospitalized older people: systematic review. 2005 , 52, 79-90		146
106	Postoperative delirium in the older patient. 2005 , 200, 767-73		58
105	["FRAGILE"-Old people and surgery]. 2005 , 76, 28-34		6
104	Een systematisch overzicht van multifactoriële interventies ter primaire preventie van delier bij ouderen. 2005 , 36, 244-250		4
103	Preventing delirium in older people. 2005 , 73-74, 25-34		16
102	Postoperative interleukin-6 and cortisol concentrations in elderly patients with postoperative confusion. 2005 , 12, 60-6		38
101	Cognitively impaired older adults: from hospital to home. 2005 , 105, 52-61; quiz 61-2		81
100	Donepezil in the Prevention and Treatment of Post-Surgical Delirium. 2005 , 13, 1100-1106		109
99	Delirio postoperatorio en pacientes neuroquirúrgicos: evaluación mediante el Test Mental Abreviado. 2006 , 17, 119-124		2

98	Is depth of anesthesia, as assessed by the Bispectral Index, related to postoperative cognitive dysfunction and recovery?. <i>Anesthesia and Analgesia</i> , 2006 , 103, 633-40	3.9	102
97	Postoperative delirium in spine surgery. 2006 , 6, 164-9		71
96	Inpatient falls and injuries in older patients treated for femoral neck fracture. <i>Archives of Gerontology and Geriatrics</i> , 2006 , 43, 389-99	4	28
95	The Organic Brain Syndrome (OBS) scale: a systematic review. <i>International Journal of Geriatric Psychiatry</i> , 2006 , 21, 210-22	3.9	42
94	Care coordination for cognitively impaired older adults and their caregivers. 2007 , 26, 57-78		58
93	Delirium in the orthopaedic patient. <i>Orthopaedic Nursing</i> , 2007 , 26, 354-63; quiz 364-5	0.9	10
92	Managing delirium in hospitalized elderly patients. 2007 , 2, 317-327		
91	[Analgesia after hip fracture repair in elderly patients: the effect of a continuous femoral nerve block: a prospective and randomised study]. 2007 , 26, 2-9		26
90	A study of a pathway to reduce pressure ulcers for patients with a hip fracture. <i>Journal of Orthopaedic Nursing</i> , 2007 , 11, 151-159		12
89	Delirium: guidelines for general hospitals. 2007 , 62, 371-83		104
88	Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. 2007 , 19, 178-86		186
87	Risk factors and characteristics of falls during inpatient rehabilitation of elderly patients. <i>Archives of Gerontology and Geriatrics</i> , 2007 , 44, 1-12	4	42
86	Cognitive function and pressure ulcers in hip fracture patients. <i>Scandinavian Journal of Caring Sciences</i> , 2007 , 21, 79-83	2.3	19
85	A multidisciplinary, multifactorial intervention program reduces postoperative falls and injuries after femoral neck fracture. <i>Osteoporosis International</i> , 2007 , 18, 167-75	5.3	170
84	How to prevent perioperative delirium in the elderly?. <i>Zeitschrift Fur Gerontologie Und Geriatrie</i> , 2008 , 41, 447-52	2.7	29
83	Quantifying the association between computerised measures of attention and confusion assessment method defined delirium: a prospective study of older orthopaedic surgical patients, free of dementia. <i>International Journal of Geriatric Psychiatry</i> , 2008 , 23, 1253-60	3.9	22
82	A comprehensive hip fracture program reduces complication rates and mortality. <i>Journal of the American Geriatrics Society</i> , 2008 , 56, 1831-8	5.6	139
81	Differences in complications and length of stay between patients with a hip fracture treated in an orthopaedic department and patients treated in other hospital departments. <i>Journal of Orthopaedic Nursing</i> , 2008 , 12, 13-25		12

80	Does femoral nerve analgesia impact the development of postoperative delirium in the elderly? A retrospective investigation. <i>Acute Pain</i> , 2008 , 10, 59-64		11
79	[Acute orthogeriatric care (II). Clinical aspects]. <i>Revista Espanola De Geriatria Y Gerontologia</i> , 2008 , 43, 316-29	1.7	13
78	Evaluation and management of the elderly patient at risk for postoperative delirium. <i>Clinics in Geriatric Medicine</i> , 2008 , 24, 667-86, viii	3.8	19
77	Comparison of three scores to screen for delirium in the recovery room. <i>British Journal of Anaesthesia</i> , 2008 , 101, 338-43	5.4	77
76	A survey of old age psychiatry consultation/liaison services in Ireland. <i>Irish Journal of Psychological Medicine</i> , 2008 , 25, 66-68	3	
75	Dehydration induced by bowel preparation in older adults does not result in cognitive dysfunction. <i>Anesthesia and Analgesia</i> , 2008 , 106, 924-9, table of contents	3.9	18
74	Delirium. 173-186		
73	The hip fracture incidence curve is shifting to the right. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009 , 80, 520-4	4.3	72
72	Effects of an interdisciplinary nurse-led Delirium Prevention and Management Program (DPMP) on nursing workload: a pilot study. <i>International Journal of Nursing Studies</i> , 2009 , 46, 804-12	5.8	36
71	Thinking clearly about postoperative delirium and perioperative medications: how concerned should we be?. <i>Canadian Journal of Anaesthesia</i> , 2009 , 56, 785-8	3	1
70	Postoperative cognitive dysfunction in the elderly. <i>Anesthesiology Clinics</i> , 2009 , 27, 485-96, table of contents	6.5	65
69	Evaluation and management of the elderly patient at risk for postoperative delirium. <i>Thoracic Surgery Clinics</i> , 2009 , 19, 363-76, vi	3.1	2
68	The role of the preoperative clinic in perioperative risk reduction. <i>International Anesthesiology Clinics</i> , 2009 , 47, 151-60	0.6	3
67	Delirium after spinal surgery in Korean population. <i>Spine</i> , 2010 , 35, 1729-32	3.3	31
66	Early postoperative delirium after open-heart cardiac surgery is associated with decreased bispectral EEG and increased cortisol and interleukin-6. <i>Intensive Care Medicine</i> , 2010 , 36, 2081-9	14.5	115
65	Medical problems in hip fracture patients. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2010 , 130, 1355-61	6.16	22
64	Delirium--awareness, observation and interventions in intensive care units: a national survey of Swedish ICU head nurses. <i>Intensive and Critical Care Nursing</i> , 2010 , 26, 296-303	3.1	40
63	Reducing delirium in elderly patients with hip fracture: a multi-factorial intervention study. <i>Acta Anaesthesiologica Scandinavica</i> , 2010 , 54, 678-88	1.9	137

62	Delirium in the older emergency department patient: a quiet epidemic. <i>Emergency Medicine Clinics of North America</i> , 2010 , 28, 611-31	1.9	73
61	Emergency orthogeriatrics: concepts and therapeutic alternatives. <i>Emergency Medicine Clinics of North America</i> , 2010 , 28, 927-49	1.9	16
60	Sedation depth during spinal anesthesia and the development of postoperative delirium in elderly patients undergoing hip fracture repair. <i>Mayo Clinic Proceedings</i> , 2010 , 85, 18-26	6.4	300
59	General anesthesia occurs frequently in elderly patients during propofol-based sedation and spinal anesthesia. <i>Journal of Clinical Anesthesia</i> , 2010 , 22, 179-83	1.9	48
58	Delirium: evaluación de tres guías clínicas con el instrumento AGREE. <i>Revista Colombiana De Psiquiatría</i> , 2011 , 40, 363-376	0.8	
57	Perioperative Medical Management. 2011 , 93-99		2
56	Postoperative cognitive disorders. <i>Current Opinion in Critical Care</i> , 2011 , 17, 376-81	3.5	139
55	The effect of sevoflurane anesthesia on cognitive function and the expression of Insulin-like Growth Factor-1 in CA1 region of hippocampus in old rats. <i>Molecular Biology Reports</i> , 2011 , 38, 1195-9	2.8	26
54	Impact of a multifaceted program to prevent postoperative delirium in the elderly: the CONFUCIUS stepped wedge protocol. <i>BMC Geriatrics</i> , 2011 , 11, 25	4.1	29
53	The comparison of hip fractures care in Lithuania and Sweden. <i>International Journal of Orthopaedic and Trauma Nursing</i> , 2012 , 16, 47-52	1.1	3
52	Does using a femoral nerve block for total knee replacement decrease postoperative delirium?. <i>BMC Anesthesiology</i> , 2012 , 12, 4	2.4	26
51	GCâMS Based Serum Metabolomic Analysis of Isoflurane-Induced Postoperative Cognitive Dysfunctional Rats: Biomarker Screening and Insight into Possible Pathogenesis. <i>Chromatographia</i> , 2012 , 75, 799-808	2.1	5
50	A multidisciplinary intervention program improved the outcome after hip fracture for people with dementia--subgroup analyses of a randomized controlled trial. <i>Archives of Gerontology and Geriatrics</i> , 2012 , 54, e284-9	4	89
49	Effectiveness of a multi-component intervention to reduce delirium incidence in elderly care wards. <i>Age and Ageing</i> , 2013 , 42, 721-7	3	54
48	What is the evidence to guide best practice for the management of older people with cognitive impairment presenting to emergency departments? A systematic review. <i>Advanced Emergency Nursing Journal</i> , 2013 , 35, 154-69	0.8	16
47	Continuous spinal anaesthesia versus ultrasound-guided combined psoas compartment-sciatic nerve block for hip replacement surgery in elderly high-risk patients: a prospective randomised study. <i>BMC Anesthesiology</i> , 2014 , 14, 99	2.4	20
46	The cost-effectiveness of multi-component interventions to prevent delirium in older people undergoing surgical repair of hip fracture. <i>European Journal of Orthopaedic Surgery and Traumatology</i> , 2014 , 24, 187-95	2.2	17
45	Insulin-like growth factor-1 and insulin-like growth factor binding protein 3 and risk of postoperative cognitive dysfunction. <i>SpringerPlus</i> , 2015 , 4, 787		4

44	Dysfonctions cognitives postop�atoires. <i>Praticien En Anesth�sie Reanimation</i> , 2015 , 19, 115-119	0	
43	Clinical outcome and predictors of adverse events of an enhanced older adult psychiatric liaison service: Rapid Assessment Interface and Discharge (Newport). <i>Clinical Interventions in Aging</i> , 2017 , 12, 29-36	4	9
42	Use of the Montreal Cognitive Assessment test to investigate the prevalence of mild cognitive impairment in the elderly elective surgical population. <i>Anaesthesia and Intensive Care</i> , 2016 , 44, 581-6	1.1	16
41	Preventing delirium in dementia: Managing risk factors. <i>Maturitas</i> , 2016 , 92, 35-40	5	16
40	Burden of Geriatric Events Among Older Adults Undergoing Major Cancer Surgery. <i>Journal of Clinical Oncology</i> , 2016 , 34, 1231-8	2.2	49
39	European Society of Anaesthesiology evidence-based and consensus-based guideline on postoperative delirium. <i>European Journal of Anaesthesiology</i> , 2017 , 34, 192-214	2.3	362
38	Sleep-Disordered Breathing, Postoperative Delirium, and Cognitive Impairment. <i>Anesthesia and Analgesia</i> , 2017 , 124, 1626-1635	3.9	16
37	Risk Factors for Delirium After Spine Surgery in Extremely Elderly Patients Aged 80 Years or Older and Review of the Literature: Japan Association of Spine Surgeons with Ambition Multicenter Study. <i>Global Spine Journal</i> , 2017 , 7, 560-566	2.7	34
36	Volatile anaesthetics and postoperative delirium in older surgical patients-A secondary analysis of prospective cohort studies. <i>Acta Anaesthesiologica Scandinavica</i> , 2019 , 63, 18-26	1.9	11
35	Recommandation sur lâ�nsth�sie du sujet ��: lâ�exemple de fracture de lâ�extr�mit�sup�rieure du f�mur. <i>Anesth�sie & R�animation</i> , 2019 , 5, 122-138	0.1	6
34	Implementing a Geriatric Fracture Program in a Mixed Practice Environment Reduces Total Cost and Length of Stay. <i>Geriatric Orthopaedic Surgery and Rehabilitation</i> , 2021 , 12, 2151459320987701	2	0
33	A bibliometric analysis of orthogeriatric care: top 50 articles. <i>European Journal of Trauma and Emergency Surgery</i> , 2021 , 1	2.3	0
32	Non-pharmacological interventions for preventing delirium in hospitalised non-ICU patients. <i>The Cochrane Library</i> , 2021 , 7, CD013307	5.2	4
31	Delirium. 2007 , 210-218		5
30	Delirium in the intensive care unit: are we helping the patient?. <i>Critical Care Nursing Quarterly</i> , 2000 , 22, 47-60	2	16
29	Managing delirium and agitation in elderly hospitalized orthopaedic patients: Part I--Theoretical aspects. <i>Orthopaedic Nursing</i> , 2001 , 20, 31-43; quiz 44-6	0.9	11
28	Orthopaedic and trauma surgery. <i>BMJ: British Medical Journal</i> , 1995 , 310, 917-9		4
27	Patient falls in stroke rehabilitation. A challenge to rehabilitation strategies. <i>Stroke</i> , 1995 , 26, 838-42	6.7	279

26	Fall prediction index for patients in stroke rehabilitation. <i>Stroke</i> , 1997 , 28, 716-21	6.7	120
25	Risk Factors of Postoperative Delirium in Elderly Patients with Hip Fractures. <i>Journal of the Korean Fracture Society</i> , 2008 , 21, 189	0.1	13
24	THE INFLUENCE OF COGNITIVE FUNCTION ON OUTCOME AFTER A HIP FRACTURE. <i>Journal of Bone and Joint Surgery - Series A</i> , 2006 , 88, 2115-2123	5.6	1
23	Delirium in elderly patients: an overview of the state of the science. <i>Journal of Gerontological Nursing</i> , 2001 , 27, 12-20	1.2	44
22	Acute confusion/delirium protocol. <i>Journal of Gerontological Nursing</i> , 2001 , 27, 21-33; quiz 62-3	1.2	28
21	Documentation of delirium in elderly patients with hip fracture. <i>Journal of Gerontological Nursing</i> , 2002 , 28, 23-9	1.2	63
20	Risk factors of postoperative delirium following liver transplantation. <i>Korean Journal of Anesthesiology</i> , 2009 , 57, 584-589	3.8	4
19	Lidocaine administration before tracheal extubation cannot reduce post-operative cognition disorders in elderly patients. <i>Advanced Biomedical Research</i> , 2013 , 2, 81	1.2	3
18	The Risk Factor of Delirium after Transurethral Resection of the Prostate. <i>Korean Journal of Urology</i> , 2006 , 47, 953		3
17	Fall prevention in the elderly: analysis and comprehensive review of methods used in the hospital and in the home. <i>Journal of the American Academy of Orthopaedic Surgeons</i> , 2011 , 19, 402-9	4.5	28
16	Care of the Injured Elderly. 2001 , 240-260		
15	Surgery in the Elderly – Outcome and Complication Pattern. 2002 , 893-901		
14	POSTOPERATIVE DELIRIUM AFTER HIP FRACTURE. <i>Journal of Bone and Joint Surgery - Series A</i> , 2006 , 88, 2060-2067	5.6	2
13	Orthopedic disorders. 2007 , 511-529		
12	Postoperative Delirium after Hip Arthroplasty in the Elderly. <i>The Journal of the Korean Hip Society</i> , 2010 , 22, 151		3
11	Risk Factors for Delirium in the Elderly. 1998 , 159-166		
10	Perioperative Risk Assessment and Decision Making. 2006 , 143-151		
9	Revisiting the O complex: urinary incontinence, delirium and polypharmacy in elderly patients. <i>Cmaj</i> , 1997 , 157, 1071-7	3.5	4

8	Effectiveness of interventions to prevent delirium in hospitalized patients: a systematic review. <i>Cmaj</i> , 1996 , 155, 1263-8	3.5	9
7	Post-operative delirium after hip fracture treatment - a review of the current literature. <i>Gms Psycho-social-medicine</i> , 2006 , 3, Doc01		8
6	Sensi-steps: Using Patient-Generated Data to Prevent Post-stroke Falls. 2017 , 2017, 2294-2298	0.7	0
5	Non-pharmacological interventions for preventing delirium in hospitalised non-ICU patients. <i>The Cochrane Library</i> , 2021 , 11, CD013307	5.2	2
4	Low geriatric nutritional risk index is a risk factor for death within 1 year following hip fracture.. <i>Journal of Orthopaedic Surgery</i> , 2022 , 30, 10225536221103360	1.4	
3	Prehabilitation of elderly frail or pre-frail patients prior to elective surgery (PRAEP-GO): study protocol for a randomized, controlled, outcome assessor-blinded trial. <i>Trials</i> , 2022 , 23,	2.8	0
2	DELIRIUM IN THE HOSPITALIZED ELDERLY. 1998 , 33, 417-439		9
1	Der fraile Patient: fiktives Fallbeispiel. 2022 , 57, 674-681		0