

# CITATION REPORT

List of articles citing

## Dobutamine digital echocardiography for detecting coronary artery disease

DOI: 10.1016/0002-9149(91)90457-v  
American Journal of Cardiology, 1991, 67, 1311-8.

**Source:** <https://exaly.com/paper-pdf/22434300/citation-report.pdf>

**Version:** 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| #   | Paper                                                                                                                                                                                                                             | IF | Citations |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|-----------|
| 211 | Evolution of stress testing. <b>1992</b> , 85, 1217-8                                                                                                                                                                             |    | 19        |
| 210 | Stress Doppler echocardiography using dobutamine in coronary patients with and without ischaemia induction. <b>1992</b> , 13, 1020-7                                                                                              |    | 14        |
| 209 | Dobutamine stress echocardiography as a predictor of cardiac events associated with aortic surgery. <b>1992</b> , 15, 831-842                                                                                                     |    | 124       |
| 208 | Dobutamine stress echocardiography for detection and assessment of coronary artery disease. <b>1992</b> , 19, 1203-11                                                                                                             |    | 255       |
| 207 | New developments in pharmacologic stress imaging. <b>1992</b> , 124, 468-85                                                                                                                                                       |    | 58        |
| 206 | On-line quantification of ventricular function during dobutamine stress echocardiography. <b>1992</b> , 13, 1669-76                                                                                                               |    | 62        |
| 205 | Clinical significance of abrupt vasodepression during dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1992</b> , 69, 1484-6                                                                        | 3  | 49        |
| 204 | Enhanced sensitivity for detection of coronary artery disease by addition of atropine to dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1992</b> , 70, 41-6                                       | 3  | 359       |
| 203 | Comparison of exercise electrocardiography and dobutamine echocardiography. <b>1992</b> , 15, 641-6                                                                                                                               |    | 12        |
| 202 | Effect of dobutamine on left ventricular relaxation and filling phase in patients with ischemic heart disease and preserved systolic function. <b>1993</b> , 7, 325-31                                                            |    | 6         |
| 201 | Comparison of dobutamine and exercise echocardiography for detecting coronary artery disease. <i>American Journal of Cardiology</i> , <b>1993</b> , 72, 1226-31                                                                   | 3  | 85        |
| 200 | Dipyridamole thallium-201 imaging versus dobutamine echocardiography for the evaluation of coronary artery disease in patients unable to exercise. <i>American Journal of Cardiology</i> , <b>1993</b> , 72, 1257-62 <sup>3</sup> |    | 30        |
| 199 | Predictive value of dobutamine echocardiography just before noncardiac vascular surgery. <i>American Journal of Cardiology</i> , <b>1993</b> , 72, 602-7                                                                          | 3  | 121       |
| 198 | Comparison of dobutamine stress echocardiography, dipyridamole stress echocardiography and exercise stress testing for diagnosis of coronary artery disease. <i>American Journal of Cardiology</i> , <b>1993</b> , 72, 865-70     | 3  | 123       |
| 197 | Dobutamine stress echocardiography: prevalence of a nonischemic response in a low-risk population. <b>1993</b> , 125, 1257-61                                                                                                     |    | 28        |
| 196 | Comparison of dobutamine stress echocardiography and stress thallium-201 single-photon emission computed tomography for detecting coronary artery disease. <b>1993</b> , 6, 593-602                                               |    | 59        |
| 195 | Dobutamine stress magnetic resonance imaging for detection of coronary artery disease. <b>1993</b> , 22, 431-9                                                                                                                    |    | 95        |

|     |                                                                                                                                                                                                                                                                                |   |     |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|-----|
| 194 | Ischemia-induced regional wall motion abnormality is improved after coronary angioplasty: demonstration by dobutamine stress echocardiography. <b>1993</b> , 21, 584-9                                                                                                         |   | 40  |
| 193 | Dobutamine stress echocardiography predicts surgical outcome in patients with an aortic aneurysm and peripheral vascular disease. <b>1993</b> , 21, 957-63                                                                                                                     |   | 161 |
| 192 | Optimal use of dobutamine stress for the detection and evaluation of coronary artery disease: combination with echocardiography or scintigraphy, or both?. <b>1993</b> , 22, 159-67                                                                                            |   | 301 |
| 191 | Selection of the optimal nonexercise stress for the evaluation of ischemic regional myocardial dysfunction and malperfusion. Comparison of dobutamine and adenosine using echocardiography and 99mTc-MIBI single photon emission computed tomography. <b>1993</b> , 87, 345-54 |   | 294 |
| 190 | Dobutamine stress echocardiography identifies hibernating myocardium and predicts recovery of left ventricular function after coronary revascularization. <b>1993</b> , 88, 430-6                                                                                              |   | 417 |
| 189 | Clinical Utility of Digital Dobutamine Stress Echocardiography in the Noninvasive Evaluation of Coronary Artery Disease. <b>1994</b> , 154, 1065                                                                                                                               |   | 7   |
| 188 | Dobutamine-atropine stress echocardiography in elderly patients unable to perform an exercise test. Hemodynamic characteristics, safety, and prognostic value. <b>1994</b> , 154, 2681-6                                                                                       |   | 41  |
| 187 | Comparative ability of dobutamine and exercise stress in inducing myocardial ischaemia in active patients. <b>1994</b> , 72, 31-8                                                                                                                                              |   | 51  |
| 186 | Myocardial perfusion imaging versus two-dimensional echocardiography: comparative value in the diagnosis of coronary artery disease. <b>1994</b> , 1, 399-414                                                                                                                  |   | 15  |
| 185 | Ischemic heart disease: assessment with gadolinium-enhanced ultrafast MR imaging and dipyridamole stress. <b>1994</b> , 4, 425-31                                                                                                                                              |   | 125 |
| 184 | Comparison of dobutamine and exercise using technetium-99m sestamibi imaging for the evaluation of coronary artery disease. <i>American Journal of Cardiology</i> , <b>1994</b> , 73, 164-9                                                                                    | 3 | 23  |
| 183 | Left ventricular volume changes during dobutamine stress echocardiography identify patients with more extensive coronary artery disease. <b>1994</b> , 24, 1268-73                                                                                                             |   | 42  |
| 182 | Abnormal left ventricular early diastolic filling during dobutamine stress Doppler echocardiography is a sensitive indicator of significant coronary artery disease. <b>1994</b> , 24, 1618-24                                                                                 |   | 52  |
| 181 | Quantitative coronary angiography in the estimation of the functional significance of coronary stenosis: correlations with dobutamine-atropine stress test. <b>1994</b> , 23, 1434-9                                                                                           |   | 101 |
| 180 | Improved endocardial border resolution during dobutamine stress echocardiography with intravenous sonicated dextrose albumin. <b>1994</b> , 23, 1440-3                                                                                                                         |   | 105 |
| 179 | How accurate is dobutamine stress electrocardiography for detection of coronary artery disease? Comparison with two-dimensional echocardiography and technetium-99m methoxyl isobutyl isonitrile (mibi) perfusion scintigraphy. <b>1994</b> , 24, 920-7                        |   | 84  |
| 178 | False positive dobutamine stress echocardiograms: characterization of clinical, echocardiographic and angiographic findings. <b>1994</b> , 24, 928-33                                                                                                                          |   | 108 |
| 177 | Dobutamine two-dimensional transesophageal echocardiographic stress testing for detection of coronary artery disease. <b>1994</b> , 128, 36-41                                                                                                                                 |   | 25  |

|     |                                                                                                                                                                                                                                                            |   |     |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|-----|
| 176 | Dobutamine stress echocardiography: clinical utility and predictive value at various infusion rates. <b>1994</b> , 128, 804-8                                                                                                                              |   | 17  |
| 175 | Dobutamine stress echocardiography: sensitivity, specificity, and predictive value for future cardiac events. <b>1994</b> , 127, 1510-5                                                                                                                    |   | 91  |
| 174 | Dobutamine-atropine stress echocardiography and clinical data for predicting late cardiac events in patients with suspected coronary artery disease. <b>1994</b> , 97, 119-25                                                                              |   | 73  |
| 173 | Safety and tolerability of dobutamine-atropine stress echocardiography: a prospective, multicentre study. Echo Dobutamine International Cooperative Study Group. <b>1994</b> , 344, 1190-2                                                                 |   | 339 |
| 172 | Dobutamine stress echocardiography compared with dipyridamole thallium-201 single-photon emission computed tomography in detecting coronary artery disease. <b>1995</b> , 16, 570-5                                                                        |   | 50  |
| 171 | Diagnosis of coronary artery disease and viable myocardium by stress echocardiography. Diagnostic accuracy of different stress modalities. <b>1995</b> , 16 Suppl J, 10-8                                                                                  |   | 3   |
| 170 | Diagnostic and prognostic value of stress testing in older persons. <b>1995</b> , 43, 190-4                                                                                                                                                                |   | 7   |
| 169 | The multicentre trial philosophy in stress echocardiography: lessons learned from the EPIC study. EPIC-EDIC Study Project. Echo Persantine International Cooperative Study. Echo-Dobutamine International Cooperative Study. <b>1995</b> , 16 Suppl J, 2-4 |   | 9   |
| 168 | Clinical and diagnostic utility of arbutamine for cardiovascular stress testing during echocardiographic monitoring. <b>1995</b> , 16 Suppl M, 24-7                                                                                                        |   | 2   |
| 167 | Dobutamine stress echocardiography predicts early wall motion improvement after elective percutaneous transluminal coronary angioplasty. <i>American Journal of Cardiology</i> , <b>1995</b> , 76, 652-6                                                   | 3 | 9   |
| 166 | Comparison of stress echocardiography and stress myocardial perfusion scintigraphy for diagnosing coronary artery disease and assessing its severity. <i>American Journal of Cardiology</i> , <b>1995</b> , 75, 25D-34D                                    | 3 | 54  |
| 165 | Dobutamine stress echocardiography: Causes and approach to false negative and false positive studies. <b>1995</b> , 4, 34-38                                                                                                                               |   | 3   |
| 164 | The influence of angiographically demonstrated coronary collaterals on the results of stress echocardiography. <b>1995</b> , 18, 205-8                                                                                                                     |   | 12  |
| 163 | Dobutamine stress echocardiography for the diagnosis and management of coronary artery disease. <b>1995</b> , 18, 252-60                                                                                                                                   |   | 9   |
| 162 | Dobutamine echocardiography: usefulness of digital image processing. <b>1995</b> , 16, 1420-4                                                                                                                                                              |   | 22  |
| 161 | Stress Echocardiography. Part II. Dobutamine Stress Echocardiography: Techniques, Implementation, Clinical Applications, and Correlations. <b>1995</b> , 70, 16-27                                                                                         |   |     |
| 160 | Cardiovascular response in patients with and without myocardial ischaemia during dobutamine echocardiography stress test for coronary artery disease. <b>1995</b> , 15, 249-63                                                                             |   | 4   |
| 159 | Electrocardiographic response during dobutamine stress echocardiography. <b>1995</b> , 129, 672-7                                                                                                                                                          |   | 26  |

|     |                                                                                                                                                                                                                                    |   |     |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|-----|
| 158 | Does hypotension during dobutamine stress echocardiography correlate with anatomic or functional cardiac impairment?. <b>1995</b> , 129, 1121-6                                                                                    |   | 24  |
| 157 | Effect of beta-adrenergic receptor blockade on the physiologic response to dobutamine stress echocardiography. <b>1995</b> , 130, 248-53                                                                                           |   | 21  |
| 156 | Intravenous albumin during dobutamine stress echocardiography: enhanced localization of left ventricular endocardial borders. <b>1995</b> , 130, 254-8                                                                             |   | 61  |
| 155 | Dipyridamole-thallium versus dobutamine echocardiographic stress testing: a clinician's viewpoint. <b>1995</b> , 130, 373-85                                                                                                       |   | 7   |
| 154 | Assessment of coronary artery disease by dobutamine stress echocardiography (DSE). <b>1995</b> , 25, 707-15                                                                                                                        |   | 5   |
| 153 | Dobutamine stress echocardiography for the identification of multivessel coronary artery disease after uncomplicated myocardial infarction: the importance of test end-point. <b>1995</b> , 50, 51-60                              |   | 13  |
| 152 | Optimal stage duration in dobutamine stress echocardiography. <b>1995</b> , 25, 605-9                                                                                                                                              |   | 26  |
| 151 | Transesophageal dobutamine stress echocardiography in the evaluation of coronary artery disease. <b>1995</b> , 25, 823-9                                                                                                           |   | 27  |
| 150 | Dobutamine stress echocardiography in orthotopic heart transplant recipients. VACOMED Research Group. <b>1995</b> , 25, 1665-72                                                                                                    |   | 75  |
| 149 | Arbutamine echocardiography: efficacy and safety of a new pharmacologic stress agent to induce myocardial ischemia and detect coronary artery disease. The International Arbutamine Study Group. <b>1995</b> , 26, 1168-75         |   | 44  |
| 148 | Stress testing with closed-loop arbutamine as an alternative to exercise. The International Arbutamine Study Group. <b>1995</b> , 26, 1151-8                                                                                       |   | 35  |
| 147 | Stress echocardiography. Part II. Dobutamine stress echocardiography: techniques, implementation, clinical applications, and correlations. <b>1995</b> , 70, 16-27                                                                 |   | 168 |
| 146 | Mechanisms of hypotension during dobutamine stress echocardiography in patients with coronary artery disease. <i>American Journal of Cardiology</i> , <b>1995</b> , 76, 26-30                                                      | 3 | 43  |
| 145 | JPEG compression of digital echocardiographic images: impact on image quality. <b>1995</b> , 8, 306-18                                                                                                                             |   | 68  |
| 144 | An analysis of the safety of performing dobutamine stress echocardiography in an ambulatory setting. <b>1995</b> , 8, 15-20                                                                                                        |   | 10  |
| 143 | Safety and efficacy of computerized closed-loop delivery of arbutamine: a new pharmacologic myocardial stress modality for the assessment of coronary artery disease. The European Arbutamine Study Group. <b>1995</b> , 8, 854-63 |   | 3   |
| 142 | Stress echocardiography, contrast echocardiography, and tissue characterization: applications for the future. <b>1996</b> , 12, 429-50                                                                                             |   | 5   |
| 141 | Prognostic value of dobutamine echocardiography in patients with left ventricular dysfunction. <b>1996</b> , 27, 132-9                                                                                                             |   | 145 |

|     |                                                                                                                                                                                                                                                                                                                                             |      |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 140 | Dobutamine stress echocardiography for detection of coronary artery stenosis in children with Kawasaki disease. <b>1996</b> , 27, 1251-6                                                                                                                                                                                                    | 66   |
| 139 | Guidelines for perioperative cardiovascular evaluation for noncardiac surgery. Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). <b>1996</b> , 27, 910-48                                            | 309  |
| 138 | Dobutamine-induced wall motion abnormalities: correlations with myocardial fractional flow reserve and quantitative coronary angiography. <b>1996</b> , 27, 1429-36                                                                                                                                                                         | 83   |
| 137 | Atropine augmentation in dobutamine stress echocardiography: role and incremental value in a clinical practice setting. <b>1996</b> , 28, 551-7                                                                                                                                                                                             | 84   |
| 136 | Dobutamine-induced ST segment elevation and ventricular fibrillation with nonsignificant coronary artery disease. <b>1996</b> , 132, 1058-60                                                                                                                                                                                                | 16   |
| 135 | Effective therapy with transthoracic video-assisted endoscopic coagulation of the left stellate ganglion and upper sympathetic trunk in congenital long-QT syndrome. <b>1996</b> , 132, 1060-3                                                                                                                                              | 12   |
| 134 | Effect of beta-blockade on dobutamine stress echocardiography. <b>1996</b> , 131, 698-703                                                                                                                                                                                                                                                   | 18   |
| 133 | Regional wall motion analysis by dobutamine stress echocardiography to distinguish between ischemic and nonischemic dilated cardiomyopathy. <b>1996</b> , 131, 537-43                                                                                                                                                                       | 40   |
| 132 | The importance of work-up (verification) bias correction in assessing the accuracy of SPECT thallium-201 testing for the diagnosis of coronary artery disease. <b>1996</b> , 49, 735-42                                                                                                                                                     | 63   |
| 131 | Transesophageal dobutamine-atropine stress echocardiography: diagnostic accuracy for coronary stenosis detection and localization. <b>1996</b> , 56, 185-92                                                                                                                                                                                 |      |
| 130 | Effects of beta-blockade and atropine on ischemic responses in left ventricular regions subtending coronary stenosis during dobutamine stress echocardiography. <b>1996</b> , 28, 1866-76                                                                                                                                                   | 20   |
| 129 | Dobutamine stress echocardiography: a new, noninvasive method for detecting ischemic heart disease. <b>1996</b> , 25, 87-97                                                                                                                                                                                                                 | 1    |
| 128 | ACC/AHA task force report. Special report: guidelines for perioperative cardiovascular evaluation for noncardiac surgery. Report of the American College of Cardiology/American Heart Association Task Force on practice guidelines (Committee on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). <b>1996</b> , 10, 540-52 | 16   |
| 127 | Comparison of adenosine echocardiography, with and without isometric handgrip, to exercise echocardiography in the detection of ischemia in patients with coronary artery disease. <b>1996</b> , 9, 33-43                                                                                                                                   | 37   |
| 126 | Sustained nonischemic ventricular tachycardia during dobutamine stress echocardiography. <b>1996</b> , 87, 82-5                                                                                                                                                                                                                             | 3    |
| 125 | Simultaneous dobutamine stress echocardiography and dobutamine scintigraphy (99mTc-MIBI-SPET) for assessment of coronary artery disease. <b>1996</b> , 12, 185-90                                                                                                                                                                           | 4    |
| 124 | Value of dobutamine stress echocardiography in determining the prognosis of patients with known or suspected coronary artery disease. <i>American Journal of Cardiology</i> , <b>1996</b> , 78, 404-8                                                                                                                                       | 3 63 |
| 123 | Stress echocardiography for the assessment of myocardial ischemia and viability. <b>1996</b> , 21, 445-520                                                                                                                                                                                                                                  | 15   |

|     |                                                                                                                                                                                                                                                                        |      |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 122 | Incremental diagnostic value of dobutamine stress echocardiography and dobutamine scintigraphy (technetium 99m-labeled sestamibi single-photon emission computed tomography) for assessment of presence and extent of coronary artery disease. <b>1996</b> , 3, 212-20 | 23   |
| 121 | The role of stress echocardiography versus stress perfusion: a view from the other side. <b>1996</b> , 3, S66-74                                                                                                                                                       | 11   |
| 120 | Impact of Beta Blockers on Dobutamine-Atropine Stress Echocardiography. <b>1996</b> , 13, 373-380                                                                                                                                                                      | 3    |
| 119 | Dipyridamole and dobutamine-atropine stress echocardiography in the diagnosis of coronary artery disease. Comparison with exercise stress test, analysis of agreement, and impact of antianginal treatment. <b>1996</b> , 110, 1248-54                                 | 66   |
| 118 | Assessment of the functional significance of coronary artery stenosis by dobutamine-atropine stress echocardiography. <b>1997</b> , 88, 386-92                                                                                                                         | 6    |
| 117 | Dobutamine stress echocardiography. <b>1997</b> , 18 Suppl D, D9-15                                                                                                                                                                                                    | 25   |
| 116 | [The predictive signs of multivessel coronary disease in echocardiography with dobutamine]. <b>1997</b> , 50, 772-7                                                                                                                                                    | 0    |
| 115 | Cardiac rupture with dobutamine stress echocardiography. <b>1997</b> , 10, 979-81                                                                                                                                                                                      | 18   |
| 114 | Long-term prognostic significance of dobutamine echocardiography in patients with suspected coronary artery disease: results of a 5-year follow-up study. <b>1997</b> , 29, 969-73                                                                                     | 57   |
| 113 | Evolution of dobutamine echocardiography protocols and indications: safety and side effects in 3,011 studies over 5 years. <b>1997</b> , 29, 1234-40                                                                                                                   | 179  |
| 112 | Methodology, feasibility, safety and diagnostic accuracy of dobutamine stress echocardiography. <b>1997</b> , 30, 595-606                                                                                                                                              | 387  |
| 111 | Significance of persistent left ventricular dysfunction during recovery after dobutamine stress echocardiography. <b>1997</b> , 30, 621-6                                                                                                                              | 44   |
| 110 | On-line quantitative assessment of left ventricular filling during dobutamine stress echocardiography: a useful addition to conventional wall motion scoring. <b>1997</b> , 59, 57-69                                                                                  | 19   |
| 109 | Dobutamine stress echocardiography in the detection of coronary artery disease in a clinical practice setting. <b>1997</b> , 62, 55-62                                                                                                                                 | 71   |
| 108 | Evaluation of signal-averaged cardiokymography for the detection of ischaemic left ventricular dysfunction. <b>1997</b> , 59, 305-12                                                                                                                                   | 4    |
| 107 | Simultaneous dobutamine stress echocardiography and thallium-201 perfusion imaging for the detection of coronary artery disease. <b>1997</b> , 88, 556-62                                                                                                              | 22   |
| 106 | Stress echocardiography: introduction, history, and methods. <b>1997</b> , 39, 499-522                                                                                                                                                                                 | 21   |
| 105 | Adenosine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1997</b> , 79, 25-9                                                                                                                                                                      | 3 19 |

|     |                                                                                                                                                                                                                                                              |   |     |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|-----|
| 104 | Impact of metoprolol on heart rate, blood pressure, and contractility in normal subjects during dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1997</b> , 80, 386-8                                                          | 3 | 6   |
| 103 | Cardiac outcomes in coronary patients with submaximum dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1997</b> , 80, 725-9                                                                                                    | 3 | 22  |
| 102 | Comparison between dobutamine echocardiography and thallium-201 scintigraphy in detecting residual stenosis, ischemia, and necrosis in patients with prior myocardial infarction. <b>1997</b> , 20, 351-6                                                    |   | 5   |
| 101 | Symptomatic and hemodynamic recovery following dobutamine stress echo: benefit of low-dose esmolol administration. <b>1997</b> , 13, 53-7                                                                                                                    |   | 2   |
| 100 | Factors influencing the diagnostic accuracy of dobutamine stress echocardiography. <b>1997</b> , 13, 493-8                                                                                                                                                   |   | 6   |
| 99  | Detection of Viability of Dysfunctional Myocardium in Coronary Heart Disease. II. Echocardiography. <b>1998</b> , 2, 207-233                                                                                                                                 |   | 4   |
| 98  | [Staged diagnosis in suspected restenosis after PTCA--clinical aspects, ergometry and then stress echocardiography?]. <b>1998</b> , 87 Suppl 2, 215-7                                                                                                        |   |     |
| 97  | Value of dobutamine stress echocardiography for the detection of multivessel coronary artery disease. <i>American Journal of Cardiology</i> , <b>1998</b> , 81, 298-301                                                                                      | 3 | 14  |
| 96  | Ischaemia monitoring with on-line vectorcardiography during dobutamine stress echocardiography in patients after an episode of unstable coronary artery disease. <b>1998</b> , 243, 61-70                                                                    |   | 2   |
| 95  | Ischaemia monitoring with on-line vectorcardiography during dobutamine stress echocardiography in patients after an episode of unstable coronary artery disease. <b>1998</b> , 244, 61-70                                                                    |   | 3   |
| 94  | Use of dobutamine echocardiography for cardiac risk stratification of patients with chronic renal failure. <b>1998</b> , 244, 155-61                                                                                                                         |   | 26  |
| 93  | Stress echocardiography: recommendations for performance and interpretation of stress echocardiography. Stress Echocardiography Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. <b>1998</b> , 11, 97-104 |   | 307 |
| 92  | Impaired systolic blood pressure response to dobutamine stress testing: a marker of more severe functional abnormalities in patients with myocardial infarction. <b>1998</b> , 11, 436-41                                                                    |   | 6   |
| 91  | Diagnostic value of 12-lead electrocardiogram during dobutamine echocardiographic studies. <b>1998</b> , 136, 1061-4                                                                                                                                         |   | 13  |
| 90  | Assessment of coronary artery disease in women by dobutamine stress echocardiography: comparison with stress thallium-201 single-photon emission computed tomography and exercise electrocardiography. <b>1998</b> , 135, 655-62                             |   | 58  |
| 89  | Measurement of haemodynamic parameters during dobutamine echocardiography by automated impedance cardiography.                                                                                                                                               |   |     |
| 88  | Risk stratification by pharmacological stress echocardiography in a primary care cardiology centre. Experience in 1082 patients. <b>1998</b> , 19, 1673-80                                                                                                   |   | 16  |
| 87  | Comparative study of dobutamine stress electron-beam computed tomography and exercise thallium scintigraphy in the diagnosis of patients with suspected coronary artery disease. <b>1998</b> , 62, 83-90                                                     |   | 3   |



|    |                                                                                                                                                                                                                      |   |    |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|----|
| 86 | Assessment of the coronary artery disease and systolic dysfunction in hypertensive patients with the dobutamine-atropine stress echocardiography: effect of the left ventricular hypertrophy. <b>1998</b> , 89, 52-8 |   | 15 |
| 85 | Evaluation of systolic performance by automated impedance cardiography. <b>1999</b> , 873, 167-73                                                                                                                    |   | 5  |
| 84 | Advances in Exercise Echocardiography Can This Technique Still Thrive in the Era of Pharmacologic Stress Testing?. <b>1999</b> , 16, 841-856                                                                         |   | 4  |
| 83 | Diagnostic and Prognostic Use of Stress Echocardiography and Radionuclide Scintigraphy. <b>1999</b> , 16, 857-877                                                                                                    |   | 1  |
| 82 | Long-term prognosis in women with normal dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>1999</b> , 83, 1127-9, A9                                                                    | 3 | 15 |
| 81 | Exercise echocardiography. Principles, methods, and clinical use. <b>1999</b> , 17, 447-60, vii                                                                                                                      |   | 10 |
| 80 | Pharmacologic stress echocardiography. Dobutamine and arbutamine stress testing. <b>1999</b> , 17, 461-79, viii                                                                                                      |   | 7  |
| 79 | Assessment of flow mismatch with pharmacologic stress test on myocardial contrast echocardiography in a model of critical stenosis: adenosine triphosphate and dipyridamole. <b>1999</b> , 12, 257-65                |   | 4  |
| 78 | A perspective on standardizing the predictive power of noninvasive cardiovascular tests by likelihood ratio computation: 2. Clinical applications. <b>1999</b> , 74, 1072-87                                         |   | 20 |
| 77 | Objective evaluation of regional left ventricular wall motion during dobutamine stress echocardiographic studies using segmental analysis of color kinesis images. <b>1999</b> , 34, 409-19                          |   | 71 |
| 76 | Can dobutamine stress echocardiography predict cardiac events in nonrevascularized diabetic patients following acute myocardial infarction?. <b>1999</b> , 116, 1224-32                                              |   | 6  |
| 75 | Safety profile and hemodynamic responses to beta-adrenergic stimulation by dobutamine in heart transplant patients. <b>1999</b> , 116, 1587-92                                                                       |   | 9  |
| 74 | Comparative physiological study of arbutamine with exercise in humans. <b>2000</b> , 98, 489-494                                                                                                                     |   | 4  |
| 73 | Comparative physiological study of arbutamine with exercise in humans. <b>2000</b> , 98, 489                                                                                                                         |   | 3  |
| 72 | The comparable diagnostic accuracies of dobutamine-stress and dipyridamole-stress echocardiographies: a meta-analysis. <b>2000</b> , 11, 151-9                                                                       |   | 57 |
| 71 | Does the lack of hyperkinesis during dobutamine stress echocardiography predict the functional significance of coronary arterial stenosis?. <b>2000</b> , 17, 229-39                                                 |   | 2  |
| 70 | Digital echocardiography. <i>American Journal of Cardiology</i> , <b>2000</b> , 86, 2G-3G                                                                                                                            | 3 | 11 |
| 69 | Usefulness of the subendocardial myocardial velocity gradient in low-dose dobutamine stress echocardiography. <b>2000</b> , 15, 11-7                                                                                 |   | 15 |

|    |                                                                                                                                                                                                                                                                                                                                                   |     |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 68 | Dobutamine-atropine stress echocardiography for the detection of coronary artery disease in patients with left ventricular hypertrophy. Importance of chamber size and systolic wall stress. <b>2000</b> , 101, 258-63                                                                                                                            | 76  |
| 67 | Non-contrast second harmonic imaging improves interobserver agreement and accuracy of dobutamine stress echocardiography in patients with impaired image quality. <b>2000</b> , 83, 133-40                                                                                                                                                        | 60  |
| 66 | Dobutamine-atropine stress echocardiography and dipyridamole sestamibi scintigraphy for the detection of coronary artery disease: limitations and concordance. <b>2000</b> , 36, 1265-73                                                                                                                                                          | 60  |
| 65 | Effect of intravenous contrast for left ventricular opacification and border definition on sensitivity and specificity of dobutamine stress echocardiography compared with coronary angiography in technically difficult patients. <b>2001</b> , 142, 908-15                                                                                      | 97  |
| 64 | Pharmacologic stress testing for coronary disease diagnosis: A meta-analysis. <b>2001</b> , 142, 934-44                                                                                                                                                                                                                                           | 157 |
| 63 | Stratification of single-vessel coronary stenosis by ischemic threshold at the onset of wall motion abnormality during continuous monitoring of left ventricular function by semisupine exercise echocardiography. <b>2001</b> , 14, 798-805                                                                                                      | 6   |
| 62 | Measurement of fractional flow reserve to assess moderately severe coronary lesions: correlation with dobutamine stress echocardiography. <b>2001</b> , 14, 499-504                                                                                                                                                                               | 30  |
| 61 | Stress echocardiography: technical considerations. <b>2001</b> , 43, 303-14                                                                                                                                                                                                                                                                       | 3   |
| 60 | Physiologic assessment of coronary artery stenoses. <b>2001</b> , 26, 423-92                                                                                                                                                                                                                                                                      | 3   |
| 59 | Native tissue second harmonic imaging improves endocardial and epicardial border definition in dobutamine stress echocardiography. <b>2001</b> , 2, 52-61                                                                                                                                                                                         | 17  |
| 58 | Preoperative cardiac risk assessment: an updated approach. <b>2001</b> , 136, 1370-6                                                                                                                                                                                                                                                              | 14  |
| 57 | Low Utility of Dobutamine Stress Echocardiograms in the Preoperative Evaluation of Patients Scheduled for Noncardiac Surgery. <b>2002</b> , 95, 512-516                                                                                                                                                                                           | 15  |
| 56 | Low utility of dobutamine stress echocardiograms in the preoperative evaluation of patients scheduled for noncardiac surgery. <b>2002</b> , 95, 512-6, table of contents                                                                                                                                                                          | 24  |
| 55 | Effect of early administration of atropine on paradoxical sinus deceleration during dobutamine stress echocardiography. <i>American Journal of Cardiology</i> , <b>2002</b> , 89, 645-7                                                                                                                                                           | 3 6 |
| 54 | Contrast and harmonic imaging improves accuracy and efficiency of novice readers for dobutamine stress echocardiography. <b>2002</b> , 19, 483-8                                                                                                                                                                                                  | 36  |
| 53 | ACC/AHA/ASE 2003 Guideline Update for the Clinical Application of Echocardiography: Summary Article. <b>2003</b> , 16, 1091-1110                                                                                                                                                                                                                  | 33  |
| 52 | Relationship between dobutamine response of dyssynergic myocardium and angiographically documented blood supply. <b>2003</b> , 16, 949-57                                                                                                                                                                                                         | 1   |
| 51 | ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography--summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). <b>2003</b> , 42, 954-70 | 392 |

|    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |     |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 50 | ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). <b>2003</b> , 103, 1145-120                                                                                                                                                                                                          | 623 |
| 49 | ACC/AHA/ASE 2003 Guideline Update for the Clinical Application of Echocardiography: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). <b>2003</b> , 103, 1145-120                                                                                                                                                                                                          | 464 |
| 48 | Transmitral pulsed-Doppler echocardiography is a more accurate technique compared with two-dimensional echocardiography using dobutamine, in patients with one vessel coronary artery disease. <b>2003</b> , 5, 63-72                                                                                                                                                                                                                                                                                                                                         | 2   |
| 47 | Intraoperative Stress Echocardiography. <b>2003</b> , 3, 87-96                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |     |
| 46 | Inducible left ventricular outflow tract gradient during dobutamine stress echocardiography: an association with intraoperative hypotension but not a contraindication to liver transplantation. <b>2004</b> , 21, 681-5                                                                                                                                                                                                                                                                                                                                      | 32  |
| 45 | Methodological analysis of diagnostic dobutamine stress echocardiography studies. <b>2004</b> , 21, 725-36                                                                                                                                                                                                                                                                                                                                                                                                                                                    | 13  |
| 44 | Sonovue improves endocardial border detection and variability in assessing wall motion score and ejection fraction during stress echocardiography. <b>2004</b> , 173, 13-7                                                                                                                                                                                                                                                                                                                                                                                    | 9   |
| 43 | The transition from an analog to a digital echocardiography laboratory: the Mayo experience. <b>2004</b> , 17, 1214-24                                                                                                                                                                                                                                                                                                                                                                                                                                        | 5   |
| 42 | The validation of fractional flow reserve in patients with coronary multivessel disease: a comparison with SPECT and contrast-enhanced dobutamine stress echocardiography. <b>2005</b> , 94, 321-7                                                                                                                                                                                                                                                                                                                                                            | 13  |
| 41 | A meta-analytic comparison of echocardiographic stressors. <b>2005</b> , 21, 189-207                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | 25  |
| 40 | ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). developed in collaboration with the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society. <b>2007</b> , 116, e418-99 | 540 |
| 39 | ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). developed in collaboration with the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society. <b>2007</b> , 116, e418-99 | 409 |
| 38 | Feasibility, safety and accuracy of dobutamine/atropine stress echocardiography for the detection of coronary artery disease in renal transplant candidates. <b>2007</b> , 88, 45-51                                                                                                                                                                                                                                                                                                                                                                          | 20  |
| 37 | Stress echocardiography expert consensus statement: European Association of Echocardiography (EAE) (a registered branch of the ESC). <b>2008</b> , 9, 415-37                                                                                                                                                                                                                                                                                                                                                                                                  | 462 |
| 36 | Exercise-induced changes in E/ESratio in patients with suspected coronary artery disease. <b>2008</b> , 19, 405-11                                                                                                                                                                                                                                                                                                                                                                                                                                            | 13  |
| 35 | 2009 ACCF/AHA focused update on perioperative beta blockade incorporated into the ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American college of cardiology foundation/American heart association task force on practice guidelines. <b>2009</b> , 100, 118-27                                                                                                                                                                                                                       | 226 |
| 34 | 2009 ACCF/AHA focused update on perioperative beta blockade incorporated into the ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery. <b>2009</b> , 54, e13-e118                                                                                                                                                                                                                                                                                                                                              | 223 |
| 33 | Factors affecting sensitivity and specificity of diagnostic testing: dobutamine stress echocardiography. <b>2009</b> , 22, 1199-208                                                                                                                                                                                                                                                                                                                                                                                                                           | 46  |

|    |                                                                                                                                                                                                                                                                                                                                                |     |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 32 | The utility of dobutamine stress echocardiography for the diagnosis of coronary artery disease in the HIV population. <b>2010</b> , 27, 1228-32                                                                                                                                                                                                | 1   |
| 31 | The value of Eb blockers administration during recovery phase of dobutamine stress echocardiography: a review. <b>2013</b> , 30, 723-9                                                                                                                                                                                                         | 1   |
| 30 | Relation between ischaemic threshold measured during dobutamine stress echocardiography and coronary angiographic features. <b>2014</b> , 72, 1380-7                                                                                                                                                                                           |     |
| 29 | Strain Imaging with a Bulls-Eye Map for Detecting Significant Coronary Stenosis during Dobutamine Stress Echocardiography. <b>2017</b> , 30, 159-167.e1                                                                                                                                                                                        | 16  |
| 28 | Diagnostic Testing for Cardiovascular Diseases. <i>Developments in Cardiovascular Medicine</i> , <b>1993</b> , 73-90                                                                                                                                                                                                                           | 1   |
| 27 | Dobutamine Stress Echocardiography. <b>2015</b> , 197-214                                                                                                                                                                                                                                                                                      | 1   |
| 26 | Dobutamine Stress Echocardiography. <b>2003</b> , 115-132                                                                                                                                                                                                                                                                                      | 1   |
| 25 | Dobutamine Stress Echocardiography. <b>1997</b> , 95-102                                                                                                                                                                                                                                                                                       | 1   |
| 24 | Combined Tc-99m sestamibi first-pass radionuclide angiography and cardiac SPECT imaging during arbutamine infusion delivered by a computerized closed-loop system. <b>1999</b> , 24, 42-7                                                                                                                                                      | 1   |
| 23 | Hemodynamic changes, plasma catecholamine responses, and echocardiographically detected contractile reserve during two different dobutamine-infusion protocols. <b>1997</b> , 29, 808-13                                                                                                                                                       | 3   |
| 22 | Two-dimensional and dobutamine stress echocardiography in the preoperative assessment of patients with end-stage liver disease prior to orthotopic liver transplantation. <b>1996</b> , 61, 1180-8                                                                                                                                             | 203 |
| 21 | Dobutamine-atropine stress echocardiography in asymptomatic healthy individuals. The relativity of stress-induced hyperkinesia. <b>1995</b> , 92, 3453-63                                                                                                                                                                                      | 54  |
| 20 | Relation between ischemic threshold measured during dobutamine stress echocardiography and known indices of poor prognosis in patients with coronary artery disease. <b>1995</b> , 92, 2095-101                                                                                                                                                | 25  |
| 19 | ACC/AHA Guidelines for the Clinical Application of Echocardiography. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Clinical Application of Echocardiography). Developed in collaboration with the American Society of Echocardiography. <b>1997</b> , 95, 1686-744 | 431 |
| 18 | Fractional Flow Reserve to Distinguish Significant Stenosis: Use at Diagnostic Catherization. <i>Developments in Cardiovascular Medicine</i> , <b>2000</b> , 203-229                                                                                                                                                                           |     |
| 17 | Performance of stress echocardiography. <i>Developments in Cardiovascular Medicine</i> , <b>2003</b> , 1-42                                                                                                                                                                                                                                    |     |
| 16 | Accuracy of Stress Echocardiography. <i>Developments in Cardiovascular Medicine</i> , <b>2003</b> , 65-104                                                                                                                                                                                                                                     |     |
| 15 | Comparison of stress echo with standard stress testing. <i>Developments in Cardiovascular Medicine</i> , <b>2003</b> , 105-118                                                                                                                                                                                                                 |     |

- 14 Clinical Usefulness and Safety of Dobutamine Stress Echocardiography for the Detection of Significant Stenosis in the Infarcted Area. **2004**, 16, 275-282
- 13 Utility of Combined Assessment of Coronary Flow Velocity and Myocardial Perfusion During Low-dose Dobutamine Stress Echocardiography in the Detection of Left Anterior Coronary Artery Disease. **2006**, 4, 69-77 1
- 12 Stress Echocardiography with Nonexercise Techniques: Principles, Protocols, Interpretation, and Clinical Applications. **2007**, 353-392
- 11 Dobutamine stress echocardiography as a prognostic tool for future cardiac events. *Annals of Saudi Medicine*, **2008**, 28, 371-3 1.6
- 10 Coronary Artery Disease. **2012**, 213-224
- 9 The Prognostic Significance of Resting Regional Left Ventricular Function in Patients With Varying Degrees of Myocardial Ischemia. *Cardiology Research*, **2013**, 4, 178-185 1.8 1
- 8 New developments in pharmacological stress imaging. *Developments in Cardiovascular Medicine*, **1992**, 503-537
- 7 Stellenwert der pharmakologischen Belastungsechokardiographie. **1993**, 85-100 1
- 6 Pharmakologische Streßechokardiographie mit Dipyridamol und Dobutamin: Praktische Durchführung. **1994**, 175-190
- 5 Comparison of Exercise and Pharmacologic Stress Echocardiography and Electrocardiography. *Developments in Cardiovascular Medicine*, **1994**, 95-111
- 4 Pharmakologische Streßechokardiographie mit Dobutamin, Dipyridamol und Arbutamin: Praktische Durchführung. **1998**, 321-367
- 3 Dobutamine stress echo has marginal utility and is context-dependent for assessing myocardial ischemia. **2015**, 1, 1
- 2 Evaluating coronary artery disease noninvasively--which test for whom?. *Western Journal of Medicine*, **1994**, 161, 173-80 7
- 1 Sustained atrial fibrillation after dobutamine stress echocardiography in an older patient with left atrial enlargement. *Western Journal of Medicine*, **1995**, 162, 268-9 5