Exercise echocardiography as a screening test for coronary artery disease and correlation with coronary arteri

DOI: 10.1016/0002-9149(91)90929-f
American Journal of Cardiology, 1991, 67, 1213-8.

#	Paper	IF	Citations
191	Evolution of stress testing. *Circulation*, **1992**, 85, 1217-8	16.7	19
190	Preoperative cardiac evaluation for noncardiac surgery: a functional approach. **1992**, 74, 586-98		55
189	Echocardiography Articles Appearing in Clinical Cardiology Journals in 1991. *Journal of the American Society of Echocardiography*, **1992**, 5, 555-568	5.8	
188	Exercise echocardiography. **1992**, 22, 610-2		1
187	Exercise echocardiography: phase II, convincing the skeptics. *Journal of the American College of Cardiology*, **1992**, 19, 82-3	15.1	8
186	On-line quantification of ventricular function during dobutamine stress echocardiography. *European Heart Journal*, **1992**, 13, 1669-76	9.5	62
185	Exercise echocardiography after coronary artery bypass grafting. *American Journal of Cardiology*, **1992**, 70, 572-6	3	47
184	Sonicated serum albumin in contrast echocardiography: improved segmental wall motion depiction and implications for stress echocardiography. *American Journal of Cardiology*, **1992**, 69, 42H-45H	3	26
183	Exercise echocardiography. *American Journal of Cardiology*, **1992**, 69, 82H-89H	3	42
182	Accuracy of dobutamine stress echocardiography in detecting coronary artery disease. *American Journal of Cardiology*, **1992**, 69, 1269-73	3	286
181	The value and promise of echocardiography in acute myocardial infarction and coronary artery disease. *Clinical Cardiology*, **1992**, 15, 401-10	3.3	3
180	Echocardiography and coronary artery disease. **1993**, 9 Suppl 2, 55-67		5
179	Comparison of dobutamine and exercise echocardiography for detecting coronary artery disease. *American Journal of Cardiology*, **1993**, 72, 1226-31	3	85
178	Comparative evaluation of bicycle and dobutamine stress echocardiography with perfusion scintigraphy and bicycle electrocardiogram for identification of coronary artery disease. *American Journal of Cardiology*, **1993**, 72, 555-9	3	91
177	Prognostic usefulness of positive or negative exercise stress echocardiography for predicting coronary events in ensuing twelve months. *American Journal of Cardiology*, **1993**, 71, 646-51	3	102
176	Supine bicycle stress echocardiography versus tomographic thallium-201 exercise imaging for the detection of coronary artery disease. *Journal of the American Society of Echocardiography*, **1993**, 6, 177-85	5.8	46
175	Supine bicycle stress echocardiography: peak exercise imaging is superior to postexercise imaging. *Journal of the American Society of Echocardiography*, **1993**, 6, 265-71	5.8	57

174	Stress echocardiography: is there an optimal type of stress?. *Journal of the American Society of Echocardiography*, **1993**, 6, 198-9	5.8	2
173	Detection of coronary artery disease with upright bicycle exercise echocardiography. *Journal of the American Society of Echocardiography*, **1993**, 6, 186-97	5.8	120
172	Comparison of dobutamine stress echocardiography and stress thallium-201 single-photon emission computed tomography for detecting coronary artery disease. *Journal of the American Society of Echocardiography*, **1993**, 6, 593-602	5.8	59
171	Paradoxic hypotension during dobutamine stress echocardiography: clinical and diagnostic implications. *Journal of the American College of Cardiology*, **1993**, 21, 1080-6	15.1	89
170	Digital supine bicycle stress echocardiography: a new technique for evaluating coronary artery disease. *Journal of the American College of Cardiology*, **1993**, 21, 950-6	15.1	83
169	Myocardial perfusion imaging versus two-dimensional echocardiography: comparative value in the diagnosis of coronary artery disease. *Journal of Nuclear Cardiology*, **1994**, 1, 399-414	2.1	15
168	A consideration of current clinical options for stress imaging in the diagnosis and evaluation of coronary artery disease. *Journal of Nuclear Cardiology*, **1994**, 1, S147-70	2.1	5
167	Left and right heart Doppler stress echo in congestive heart failure. **1994**, 10, 289-97		3
166	Limited usefulness of exercise testing and thallium scintigraphy in evaluation of ambulatory patients several months after recovery from an acute coronary event: implications for management of stable coronary heart disease. Multicenter Myocardial Ischemia Research Group. *Journal of the American College of Cardiology*, **1994**, 24, 1274-81	15.1	20
165	Identification of multivessel coronary artery disease by exercise echocardiography. *Journal of the American College of Cardiology*, **1994**, 24, 109-14	15.1	90
164	Exercise echocardiography: coming of age. *Journal of the American College of Cardiology*, **1994**, 24, 115-6	15.1	12
163	Color Doppler myocardial imaging: a new technique for the assessment of myocardial function. *Journal of the American Society of Echocardiography*, **1994**, 7, 441-58	5.8	517
162	Clinical use of stress echocardiography: factors affecting diagnostic accuracy. *European Heart Journal*, **1994**, 15, 672-80	9.5	23
161	Exercise echocardiography performed early and late after percutaneous transluminal coronary angioplasty for prediction of restenosis. *European Heart Journal*, **1995**, 16, 1872-9	9.5	10
160	Functional evaluation of patients with coronary artery disease: selection of appropriate pharmacologic agents and imaging modalities. *European Heart Journal*, **1995**, 16 Suppl M, 11-6	9.5	3
159	Diagnosis of coronary artery disease and viable myocardium by stress echocardiography. Diagnostic accuracy of different stress modalities. *European Heart Journal*, **1995**, 16 Suppl J, 10-8	9.5	3
158	Diagnostic and prognostic value of stress testing in older persons. **1995**, 43, 190-4		7
157	Stress imaging. Current clinical options for the diagnosis, localization, and evaluation of coronary artery disease. **1995**, 79, 1025-61		6

156	Emerging technology in stress echocardiography. *European Heart Journal*, **1995**, 16 Suppl J, 5-9	9.5	2
155	Detection of myocardial ischemia. **1995**, 20, 773-824		3
154	Comparison of stress echocardiography and stress myocardial perfusion scintigraphy for diagnosing coronary artery disease and assessing its severity. *American Journal of Cardiology*, **1995**, 75, 25D-34D	3	54
153	Digital high frame rate stress echocardiography for detection of coronary artery stenosis by high dose dipyridamole stress testing. **1995**, 11, 163-70		5
152	The influence of angiographically demonstrated coronary collaterals on the results of stress echocardiography. *Clinical Cardiology*, **1995**, 18, 205-8	3.3	12
151	Correlation of coronary stenosis by quantitative coronary arteriography with exercise echocardiography. *American Journal of Cardiology*, **1995**, 75, 287-90	3	14
150	Comparison of digital dipyridamole stress echocardiography and upright bicycle stress echocardiography for identification of coronary artery stenosis. **1995**, 86, 514-20		22
149	Stress echocardiography. Part I. Exercise echocardiography: techniques, implementation, clinical applications, and correlations. **1995**, 70, 5-15		88
148	Application and perspectives of transesophageal stress echocardiography using monoplane and biplane instruments. *Echocardiography*, **1995**, 12, 317-24	1.5	1
147	Exercise echocardiography after coronary artery bypass surgery: correlation with coronary angiography. *Journal of the American College of Cardiology*, **1995**, 25, 1019-23	15.1	38
146	Exercise echocardiography is an accurate and cost-efficient technique for detection of coronary artery disease in women. *Journal of the American College of Cardiology*, **1995**, 26, 335-41	15.1	215
145	Influence of left ventricular hypertrophy on detection of coronary artery disease using exercise echocardiography. *Journal of the American College of Cardiology*, **1995**, 26, 1180-6	15.1	64
144	Stress Echocardiography. Part I. Exercise Echocardiography: Techniques, Implementation, Clinical Applications, and Correlations. **1995**, 70, 5-15		82
143	Safety and efficacy of computerized closed-loop delivery of arbutamine: a new pharmacologic myocardial stress modality for the assessment of coronary artery disease. The European Arbutamine Study Group. *Journal of the American Society of Echocardiography*, **1995**, 8, 854-63	5.8	3
142	Sonicated albumin in exercise echocardiography: technique and feasibility to enhance endocardial visualization. *Journal of the American Society of Echocardiography*, **1996**, 9, 462-9	5.8	20
141	Interobserver and intraobserver variation for analysis of left ventricular wall motion at baseline and during low- and high-dose dobutamine stress echocardiography in patients with high prevalence of wall motion abnormalities at rest. *Journal of the American Society of Echocardiography*, **1996**, 9, 320-8	5.8	17
140	Comparison of ischemic response during exercise and dobutamine echocardiography in patients with left main coronary artery disease. *Journal of the American College of Cardiology*, **1996**, 27, 1171-7	15.1	50
139	The importance of work-up (verification) bias correction in assessing the accuracy of SPECT thallium-201 testing for the diagnosis of coronary artery disease. **1996**, 49, 735-42		63

138	Arbutamine vs. exercise stress testing in patients with coronary artery disease: evaluation by echocardiography and electrocardiography. *International Journal of Cardiology*, **1996**, 57, 81-9	3.2	1
137	Detection of coronary artery disease in the presence of left ventricular atrophy. *International Journal of Cardiology*, **1996**, 57, 245-55	3.2	5
136	Exercise echocardiography in the diagnosis of coronary artery disease. **1996**, 28, 73-7		25
135	Stress Testing for Coronary Artery Disease in the Elderly. **1996**, 12, 101-119		6
134	Use of exercise echocardiography to evaluate patients after coronary angioplasty. *American Journal of Cardiology*, **1996**, 78, 1163-6	3	5
133	Exercise echocardiography, angiography, and intracoronary ultrasound after cardiac transplantation. *American Journal of Cardiology*, **1996**, 77, 1216-9	3	25
132	Stress echocardiography for the assessment of myocardial ischemia and viability. **1996**, 21, 445-520		15
131	Preoperative assessment and perioperative management of cardiac ischemic risk in noncardiac surgery. **1996**, 21, 291-382		9
130	The role of stress echocardiography versus stress perfusion: a view from the other side. *Journal of Nuclear Cardiology*, **1996**, 3, S66-74	2.1	11
129	Stress echocardiography for assessing myocardial ischaemia and viable myocardium. *Heart*, **1997**, 78 Suppl 1, 12-8	5.1	9
128	Treadmill exercise echocardiography: methodology and clinical role. *European Heart Journal*, **1997**, 18 Suppl D, D2-8	9.5	38
127	Amlodipine Improves the Anti-Ischaemic Effect of Atenolol in Postinfarction Patients with Effort-Induced Ischaemia. **1997**, 13, 22-28		
126	[II. Role of Doppler echocardiography in the management of chronic ischemic cardiopathy]. *Revista Espanola De Cardiologia*, **1997**, 50, 15-25	1.5	3
125	Is review of videotape necessary after review of digitized cine-loop images in stress echocardiography? A prospective study in 306 patients. *Journal of the American Society of Echocardiography*, **1997**, 10, 179-84	5.8	39
124	The Five View Technique for Stress Echocardiography: A Description and Evaluation of Segmental Imaging and Reported Angiographic Data. *Echocardiography*, **1997**, 14, 231-242	1.5	
123	Prognostic issues in stress echocardiography. *Progress in Cardiovascular Diseases*, **1997**, 39, 533-42	8.5	19
122	Comparison between dobutamine echocardiography and thallium-201 scintigraphy in detecting residual stenosis, ischemia, and necrosis in patients with prior myocardial infarction. *Clinical Cardiology*, **1997**, 20, 351-6	3.3	5
121	How does computer-assisted digital wall motion analysis influence observer agreement and diagnostic accuracy during stress echocardiography?. **1997**, 13, 105-14		10

120	The Sensitivity of Exercise Echocardiography Can Be Improved by Taking the Delayed Myocardial Contraction into Account. *Echocardiography*, **1998**, 15, 345-352	1.5	1
119	Stress Echocardiography: A Review of the Principles and Practice. *Echocardiography*, **1998**, 15, 669-692	1.5	4
118	Prognostic value of stress echocardiography in the evaluation of atypical chest pain patients without known coronary artery disease. *American Journal of Cardiology*, **1998**, 81, 545-51	3	34
117	Usefulness of exercise echocardiography in predicting cardiac events in an outpatient population. *American Journal of Cardiology*, **1998**, 82, 569-73	3	11
116	Risk stratification of patients with medically treated unstable angina using exercise echocardiography. *American Journal of Cardiology*, **1998**, 82, 720-4	3	16
115	Utility of stress echocardiography in the triage of patients with atypical chest pain from the emergency department. *American Journal of Cardiology*, **1998**, 82, 1282-4, A10	3	26
114	Color Doppler imaging of the myocardium: current status and potential clinical applications. **1998**, 24, 177-85		6
113	Critical analysis of coronary artery bypass graft surgery: a 30-year journey. *Journal of the American College of Cardiology*, **1998**, 31, 1B-63B	15.1	82
112	Stress echocardiography: recommendations for performance and interpretation of stress echocardiography. Stress Echocardiography Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. *Journal of the American Society of Echocardiography*, **1998**, 11, 97-104	5.8	307
111	Exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance. **1998**, 280, 913-20		324
110	Use of Exercise Echocardiography to Evaluate Patients With Chest Pain. *American Journal of the Medical Sciences*, **1998**, 316, 345-350	2.2	
109	Fuzzy cluster analysis of positive stress tests: comparison with stress echocardiography and nuclear perfusion imaging in patients with triple vessel and left main coronary disease. **1999**, 91, 66-8		6
108	Overall and Segmental Agreement of Stress Echocardiography. *Echocardiography*, **1999**, 16, 531-538	1.5	1
107	Advances in Exercise Echocardiography Can This Technique Still Thrive in the Era of Pharmacologic Stress Testing?. *Echocardiography*, **1999**, 16, 841-856	1.5	4
106	Diagnostic and Prognostic Use of Stress Echocardiography and Radionuclide Scintigraphy. *Echocardiography*, **1999**, 16, 857-877	1.5	1
105	Long-Term Value of Stress Echocardiography in the Triage of Patients with Atypical Chest Pain Presenting to the Emergency Department. *Echocardiography*, **1999**, 16, 171-177	1.5	13
104	Comparison of bicycle, heavy isometric, dipyridamole-atropine and dobutamine stress echocardiography for diagnosis of myocardial ischemia. *American Journal of Cardiology*, **1999**, 84, 1396-400	3	22
103	Exercise echocardiography. Principles, methods, and clinical use. **1999**, 17, 447-60, vii		10

102	Stress echocardiography in women. **1999**, 17, 573-82		13
101	Comparison of treadmill exercise echocardiography before and after exercise in the evaluation of patients with known or suspected coronary artery disease. *Journal of the American Society of Echocardiography*, **1999**, 12, 1073-9	5.8	35
100	A perspective on standardizing the predictive power of noninvasive cardiovascular tests by likelihood ratio computation: 2. Clinical applications. **1999**, 74, 1072-87		20
99	Supine bicycle versus post-treadmill exercise echocardiography in the detection of myocardial ischemia: a randomized single-blind crossover trial. *Journal of the American College of Cardiology*, **1999**, 33, 1485-90	15.1	74
98	ACC/AHA/ACP-ASIM guidelines for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Patients With Chronic Stable Angina). *Journal of the American College of Cardiology*, **1999**, 33, 2092-197	15.1	482
97	Objective evaluation of regional left ventricular wall motion during dobutamine stress echocardiographic studies using segmental analysis of color kinesis images. *Journal of the American College of Cardiology*, **1999**, 34, 409-19	15.1	71
96	Palpable cardiac impulse predicts adequate acoustic windows. *Echocardiography*, **2000**, 17, 1-6	1.5	1
95	Diagnostic and prognostic use of stress echocardiography in stable patients. *Echocardiography*, **2000**, 17, 465-77	1.5	3
94	Cost-effectiveness of stress echocardiography and nuclear perfusion imaging. *Progress in Cardiovascular Diseases*, **2000**, 43, 197-214	8.5	1
93	Accuracy of exercise echocardiography to detect coronary artery disease in left bundle branch block unassociated with either acute or healed myocardial infarction. *American Journal of Cardiology*, **2000**, 85, 890-3, A9	3	13
92	Anatomy of a meta-analysis: a critical review of "exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance". *Journal of Nuclear Cardiology*, **2000**, 7, 599-615	2.1	13
91	Current status of stress echocardiography. *Clinical Cardiology*, **2000**, 23, 242-6	3.3	5
90	Mean myocardial velocity mapping in quantifying regional myocardial contractile reserve in patients with impaired left ventricular systolic function: Doppler myocardial imaging study. *Journal of the American Society of Echocardiography*, **2000**, 13, 96-107	5.8	4
89	Influence of age on the electrophysiological mechanism of paroxysmal supraventricular tachycardias. *International Journal of Cardiology*, **2001**, 78, 293-8	3.2	31
88	Pharmacological stress agents for evaluation of ischemic heart disease. *International Journal of Cardiology*, **2001**, 81, 157-67	3.2	19
87	The clinical use of imaging techniques with exercise testing. *Primary Care - Clinics in Office Practice*, **2001**, 28, 181-98 ,viii	2.2	1
86	Stratification of single-vessel coronary stenosis by ischemic threshold at the onset of wall motion abnormality during continuous monitoring of left ventricular function by semisupine exercise echocardiography. *Journal of the American Society of Echocardiography*, **2001**, 14, 798-805	5.8	6
85	Stress echocardiography and its role in performance assessment. *Veterinary Clinics of North America Equine Practice*, **2001**, 17, 179-89, viii	1.9	18

84	Stress echocardiography: technical considerations. *Progress in Cardiovascular Diseases*, **2001**, 43, 303-14	8.5	3
83	Stress testing in the elderly. *The American Journal of Geriatric Cardiology*, **2001**, 10, 308-13; quiz 313-5		7
82	Hypertensive response to exercise: a potential cause for new wall motion abnormality in the absence of coronary artery disease. *Journal of the American College of Cardiology*, **2002**, 39, 323-7	15.1	66
81	Contribution of stress echocardiography to clinical decision making in unselected ambulatory patients with known or suspected coronary artery disease. *International Journal of Cardiology*, **2002**, 84, 179-85	3.2	5
80	Low-level exercise echocardiography identifies contractile reserve in patients with a recent myocardial infarction: comparison with dobutamine stress echocardiography. *Journal of the American Society of Echocardiography*, **2002**, 15, 671-7	5.8	2
79	Effect of left ventricular global systolic function, mitral regurgitation, and left ventricular inflow pattern on exercise echocardiography results. *Echocardiography*, **2002**, 19, 115-23	1.5	5
78	Feasibility of pulsed-Doppler tissue imaging of the interventricular septum during exercise echocardiography. *Echocardiography*, **2002**, 19, 299-305	1.5	1
77	[Exercise echocardiography to differentiate dilated cardiomyopathy from ischemic left ventricular dysfunction]. *Revista Espanola De Cardiologia*, **2003**, 56, 57-64	1.5	3
76	ACC/AHA/ASE 2003 Guideline Update for the Clinical Application of Echocardiography: Summary Article. *Journal of the American Society of Echocardiography*, **2003**, 16, 1091-1110	5.8	33
75	ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography--summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). *Journal of the American College of Cardiology*, **2003**, 42, 954-70	15.1	392
74	Clinical techniques for diagnosing cardiovascular abnormalities in performance horses. *Clinical Techniques in Equine Practice*, **2003**, 2, 266-277		6
73	Biplane stress echocardiography using a prototype matrix-array transducer. *Journal of the American Society of Echocardiography*, **2003**, 16, 937-41	5.8	29
72	ACC/AHA/ASE 2003 guideline update for the clinical application of echocardiography: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). *Circulation*, **2003**, 108, 1146-62	16.7	623
71	Accuracy of peak treadmill exercise echocardiography to detect multivessel coronary artery disease: comparison with post-exercise echocardiography. *European Journal of Echocardiography*, **2003**, 4, 182-90		28
70	Intraoperative Stress Echocardiography. *Cardiology*, **2003**, 3, 87-96		
69	Novel stress echocardiographic model incorporating the extent and severity of wall motion abnormality for risk stratification and prognosis. *American Journal of Cardiology*, **2004**, 94, 715-9	3	30
68	Échocardiographie de stress. *EMC - Cardiologie-Angeiologie*, **2004**, 1, 38-48		0
67	Detection of exercise-induced ischemia by changes in B-type natriuretic peptides. *Journal of the American College of Cardiology*, **2004**, 44, 1980-7	15.1	112

66	Relationship between the ischemic threshold at the onset of wall-motion abnormality on semisupine exercise echocardiography and the extent of coronary artery disease. *Journal of the American Society of Echocardiography*, **2004**, 17, 121-5	5.8	1
65	Prognostic value of contrast stress echocardiography in patients with image quality too limited for traditional noncontrast harmonic echocardiography. *Journal of the American Society of Echocardiography*, **2004**, 17, 15-20	5.8	5
64	Comparing stress testing methods. Available techniques and their use in CAD evaluation. *Postgraduate Medicine*, **2004**, 115, 61-70	3.7	10
63	Comparison of prognostic value of stress echocardiography versus stress electrocardiography in patients with suspected coronary artery disease. *American Journal of Cardiology*, **2005**, 96, 628-34	3	23
62	A meta-analytic comparison of echocardiographic stressors. *International Journal of Cardiovascular Imaging*, **2005**, 21, 189-207	2.5	25
61	Myocardial blood volume and perfusion reserve responses to combined dipyridamole and exercise stress: a quantitative approach to contrast stress echocardiography. *Journal of the American Society of Echocardiography*, **2005**, 18, 1187-93	5.8	18
60	Emergency nontraumatic chest pain: use of stress echocardiography to detect significant coronary artery stenosis. *Journal of the American Society of Echocardiography*, **2005**, 18, 1181-6	5.8	7
59	Stress echocardiography: current methodology and clinical applications. *Journal of the American College of Cardiology*, **2005**, 45, 1739-47	15.1	101
58	Right ventricular pressure dynamics and stress echocardiography in pharmacological and exercise stress testing. *Equine Veterinary Journal*, **2006**, 38, 183-92	2.4	11
57	E□chocardiographie de stress. *EMC - Cardiologie*, **2006**, 1, 1-7		
56	Usefulness of high functional capacity in patients with exercise-induced ST-depression to predict a negative result on exercise echocardiography and low prognostic risk. *American Journal of Cardiology*, **2008**, 101, 1541-3	3	15
55	Stress echocardiography expert consensus statement: European Association of Echocardiography (EAE) (a registered branch of the ESC). *European Journal of Echocardiography*, **2008**, 9, 415-37		462
54	Stress echocardiography from 1979 to present. *Journal of the American Society of Echocardiography*, **2008**, 21, 22-8	5.8	31
53	Arterial Hypertension and Cardiac Damage. *High Blood Pressure and Cardiovascular Prevention*, **2008** , 15, 141-170	2.9	9
52	Stress Echocardiography Expert Consensus Statement--Executive Summary: European Association of Echocardiography (EAE) (a registered branch of the ESC). *European Heart Journal*, **2009**, 30, 278-89	9.5	211
51	Gender differences in onset of symptoms in AV nodal re-entrant and accessory pathway-mediated re-entrant tachycardia. *Herzschrittmachertherapie Und Elektrophysiologie*, **2009**, 20, 33-8	0.8	11
50	Grading of Ischemic Response. **2009**, 247-258		
49	Coronary Artery Disease. **2009**, 255-317		

48	Diagnosis and Management of Coronary Artery Disease. **2010**, 286-294		
47	Independent and incremental value of stress echocardiography over clinical and stress electrocardiographic parameters for the prediction of hard cardiac events in new-onset suspected angina with no history of coronary artery disease. *European Journal of Echocardiography*, **2010**, 11, 875-82		26
46	Right ventricular pressure dynamics during exercise: relationship to stress echocardiography. *Equine Veterinary Journal*, **2002**, 34, 472-7	2.4	14
45	Stress echocardiography for the detection and assessment of coronary artery disease. *Journal of Nuclear Cardiology*, **2011**, 18, 501-15	2.1	7
44	Transthoracic detection of coronary flow in left and right coronary descending arteries during supine bicycle stress echocardiography. *Coronary Artery Disease*, **2012**, 23, 337-47	1.4	3
43	Utility of exercise stress echocardiography in pediatric cardiac transplant recipients: a single-center experience. *Journal of Heart and Lung Transplantation*, **2012**, 31, 517-23	5.8	26
42	Echocardiographic Evaluation of Coronary Artery Disease. *Cardiovascular Medicine*, **2015**, 217-252	0.1	1
41	Chronic obstructive pulmonary disease and coronary disease: COPDCoRi, a simple and effective algorithm for predicting the risk of coronary artery disease in COPD patients. *Respiratory Medicine*, **2015**, 109, 1019-25	4.6	16
40	The additive prognostic value of coronary flow velocity reserve during exercise echocardiography. *European Heart Journal Cardiovascular Imaging*, **2017**, 18, 1179-1184	4.1	13
39	The Significance of Equivocal Exercise Treadmill ECG for Intermediate Risk Chest Pain Assessment - Insight From Coronary CT Angiography Data. *Heart Lung and Circulation*, **2018**, 27, 50-57	1.8	2
38	Usefulness of Excellent Functional Capacity in Men and Women With Ischemic Exercise Electrocardiography to Predict a Negative Stress Imaging Test and Very Low Late Mortality. *American Journal of Cardiology*, **2019**, 124, 661-665	3	4
37	Management of cardiac conduction abnormalities and arrhythmia in aircrew. *Heart*, **2019**, 105, s38-s49	5.1	9
36	Effect of random deferral of percutaneous coronary intervention in patients with diabetes and stable ischaemic heart disease. *Heart*, **2020**, 106, 1651-1657	5.1	1
35	Stress Echocardiography: Instructions for Use. **1994**, 81-97		2
34	Grading of Ischemic Response. **1997**, 139-148		1
33	Exercise Echocardiography. *Developments in Cardiovascular Medicine*, **1994**, 39-59		1
32	Feasibility of exercise stress echocardiography for the follow-up of children with coronary involvement secondary to Kawasaki disease. *Circulation*, **1995**, 91, 122-8	16.7	41
31	Sex and test verification bias. Impact on the diagnostic value of exercise echocardiography. *Circulation*, **1997**, 95, 405-10	16.7	126

30	ACC/AHA Guidelines for the Clinical Application of Echocardiography. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Clinical Application of Echocardiography). Developed in collaboration with the American Society of Echocardiography. *Circulation*, **1997**, 95, 1686-744	16.7	431
29	Grading of Ischemic Response. **2003**, 189-198		
28	Accuracy of Stress Echocardiography. *Developments in Cardiovascular Medicine*, **2003**, 65-104		
27	Exercise Echocardiography. **2003**, 103-114		
26	Comparison of stress echo with standard stress testing. *Developments in Cardiovascular Medicine*, **2003**, 105-118		
25	Echocardiographic Evaluation of Coronary Artery Disease. **2007**, 811-839		
24	Exercise Echocardiography. **2007**, 326-352		
23	□hocardiographie en pr□op□atoire de chirurgie non cardiaque. **2008**, 395-415		
22	Diagnosis of Coronary Heart Diseaseinthe Elderly. *Fundamental and Clinical Cardiology*, **2008**, 243-268		
21	Stress test: a primer for primary care physicians. *Southern Medical Journal*, **2008**, 101, 806-14	0.6	
20	Stress Echocardiography Versus Radionuclide Imaging. **2010**, 159-176		
19	Coronary Artery Disease. **2012**, 213-224		
18	Stellenwert der Stressechokardiographie in der kardiologischen Rehabilitation. **1993**, 133-146		5
17	Indikationen und Differentialindikationen stre□chokardiographischer Routinetechniken. **1994**, 99-111		
16	Grading of Ischemic Response in Stress Echocardiography. **1994**, 98-109		
15	Stre□chokardiographie-Methoden. **1994**, 25-60		1
14	Comparison of Exercise and Pharmacologic Stress Echocardiography and Electrocardiography. *Developments in Cardiovascular Medicine*, **1994**, 95-111		
13	THE ROLE OF IMAGING TECHNIQUES IN STRESS TESTING. *Primary Care - Clinics in Office Practice*, **1994**, 21, 535-555	2.2	4

12	Diagnostic accuracy of stress-echocardiography for the detection of significant coronary artery disease. *Developments in Cardiovascular Medicine*, **1996**, 105-116	
11	Exercise Echocardiography. **1997**, 85-94	
10	Stellenwert der Stre echokardiographie in der kardiologischen Rehabilitation. **1998**, 155-164	
9	Methodik der Stre echokardiographie. **1998**, 101-110	
8	Stre echokardiographie-Methoden. **1998**, 35-89	
7	Use of exercise echocardiography to evaluate patients with chest pain. *American Journal of the Medical Sciences*, **1998**, 316, 345-50	2.2
6	Echocardiography in the Evaluation of Coronary Artery Disease. *Developments in Cardiovascular Medicine*, **1999**, 397-420	
5	Grading of Ischemic Response. **2015**, 291-302	
4	Cardiovascular disease in COPD. **2020**, 47-65	
3	Evaluating coronary artery disease noninvasively--which test for whom?. *Western Journal of Medicine*, **1994**, 161, 173-80	7
2	Ischemic Heart Disease. **2016**, 209-235	
1	STRESS ECHOCARDIOGRAPHY. **1997**, 51, 41-46	0