

Cocaine induces striatal c-fos-immunoreactive proteins

Proceedings of the National Academy of Sciences of the United States of America
88, 1291-1295

DOI: [10.1073/pnas.88.4.1291](https://doi.org/10.1073/pnas.88.4.1291)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Basal ganglia input, neural activity, and relation to the cortex. <i>Current Opinion in Neurobiology</i> , 1991, 1, 644-651.	2.0	68
2	Dynamic gene regulation in striatal neurons: perspectives in memorial to Dr. Kyoji Maekawa. <i>Neuroscience Research</i> , 1991, 12, 41-44.	1.0	0
3	Stress Selectively Increases Fos Protein in Dopamine Neurons Innervating the Prefrontal Cortex. <i>Cerebral Cortex</i> , 1991, 1, 273-292.	1.6	136
4	Sexual behavior increases c-fos expression in the forebrain of the male rat. <i>Brain Research</i> , 1991, 564, 352-357.	1.1	162
5	The molecular biology of addictive drugs. <i>Molecular Neurobiology</i> , 1991, 5, 45-58.	1.9	31
6	D1-dopamine receptors activate c-fos expression in the fetal suprachiasmatic nuclei.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 9201-9204.	3.3	93
7	Differential expression of c-fos and zif268 in rat striatum after haloperidol, clozapine, and amphetamine.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 4270-4274.	3.3	314
8	Transduction of psychosocial stress into the neurobiology of recurrent affective disorder. <i>American Journal of Psychiatry</i> , 1992, 149, 999-1010.	4.0	1,598
9	Regulation of immediate early gene expression and AP-1 binding in the rat nucleus accumbens by chronic cocaine.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 5764-5768.	3.3	421
10	Preproenkephalin promoter "cassette" confers brain expression and synaptic regulation in transgenic mice.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 2345-2349.	3.3	27
11	Activation of the zinc finger encoding gene krox-20 in adult rat brain: comparison with zif268. <i>Molecular Brain Research</i> , 1992, 13, 263-266.	2.5	49
12	In situ DNA-protein binding: A novel method for detecting DNA-binding activity of transcription factor in brain. <i>Neuroscience Letters</i> , 1992, 146, 25-28.	1.0	17
13	Fos-lacZ transgenic mice: Mapping sites of gene induction in the central nervous system. <i>Neuron</i> , 1992, 8, 13-23.	3.8	239
14	Late onset and long-lasting suppressive effects of ceruletide, an analogue of cholecystokinin, on c-fos mRNA expression in the rat striatum. <i>Neuroscience Letters</i> , 1992, 138, 233-236.	1.0	4
15	Conditioned Sensitization to the Psychomotor Stimulant Cocaine. <i>Annals of the New York Academy of Sciences</i> , 1992, 654, 386-399.	1.8	92
16	A 35 kDa Fos-related antigen is co-localized with substance P and dynorphin in striatal neurons. <i>Brain Research</i> , 1992, 577, 312-317.	1.1	28
17	Expression of c-fos mRNA in acute and kindled cocaine seizures in rats. <i>Brain Research</i> , 1992, 582, 101-106.	1.1	33
18	Differential effects of intermittent or continuous exposure to cocaine on the hypothalamic-pituitary-adrenal axis and c-fos expression. <i>Brain Research</i> , 1992, 571, 204-211.	1.1	61

#	ARTICLE	IF	CITATIONS
19	C-fos immunoreactivity in the brain following unilateral electrical stimulation of the dorsal periaqueductal gray in freely moving rats. <i>Brain Research</i> , 1992, 573, 276-283.	1.1	47
20	Synergistic Interactions of D1- and D2-Selective Dopamine Agonists in Animal Models for Parkinson's Disease: Sites of Action and Implications for the Pathogenesis of Dyskinesias. <i>Canadian Journal of Neurological Sciences</i> , 1992, 19, 147-152.	0.3	21
21	Both phasic sensory stimulation and tonic pharmacological activation increase Fos-like immunoreactivity in the rat locus coeruleus. <i>Synapse</i> , 1992, 12, 112-118.	0.6	14
22	Characterization of Gene Organization and Promoter Region of the Rat Dopamine D1 Receptor Gene. <i>Journal of Neurochemistry</i> , 1992, 59, 1875-1883.	2.1	69
23	D1 Dopamine Receptor Activation of Multiple Transcription Factor Genes in Rat Striatum. <i>Journal of Neurochemistry</i> , 1992, 58, 1420-1426.	2.1	193
24	Colocalization of Fos- and Glucocorticoid Receptor-Like Immunoreactivities in the Rat Amygdaloid Complex After Immobilization Stress. <i>Journal of Neuroendocrinology</i> , 1992, 4, 547-555.	1.2	23
25	Opposite effects of rough and gentle handling with repeated saline administration on c-fos mRNA expression in the rat brain. <i>Journal of Neural Transmission</i> , 1992, 90, 163-169.	1.4	31
26	Muscarinic cholinergic receptor-mediated modulation on striatal c-fos mRNA expression induced by levodopa in rat brain. <i>Journal of Neural Transmission</i> , 1992, 90, 171-181.	1.4	19
27	Developmental regulation of Fos and Fos-related antigens in cerebral cortex, striatum, hippocampus, and cerebellum of the rat. <i>Journal of Comparative Neurology</i> , 1993, 334, 75-85.	0.9	52
28	Selective induction of Fos and FRA immunoreactivity within the mesolimbic and mesostriatal dopamine terminal fields. <i>Synapse</i> , 1993, 13, 251-263.	0.6	69
29	Distribution of dopamine- and cAMP-dependent phosphoprotein (DARPP-32) in the developing and mature kidney. <i>Kidney International</i> , 1993, 44, 495-502.	2.6	13
30	Transection of Corticostriatal Afferents Reduces Amphetamine- and Apomorphine-induced Striatal Fos Expression and Turning Behaviour in Unilaterally 6-Hydroxydopamine-lesioned Rats. <i>European Journal of Neuroscience</i> , 1993, 5, 1062-1070.	1.2	85
31	Fos Immunoreactivity after Stimulation or Inhibition of Muscarinic Receptors Indicates Anatomical Specificity for Cholinergic Control of Striatal Efferent Neurons and Cortical Neurons in the Rat. <i>European Journal of Neuroscience</i> , 1993, 5, 1218-1225.	1.2	73
32	Kainate-Induced Changes in Opioid Peptide Genes and AP-1 Protein Expression in the Rat Hippocampus. <i>Journal of Neurochemistry</i> , 1993, 60, 204-211.	2.1	63
33	Co-localization of peptide-like immunoreactivities with glucocorticoid receptor- and Fos-like immunoreactivities in the rat parabrachial nucleus. <i>Brain Research</i> , 1993, 615, 245-251.	1.1	23
34	Colocalization of Fos- and glucocorticoid receptor-immunoreactivities is present only in a very restricted population of dorsal horn neurons of the rat spinal cord after nociceptive stimulation. <i>Brain Research</i> , 1993, 632, 334-338.	1.1	34
35	c-fos and NGFI-A mRNA of rat retina: evidence for light-induced augmentation and a role for cholinergic and glutamate receptors. <i>Brain Research</i> , 1993, 631, 77-82.	1.1	23
36	Sensitization of c-fos expression in rat striatum following multiple challenges with d-amphetamine. <i>Brain Research</i> , 1993, 603, 125-128.	1.1	30

#	ARTICLE	IF	CITATIONS
37	Destruction of the nigrostriatal pathway increases Fos-like immunoreactivity predominantly in striatopallidal neurons. <i>Molecular Brain Research</i> , 1993, 19, 156-160.	2.5	40
38	Elevated basal AP-1 DNA binding activity in developing rat brain. <i>Molecular Brain Research</i> , 1993, 19, 349-352.	2.5	25
39	Acute nicotine injections induce c-fos mostly in non-dopaminergic neurons of the midbrain of the rat. <i>Molecular Brain Research</i> , 1993, 20, 162-170.	2.5	53
40	Brain transcription factor expression: effects of acute and chronic amphetamine and injection stress. <i>Molecular Brain Research</i> , 1993, 20, 91-100.	2.5	102
41	Induction of immediate early genes by cyclic AMP in primary cultures of neurons from rat cerebral cortex. <i>Molecular Brain Research</i> , 1993, 19, 76-82.	2.5	44
42	Chapter 13 Neurochemically specialized projection neurons of the striatum respond differentially to psychomotor stimulants. <i>Progress in Brain Research</i> , 1993, 99, 201-205.	0.9	23
43	Chapter 9 Dopamine receptors. <i>New Comprehensive Biochemistry</i> , 1993, 24, 251-265.	0.1	2
44	Striatal Fos expression is indicative of dopamine D1/D2 synergism and receptor supersensitivity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 7451-7455.	3.3	158
45	Coordinate expression of c-fos and jun B is induced in the rat striatum by cocaine. <i>Journal of Neuroscience</i> , 1993, 13, 423-433.	1.7	145
46	Influence of Male Rats on the Luteinizing Hormone-Releasing Hormone Neuronal System in Female Rats: Role of the Vomeronasal Organ. <i>Neuroendocrinology</i> , 1993, 57, 898-906.	1.2	34
47	Chronic electroconvulsive seizure (ECS) treatment results in expression of a long-lasting AP-1 complex in brain with altered composition and characteristics. <i>Journal of Neuroscience</i> , 1994, 14, 4318-4328.	1.7	159
48	Amphetamine regulates gene expression in rat striatum via transcription factor CREB. <i>Journal of Neuroscience</i> , 1994, 14, 5623-5634.	1.7	346
49	Pharmacological regulation of AP-1 transcription factor DNA binding activity 1. <i>FASEB Journal</i> , 1994, 8, 475-478.	0.2	63
50	Molecular Neurobiology of Drug Addiction. <i>Neuropsychopharmacology</i> , 1994, 11, 77-87.	2.8	99
51	Pitfalls in Assessment of c-fos mRNA Expression in the Brain: Effects of Animal Handling. <i>Reviews in the Neurosciences</i> , 1994, 5, 171-8.	1.4	26
52	Cocaine behavioral sensitization and the excitatory amino acids. <i>Psychopharmacology</i> , 1994, 115, 305-310.	1.5	136
53	Acute and chronic cocaine administration differentially alters striatal opioid and nuclear transcription factor mRNAs. <i>Synapse</i> , 1994, 18, 35-45.	0.6	119
54	Acute administration of alcohol blocks cocaine-induced striatal c-fos immunoreactivity protein in the rat. <i>Synapse</i> , 1994, 18, 161-167.	0.6	20

#	ARTICLE	IF	CITATIONS
55	Amphetamine- and cocaine-induced fos in the rat striatum depends on D2 dopamine receptor activation. <i>Synapse</i> , 1994, 18, 233-240.	0.6	78
56	Tactile sensory input regulates basal and apomorphine-induced immediate-early gene expression in rat barrel cortex. <i>Journal of Comparative Neurology</i> , 1994, 344, 297-304.	0.9	55
57	NMDA receptor-mediated expression of Fos protein in the rat striatum following methamphetamine administration: relation to behavioral sensitization. <i>Brain Research</i> , 1994, 665, 135-140.	1.1	43
58	Dopamine D1 receptor mutant mice are deficient in striatal expression of dynorphin and in dopamine-mediated behavioral responses. <i>Cell</i> , 1994, 79, 729-742.	13.5	474
59	Persistent reduction of immediate early gene mRNA in rat forebrain following single or multiple doses of cocaine. <i>Molecular Brain Research</i> , 1994, 26, 106-112.	2.5	65
60	Coexpression of c-fos and hsp70 mRNAs in gerbil brain after ischemia: Induction threshold, distribution and time course evaluated by in situ hybridization. <i>Molecular Brain Research</i> , 1994, 26, 249-258.	2.5	66
61	Induction of multiple immediate early genes in rat hypothalamic paraventricular nucleus after stress. <i>Molecular Brain Research</i> , 1994, 25, 234-241.	2.5	72
62	Differential induction of Fos protein and a Fos-related antigen following acute and repeated cocaine administration. <i>Molecular Brain Research</i> , 1994, 25, 168-172.	2.5	66
63	Comparison between normal developing striatum and developing striatal grafts using drug-induced fos expression and neuron-specific enolase immunohistochemistry. <i>Neuroscience</i> , 1994, 60, 399-415.	1.1	18
64	Cortical stimulation induces fos expression in intrastriatal striatal grafts. <i>Brain Research</i> , 1994, 652, 87-97.	1.1	25
65	Induction of c-fos in rat brain by acute cocaine and fenfluramine exposure: a comparison study. <i>Brain Research</i> , 1994, 647, 1-9.	1.1	46
66	Induction of c-fos and CRF mRNA by MK-801 in the parvocellular paraventricular nucleus of the rat hypothalamus. <i>Molecular Brain Research</i> , 1994, 24, 192-198.	2.5	32
67	Interactive effects of stimulation of D1 and D2 dopamine receptors on fos-like immunoreactivity in the normosensitive rat striatum. <i>Brain Research Bulletin</i> , 1994, 35, 85-91.	1.4	58
68	Morphine induces c-fos and junB in striatum and nucleus accumbens via D1 and N-methyl-D-aspartate receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 8537-8541.	3.3	175
69	Sphingosylphosphocholine, a signaling molecule which accumulates in Niemann-Pick disease type A, stimulates DNA-binding activity of the transcription activator protein AP-1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 5885-5889.	3.3	45
70	Screening the dopamine D1 receptor gene in 131 schizophrenics and eight alcoholics: Identification of polymorphisms but lack of functionally significant sequence changes. <i>American Journal of Medical Genetics Part A</i> , 1995, 60, 165-171.	2.4	45
71	Comparison of c-jun, junB, and junD mRNA expression and protein in the rat dorsal root ganglia following sciatic nerve transection. <i>Journal of Neuroscience Research</i> , 1995, 42, 391-401.	1.3	31
72	Fos: An immediate-early transcription factor in neurons. <i>Journal of Neurobiology</i> , 1995, 26, 403-412.	3.7	329

#	ARTICLE	IF	CITATIONS
73	Postnatal age defines specificity of immediate early gene induction by cocaine in developing rat brain. <i>Journal of Comparative Neurology</i> , 1995, 351, 27-40.	0.9	45
74	Substance P phenotype defines specificity of c-fos induction by cocaine in developing rat striatum. <i>Journal of Comparative Neurology</i> , 1995, 351, 41-50.	0.9	45
75	Dynorphin opioid inhibition of cocaine-induced, D1 dopamine receptor-mediated immediate-early gene expression in the striatum. <i>Journal of Comparative Neurology</i> , 1995, 353, 200-212.	0.9	96
76	Cocaine and d-amphetamine increase c-fos expression in the rat cerebellum. <i>Synapse</i> , 1995, 19, 29-36.	0.6	25
77	Brain transcription factor gene expression, neurotransmitter levels, and novelty response behaviors: Alterations during rat amphetamine withdrawal and following chronic injection stress. <i>Synapse</i> , 1995, 19, 212-227.	0.6	56
78	Time-dependent changes in dopamine agonist-induced striatal fos immunoreactivity are related to sensory neglect and its recovery after unilateral prefrontal cortex injury. <i>Synapse</i> , 1995, 20, 305-315.	0.6	38
79	Cholecystokinin octapeptide and the D2 antagonist raclopride induce Fos-like immunoreactivity in the shell part of the rat nucleus accumbens via different mechanisms. <i>Brain Research</i> , 1995, 684, 225-229.	1.1	7
80	Alterations in immediate-early gene proteins in the rat forebrain induced by acute morphine injection. <i>Brain Research</i> , 1995, 692, 23-40.	1.1	82
81	Amphetamine induction of c-fos in the nucleus accumbens is not inhibited by glutamate antagonists. <i>Brain Research</i> , 1995, 694, 299-307.	1.1	24
82	Neuronal regulation of c-fos, c-jun, and junB immediate-early genes in rat adrenal medulla. <i>Journal of Neuroscience</i> , 1995, 15, 1854-1868.	1.7	29
83	D1 and D2 dopamine receptor function in the striatum: coactivation of D1- and D2-dopamine receptors on separate populations of neurons results in potentiated immediate early gene response in D1-containing neurons. <i>Journal of Neuroscience</i> , 1995, 15, 8167-8176.	1.7	252
84	PCR differential display identifies a rat brain mRNA that is transcriptionally regulated by cocaine and amphetamine. <i>Journal of Neuroscience</i> , 1995, 15, 2471-2481.	1.7	620
85	Circadian Expression of Transcription Factor Fra-2 in the Rat Pineal Gland. <i>Journal of Biological Chemistry</i> , 1995, 270, 27319-27325.	1.6	90
86	Dopamine Regulation of Transcription Factor-Target Interactions in Rat Striatum. <i>Chemical Senses</i> , 1995, 20, 257-260.	1.1	26
87	Review : Molecular Basis of Addictive States. <i>Neuroscientist</i> , 1995, 1, 212-220.	2.6	16
88	Neuronal adaptation to amphetamine and dopamine: Molecular mechanisms of prodynorphin gene regulation in rat striatum. <i>Neuron</i> , 1995, 14, 813-823.	3.8	342
89	D1-class dopamine receptor involvement in the behavioral recovery from prefrontal cortical injury. <i>Behavioural Brain Research</i> , 1995, 72, 39-48.	1.2	20
90	Dopaminergic control of gene transcription during striatal ontogeny: c-fos induction by D1 receptor activation in the developing striosomes. <i>Molecular Brain Research</i> , 1995, 30, 223-232.	2.5	23

#	ARTICLE	IF	CITATIONS
91	The human D1A dopamine receptor gene promoter directs expression of a reporter gene to the central nervous system in transgenic mice. <i>Molecular Brain Research</i> , 1995, 30, 336-346.	2.5	12
92	Basal expression of 35 kDa fos-related antigen in olfactory bulb. <i>Molecular Brain Research</i> , 1995, 34, 161-165.	2.5	4
93	Differential expression of inducible transcription factors in basal ganglia neurons. <i>Molecular Brain Research</i> , 1995, 34, 294-302.	2.5	26
94	Pretreatment with a dopamine D1-receptor antagonist prevents metabolic activation by cocaine. <i>Neuroscience Letters</i> , 1995, 196, 161-164.	1.0	5
95	Novel Dopamine Receptor Subtypes as Targets for Antipsychotic Drugs. <i>Annals of the New York Academy of Sciences</i> , 1995, 757, 278-292.	1.8	16
96	Stress, Conditioning, and the Temporal Aspects of Affective Disorders. <i>Annals of the New York Academy of Sciences</i> , 1995, 771, 677-696.	1.8	35
97	The role of dopamine in drug abuse viewed from the perspective of its role in motivation. <i>Drug and Alcohol Dependence</i> , 1995, 38, 95-137.	1.6	605
98	Molecular Mechanisms of Drug Reinforcement and Addiction. <i>Annual Review of Neuroscience</i> , 1995, 18, 463-495.	5.0	551
99	Roles of dopamine D1 receptors in striatal fos protein induction associated with methamphetamine behavioral sensitization in rats. <i>Brain Research Bulletin</i> , 1995, 38, 393-397.	1.4	21
100	NMDA and D1 receptors mediate induction of c-fos and junB genes in striatum following morphine administration: implications for studies of memory. <i>Behavioural Brain Research</i> , 1995, 66, 225-230.	1.2	61
101	C-fos expression in the rat brain after pharmacological stimulation of the rat <i>œ</i> mediodorsal <i>œ</i> thalamus by means of microdialysis. <i>Neuroscience</i> , 1995, 66, 115-131.	1.1	21
102	Synthesis and Biological Evaluation of Certain Alkenyldiarylmethanes as Anti-HIV-1 Agents Which Act as Non-Nucleoside Reverse Transcriptase Inhibitors. <i>Journal of Medicinal Chemistry</i> , 1996, 39, 3217-3227.	2.9	48
103	Novel Transcription Factors Are Induced by Chronic Cocaine Treatment. <i>Annals of the New York Academy of Sciences</i> , 1996, 801, 1-12.	1.8	9
104	Antisense oligodeoxynucleotide attenuates in vivo expression of c-fos in the paraventricular hypothalamic nucleus of the rat brain. <i>Neuroscience Letters</i> , 1996, 209, 89-92.	1.0	8
105	A nitric oxide synthesis inhibitor (L-NAME) reduces licking behavior and Fos-labeling in the spinal cord of rats during formalin-induced inflammation. <i>Pain</i> , 1996, 66, 331-341.	2.0	85
106	Changes in striatal immediate early gene expression following neonatal dopaminergic lesion and effects of intrastriatal dopaminergic transplants. <i>Neuroscience</i> , 1996, 73, 145-159.	1.1	11
107	D1 dopamine receptor-deficient mouse: cocaine-induced regulation of immediate-early gene and substance P expression in the striatum. <i>Neuroscience</i> , 1996, 74, 813-823.	1.1	143
108	The effects of D1 or D2 dopamine receptor blockade on zif/268 and preprodynorphin gene expression in rat forebrain following a short-term cocaine binge. <i>Molecular Brain Research</i> , 1996, 35, 237-248.	2.5	49

#	ARTICLE	IF	CITATIONS
109	Fenfluramine-induced activation of the immediate-early gene c-fos in the striatum: possible interaction between serotonin and dopamine. <i>Molecular Brain Research</i> , 1996, 37, 105-115.	2.5	40
110	Transcription factors in primary glial cultures: changes with neuronal interactions. <i>Molecular Brain Research</i> , 1996, 37, 224-230.	2.5	21
111	Individual and combined effects of ethanol and cocaine on intracellular signals and gene expression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1996, 20, 561-596.	2.5	31
112	Cardiovascular hypertrophy and increased vascular contractile responsiveness following repeated cocaine administration in rabbits. <i>Life Sciences</i> , 1996, 58, 675-682.	2.0	13
113	Activation of in the brain. <i>Progress in Neurobiology</i> , 1996, 50, 83-107.	2.8	610
114	Cocaethylene stimulates the secretion of ACTH and corticosterone and the transcriptional activation of hypothalamic NGFI-B. <i>Molecular Brain Research</i> , 1996, 43, 225-232.	2.5	12
115	Network-Level Changes in Expression of Inducible Fos/Jun Proteins in the Striatum during Chronic Cocaine Treatment and Withdrawal. <i>Neuron</i> , 1996, 17, 147-156.	3.8	256
116	Methamphetamine-Induced Changes in AP-1 Binding and Dynorphin in the Striatum: Correlated, Not Causally Related Events?. <i>NeuroSignals</i> , 1996, 5, 317-333.	0.5	6
117	Influence of Cocaine on the JAK/STAT Pathway in the Mesolimbic Dopamine System. <i>Journal of Neuroscience</i> , 1996, 16, 8019-8026.	1.7	50
118	D1-class dopamine receptors influence cocaine-induced persistent expression of Fos-related proteins in striatum. <i>NeuroReport</i> , 1996, 8, 1-5.	0.6	55
119	Potential brain neuronal targets for amphetamine-, methylphenidate-, and modafinil-induced wakefulness, evidenced by c-fos immunocytochemistry in the cat. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 14128-14133.	3.3	294
120	Cellular responses to psychomotor stimulant and neuroleptic drugs are abnormal in mice lacking the D1 dopamine receptor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 14928-14933.	3.3	178
121	Amphetamine and Dopamine-Induced Immediate Early Gene Expression in Striatal Neurons Depends on Postsynaptic NMDA Receptors and Calcium. <i>Journal of Neuroscience</i> , 1996, 16, 4231-4239.	1.7	280
122	Developmental expression of Fos-lacZ in the brains of postnatal transgenic rats. <i>Developmental Brain Research</i> , 1996, 93, 191-197.	2.1	14
123	Nicotine injections into the ventral tegmental area increase locomotion and Fos-like immunoreactivity in the nucleus accumbens of the rat. <i>Brain Research</i> , 1996, 730, 133-142.	1.1	96
124	Locating Reward Cue at Response Manipulandum (CAM) Induces Symptoms of Drug Abuse. <i>Neuroscience and Biobehavioral Reviews</i> , 1996, 20, 505-535.	2.9	129
125	Combined effects of ethanol and cocaine on FOS-like protein and cocaethylene biosynthesis in the rat. <i>Psychopharmacology</i> , 1996, 128, 105-114.	1.5	7
126	Psychomotor stimulant- and opiate-induced c-fos mRNA expression patterns in the rat forebrain: Comparisons between acute drug treatment and a drug challenge in sensitized animals. <i>Neurochemical Research</i> , 1996, 21, 1425-1435.	1.6	105

#	ARTICLE	IF	CITATIONS
127	Development of striatal dopaminergic function. II: Dopaminergic regulation of transcription of the immediate early gene zif268 and of D1 (D1a) and D2 (D2a) receptors during pre- and postnatal development. <i>Developmental Brain Research</i> , 1996, 94, 121-132.	2.1	20
128	Increased expression of neuronal nitric oxide synthase in bladder afferent pathways following chronic bladder irritation. , 1996, 370, 191-202.		111
129	Atypical and typical neuroleptic treatments induce distinct programs of transcription factor expression in the striatum. , 1996, 374, 70-83.		95
130	Dynorphin regulates D1 dopamine receptor-mediated responses in the striatum: Relative contributions of pre- and postsynaptic mechanisms in dorsal and ventral striatum demonstrated by altered immediate-early gene induction. , 1996, 376, 530-541.		72
131	Dopamine receptors mediate differential morphological effects on cerebral cortical neurons in vitro. , 1996, 43, 439-453.		107
132	Involvement of adenosine and glutamate receptors in the induction of c-fos in the striatum by haloperidol. , 1996, 22, 70-77.		82
133	Frontal cortex ablation reversibly decreases striatal zif/268 and junB expression: Temporal correspondence with sensory neglect and its spontaneous recovery. , 1996, 22, 291-303.		22
134	Chronic Alterations in Dopaminergic Neurotransmission Produce a Persistent Elevation of FosB-like Protein(s) in both the Rodent and Primate Striatum. <i>European Journal of Neuroscience</i> , 1996, 8, 365-381.	1.2	178
135	Phenotype of Striatal Cells Expressing c-Fos Following Amphetamine Treatment of Rats with Intrastratial Dopaminergic Grafts. <i>European Journal of Neuroscience</i> , 1996, 8, 2521-2529.	1.2	4
136	Chapter II The basal ganglia. <i>Handbook of Chemical Neuroanatomy</i> , 1996, , 371-468.	0.3	202
137	Initiation and adaptation: a paradigm for understanding psychotropic drug action. <i>American Journal of Psychiatry</i> , 1996, 153, 151-162.	4.0	442
138	Two Proximal Activating Protein-1-Binding Sites Are Sufficient to Stimulate Transcription of the Ovine Follicle-Stimulating Hormone- β Gene*. <i>Endocrinology</i> , 1997, 138, 2621-2631.	1.4	55
139	A single dose of kainic acid elevates the levels of enkephalins and activator protein-1 transcription factors in the hippocampus for up to 1 year. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 9422-9427.	3.3	62
140	Cycloheximide phase-shifts, but does not prevent, de novo Krox-24 protein expression. <i>NeuroReport</i> , 1997, 8, 3263-3266.	0.6	38
141	5-Hydroxytryptamine _{1B} Receptors Modulate the Effect of Cocaine on c-fos Expression: Converging Evidence Using 5-Hydroxytryptamine _{1B} Knockout Mice and the 5-Hydroxytryptamine _{1B/1D} Antagonist GR127935. <i>Molecular Pharmacology</i> , 1997, 51, 755-763.	1.0	90
142	FosB mutant mice: Loss of chronic cocaine induction of Fos-related proteins and heightened sensitivity to cocaine's psychomotor and rewarding effects. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 10397-10402.	3.3	274
143	Common Neural Substrates for the Addictive Properties of Nicotine and Cocaine. <i>Science</i> , 1997, 275, 83-86.	6.0	376
144	Expression of Fos, Jun, and Krox Family Proteins in Alzheimer's Disease. <i>Experimental Neurology</i> , 1997, 147, 316-332.	2.0	84

#	ARTICLE	IF	CITATIONS
145	A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. <i>Brain Research Reviews</i> , 1997, 25, 192-216.	9.1	1,122
146	Neural mechanisms of tolerance to the effects of cocaine. <i>Behavioural Brain Research</i> , 1997, 84, 225-239.	1.2	41
147	Changes in expression of 35 S-FosB and the Fos family proteins following NMDA receptor activation in the rat striatum. <i>Molecular Brain Research</i> , 1997, 47, 31-43.	2.5	9
148	Enhanced serum response element binding activity correlates with down-regulation of c-fos mRNA expression in the rat brain following repeated cortical lesions. <i>Molecular Brain Research</i> , 1997, 52, 62-70.	2.5	5
149	Dopaminergic function in the neostriatum and nucleus accumbens of young and aged fischer 344 rats. <i>Neurobiology of Aging</i> , 1997, 18, 57-66.	1.5	29
150	Modulation of c-fos expression in the rat striatum by diazepam. <i>Neuroscience Letters</i> , 1997, 236, 5-8.	1.0	15
151	Ventral pallidum self-stimulation induces stimulus dependent increase in c-fos expression in reward-related brain regions. <i>Neuroscience</i> , 1997, 77, 175-186.	1.1	60
152	Temporal pattern of AP-1 DNA-binding activity in the rat hippocampus following a kindled seizure. <i>Neuroscience</i> , 1997, 80, 753-761.	1.1	10
153	Dopaminergic and Glutamatergic Mechanisms Mediate the Induction of FOS-Like Protein by Cocaethylene. <i>Brain Research Bulletin</i> , 1997, 42, 393-398.	1.4	12
154	Effect of Pertussis Toxin Injected Into the Ventral Tegmental Area on Amphetamine-Induced c-Fos in the Nucleus Accumbens. <i>Brain Research Bulletin</i> , 1997, 43, 435-439.	1.4	1
155	Systemic Morphine-Induced Fos Protein in the Rat Striatum and Nucleus Accumbens Is Regulated by $\frac{1}{4}$ Opioid Receptors in the Substantia Nigra and Ventral Tegmental Area. <i>Journal of Neuroscience</i> , 1997, 17, 8596-8612.	1.7	111
156	NAC-1, a Rat Brain mRNA, Is Increased in the Nucleus Accumbens Three Weeks after Chronic Cocaine Self-Administration. <i>Journal of Neuroscience</i> , 1997, 17, 6864-6871.	1.7	82
157	Dopamine-Adenosine Interactions in the Striatum and the Globus Pallidus: Inhibition of Striatopallidal Neurons through Either D ₂ or A _{2A} Receptors Enhances D ₁ Receptor-Mediated Effects on c-fos Expression. <i>Journal of Neuroscience</i> , 1997, 17, 8038-8048.	1.7	123
158	Glutamate, But Not Dopamine, Stimulates Stress-Activated Protein Kinase and AP-1-Mediated Transcription in Striatal Neurons. <i>Journal of Neuroscience</i> , 1997, 17, 3455-3466.	1.7	156
159	Absence of a Persistently Elevated 37 kDa Fos-Related Antigen and AP-1-Like DNA-Binding Activity in the Brains of Kainic Acid-Treated c-fos/B Null Mice. <i>Journal of Neuroscience</i> , 1997, 17, 5407-5415.	1.7	56
160	Increased Expression of NGFI-A mRNA in the Rat Striatum Following Burst Stimulation of the Medial Forebrain Bundle. <i>European Journal of Neuroscience</i> , 1997, 9, 2370-2382.	1.2	44
161	Genetics, Ethanol and the Fos Response: A Comparison of the C57BL/6J and DBA/2J Inbred Mouse Strains. <i>Alcoholism: Clinical and Experimental Research</i> , 1997, 21, 1497-1507.	1.4	58
162	Chronic Nicotine Enhances Basal and Nicotine-Induced Fos Immunoreactivity Preferentially in the Medial Prefrontal Cortex of the Rat. <i>Neuropsychopharmacology</i> , 1997, 17, 151-161.	2.8	48

#	ARTICLE	IF	CITATIONS
163	Modulatory role of catecholamines in the transsynaptic expression of c-fos in the rat medial prefrontal cortex induced by disinhibition of the mediodorsal thalamus: a study employing microdialysis and immunohistochemistry. <i>Brain Research</i> , 1997, 749, 214-225.	1.1	18
164	Desensitization of Fos protein induction in rat striatum and nucleus accumbens following repeated administration of δ^9 -tetrahydrocannabinol. <i>Brain Research</i> , 1997, 763, 137-140.	1.1	6
165	Characterization of the long-lasting activator protein-1 complex induced by kainic acid treatment. <i>Brain Research</i> , 1997, 770, 53-59.	1.1	18
166	Protein kinases A and C are involved in the mechanisms underlying consolidation of cocaine place conditioning. <i>Brain Research</i> , 1997, 775, 30-36.	1.1	84
167	SCH 23390-induced behavioral supersensitivity is not related to striatal c-fos levels. <i>Brain Research</i> , 1997, 774, 51-54.	1.1	0
168	Widespread expression of functional D1-dopamine receptors in fetal rat brain. <i>Developmental Brain Research</i> , 1997, 102, 105-115.	2.1	34
169	Treadmill running induces striatal Fos expression via NMDA glutamate and dopamine receptors. <i>Experimental Brain Research</i> , 1997, 115, 458-469.	0.7	61
170	NMDA and D1 receptors regulate the phosphorylation of CREB and the induction of c-fos in striatal neurons in primary culture. , 1997, 25, 227-233.		91
171	Regulation of c-Fos and NGF1-A by antidepressant treatments. <i>Synapse</i> , 1997, 25, 313-320.	0.6	63
172	Increased vulnerability to cocaine in mice lacking the serotonin-1B receptor. <i>Nature</i> , 1998, 393, 175-178.	13.7	309
173	Frizzled signalling controls orientation of asymmetric sense organ precursor cell divisions in <i>Drosophila</i> . <i>Nature</i> , 1998, 393, 178-181.	13.7	207
174	Altered synaptic physiology and reduced susceptibility to kainate-induced seizures in GluR6-deficient mice. <i>Nature</i> , 1998, 392, 601-605.	13.7	450
175	Repeated Alcohol Administration Differentially Affects c-Fos and FosB Protein Immunoreactivity in DBA/2J Mice. <i>Alcoholism: Clinical and Experimental Research</i> , 1998, 22, 1646-1654.	1.4	48
176	Cocaine and the AP-1 Transcription Factor Complex. <i>Annals of the New York Academy of Sciences</i> , 1998, 844, 1-6.	1.8	42
177	Cocaine-Induced Activation of c-fos Gene Expression Is Attenuated in Prenatal Cocaine-Exposed Rabbits. <i>Annals of the New York Academy of Sciences</i> , 1998, 846, 368-370.	1.8	1
178	Stimulant-mediated c-fos induction in striatum as a function of age, sex, and prenatal cocaine exposure. <i>Brain Research</i> , 1998, 794, 88-95.	1.1	24
179	Behavioral sensitization to cocaine after a brief social stress is accompanied by changes in Fos expression in the murine brainstem. <i>Brain Research</i> , 1998, 810, 200-210.	1.1	57
180	Neural substrate of nicotine addiction as defined by functional brain maps of gene expression. <i>Journal of Physiology (Paris)</i> , 1998, 92, 225-228.	2.1	8

#	ARTICLE	IF	CITATIONS
181	Effect of cocaine self-administration on striatal dopamine D1 receptors in rhesus monkeys. <i>Synapse</i> , 1998, 28, 1-9.	0.6	58
182	D1 dopamine receptors mediate neuroleptic-induced Fos expression in the islands of Calleja. , 1998, 28, 154-159.		10
183	Neurotransmitter regulation of MAP kinase signaling in striatal neurons in primary culture. <i>Synapse</i> , 1998, 29, 29-36.	0.6	66
184	5-HT3 receptor activation is required for induction of striatal c-Fos and phosphorylation of ATF-1 by amphetamine. , 1998, 30, 71-78.		24
185	Effect of cocaine self-administration on dopamine D2 receptors in rhesus monkeys. , 1998, 30, 88-96.		113
186	Drugs of abuse and immediate-early genes in the forebrain. <i>Molecular Neurobiology</i> , 1998, 16, 221-267.	1.9	189
187	Role of dynorphin and enkephalin in the regulation of striatal output pathways and behavior. <i>Experimental Brain Research</i> , 1998, 123, 60-76.	0.7	361
188	Δ ⁹ -Tetrahydrocannabinol increases sequence-specific AP-1 DNA-binding activity and Fos-related antigens in the rat brain. <i>European Journal of Neuroscience</i> , 1998, 10, 1743-1751.	1.2	35
189	MPTP-Parkinsonism is accompanied by persistent expression of a Δ ⁹ -FosB-like protein in dopaminergic pathways. <i>Molecular Brain Research</i> , 1998, 53, 41-52.	2.5	41
190	Subtracted differential display: Genes with amphetamine-altered expression patterns include calcineurin. <i>Molecular Brain Research</i> , 1998, 53, 344-347.	2.5	13
191	Differential effects of mood stabilizers on Fos/Jun proteins and AP-1 DNA binding activity in human neuroblastoma SH-SY5Y cells. <i>Molecular Brain Research</i> , 1998, 58, 95-102.	2.5	75
192	Conditional coupling of striatal dopamine D1 receptor to transcription factors: ontogenic and regional differences in CREB activation. <i>Molecular Brain Research</i> , 1998, 60, 127-132.	2.5	12
193	C-fos Mediates Cocaine Inhibition of NGF-Induced PC12 Cell Differentiation. <i>Molecular Genetics and Metabolism</i> , 1998, 64, 62-69.	0.5	7
194	Ap-1 Transcription Factors: Short- and Long-Term Modulators of Gene Expression in The Brain. <i>International Review of Neurobiology</i> , 1998, 42, 169-197.	0.9	30
195	Dopamine Receptors and Dopamine Transporter in Brain Function and Addictive Behaviors: Insights from Targeted Mouse Mutants. <i>Developmental Neuroscience</i> , 1998, 20, 188-203.	1.0	88
196	MK-801 reverses Fos expression induced by the full dopamine D1 receptor agonist SKF-82958 in the rat striatum. <i>European Journal of Pharmacology</i> , 1998, 342, 209-212.	1.7	15
197	Transcriptional Activation of the Ovine Follicle-Stimulating Hormone β ₂ -Subunit Gene by Gonadotropin-Releasing Hormone: Involvement of Two Activating Protein-1-Binding Sites and Protein Kinase C**This work was supported by the North Carolina State University Agricultural Research Service, NICHD Grant 34863, and the Mellon Foundation.. <i>Endocrinology</i> , 1998, 139, 4455-4465.	1.4	63
198	Light deprivation accelerates recovery from frontal cortical neglect: Relation to locomotion and striatal fos expression.. <i>Behavioral Neuroscience</i> , 1998, 112, 387-398.	0.6	18

#	ARTICLE	IF	CITATIONS
199	Essential Role of the <i>c-fos</i> Gene in Molecular, Cellular, and Behavioral Actions of Chronic Electroconvulsive Seizures. <i>Journal of Neuroscience</i> , 1998, 18, 6952-6962.	1.7	115
200	Internalization of D1 Dopamine Receptor in Striatal Neurons <i>In Vivo</i> as Evidence of Activation by Dopamine Agonists. <i>Journal of Neuroscience</i> , 1998, 18, 1650-1661.	1.7	187
201	A Complex Program of Striatal Gene Expression Induced by Dopaminergic Stimulation. <i>Journal of Neuroscience</i> , 1998, 18, 5301-5310.	1.7	320
202	An Antisense Oligonucleotide Reverses the Footshock-Induced Expression of Fos in the Rat Medial Prefrontal Cortex and the Subsequent Expression of Conditioned Fear-Induced Immobility. <i>Journal of Neuroscience</i> , 1999, 19, 5666-5673.	1.7	100
203	Decreased <i>c-fos</i> Responses to Dopamine D1 Receptor Agonist Stimulation in Mice Deficient for D3 Receptors. <i>Journal of Biological Chemistry</i> , 1999, 274, 29406-29412.	1.6	35
204	Blockade of neurotensin receptors suppresses the dopamine D1/D2 synergism on immediate early gene expression in the rat brain. <i>European Journal of Neuroscience</i> , 1999, 11, 967-974.	1.2	23
205	Functional heterogeneity in dopamine release and in the expression of Fos-like proteins within the rat striatal complex. <i>European Journal of Neuroscience</i> , 1999, 11, 1155-1166.	1.2	72
206	Neuronal and behavioural abnormalities in striatal function in DARPP-32-mutant mice. <i>European Journal of Neuroscience</i> , 1999, 11, 1114-1118.	1.2	73
207	Caffeine-mediated induction of <i>c-fos</i> , <i>zif-268</i> and <i>arc</i> expression through A1 receptors in the striatum: different interactions with the dopaminergic system. <i>European Journal of Neuroscience</i> , 1999, 11, 3101-3114.	1.2	38
208	Selective Effects of Alcohol Drinking on Restraint-Induced Expression of Immediate Early Genes in Mouse Brain. <i>Alcoholism: Clinical and Experimental Research</i> , 1999, 23, 1272-1280.	1.4	35
209	Behavioral sensitization to cocaine after a brief social defeat stress: <i>c-fos</i> expression in the PAG. <i>Psychopharmacology</i> , 1999, 141, 225-234.	1.5	82
210	Dextromethorphan modulates the AP-1 DNA-binding activity induced by kainic acid. <i>Brain Research</i> , 1999, 824, 125-132.	1.1	37
211	Compartmental changes in expression of <i>c-Fos</i> and <i>FosB</i> proteins in intact and dopamine-depleted striatum after chronic apomorphine treatment. <i>Brain Research</i> , 1999, 825, 104-114.	1.1	31
212	Effects of a D1 antagonist and of sexual experience on copulation-induced Fos-like immunoreactivity in the medial preoptic nucleus. <i>Brain Research</i> , 1999, 829, 55-68.	1.1	70
213	<i>c-fos</i> Gene expression parallels auditory adaptation in the adult rat. <i>Brain Research</i> , 1999, 839, 292-297.	1.1	16
214	Alcohol drinking produces brain region-selective changes in expression of inducible transcription factors. <i>Brain Research</i> , 1999, 847, 157-165.	1.1	129
215	Region-specific induction of <i>FosB</i> by repeated administration of typical versus atypical antipsychotic drugs. <i>Synapse</i> , 1999, 33, 118-128.	0.6	89
216	MK-801-induced expression of Fos protein family members in the rat retrosplenial granular cortex. <i>Journal of Neuroscience Research</i> , 1999, 57, 719-729.	1.3	7

#	ARTICLE	IF	CITATIONS
217	Actions of intrathecal diphtheria toxin-substance P fusion protein on models of persistent pain. <i>Pain</i> , 1999, 79, 243-253.	2.0	48
218	Enkephalin regulates acute D2 dopamine receptor antagonist-induced immediate-early gene expression in striatal neurons. <i>Neuroscience</i> , 1999, 88, 795-810.	1.1	69
219	Characterization of CArG-binding protein A initially identified by differential display. <i>Neuroscience</i> , 1999, 94, 637-649.	1.1	16
220	Distribution, biochemistry and function of striatal adenosine A2A receptors. <i>Progress in Neurobiology</i> , 1999, 59, 355-396.	2.8	468
221	Acute ibogaine injection induces expression of the immediate early genes, egr-1 and c-fos, in mouse brain. <i>Molecular Brain Research</i> , 1999, 74, 237-241.	2.5	13
222	Pseudoephedrine, a sympathomimetic agent, induces Fos-like immunoreactivity in rat nucleus accumbens and striatum. <i>Neuropharmacology</i> , 1999, 38, 1381-1387.	2.0	15
223	Effects of Amphetamine and Cocaine Treatment on c-Fos, Jun-B, and Krox-24 Expression in Rats with Intrastratial Dopaminergic Grafts. <i>Experimental Neurology</i> , 1999, 159, 139-152.	2.0	8
224	Drugs of Abuse and Brain Gene Expression. <i>Psychosomatic Medicine</i> , 1999, 61, 630-650.	1.3	52
225	3,4-Methylenedioxyamphetamine (ecstasy) induces c-fos-like protein and mRNA in rat organotypic dorsal striatal slices. , 2000, 36, 75-83.		15
226	Suppression of c-fos induction in the nucleus accumbens prevents acquisition but not expression of morphine-conditioned place preference. <i>European Journal of Neuroscience</i> , 2000, 12, 3399-3406.	1.2	40
227	The dopaminergic hyper-responsiveness of the shell of the nucleus accumbens is hormone-dependent. <i>European Journal of Neuroscience</i> , 2000, 12, 973-979.	1.2	190
228	Structure and function of dopamine receptors. <i>Neuroscience and Biobehavioral Reviews</i> , 2000, 24, 125-132.	2.9	607
229	Chronic tooth pulp inflammation causes transient and persistent expression of Fos in dynorphin-rich regions of rat brainstem. <i>Brain Research</i> , 2000, 861, 191-207.	1.1	29
230	Cocaine- and alcohol-mediated expression of inducible transcription factors is blocked by pentobarbital anesthesia. <i>Brain Research</i> , 2000, 877, 251-261.	1.1	32
231	Leftward shift in the acquisition of cocaine self-administration in isolation-reared rats: relationship to extracellular levels of dopamine, serotonin and glutamate in the nucleus accumbens and amygdala-striatal FOS expression. <i>Psychopharmacology</i> , 2000, 151, 55-63.	1.5	144
232	Margatoxin and iberiotoxin, two selective potassium channel inhibitors, induce c-fos like protein and mRNA in rat organotypic dorsal striatal slices. <i>Amino Acids</i> , 2000, 19, 23-31.	1.2	4
233	Fos Protein Expression and Cocaine-Seeking Behavior in Rats after Exposure to a Cocaine Self-Administration Environment. <i>Journal of Neuroscience</i> , 2000, 20, 798-805.	1.7	417
234	Involvement of the Extracellular Signal-Regulated Kinase Cascade for Cocaine-Rewarding Properties. <i>Journal of Neuroscience</i> , 2000, 20, 8701-8709.	1.7	500

#	ARTICLE	IF	CITATIONS
235	G _{OLF±} Mediates Dopamine D ₁ Receptor Signaling. <i>Journal of Neuroscience</i> , 2000, 20, RC91-RC91.	1.7	177
236	Fos-Related Antigen 2: Potential Mediator of the Transcriptional Activation in Rat Adrenal Medulla Evoked by Repeated Immobilization Stress. <i>Journal of Neuroscience</i> , 2000, 20, 5647-5653.	1.7	57
237	Regulation of the phosphorylation of the dopamine- and cAMP-regulated phosphoprotein of 32 kDa in vivo by dopamine D1, dopamine D2, and adenosine A2A receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 1856-1860.	3.3	190
238	Changes in CARG-binding protein A expression levels following injection(s) of the D1-dopamine agonist SKF-82958 in the intact and 6-hydroxydopamine-lesioned rat. <i>Neuroscience</i> , 2000, 98, 69-78.	1.1	8
239	Co-stimulation of D1/D5 and D2 dopamine receptors leads to an increase in c-fos messenger RNA in cholinergic interneurons and a redistribution of c-fos messenger RNA in striatal projection neurons. <i>Neuroscience</i> , 2000, 98, 749-757.	1.1	37
240	NGFI-B and Nor1 mRNAs are upregulated in brain reward pathways by drugs of abuse: different effects in Fischer and Lewis rats. <i>Molecular Brain Research</i> , 2000, 76, 18-24.	2.5	47
241	Dual effects of dextromethorphan on cocaine-induced conditioned place preference in mice. <i>Neuroscience Letters</i> , 2000, 288, 76-80.	1.0	36
242	Addiction, Dopamine, and the Molecular Mechanisms of Memory. <i>Neuron</i> , 2000, 25, 515-532.	3.8	1,054
243	Molecular effects of dopamine on striatal-projection pathways. <i>Trends in Neurosciences</i> , 2000, 23, S64-S70.	4.2	271
244	Cocaine Activation Discriminates Dopaminergic Projections by Temporal Response: An fMRI Study in Rat. <i>NeuroImage</i> , 2000, 11, 13-23.	2.1	162
245	Spatiotemporal Analysis of Fos Expression Associated with Cocaine- and PTZ-Induced Seizures in Prenatally Cocaine-Treated Rats. <i>Experimental Neurology</i> , 2001, 170, 109-120.	2.0	9
246	Parafascicular thalamic nucleus deafferentation reduces c-fos expression induced by dopamine D-1 receptor stimulation in rat striatum. <i>Neuroscience</i> , 2001, 103, 653-661.	1.1	15
247	Expression of c-Fos, Fos B, Jun B, and Zif268 transcription factor proteins in rat barrel cortex following apomorphine-evoked whisking behavior. <i>Neuroscience</i> , 2001, 106, 679-688.	1.1	23
248	Time course of Fos and Fras expression in the hypothalamic supraoptic neurons during chronic osmotic stimulation. <i>Molecular Brain Research</i> , 2001, 90, 39-47.	2.5	48
249	Carbetapentane attenuates kainate-induced seizures via I_f -1 receptor modulation. <i>Life Sciences</i> , 2001, 69, 915-922.	2.0	16
250	Influence of glucocorticoids on dopaminergic transmission in the rat dorsolateral striatum. <i>European Journal of Neuroscience</i> , 2001, 13, 812-818.	1.2	49
251	Analysis of Early Gene Responses. , 2001, Chapter 2, Unit2.5.		0
252	Amphetamine and cocaine induce different patterns of c-fosmRNA expression in the striatum and subthalamic nucleus depending on environmental context. <i>European Journal of Neuroscience</i> , 2001, 13, 1977-1983.	1.2	105

#	ARTICLE	IF	CITATIONS
253	Attenuation of cocaine-induced genomic and functional responses in prenatal cocaine-exposed rabbits. <i>Pharmacology Biochemistry and Behavior</i> , 2001, 69, 225-232.	1.3	9
254	The distribution of cannabinoid-induced Fos expression in rat brain: differences between the Lewis and Wistar strain. <i>Brain Research</i> , 2001, 921, 240-255.	1.1	75
255	ΔFosB: A sustained molecular switch for addiction. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 11042-11046.	3.3	533
256	Male Sexual Behavior. , 2002, , 3-137.		90
257	Chapter I Methods used in inducible transcription factor studies: focus on mRNA. <i>Handbook of Chemical Neuroanatomy</i> , 2002, 19, 1-38.	0.3	6
258	Cloning and characterization of guanine deaminase from mouse and rat brain. <i>Neuroscience</i> , 2002, 109, 15-26.	1.1	37
259	Withdrawal duration differentially affects c-fos expression in the medial prefrontal cortex and discrete subregions of the nucleus accumbens in cocaine-sensitized rats. <i>Neuroscience</i> , 2002, 114, 1061-1069.	1.1	49
260	Effects of cocaine on c-fos and preprodynorphin mRNA levels in intact and ovariectomized Fischer rats. <i>Brain Research Bulletin</i> , 2002, 58, 295-299.	1.4	16
261	Contrasting effects of dopamine antagonists and frequency reduction on Fos expression induced by lateral hypothalamic stimulation. <i>Behavioural Brain Research</i> , 2002, 132, 187-201.	1.2	20
262	Locomotor sensitization to cocaine is associated with increased Fos expression in the accumbens, but not in the caudate. <i>Behavioural Brain Research</i> , 2002, 136, 455-462.	1.2	70
263	Age-related distribution of c-fos expression in the striatum of CD-1 mice after acute methylphenidate administration. <i>Developmental Brain Research</i> , 2002, 135, 71-77.	2.1	37
264	Heroin sensitization as mapped by c-Fos immunoreactivity in the rat striatum. <i>Brain Research</i> , 2002, 933, 144-149.	1.1	26
265	Anatomical Markers of Activity in Neuroendocrine Systems: Are we all "Fos-ed out"? <i>Journal of Neuroendocrinology</i> , 2002, 14, 259-268.	1.2	200
266	The dopamine D1 receptor is a critical mediator for cocaine-induced gene expression. <i>Journal of Neurochemistry</i> , 2002, 82, 1453-1464.	2.1	113
267	Activation of <i>arc</i> , a Putative "Effector" Immediate Early Gene, by Cocaine in Rat Brain. <i>Journal of Neurochemistry</i> , 1995, 64, 2377-2380.	2.1	122
268	Acute methamphetamine administration upregulates NGFI-B mRNA expression in the striatum: Co-localization with C-FOS immunoreactivity. <i>Synapse</i> , 2002, 44, 158-165.	0.6	14
269	Dopamine transporter-dependent induction of C-Fos in HEK cells. <i>Synapse</i> , 2002, 45, 52-65.	0.6	28
270	Repeated ventral tegmental area amphetamine administration alters dopamine D1 receptor signaling in the nucleus accumbens. <i>Synapse</i> , 2002, 45, 159-170.	0.6	40

#	ARTICLE	IF	CITATIONS
271	Effects of <i>Coptis japonica</i> on morphine-induced conditioned place preference in mice. <i>Archives of Pharmacal Research</i> , 2003, 26, 540-544.	2.7	15
272	Inducible, brain region-specific expression of a dominant negative mutant of c-Jun in transgenic mice decreases sensitivity to cocaine. <i>Brain Research</i> , 2003, 970, 73-86.	1.1	87
273	Firing regulation in dopaminergic cells: effect of the partial degeneration of nigrostriatal system in surviving neurons. <i>European Journal of Neuroscience</i> , 2003, 18, 53-60.	1.2	22
274	Repeated methylphenidate treatment in adolescent rats alters gene regulation in the striatum. <i>European Journal of Neuroscience</i> , 2003, 18, 1584-1592.	1.2	95
275	Metabolism to dextrorphan is not essential for dextromethorphan's anticonvulsant activity against kainate in mice. <i>Life Sciences</i> , 2003, 72, 769-783.	2.0	33
276	Molecular and Behavioral Interactions Between Central Melanocortins and Cocaine. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003, 304, 391-399.	1.3	50
277	Effects of Cocaine on c-Fos and NGFI-B mRNA Expression in Transgenic Mice Underexpressing Glucocorticoid Receptors. <i>Neuropsychopharmacology</i> , 2003, 28, 478-489.	2.8	24
278	The Rate of Cocaine Administration Alters Gene Regulation and Behavioral Plasticity: Implications for Addiction. <i>Journal of Neuroscience</i> , 2004, 24, 6362-6370.	1.7	107
279	Inorganic Lead Exposure in the Rat Activates Striatal cFOS Expression at Lower Blood Levels and Inhibits Amphetamine-Induced cFOS Expression at Higher Blood Levels. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004, 310, 815-820.	1.3	6
280	The Type IV phosphodiesterase inhibitor rolipram interferes with drug-induced conditioned place preference but not immediate early gene induction in mice. <i>European Journal of Neuroscience</i> , 2004, 19, 2561-2568.	1.2	38
281	The transition from controlled to compulsive drug use is associated with a loss of sensitization. <i>Brain Research</i> , 2004, 995, 46-54.	1.1	98
282	Caffeine as a psychomotor stimulant: mechanism of action. <i>Cellular and Molecular Life Sciences</i> , 2004, 61, 857-872.	2.4	425
283	Fos immunocytochemical studies on the neuroanatomical sites of action of acute tyrosine depletion in the rat brain. <i>Psychopharmacology</i> , 2004, 171, 435-440.	1.5	15
284	Amphetamine-evoked c-fos mRNA expression in the caudate-putamen: the effects of DA and NMDA receptor antagonists vary as a function of neuronal phenotype and environmental context. <i>Journal of Neurochemistry</i> , 2004, 86, 33-44.	2.1	38
285	Molecular Neurobiology of Drug Addiction. <i>Annual Review of Medicine</i> , 2004, 55, 113-132.	5.0	296
286	Basal Ganglia. , 2004, , 455-508.		83
287	Dimemorfan prevents seizures induced by the L-type calcium channel activator BAY k-8644 in mice. <i>Behavioural Brain Research</i> , 2004, 151, 267-276.	1.2	20
288	Historical review: Molecular and cellular mechanisms of opiate and cocaine addiction. <i>Trends in Pharmacological Sciences</i> , 2004, 25, 210-218.	4.0	376

#	ARTICLE	IF	CITATIONS
289	DARPP-32 and modulation of cAMP signaling: involvement in motor control and levodopa-induced dyskinesia. <i>Parkinsonism and Related Disorders</i> , 2004, 10, 281-286.	1.1	48
290	Molecular mechanisms of drug addiction. <i>Neuropharmacology</i> , 2004, 47, 24-32.	2.0	538
291	Long-term behavioral and neuronal cross-sensitization to amphetamine induced by repeated brief social defeat stress: Fos in the ventral tegmental area and amygdala. <i>Neuroscience</i> , 2004, 123, 857-865.	1.1	146
292	Cocaine-induced Fos expression in rat striatum is blocked by chloral hydrate or urethane. <i>Neuroscience</i> , 2004, 127, 233-242.	1.1	40
293	Selegiline potentiates the effects of EGb 761 in response to ischemic brain injury. <i>Neurochemistry International</i> , 2004, 45, 157-170.	1.9	28
294	PAP 9704, a Korean Herbal Medicine Attenuates Methamphetamine-Induced Hyperlocomotion via Adenosine A2A Receptor Stimulation in Mice. <i>Biological and Pharmaceutical Bulletin</i> , 2004, 27, 906-909.	0.6	14
295	Neural Substrates of Coping Behavior in the Rat: Possible Importance of Mesocorticolimbic Dopamine System.. <i>Behavioral Neuroscience</i> , 2005, 119, 429-445.	0.6	18
296	Relationship of cocaine-induced c-Fos expression to behaviors and the role of serotonin 5-HT2A receptors in cocaine-induced c-Fos expression.. <i>Behavioral Neuroscience</i> , 2005, 119, 1173-1183.	0.6	21
297	Î ^γ FosB accumulates in a GABAergic cell population in the posterior tail of the ventral tegmental area after psychostimulant treatment. <i>European Journal of Neuroscience</i> , 2005, 21, 2817-2824.	1.2	153
298	Acute Alcohol Withdrawal is Associated with c-Fos Expression in the Basal Ganglia and Associated Circuitry: C57BL/6J and DBA/2J Inbred Mouse Strain Analyses. <i>Alcoholism: Clinical and Experimental Research</i> , 2005, 29, 1939-1948.	1.4	39
299	A48G polymorphism in the D1 receptor genes associated with bipolar I disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2005, 134B, 37-38.	1.1	42
300	Dopamine and norepinephrine transporter-dependent c-Fos production in vitro: relevance to neuroadaptation. <i>Journal of Neuroscience Methods</i> , 2005, 143, 69-78.	1.3	5
301	Increased activity of cyclin-dependent kinase 5 leads to attenuation of cocaine-mediated dopamine signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 1737-1742.	3.3	81
302	Cocaine Up-Regulates Fra-2 and Î ^γ -1 Receptor Gene and Protein Expression in Brain Regions Involved in Addiction and Reward. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005, 314, 770-779.	1.3	48
303	Chapter III The use of dopamine receptor knockout mice in understanding brain dopamine neurotransmission and sprouting in the nigrostriatal pathway. <i>Handbook of Chemical Neuroanatomy</i> , 2005, 21, 153-197.	0.3	1
304	Gene expression induced by drugs of abuse. <i>Current Opinion in Pharmacology</i> , 2005, 5, 26-33.	1.7	48
305	Why does the rapid delivery of drugs to the brain promote addiction?. <i>Trends in Pharmacological Sciences</i> , 2005, 26, 82-87.	4.0	184
306	Chromatin Remodeling Is a Key Mechanism Underlying Cocaine-Induced Plasticity in Striatum. <i>Neuron</i> , 2005, 48, 303-314.	3.8	692

#	ARTICLE	IF	CITATIONS
307	Preference for cocaine- versus pup-associated cues differentially activates neurons expressing either Fos or cocaine- and amphetamine-regulated transcript in lactating, maternal rodents. <i>Neuroscience</i> , 2005, 135, 315-328.	1.1	93
308	Inhibition of methylphenidate-induced gene expression in the striatum by local blockade of D1 dopamine receptors: Interhemispheric effects. <i>Neuroscience</i> , 2006, 140, 699-709.	1.1	42
309	Cocaine-induced locomotor activity and Fos expression in nucleus accumbens are sensitized for 6 months after repeated cocaine administration outside the home cage. <i>European Journal of Neuroscience</i> , 2006, 24, 867-875.	1.2	60
310	Neurochemical Changes and Neurotoxic Effects of an Acute Treatment with Sydnocarb, A Novel Psychostimulant. <i>Annals of the New York Academy of Sciences</i> , 2002, 965, 180-192.	1.8	11
311	Major affective disorders and schizophrenia: a common molecular signature?. <i>Human Mutation</i> , 2006, 27, 833-853.	1.1	40
312	Motor-Skill Learning-Associated Gene Regulation in the Striatum: Effects of Cocaine. <i>Neuropsychopharmacology</i> , 2006, 31, 2669-2682.	2.8	33
313	Plasticity-Associated Gene Krox24/Zif268 Is Required for Long-Lasting Behavioral Effects of Cocaine. <i>Journal of Neuroscience</i> , 2006, 26, 4956-4960.	1.7	111
314	Psychostimulants. , 2006, , 69-120.		3
315	GABAB Receptor-Positive Modulation Decreases Selective Molecular and Behavioral Effects of Cocaine. <i>Neuropsychopharmacology</i> , 2007, 32, 388-398.	2.8	59
316	Exposure to Extremely Low Frequency Magnetic Fields Enhances Locomotor Activity via Activation of Dopamine D1-Like Receptors in Mice. <i>Journal of Pharmacological Sciences</i> , 2007, 105, 367-371.	1.1	25
317	Cocaine increases Ras-guanine nucleotide-releasing factor 1 protein expression in the rat striatum in vivo. <i>Neuroscience Letters</i> , 2007, 427, 117-121.	1.0	14
319	c-fos gene expression is increased in the paraventricular hypothalamic nucleus of Sprague-Dawley rats With visceral pain induced by acetic acid without detectable changes of corticotrophin-releasing factor mRNA: A quantitative approach with an image analysis system. <i>Anatomical Record</i> , 2007, 290, 406-413.	0.8	9
320	Haplotype association study between DRD1 gene and bipolar type I affective disorder in two samples from Canada and Sardinia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2007, 144B, 237-241.	1.1	13
321	Context-specific sensitization of cocaine-induced locomotor activity and associated neuronal ensembles in rat nucleus accumbens. <i>European Journal of Neuroscience</i> , 2008, 27, 202-212.	1.2	59
322	Changes in levels of D1, D2, or NMDA receptors during withdrawal from brief or extended daily access to IV cocaine. <i>Brain Research</i> , 2007, 1131, 220-228.	1.1	59
323	Morphine-induced place preference: Involvement of the central amygdala NMDA receptors. <i>Brain Research</i> , 2007, 1133, 34-41.	1.1	47
324	Regulation of fosB and \hat{I} fosB mRNA expression: In vivo and in vitro studies. <i>Brain Research</i> , 2007, 1143, 22-33.	1.1	67
325	Differential gene expression in the striatum of mice with very low expression of the vesicular monoamine transporter type 2 gene. <i>Brain Research</i> , 2007, 1152, 10-16.	1.1	9

#	ARTICLE	IF	CITATIONS
326	Effects of bilateral adrenalectomy on systemic kainate-induced activation of the nucleus of the solitary tract. Regulation of blood pressure and local neurotransmitters. <i>Journal of Molecular Histology</i> , 2008, 39, 253-263.	1.0	3
327	Relapse to cocaine seeking increases activity-regulated gene expression differentially in the prefrontal cortex of abstinent rats. <i>Psychopharmacology</i> , 2008, 198, 77-91.	1.5	68
328	Relapse to cocaine-seeking increases activity-regulated gene expression differentially in the striatum and cerebral cortex of rats following short or long periods of abstinence. <i>Brain Structure and Function</i> , 2008, 213, 215-227.	1.2	49
329	Behavioral and biochemical responses to <i>l</i> -amphetamine in MCH ¹ receptor knockout mice. <i>Synapse</i> , 2008, 62, 128-136.	0.6	27
330	Distinct patterns of ³⁵ S-FosB induction in brain by drugs of abuse. <i>Synapse</i> , 2008, 62, 358-369.	0.6	183
331	Amphetamine and pseudoephedrine cross-tolerance measured by c-Fos protein expression in brains of chronically treated rats. <i>BMC Neuroscience</i> , 2008, 9, 99.	0.8	8
332	The rate of intravenous cocaine administration alters c-fos mRNA expression and the temporal dynamics of dopamine, but not glutamate, overflow in the striatum. <i>Brain Research</i> , 2008, 1209, 151-156.	1.1	32
333	Environmental modulation of ethanol-induced locomotor activity: Correlation with neuronal activity in distinct brain regions of adolescent and adult Swiss mice. <i>Brain Research</i> , 2008, 1239, 127-140.	1.1	60
334	Effects of dopamine and NMDA receptors on cocaine-induced Fos expression in the striatum of Fischer rats. <i>Brain Research</i> , 2008, 1243, 1-9.	1.1	25
335	Motor-skill learning in a novel running-wheel task is dependent on D1 dopamine receptors in the striatum. <i>Neuroscience</i> , 2008, 153, 249-258.	1.1	78
336	Inhibiting activator protein-1 activity alters cocaine-induced gene expression and potentiates sensitization. <i>Neuroscience</i> , 2008, 152, 1040-1053.	1.1	12
337	Transcriptional mechanisms of addiction: role of ³⁵ S-FosB. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2008, 363, 3245-3255.	1.8	329
338	Amphetamine alters Ras-guanine nucleotide-releasing factor expression in the rat striatum in vivo. <i>European Journal of Pharmacology</i> , 2009, 619, 50-56.	1.7	11
339	The impact of early environmental rearing condition on the discriminative stimulus effects and Fos expression induced by cocaine in adult male and female rats. <i>Psychopharmacology</i> , 2009, 203, 383-397.	1.5	15
340	Cocaine activates Homer1 immediate early gene transcription in the mesocorticolimbic circuit: Differential regulation by dopamine and glutamate signaling. <i>Synapse</i> , 2009, 63, 42-53.	0.6	37
341	c-Fos expression associated with reinstatement of cocaine-seeking behavior by response-contingent conditioned cues. <i>Synapse</i> , 2009, 63, 823-835.	0.6	87
342	Controlled cortical impact injury influences methylphenidate-induced changes in striatal dopamine neurotransmission. <i>Journal of Neurochemistry</i> , 2009, 110, 801-810.	2.1	29
343	Cocaine-induced Fos expression is detectable in the frontal cortex and striatum of rats under isoflurane but not \pm -chloralose anesthesia: Implications for fMRI. <i>Journal of Neuroscience Methods</i> , 2009, 181, 241-248.	1.3	19

#	ARTICLE	IF	CITATIONS
344	Differential activation of limbic circuitry associated with chronic ethanol withdrawal in DBA/2J and C57BL/6J mice. <i>Alcohol</i> , 2009, 43, 411-420.	0.8	32
345	Maturation of coordinated immediate early gene expression by cocaine during adolescence. <i>Neuroscience</i> , 2009, 160, 13-31.	1.1	21
346	PKA-mediated responses in females' estrous cycle affect cocaine-induced responses in dopamine-mediated intracellular cascades. <i>Neuroscience</i> , 2009, 161, 865-876.	1.1	10
347	Cocaine addiction and personality: A mathematical model. <i>British Journal of Mathematical and Statistical Psychology</i> , 2010, 63, 449-480.	1.0	22
348	The effects of adolescent methylphenidate self-administration on responding for a conditioned reward, amphetamine-induced locomotor activity, and neuronal activation. <i>Psychopharmacology</i> , 2010, 208, 455-468.	1.5	19
349	Interaction of the rostral basolateral amygdala and prelimbic prefrontal cortex in regulating reinstatement of cocaine-seeking behavior. <i>Pharmacology Biochemistry and Behavior</i> , 2010, 96, 347-353.	1.3	47
350	Differential effects of methamphetamine and SCH23390 on the expression of members of IEG families of transcription factors in the rat striatum. <i>Brain Research</i> , 2010, 1318, 1-10.	1.1	36
351	Region-specific Induction of FosB/Δ FosB by Voluntary Alcohol Intake: Effects of Naltrexone. <i>Alcoholism: Clinical and Experimental Research</i> , 2010, 34, 1742-1750.	1.4	52
352	Striatal regulation of Δ FosB, FosB, and cFos during cocaine self-administration and withdrawal. <i>Journal of Neurochemistry</i> , 2010, 115, 112-122.	2.1	68
353	Identification of Brain Nuclei Implicated in Cocaine-Primed Reinstatement of Conditioned Place Preference: A Behaviour Dissociable from Sensitization. <i>PLoS ONE</i> , 2010, 5, e15889.	1.1	58
354	Psychostimulant-Induced Gene Regulation in Corticostriatal Circuits. <i>Handbook of Behavioral Neuroscience</i> , 2010, , 501-525.	0.7	5
355	Fos After Single and Repeated Self-Administration of Cocaine and Saline in the Rat: Emphasis on the Basal Forebrain and Recalibration of Expression. <i>Neuropsychopharmacology</i> , 2010, 35, 445-463.	2.8	70
356	Ablation of Rat TRPV1-Expressing Aδ/C-Fibers with Resiniferatoxin: Analysis of Withdrawal Behaviors, Recovery of Function and Molecular Correlates. <i>Molecular Pain</i> , 2010, 6, 1744-8069-6-94.	1.0	67
357	Exposure to Extremely Low Frequency Magnetic Fields Induces Fos-Related Antigen-Immunoreactivity Via Activation of Dopaminergic D1 Receptor. <i>Experimental Neurobiology</i> , 2011, 20, 130-136.	0.7	11
358	The Striatal Balancing Act in Drug Addiction: Distinct Roles of Direct and Indirect Pathway Medium Spiny Neurons. <i>Frontiers in Neuroanatomy</i> , 2011, 5, 41.	0.9	301
359	Chronic methamphetamine exposure suppresses the striatal expression of members of multiple families of immediate early genes (IEGs) in the rat: normalization by an acute methamphetamine injection. <i>Psychopharmacology</i> , 2011, 215, 353-365.	1.5	47
360	Liquiritigenin Decreases Selective Molecular and Behavioral Effects of Cocaine in Rodents. <i>Current Neuropharmacology</i> , 2011, 9, 30-34.	1.4	8
361	Role of Dorsal Medial Prefrontal Cortex Dopamine D1-Family Receptors in Relapse to High-Fat Food Seeking Induced by the Anxiogenic Drug Yohimbine. <i>Neuropsychopharmacology</i> , 2011, 36, 497-510.	2.8	80

#	ARTICLE	IF	CITATIONS
362	Dextromethorphan-induced psychotoxic behaviors cause sexual dysfunction in male mice via stimulation of 5-HT _{2A} receptors. <i>Neurochemistry International</i> , 2012, 61, 913-922.	1.9	17
363	Upregulation of Npas4 protein expression by chronic administration of amphetamine in rat nucleus accumbens in vivo. <i>Neuroscience Letters</i> , 2012, 528, 210-214.	1.0	10
364	Nucleus accumbens D2- and D1-receptor expressing medium spiny neurons are selectively activated by morphine withdrawal and acute morphine, respectively. <i>Neuropharmacology</i> , 2012, 62, 2463-2471.	2.0	50
365	5-HT _{2A} receptor blockade and 5-HT _{2C} receptor activation interact to reduce cocaine hyperlocomotion and fos protein expression in the caudate-putamen. <i>Synapse</i> , 2012, 66, 989-1001.	0.6	24
366	Forced Abstinence from Cocaine Self-Administration is Associated with DNA Methylation Changes in Myelin Genes in the Corpus Callosum: a Preliminary Study. <i>Frontiers in Psychiatry</i> , 2012, 3, 60.	1.3	29
367	Neuroplasticity in addiction: cellular and transcriptional perspectives. <i>Frontiers in Molecular Neuroscience</i> , 2012, 5, 99.	1.4	31
368	Dopamine D ₁ Receptor-Mediated Intracellular Responses in the Hypothalamus after Co-Administration of Caffeine with MDMA. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2012, 110, 283-289.	1.2	6
369	Optogenetic insights into striatal function and behavior. <i>Behavioural Brain Research</i> , 2013, 255, 44-54.	1.2	87
370	Addiction-related gene regulation: Risks of exposure to cognitive enhancers vs. other psychostimulants. <i>Progress in Neurobiology</i> , 2013, 100, 60-80.	2.8	51
371	The medial preoptic area modulates cocaine-induced activity in female rats. <i>Behavioral Neuroscience</i> , 2013, 127, 293-302.	0.6	40
372	Electroacupuncture Reduces Cocaine-Induced Seizures and Mortality in Mice. <i>Evidence-based Complementary and Alternative Medicine</i> , 2013, 2013, 1-13.	0.5	10
373	Sex differences in the neurobiology of drug addiction. <i>Experimental Neurology</i> , 2014, 259, 64-74.	2.0	198
374	Histidine Decarboxylase Deficiency Causes Tourette Syndrome: Parallel Findings in Humans and Mice. <i>Neuron</i> , 2014, 81, 77-90.	3.8	212
375	Molecular neurobiology of addiction: what's all the (i)FosB about?. <i>American Journal of Drug and Alcohol Abuse</i> , 2014, 40, 428-437.	1.1	68
376	Neuromodulatory Effect of G _i - or G _q -Coupled G-Protein-Coupled Receptor on NMDA Receptor Selectively Activates the NMDA Receptor/Ca ²⁺ /Calcineurin/cAMP Response Element-Binding Protein-Regulated Transcriptional Coactivator 1 Pathway to Effectively Induce Brain-Derived Neurotrophic Factor Expression in Neurons. <i>Journal of Neuroscience</i> , 2015, 35, 5606-5624.	1.7	53
377	Methyl Supplementation Attenuates Cocaine-Seeking Behaviors and Cocaine-Induced c-Fos Activation in a DNA Methylation-Dependent Manner. <i>Journal of Neuroscience</i> , 2015, 35, 8948-8958.	1.7	101
378	Leptin in the hindbrain facilitates phosphorylation of STAT3 in the hypothalamus. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2015, 308, E351-E361.	1.8	13
379	A Genetic Mouse Model of Parkinson's Disease Shows Involuntary Movements and Increased Postsynaptic Sensitivity to Apomorphine. <i>Molecular Neurobiology</i> , 2015, 52, 1152-1164.	1.9	16

#	ARTICLE	IF	CITATIONS
380	A previous history of repeated amphetamine exposure modifies brain angiotensin II AT1 receptor functionality. <i>Neuroscience</i> , 2015, 307, 1-13.	1.1	11
381	Region-specific effects of isoflurane anesthesia on Fos immunoreactivity in response to intravenous cocaine challenge in rats with a history of repeated cocaine administration. <i>Brain Research</i> , 2015, 1594, 256-266.	1.1	6
382	Using c-fos to study neuronal ensembles in corticostriatal circuitry of addiction. <i>Brain Research</i> , 2015, 1628, 157-173.	1.1	128
383	Serotonin _{2C} receptors modulate dopamine transmission in the nucleus accumbens independently of dopamine release: behavioral, neurochemical and molecular studies with cocaine. <i>Addiction Biology</i> , 2015, 20, 445-457.	1.4	30
384	Serotonin _{2C} receptor stimulation inhibits cocaine-induced Fos expression and DARPP-32 phosphorylation in the rat striatum independently of dopamine outflow. <i>Neuropharmacology</i> , 2015, 89, 375-381.	2.0	14
386	Psychostimulant-Induced Gene Regulation in Striatal Circuits. <i>Handbook of Behavioral Neuroscience</i> , 2016, , 639-672.	0.7	2
387	Colocalization of Mating-Induced Fos and D2-Like Dopamine Receptors in the Medial Preoptic Area: Influence of Sexual Experience. <i>Frontiers in Behavioral Neuroscience</i> , 2016, 10, 75.	1.0	20
388	Dopamine D2 receptors in striatal output neurons enable the psychomotor effects of cocaine. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 11609-11614.	3.3	30
389	Prelimbic to Accumbens Core Pathway Is Recruited in a Dopamine-Dependent Manner to Drive Cued Reinstatement of Cocaine Seeking. <i>Journal of Neuroscience</i> , 2016, 36, 8700-8711.	1.7	110
390	Molecular Mechanisms of Cannabis Signaling in the Brain. <i>Progress in Molecular Biology and Translational Science</i> , 2016, 137, 123-147.	0.9	12
391	Differential regulation of the phosphorylation of Trimethyl-lysine ²⁷ histone H3 at serine 28 in distinct populations of striatal projection neurons. <i>Neuropharmacology</i> , 2016, 107, 89-99.	2.0	10
392	Exposure of adolescent mice to 3,4-methylenedioxypyrovalerone increases the psychostimulant, rewarding and reinforcing effects of cocaine in adulthood. <i>British Journal of Pharmacology</i> , 2017, 174, 1161-1173.	2.7	24
393	Cocaine Abuse, Traumatic Brain Injury, and Preexisting Brain Lesions as Risk Factors for Bupropion-Associated Psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2017, 37, 459-463.	0.7	5
394	Post-sensitization treatment with rimonabant blocks the expression of cocaine-induced behavioral sensitization and c-Fos protein in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2017, 156, 16-23.	1.3	6
395	Effects of alkaloid-rich extract from <i>Mitragyna speciosa</i> (Korth.) Havil. on naloxone-precipitated morphine withdrawal symptoms and local field potential in the nucleus accumbens of mice. <i>Journal of Ethnopharmacology</i> , 2017, 208, 129-137.	2.0	22
396	Molecular Mechanisms of Drug-Induced Plasticity. , 2017, , .		0
397	Alcohol's Effects on Pair-Bond Maintenance in Male Prairie Voles. <i>Frontiers in Psychiatry</i> , 2017, 8, 226.	1.3	16
398	Beyond Neuronal Activity Markers: Select Immediate Early Genes in Striatal Neuron Subtypes Functionally Mediate Psychostimulant Addiction. <i>Frontiers in Behavioral Neuroscience</i> , 2017, 11, 112.	1.0	42

#	ARTICLE	IF	CITATIONS
399	Common Hepatic Branch of Vagus Nerve-Dependent Expression of Immediate Early Genes in the Mouse Brain by Intraportal L-Arginine: Comparison with Cholecystokinin-8. <i>Frontiers in Neuroscience</i> , 2017, 11, 366.	1.4	3
400	The Hyperactive Rat: An Animal Model for Bipolar Disorder. <i>Current Psychiatry Reviews</i> , 2017, 13, .	0.9	2
401	Estrogen Receptor β in the Nucleus Accumbens Regulates the Rewarding Properties of Cocaine in Female Mice. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 382-392.	1.0	33
402	Dorsal Hippocampus Drives Context-Induced Cocaine Seeking via Inputs to Lateral Septum. <i>Neuropsychopharmacology</i> , 2018, 43, 987-1000.	2.8	80
403	Leveraging calcium imaging to illuminate circuit dysfunction in addiction. <i>Alcohol</i> , 2019, 74, 47-63.	0.8	43
404	Decoding the transcriptional programs activated by psychotropic drugs in the brain. <i>Genes, Brain and Behavior</i> , 2019, 18, e12511.	1.1	11
405	Glutathione peroxidase-1 gene rescues cocaine-induced conditioned place preference in mice by inhibiting β -1 receptor expression. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2019, 46, 791-797.	0.9	6
406	Effects of Muscarinic Acetylcholine m1 and m4 Receptor Blockade on Dyskinesia in the Hemi-Parkinsonian Rat. <i>Neuroscience</i> , 2019, 409, 180-194.	1.1	38
407	Fos-Expressing Neuronal Ensembles in Addiction Research. , 2019, , 75-88.		1
408	Adolescent cocaine exposure enhanced negative affect following drug re-exposure in adult rats: Attenuation of c-Fos activation. <i>Journal of Psychopharmacology</i> , 2019, 33, 154-162.	2.0	19
409	Selective Regulation of 5-HT1B Serotonin Receptor Expression in the Striatum by Dopamine Depletion and Repeated L-DOPA Treatment: Relationship to L-DOPA-Induced Dyskinesias. <i>Molecular Neurobiology</i> , 2020, 57, 736-751.	1.9	15
410	Glutathione peroxidase-1 knockout potentiates behavioral sensitization induced by cocaine in mice via β -1 receptor-mediated ERK signaling: A comparison with the case of glutathione peroxidase-1 overexpressing transgenic mice. <i>Brain Research Bulletin</i> , 2020, 164, 107-120.	1.4	5
411	Neuroadaptations in the dorsal hippocampus underlie cocaine seeking during prolonged abstinence. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 26460-26469.	3.3	24
412	Taking Rapid and Intermittent Cocaine Infusions Enhances Both Incentive Motivation for the Drug and Cocaine-induced Gene Regulation in Corticostriatal Regions. <i>Neuroscience</i> , 2020, 442, 314-328.	1.1	9
413	Psychostimulants. , 2020, , 1-245.		1
414	Vulnerability to helpless behavior is regulated by the circadian clock component CRYPTOCHROME in the mouse nucleus accumbens. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 13771-13782.	3.3	29
415	Dopaminergic Control of Striatal Cholinergic Interneurons Underlies Cocaine-Induced Psychostimulation. <i>Cell Reports</i> , 2020, 31, 107527.	2.9	23
416	Role of prefrontal cortex projections to the nucleus accumbens core in mediating the effects of ceftriaxone on cue-induced cocaine seeking. <i>Addiction Biology</i> , 2021, 26, e12928.	1.4	14

#	ARTICLE	IF	CITATIONS
417	Cocaine induces paradigm-specific changes to the transcriptome within the ventral tegmental area. <i>Neuropsychopharmacology</i> , 2021, 46, 1768-1779.	2.8	14
418	Cocaine-induced Fos expression in the rat brain: Modulation by prior δ^9 -tetrahydrocannabinol exposure during adolescence and sex-specific effects. <i>Brain Research</i> , 2021, 1764, 147480.	1.1	2
419	Drug-activated cells: From immediate early genes to neuronal ensembles in addiction. <i>Advances in Pharmacology</i> , 2021, 90, 173-216.	1.2	25
420	D1/D2 Dopamine Receptor Interactions in Basal Ganglia Functions. , 1997, , 193-219.		13
422	Dependence, Tolerance, and Alteration in Gene Expression. , 1999, , 207-211.		1
423	Understanding the Function of the Dopamine D2 Receptor: A Knockout Animal Approach. <i>Handbook of Experimental Pharmacology</i> , 2002, , 159-183.	0.9	1
424	Sites and Mechanisms of Action of Antipsychotic Drugs as Revealed by Immediate-Early Gene Expression. <i>Handbook of Experimental Pharmacology</i> , 1996, , 117-161.	0.9	14
425	Sphingosine 1-phosphate, a novel signaling molecule, stimulates DNA binding activity of AP-1 in quiescent Swiss 3T3 fibroblasts.. <i>Journal of Biological Chemistry</i> , 1994, 269, 16512-16517.	1.6	82
426	Constitutive expression of AP-1 transcription factors in the rat adrenal. Effects of nicotine.. <i>Journal of Biological Chemistry</i> , 1992, 267, 20148-20152.	1.6	34
427	Light deprivation accelerates recovery from frontal cortical neglect: relation to locomotion and striatal Fos expression. <i>Behavioral Neuroscience</i> , 1998, 112, 387-98.	0.6	5
429	Caveats in the use of the Kindling Model of Affective Disorders. <i>Toxicology and Industrial Health</i> , 1994, 10, 421-447.	0.6	30
430	D1 and D2Dopamine Receptors in Perinatal and Adult Basal Ganglia. <i>Pediatric Research</i> , 1997, 41, 822-831.	1.1	24
431	A α_2A receptor and striatal cellular functions. <i>Neurology</i> , 2003, 61, S24-9.	1.5	26
432	Repeated Methamphetamine Administration Differentially Alters Fos Expression in Caudate-Putamen Patch and Matrix Compartments and Nucleus Accumbens. <i>PLoS ONE</i> , 2012, 7, e34227.	1.1	29
433	Prolonged and Extrasynaptic Excitatory Action of Dopamine Mediated by D1 Receptors in the Rat Striatum <i>In Vivo</i> . <i>Journal of Neuroscience</i> , 1997, 17, 5972-5978.	1.7	273
434	Involvement of pCREB Expression in Inhibitory Effects of <i>Coptis japonica</i> on Morphine-induced Psychological Dependence. <i>Biomolecules and Therapeutics</i> , 2008, 16, 113-117.	1.1	4
435	δ^9 FosB: a Molecular Switch for Reward. <i>Journal of Drug and Alcohol Research</i> , 2013, 2, 1-11.	0.9	7
436	Dopamine and Gene Expression. <i>Handbook of Experimental Pharmacology</i> , 2002, , 321-337.	0.9	0

#	ARTICLE	IF	CITATIONS
437	Basal Ganglia & Cortex Interactions: Regulation of Cortical Function by D1 Dopamine Receptors in the Striatum. , 2007, , 261-282.		1
438	The Neurobiology of Cocaine-Induced Reinforcement. Novartis Foundation Symposium, 1992, 166, 96-124.	1.2	23
439	Nicotine, Cocaine and Amphetamine Share Several Common Mechanisms of c-fos Induction in the Striatum: Implications for the Functions of the Striatum. Advances in Behavioral Biology, 1994, , 465-473.	0.2	0
440	Dopamine Receptor Modulation of Gene Expression in the Brain. , 1997, , 305-342.		1
441	Cocaine and a Mechanism for Long-Term Changes in Gene Expression. , 1999, , 213-222.		0
443	Role of monoamine systems in activation of zif268 by cocaine. Journal of Psychiatry and Neuroscience, 1992, 17, 94-102.	1.4	24
444	Dopamine receptor genes: new tools for molecular psychiatry. Journal of Psychiatry and Neuroscience, 1992, 17, 158-80.	1.4	38
445	Acute treatment with antidepressant drugs selectively increases the expression of c-fos in the rat brain. Journal of Psychiatry and Neuroscience, 1995, 20, 25-32.	1.4	47
448	Convergent dopamine and <i>ALK4</i> signaling to <i>PCBP1</i> controls <i>FosB</i> alternative splicing and cocaine behavioral sensitization. EMBO Journal, 2022, 41, .	3.5	5
450	Alcohol consumption modulates prelimbic cortex response to cocaine following sequential cocaine and alcohol polysubstance use in the rat. Frontiers in Pharmacology, 0, 14, .	1.6	1