Linear mixture modelling applied to AVHRR data for cr

International Journal of Remote Sensing 13, 415-425 DOI: 10.1080/01431169208904046

Citation Report

#	Article	IF	CITATIONS
1	Relationships between reflectances at coarse and fine resolution on heterogeneous landscapes. , 0, , .		4
2	Linear mixing model applied to coarse spatial resolution data from multispectral satellite sensors. International Journal of Remote Sensing, 1993, 14, 2231-2240.	2.9	73
3	Linear mixture modelling applied to AVHRR data for monitoring vegetation. , 0, , .		1
4	Validation of the use of multiple linear regression as a tool for unmixing coarse spatial resolution images. Remote Sensing of Environment, 1994, 49, 155-166.	11.0	39
5	AVHRR bidirectional reflectance effects and compositing. Remote Sensing of Environment, 1994, 48, 77-88.	11.0	117
6	Forest canopy closure from classification and spectral unmixing of scene components-multisensor evaluation of an open canopy. IEEE Transactions on Geoscience and Remote Sensing, 1994, 32, 1067-1080.	6.3	72
7	Sub-pixel land cover composition estimation using a linear mixture model and fuzzy membership functions. International Journal of Remote Sensing, 1994, 15, 619-631.	2.9	285
8	The improvement of net primary productivity estimation taking into account the heterogeneity of Sahelian region. , 0, , .		1
9	Status of remote sensing algorithms for estimation of land surface state parameters. Remote Sensing of Environment, 1995, 51, 138-156.	11.0	210
10	Unmixing multiple land-cover type reflectances from coarse spatial resolution satellite data. Remote Sensing of Environment, 1995, 54, 98-112.	11.0	61
11	Linear and correlation analysis for computerized identification of categories in Landsat images. International Journal of Remote Sensing, 1995, 16, 2277-2284.	2.9	0
12	NOAA-AVHRR NDVI decomposition and subpixel classification using linear mixing in the Argentinean Pampa. International Journal of Remote Sensing, 1995, 16, 1303-1325.	2.9	88
13	A review of mixture modeling techniques for subâ€pixel land cover estimation. International Journal of Remote Sensing, 1996, 13, 161-186.	1.0	121
14	A METHOD FOR SYNTHESIZING IMAGES OF DIVERSE SPATIAL RESOLUTION. Mapping Sciences and Remote Sensing, 1996, 33, 196-206.	0.0	0
15	Point accuracy of a non-parametric method in estimation of forest characteristics with different satellite materials. International Journal of Remote Sensing, 1996, 17, 2333-2351.	2.9	124
16	Woodland area estimation by spectral mixing: applying a goodness-of-fit solution method. International Journal of Remote Sensing, 1996, 17, 291-301.	2.9	17
17	A knowledge-based system for the computation of land cover mixing and the classification of multi-spectral satellite imagery. International Journal of Remote Sensing, 1996, 17, 1483-1492.	2.9	5
18	Monitoring regional drought using the Vegetation Condition Index. International Journal of Remote Sensing, 1996, 17, 2761-2782.	2.9	298

TATION REDO

#	Article	IF	CITATIONS
19	Extraction of the pure spectral response of the landscape components in NOAA-AVHRR mixed pixels—application to the HAPEX-Sahel degree square. International Journal of Remote Sensing, 1996, 17, 2259-2280.	2.9	14
20	Impacts of upscaling techniques on land cover representation in Nebraska, U.S.A Geocarto International, 1997, 12, 27-39.	3.5	17
21	The pixel: A snare and a delusion. International Journal of Remote Sensing, 1997, 18, 679-685.	2.9	387
22	Mapping tropical forest fractional cover from coarse spatial resolution remote sensing imagery. Plant Ecology, 1997, 131, 143-154.	1.6	38
23	Multitemporal, multichannel AVHRR data sets for land biosphere studies—Artifacts and corrections. Remote Sensing of Environment, 1997, 60, 35-57.	11.0	202
24	Generation of fraction images from AVHRR data using linear mixing model. Journal of the Indian Society of Remote Sensing, 1998, 26, 23-27.	2.4	2
25	Subpixel Land Use Classification and Retrieval of Forest Stem Volume in the Boreal Forest Zone by Employing SSM/I Data. Remote Sensing of Environment, 1998, 63, 140-154.	11.0	5
26	Quantification of Subpixel Cover Fractions Using Principal Component Analysis and a Linear Programming Method: Application to the Coastal Zone of Roscoff (France). Remote Sensing of Environment, 1998, 64, 153-165.	11.0	31
27	Using Landscape Spatial Relationships to Improve Estimates of Land-Cover Area from Coarse Resolution Remote Sensing. Remote Sensing of Environment, 1998, 64, 202-220.	11.0	29
28	Higher resolution Earth surface features from repeat moderate resolution satellite imagery. IEEE Transactions on Geoscience and Remote Sensing, 1998, 36, 244-255.	6.3	13
29	Multiclass spectral decomposition of remotely sensed scenes by selective pixel unmixing. IEEE Transactions on Geoscience and Remote Sensing, 1998, 36, 1809-1820.	6.3	39
30	Remote sensing segmentation through a filter bank based on Gabor functions. , 0, , .		2
31	Using textural and statistical data for almond and olive fields classification. , 1998, , .		0
32	Optimal linear spectral unmixing. IEEE Transactions on Geoscience and Remote Sensing, 1999, 37, 639-644.	6.3	101
33	Global land cover characterization from satellite data: from research to operational implementation?. Global Ecology and Biogeography, 1999, 8, 367-379.	5.8	74
34	Soft classification for sub-pixel land cover extraction. Journal of the Indian Society of Remote Sensing, 1999, 27, 225-234.	2.4	2
35	Estimating lake area in an Arctic landscape using linear mixture modelling with AVHRR data. International Journal of Remote Sensing, 1999, 20, 829-835.	2.9	16
36	Continuous fields of vegetation characteristics at the global scale at 1â€km resolution. Journal of Geophysical Research, 1999, 104, 16911-16923.	3.3	204

#	Article	IF	CITATIONS
37	Land cover characterization and change detection for environmental monitoring of pan-Europe. International Journal of Remote Sensing, 2000, 21, 1159-1181.	2.9	132
38	Global continuous fields of vegetation characteristics: A linear mixture model applied to multi-year 8 km AVHRR data. International Journal of Remote Sensing, 2000, 21, 1389-1414.	2.9	227
39	Assessing positional accuracy and its effects on rice crop area measurement: an application at Coleambally Irrigation Area. Australian Journal of Experimental Agriculture, 2001, 41, 557.	1.0	10
40	Sub-pixel Classification of Land Use Using Temporal Profile of NDVI Journal of the Japan Society of Photogrammetry and Remote Sensing, 2001, 40, 43-54.	0.0	4
41	On the Study of China's Grain Output Prediction. International Transactions in Operational Research, 2001, 8, 429-437.	2.7	5
42	NOAA-AVHRR based flood monitoring. , 0, , .		5
43	Landscape-scale characterization of cropland in China using Vegetation and Landsat TM images. International Journal of Remote Sensing, 2002, 23, 3579-3594.	2.9	71
44	Temporal dynamics of soil and vegetation spectral responses in a semi-arid environment. International Journal of Remote Sensing, 2002, 23, 4073-4087.	2.9	51
45	Class-prioritized compression of multispectral imagery data. Journal of Electronic Imaging, 2002, 11, 246.	0.9	4
46	An evaluation of spectral mixture modelling applied to a semi-arid environment. International Journal of Remote Sensing, 2002, 23, 687-700.	2.9	53
47	Areal estimates of fragmented land cover: Effects of pixel size and model-based corrections. International Journal of Remote Sensing, 2002, 23, 711-724.	2.9	16
48	Linear mixing model based on optimization method for land cover mapping using LANDSAT ETM+ and SPOT-HRV data. , 0, , .		1
49	Sub-pixel habitat mapping of a costal dune ecosystem. Applied Geography, 2002, 22, 253-270.	3.7	27
50	Development of an integrated geographical information system prototype for coastal habitat monitoring. Computers and Geosciences, 2002, 28, 129-141.	4.2	20
51	Linear mixture model applied to Amazonian vegetation classification. Remote Sensing of Environment, 2003, 87, 456-469.	11.0	207
52	IKONOS imagery for the Large Scale Biosphere–Atmosphere Experiment in Amazonia (LBA). Remote Sensing of Environment, 2003, 88, 111-127.	11.0	44
53	A stepwise regression tree for nonlinear approximation: Applications to estimating subpixel land cover. International Journal of Remote Sensing, 2003, 24, 75-90.	2.9	122
54	Mesoscale water balance modelling in the Upper Danube watershed using sub-scale land cover information derived from NOAA-AVHRR imagery and GIS-techniques. Physics and Chemistry of the Earth, 2003, 28, 1351-1364.	2.9	21

# 55	ARTICLE An approach for land cover change detection using low spatial resolution data. , 0, , .	IF	CITATIONS 0
56	Spectral Mixture Analysis of the Urban Landscape in Indianapolis with Landsat ETM+ Imagery. Photogrammetric Engineering and Remote Sensing, 2004, 70, 1053-1062.	0.6	299
57	Spectral and temporal linear mixing model for vegetation classification. International Journal of Remote Sensing, 2004, 25, 4203-4218.	2.9	20
58	Cropland distributions from temporal unmixing of MODIS data. Remote Sensing of Environment, 2004, 93, 412-422.	11.0	272
59	Accuracy of land cover area estimated from coarse spatial resolution images using an unmixing method. International Journal of Remote Sensing, 2004, 25, 1673-1683.	2.9	15
60	Remote sensing and land cover area estimation. International Journal of Remote Sensing, 2004, 25, 3019-3047.	2.9	279
61	Current and potential uses of optical remote sensing in rice-based irrigation systems: a review. Australian Journal of Agricultural Research, 2004, 55, 155.	1.5	62
62	Comparison of non-linear mixture models: sub-pixel classification. Remote Sensing of Environment, 2005, 94, 145-154.	11.0	113
63	Fusion of NOAA-AVHRR imagery and geographical information system techniques to derive subscale land cover information for the upper Danube watershed. Hydrological Processes, 2005, 19, 2407-2418.	2.6	12
64	Cotton Canopy Reflectance at Landscape Scale as Affected by Nitrogen Fertilization. Agronomy Journal, 2005, 97, 654-660.	1.8	32
65	A multiscale multitemporal land cover classification method using a Bayesian approach. , 2005, , .		5
66	Irrigated area mapping in heterogeneous landscapes with MODIS time series, ground truth and census data, Krishna Basin, India. International Journal of Remote Sensing, 2006, 27, 4245-4266.	2.9	137
67	Estimation of Inter-annual Crop Area Variation by the Application of Spectral Angle Mapping to Low Resolution Multitemporal NDVI Images. Photogrammetric Engineering and Remote Sensing, 2006, 72, 55-62.	0.6	25
68	Land-cover Mapping in the Brazilian Amazon Using SPOT-4 Vegetation Data and Machine Learning Classification Methods. Photogrammetric Engineering and Remote Sensing, 2006, 72, 897-910.	0.6	32
70	Spectral mixture analysis of ASTER images for examining the relationship between urban thermal features and biophysical descriptors in Indianapolis, Indiana, USA. Remote Sensing of Environment, 2006, 104, 157-167.	11.0	157
71	Estimation of yellow starthistle abundance through CASI-2 hyperspectral imagery using linear spectral mixture models. Remote Sensing of Environment, 2006, 101, 329-341.	11.0	79
72	Resolution dependent errors in remote sensing of cultivated areas. Remote Sensing of Environment, 2006, 103, 203-217.	11.0	140
73	A combined approach for estimating vegetation cover in urban/suburban environments from remotely sensed data. Computers and Geosciences, 2006, 32, 1299-1309.	4.2	23

#	Article	IF	CITATIONS
74	A comparison of the classification of wetland characteristics by linear spectral mixture modelling and traditional hard classifiers on multispectral remotely sensed imagery in southern India. Ecological Modelling, 2006, 194, 379-394.	2.5	98
75	Residential population estimation using a remote sensing derived impervious surface approach. International Journal of Remote Sensing, 2006, 27, 3553-3570.	2.9	106
76	Nonlinear estimation of subpixel proportion via kernel least square regression. International Journal of Remote Sensing, 2007, 28, 4157-4172.	2.9	19
77	Corn and Soybean Mapping in the United States Using MODIS Time‣eries Data Sets. Agronomy Journal, 2007, 99, 1654-1664.	1.8	153
78	Representation of an alpine treeline ecotone in SPOT 5 HRG data. Remote Sensing of Environment, 2007, 110, 458-467.	11.0	32
79	Crop area estimation using high and medium resolution satellite imagery in areas with complex topography. Journal of Geophysical Research, 2008, 113, .	3.3	33
80	Yearly Grain Output Predictions in China 1980–2004. Economic Systems Research, 2008, 20, 139-150.	2.7	2
81	Comparison of Winter Wheat Classification using Multi-Temporal IRS-P6 Images. , 2008, , .		0
82	Accuracy assessment of linear spectral mixture model due to terrain undulation. Proceedings of SPIE, 2008, , .	0.8	0
83	Imagens de sensoriamento remoto no monitoramento da colheita da cana-de-açúcar. Engenharia Agricola, 2009, 29, 440-451.	0.7	19
84	Research on New Classification Methods of Remote Sensing of Mass Ingredient without Vegetation of Hei Shan Gorge in Yellow River Basin. , 2009, , .		0
85	Estimation of inter-annual winter crop area variation and spatial distribution with low resolution NDVI data by using neural networks trained on high resolution images. Proceedings of SPIE, 2009, , .	0.8	3
86	Derivation of the green vegetation fraction from TM data of three gorges area. Procedia Earth and Planetary Science, 2009, 1, 1152-1157.	0.6	5
87	Dynamic monitoring of vegetation coverage change in mining area based on CBERS-02 imagery. , 2010, , .		1
88	The spatial distribution of crop types from MODIS data: Temporal unmixing using Independent Component Analysis. Remote Sensing of Environment, 2010, 114, 1190-1204.	11.0	136
89	Evaluation of a linear spectral mixture model and vegetation indices (NDVI and EVI) in a study of schistosomiasis mansoni and Biomphalaria glabrata distribution in the state of Minas Gerais, Brazil. Memorias Do Instituto Oswaldo Cruz, 2010, 105, 512-518.	1.6	9
90	Detecting drought induced environmental changes in a Mediterranean wetland by remote sensing. Applied Geography, 2010, 30, 254-262.	3.7	44
91	Multi-resolution and temporal characterization of land-use classes in a Mediterranean wetland with land-cover fractions. International Journal of Remote Sensing, 2010, 31, 5365-5389.	2.9	10

#	Article	IF	CITATIONS
92	ARTMAP Neural Network Multisensor Fusion Model for Multiscale Land Cover Characterization. , 2010, , 529-543.		1
93	Estimation of wheat production by the integration of MODIS and ground data. International Journal of Remote Sensing, 2011, 32, 1105-1123.	2.9	9
94	Increasing the Resolution of Wide-Area Situational Awareness of the Power Grid through Event Unmixing. , 2011, , .		2
95	Maize acreage estimation using ENVISAT MERIS and CBERS-02B CCD data in the North China Plain. Computers and Electronics in Agriculture, 2011, 78, 208-214.	7.7	19
96	Endmember variability in Spectral Mixture Analysis: A review. Remote Sensing of Environment, 2011, 115, 1603-1616.	11.0	536
97	Reservoir capacity estimation of Singoor reservoir, India, by per-pixel and sub-pixel processing of multi-date satellite images. Geocarto International, 2011, 26, 305-320.	3.5	9
98	Interpreting temperature evolution of a bulk-metallic glass during cyclic loading through spatial–temporal modeling. Intermetallics, 2012, 29, 1-13.	3.9	4
99	Automatic unmixing of MODIS multi-temporal data for inter-annual monitoring of land use at a regional scale (Tensift, Morocco). International Journal of Remote Sensing, 2012, 33, 1325-1348.	2.9	16
100	Spatiotemporal dimensionality and Time-Space characterization of multitemporal imagery. Remote Sensing of Environment, 2012, 124, 793-809.	11.0	41
101	Using high resolution satellite imagery to estimate cropped area in Guatemala and Haiti. Applied Geography, 2012, 32, 433-440.	3.7	18
102	Spectral unmixing. International Journal of Remote Sensing, 2012, 33, 5307-5340.	2.9	128
103	Portability of neural nets modelling regional winter crop acreages using AVHRR time series. European Journal of Remote Sensing, 2012, 45, 371-392.	3.5	4
104	Mapping Soil Salinization of Agricultural Coastal Areas in Southeast Spain. , 2012, , .		4
105	Mapping cropping intensity of smallholder farms: A comparison of methods using multiple sensors. Remote Sensing of Environment, 2013, 134, 210-223.	11.0	118
106	Sub-pixel land-cover mapping with improved fraction images upon multiple-point simulation. International Journal of Applied Earth Observation and Geoinformation, 2013, 22, 115-126.	2.8	52
107	Fractional vegetation cover estimation in arid and semi-arid environments using HJ-1 satellite hyperspectral data. International Journal of Applied Earth Observation and Geoinformation, 2013, 21, 506-512.	2.8	148
108	Integration of in situ measured soil status and remotely sensed hyperspectral data to improve plant production system monitoring: Concept, perspectives and limitations. Remote Sensing of Environment, 2013, 128, 197-211.	11.0	26
109	Vegetation landscape change analysis based on remote sensing in northwest of Beijing. , 2013, , .		1

#	Article	IF	CITATIONS
110	Predicting water content using linear spectral mixture model on soil spectra. Journal of Applied Remote Sensing, 2013, 7, 073539.	1.3	4
111	Mapping the Spatial Distribution of Winter Crops at Sub-Pixel Level Using AVHRR NDVI Time Series and Neural Nets. Remote Sensing, 2013, 5, 1335-1354.	4.0	71
112	Daily Evaporative Fraction Parameterization Scheme Driven by Day–Night Differences in Surface Parameters: Improvement and Validation. Remote Sensing, 2014, 6, 4369-4390.	4.0	8
113	Temporal and Spatial Distribution Patterns of Grain Crops in the West Liaohe River Basin. Journal of Resources and Ecology, 2014, 5, 244-252.	0.4	6
114	County-level rice area estimation in southern China using remote sensing data. Journal of Applied Remote Sensing, 2014, 8, 083657.	1.3	15
115	Spectral unmixing of MODIS data based on improved endmember purification model: application to forest type identification. , 2014, , .		0
116	Incorporating spatial information in spectral unmixing: A review. Remote Sensing of Environment, 2014, 149, 70-87.	11.0	169
117	Mapping the human footprint from satellite measurements in Japan. ISPRS Journal of Photogrammetry and Remote Sensing, 2014, 88, 80-90.	11.1	21
118	Characterizing changes in cropping patterns using sequential Landsat imagery: an adaptive threshold approach and application to Phoenix, Arizona. International Journal of Remote Sensing, 2014, 35, 7263-7278.	2.9	25
119	Crop Type Classification by Simultaneous Use of Satellite Images of Different Resolutions. IEEE Transactions on Geoscience and Remote Sensing, 2014, 52, 3637-3649.	6.3	29
120	Obtaining crop-specific time profiles of NDVI: the use of unmixing approaches for serving the continuity between SPOT-VGT and PROBA-V time series. International Journal of Remote Sensing, 2014, 35, 2615-2638.	2.9	19
121	Information Content. , 2014, , 241-278.		1
122	Fallow land mapping for better crop monitoring in Huang-Huai-Hai Plain using HJ-1 CCD data. IOP Conference Series: Earth and Environmental Science, 2014, 17, 012048.	0.3	3
123	Determination of urban change in the vicinity of the gulf of izmit using linear spectral unmixing method and landsat data. Journal of Aeronautics and Space Technologies (Havacilik Ve Uzay) Tj ETQq1 1 0.7843	14org/BT /C	overlock 10 Ti
124	Unmixing: A new direction from classical tidal harmonic analysis?. , 2015, , .		1
125	Satellite Hyperspectral Imagery to Support Tick-Borne Infectious Diseases Surveillance. PLoS ONE, 2015, 10, e0143736.	2.5	19
126	Detection of Urban Expansion by using DMSP-OLS, Landsat Data and Linear Spectral Unmixing Method. International Journal of 3-D Information Modeling, 2015, 4, 58-67.	0.2	0
127	Optimization of spatial sampling schemes for maize acreage estimation. Journal of Applied Remote Sensing, 2015, 9, 097092.	1.3	0

#	Article	IF	CITATIONS
128	Assessment of Multiple Scattering in the Reflectance of Semiarid Shrublands. IEEE Transactions on Geoscience and Remote Sensing, 2015, 53, 4910-4921.	6.3	14
129	Estimation of Fractional Vegetation Cover in Semiarid Areas by Integrating Endmember Reflectance Purification Into Nonlinear Spectral Mixture Analysis. IEEE Geoscience and Remote Sensing Letters, 2015, 12, 1175-1179.	3.1	19
130	Estimation of winter wheat acreage via a combination of remotely sensed data and an optimized spatial sampling scheme. International Journal of Remote Sensing, 2015, 36, 5208-5221.	2.9	5
131	Mapping Rice Cropping Systems in Vietnam Using an NDVI-Based Time-Series Similarity Measurement Based on DTW Distance. Remote Sensing, 2016, 8, 19.	4.0	127
132	Using Moderate-Resolution Temporal NDVI Profiles for High-Resolution Crop Mapping in Years of Absent Ground Reference Data: A Case Study of Bole and Manas Counties in Xinjiang, China. ISPRS International Journal of Geo-Information, 2016, 5, 67.	2.9	28
133	A Framework for Large-Area Mapping of Past and Present Cropping Activity Using Seasonal Landsat Images and Time Series Metrics. Remote Sensing, 2016, 8, 312.	4.0	45
134	A Method to Determine Appropriate Spatial Resolution for Hard Image Classification. Journal of the Indian Society of Remote Sensing, 2016, 44, 11-19.	2.4	5
135	Driving Force Analysis of the Temporal and Spatial Distribution of Flash Floods in Sichuan Province. Sustainability, 2017, 9, 1527.	3.2	33
136	Multitemporal Remote Sensing Data Analysis for Agricultural Application. , 2018, , 29-38.		3
137	Remote Sensing of Croplands. , 2018, , 78-95.		11
138	Sub-Pixel Crop Type Classification Using PROBA-V 100 m NDVI Time Series and Reference Data from Sentinel-2 Classifications. Remote Sensing, 2019, 11, 1370.	4.0	13
139	Estimation of copper concentration of rocks using hyperspectral technology. Frontiers of Earth Science, 2019, 13, 563-574.	2.1	3
140	A Crop Group-Specific Pure Pixel Time Series for Europe. Remote Sensing, 2019, 11, 2668.	4.0	5
141	Mapping cropping intensity in China using time series Landsat and Sentinel-2 images and Google Earth Engine. Remote Sensing of Environment, 2020, 239, 111624.	11.0	187
142	Vegetation Growth Status and Topographic Effects in the Pisha Sandstone Area of China. Remote Sensing, 2020, 12, 2759.	4.0	13
143	Scalable mapping and monitoring of Mediterranean-climate oak landscapes with temporal mixture models. Remote Sensing of Environment, 2020, 247, 111937.	11.0	16
144	Constrained Linear Spectral Unmixing Technique for Regional Land Cover Mapping Using MODIS Data. , 2008, , 416-423.		10
145	Integration of AVHRR and Fine Spatial Resolution Imagery for Tropical Forest Monitoring. , 1996, , 377-394.		1

#	Article	IF	CITATIONS
146	Linear Mixture Model Applied to the Land-Cover Classification in an Alluvial Plain Using Landsat TM Data. Journal of Environmental Informatics, 2006, 7, 95-101.	6.0	5
147	Detection of ecotone environment based on satellite and field techniques (A case study; Northern) Tj ETQq1 1 0.	784314 rg 0.4	gBŢ /Overlo <mark>c</mark> i
148	Physical characteristics of spectrums and field techniques in detection of ecotone (a case study of) Tj ETQq0 0 0	rgBT /Ove 0.4	rlock 10 Tf 5 1
149	Geo-Big Data in Digital Augmentation and Accelerating Sustainable Agroecosystems. Studies in Big Data, 2022, , 221-242.	1.1	2
150	Determination of Treeline Using Satellite Data and Field Techniques (A Case Study; Western Nojomeh,) Tj ETQq0	0 0 rgBT /	Oyerlock 10
151	Analysis of Hyperspectral Remote Sensing Images. , 2010, , 235-269.		1
152	Sensing Uncertainty. , 2012, , 111-134.		0
153	The BRDF Model Construction and Application in Urban Areas. Advances in Intelligent and Soft Computing, 2012, , 379-386.	0.2	0
154	Applications of Space-Borne Remote Sensing Technology on Agricultural Research. Large Area Crop Inventory and Quantitative Approach for Growth Monitoring Seibutsu Kankyo Chosetsu [Environment Control in Biology, 1994, 32, 145-154.	0.2	0
155	A Study on Estimating Production of the Main Cereal Crops By Remote Sensing and GIS in China. J Agricultural Meteorology, 1997, 52, 545-554.	1.5	0
156	Subpixel mixture modeling applied for vegetation monitoring. IFIP Advances in Information and Communication Technology, 1997, , 41-48.	0.7	2
157	An Overview of Research on Remote Sensing Technology for Estimating Yield of Crop in the Arid Region. Hans Journal of Agricultural Sciences, 2015, 05, 6-13.	0.1	0
158	Detection of Urban Expansion by Using DMSP-OLS, Landsat Data and Linear Spectral Unmixing Method. , 2019, , 1372-1382.		0
159	Validation and refinement of cropland data layer using a spatial-temporal decision tree algorithm. Scientific Data, 2022, 9, 63.	5.3	34
160	Predictive Ecological Land Classification From Multi-Decadal Satellite Imagery. Frontiers in Forests and Global Change, 2022, 5, .	2.3	0
161	Grain-size composition of the surface sediments in Chinese deserts and the associated dust emission. Catena, 2022, 219, 106615.	5.0	7
162	Development of an Adaptive Linear Mixture Model for Decomposition of Mixed Pixels to Improve Crop Area Estimation Using Artificial Neural Network. IEEE Access, 2023, 11, 5714-5723.	4.2	0
163	Agricultural crop pattern mapping and change analysis at a sub-district level in South-eastern region, Bangladesh using Landsat satellite data from 2010 to 2019 2022. 116. 5-38.		1

#	Article	IF	CITATIONS
164	Fraction Images Derived from Landsat Mss, TM and Oli Images for Monitoring Forest Cover at the Rondônia State, Brazilian Amazon. , 2023, , .		0
165	Mapping multi-decadal wetland loss: Comparative analysis of linear and nonlinear spatiotemporal characterization. Remote Sensing of Environment, 2024, 302, 113969.	11.0	0