## Deposition of moss spores in relation to distance from p

Journal of Bryology 17, 355-368 DOI: 10.1179/jbr.1992.17.2.355

Citation Report

#	Article	IF	CITATIONS
1	Determination of thymidine incorporation activity in the sera from patients with short stature. European Journal of Endocrinology, 1981, 98, 352-357.	1.9	1
2	Spore Dispersal Distances in Atrichum angustatum (Polytrichaceae). Bryologist, 1992, 95, 324.	0.1	39
3	Investment in Sexual Reproduction by Antarctic Mosses. Oikos, 1993, 68, 293.	1.2	53
4	Recent Bryological Literature, 79. Journal of Bryology, 1993, 17, 523-537.	0.4	0
5	Modelling reproductive effort in sub-and maritime Antarctic mosses. Oecologia, 1994, 100-100, 45-53.	0.9	20
6	Local rate of spreading and patch dynamics of an invasive moss species,Orthodontiumlineare. Journal of Bryology, 1994, 18, 115-125.	0.4	12
8	THE INFLUENCE OF ENVIRONMENTAL CHARACTERISTICS ON LIFE HISTORY ATTRIBUTES OF ANTARCTIC TERRESTRIAL BIOTA. Biological Reviews, 1996, 71, 191-225.	4.7	257
9	Aerial dispersal of lichen soredia in the maritime Antarctic. New Phytologist, 1996, 134, 523-530.	3.5	99
10	Dispersal of moss propagules on Signy Island, maritime Antarctic. Polar Biology, 1997, 18, 376-383.	0.5	53
11	Presence and abundance of four epiphytic bryophytes in relation to density of aspen (Populus tremula) and other stand characteristics. Forest Ecology and Management, 1998, 107, 147-158.	1.4	56
12	Spore number in <i>Sphagnum</i> and its dependence on spore and capsule size. Journal of Bryology, 1998, 20, 1-16.	0.4	33
13	Levels of Genetic Variation and Its Partitioning in the Wide-Ranging Moss Polytrichum commune. Systematic Botany, 1999, 24, 512.	0.2	14
14	Vascular Plant Diversity as a Surrogate for Bryophyte and Lichen Diversity. Conservation Biology, 1999, 13, 282-292.	2.4	138
15	Spatial pattern of the threatened epiphytic bryophyte Neckera pennata at two scales in a fragmented boreal forest. Ecography, 1999, 22, 729-735.	2.1	40
16	Dispersal Patterns of Terricolous Lichens by Thallus Fragments. Lichenologist, 1999, 31, 603-612.	0.5	52
17	Are Enzyme Loci Selectively Neutral in Haploid Populations of Nonvascular Plants?. Evolution; International Journal of Organic Evolution, 1999, 53, 1050.	1.1	4
18	ARE ENZYME LOCI SELECTIVELY NEUTRAL IN HAPLOID POPULATIONS OF NONVASCULAR PLANTS?. Evolution; International Journal of Organic Evolution, 1999, 53, 1050-1059.	1.1	10
19	Experimental evidence for a persistent spore bank in Sphagnum. New Phytologist, 2000, 148, 105-116.	3.5	60

#	Article	IF	CITATIONS
20	2. <i>Pogonatum dentatum</i> (Brid.) Brid. (Bryopsida: Polytrichaceae). Journal of Bryology, 2000, 22, 55-60.	0.4	6
21	Population Structure and Taxonomy ofSphagnum cuspidatumandS. viride. Bryologist, 2000, 103, 93-103.	0.1	22
22	Epiphytic bryophytes on European aspen <i>Populus tremula</i> in old-growth forests in northeastern Finland and in adjacent sites in Russia. Canadian Journal of Botany, 2000, 78, 529-536.	1.2	11
23	Spore liberation in <i>Grimmia orbicularis</i> and <i>Tortula inermis:</i> two patterns from the Mojave desert. Journal of Bryology, 2001, 23, 83-90.	0.4	4
24	Bryological Notes. Journal of Bryology, 2001, 23, 139-152.	0.4	8
25	Environmental relationships of perichaetial and sporophyte production in <i>Andreaea</i> spp in western Norway. Journal of Bryology, 2001, 23, 97-108.	0.4	6
26	Forest age and management effects on epiphytic bryophyte communities in Adirondack northern hardwood forests, New York, U.S.A Canadian Journal of Forest Research, 2002, 32, 1562-1576.	0.8	68
27	The Association Between Substrate Variability and Bryophyte and Lichen Diversity in Eastern Australian Forests. Bryologist, 2002, 105, 11-26.	0.1	48
28	Enhanced sexual reproduction in bryophytes at high latitudes in the maritime Antarctic. Journal of Bryology, 2002, 24, 107-117.	0.4	40
29	Sporophyte production and spore dispersal phenology in Sphagnum: the importance of summer moisture and patch characteristics. Canadian Journal of Botany, 2002, 80, 543-556.	1.2	43
30	Spatial occurrence and colonisations in patch-tracking metapopulations: local conditions versus dispersal. Oikos, 2003, 103, 566-578.	1.2	167
31	Determining the immigration potential of plants colonizing disturbed environments: the case of milled peatlands in Quebec. Journal of Applied Ecology, 2003, 40, 78-91.	1.9	94
32	PLANT SPECIES COMPOSITION OF BOREAL SPRUCE SWAMP FORESTS: CLOSED DOORS AND WINDOWS OF OPPORTUNITY. Ecology, 2003, 84, 1909-1919.	1.5	42
33	Bryophyte dispersal inferred from colonization of an introduced substratum on Whiteface Mountain, New York. American Journal of Botany, 2004, 91, 1173-1182.	0.8	51
34	The effects of large-scale fragmentation on bryophytes in temperate forests. Journal of Applied Ecology, 2004, 41, 910-921.	1.9	47
35	Propagule Sources of Forest Floor Bryophytes: Spatiotemporal Compositional Patterns. Bryologist, 2004, 107, 88-97.	0.1	57
36	Size of Acer saccharum Hosts Does Not Influence Growth of Mature Bryophyte Gametophytes in Adirondack Northern Hardwood Forests. Bryologist, 2004, 107, 302-311.	0.1	7
37	Predicting bryophyte diversity in grassland and eucalypt-dominated remnants in subhumid Tasmania. Journal of Biogeography, 2005, 32, 2015-2024.	1.4	35

#	Article	IF	CITATIONS
38	Larger capsules enhance short-range spore dispersal inSphagnum, but what happens further away?. Oikos, 2005, 108, 115-124.	1.2	97
39	Niche breadth and niche overlap in three epixylic hepatics in a boreal old-growth forest, southern Finland. Journal of Bryology, 2005, 27, 119-127.	0.4	16
40	Bryophyte rarity viewed from the perspectives of life history strategy and metapopulation dynamics. Journal of Bryology, 2005, 27, 261-268.	0.4	49
41	Bryophyte recolonization on burnt soil and logs. Scandinavian Journal of Forest Research, 2005, 20, 5-16.	0.5	24
42	Phytosociology and life strategies of a new loess slope bryophyte community from N China (Gansu), including Crossidium laxefilamentosum new to China. Nova Hedwigia, 2005, 81, 229-246.	0.2	2
43	Short-term effects of timber harvest and forest edges on ground-layer mosses and liverworts. Canadian Journal of Botany, 2005, 83, 610-620.	1.2	96
44	Persistence of ground-layer bryophytes in a structural retention experiment: initial effects of level and pattern of overstory retention. Canadian Journal of Forest Research, 2006, 36, 3039-3052.	0.8	37
45	Sphagnumspore availability in boreal forests. Bryologist, 2006, 109, 173-181.	0.1	11
46	Dispersal potential of spores and asexual propagules in the epixylic hepatic Anastrophyllum hellerianum. Evolutionary Ecology, 2006, 20, 415-430.	0.5	86
47	Community Structure of In-Stream Bryophytes in English and Welsh Rivers. Hydrobiologia, 2006, 553, 143-152.	1.0	41
48	Life form and life strategy analyses of the epiphytic bryophyte communities of Madeira's laurel and ericaceous forests. Botanische Jahrbücher Für Systematik, Pflanzengeschichte Und Pflanzengeographie, 2007, 127, 151-164.	0.4	5
49	Rehabilitating boreal forest structure and species composition in Finland through logging, dead wood creation and fire: The EVO experiment. Forest Ecology and Management, 2007, 250, 77-88.	1.4	141
50	Bryophytes in a changing landscape: The hierarchical effects of habitat fragmentation on ecological and evolutionary processes. Biological Conservation, 2007, 135, 315-325.	1.9	89
51	Bryophyte responses to fragmentation in temperate coastal rainforests: A functional group approach. Biological Conservation, 2007, 136, 408-422.	1.9	45
52	Spatial distribution patterns of Rhynchostegium megapolitanum at the landscape scale ―an expanding species?. Applied Vegetation Science, 2007, 10, 111-120.	0.9	6
53	How far and how fast do bryophytes travel at the landscape scale?. Diversity and Distributions, 2008, 14, 483-492.	1.9	69
54	Age Structure and Ecological Characteristics of Some Epiphytic Liverworts (Frullania Dilatata,) Tj ETQq0 0 0 rgBT	/Overlock	10 Tf 50 102

55	Bryophyte diaspore bank: a genetic memory? Genetic structure and genetic diversity of surface populations and diaspore bank in the liverwort <i>Mannia fragrans</i> (Aytoniaceae). American Journal of Botany, 2008, 95, 542-548.	0.8	27	
----	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----	----	--

#	Article	IF	CITATIONS
56	Population and community ecology of bryophytes. , 0, , 393-444.		16
58	Clonal diversity and geographic structure in PleurochaeteÂsquarrosa (Pottiaceae): different sampling scale approach. Journal of Plant Research, 2009, 122, 161-170.	1.2	10
59	Niche assembly of epiphytic bryophyte communities in the Guianas: a regional approach. Journal of Biogeography, 2009, 36, 2076-2084.	1.4	74
60	Multiple paternity and sporophytic inbreeding depression in a dioicous moss species. Heredity, 2009, 103, 394-403.	1.2	40
61	CORRELATED EVOLUTION OF SEXUAL SYSTEM AND LIFE-HISTORY TRAITS IN MOSSES. Evolution; International Journal of Organic Evolution, 2009, 63, 1129-1142.	1.1	59
62	Epiphytic bryophytes in Canarian subtropical montane cloud forests: the importance of the time since disturbance and host identity. Canadian Journal of Forest Research, 2009, 39, 48-63.	0.8	13
63	Effects of light and nutrients on different germination phases of the Cosmopolitan moss Bryum argenteum Hedw. (Bryaceae). Brazilian Archives of Biology and Technology, 2010, 53, 763-769.	0.5	6
64	Potential Impacts of Climate Change on <i>Sphagnum</i> Bogs of the Southern Appalachian Mountains. Natural Areas Journal, 2010, 30, 417-424.	0.2	8
65	Modelling the distribution of the moss species Hypopterygium tamarisci (Hypopterygiaceae, Bryophyta) in Central and South America. Nova Hedwigia, 2010, 91, 399-420.	0.2	14
66	Introduction and recent range expansion in the moss Ptychomitrium serratum (Ptychomitriaceae) in the Southern and Eastern United StatesThis paper is one of a selection of papers published as part of the special Schofield Gedenkschrift Botany, 2010, 88, 336-344.	0.5	1
67	Bryophyte diversity and evolution: Windows into the early evolution of land plants. American Journal of Botany, 2011, 98, 352-369.	0.8	169
68	Fine-scale spatial genetic structure of a liverwort (Barbilophozia attenuata) within a network of ant trails. Evolutionary Ecology, 2011, 25, 45-57.	0.5	21
69	Bryophyte dispersal by sheep on dry grassland. Nova Hedwigia, 2011, 92, 327-341.	0.2	21
70	The contributions of forest structure and substrate to bryophyte diversity and abundance in mature coniferous forests of the Pacific Northwest. Bryologist, 2012, 115, 278.	0.1	28
71	Diaspore bank of bryophytes in tropical rain forests: the importance of breeding system, phylum and microhabitat. Oecologia, 2012, 168, 321-333.	0.9	12
72	Linking the biological traits of boreal bryophytes to forest habitat change after partial harvesting. Forest Ecology and Management, 2013, 303, 184-194.	1.4	34
73	Bryophyte persistence following major fire in eucalypt forest of southern Australia. Forest Ecology and Management, 2013, 296, 24-32.	1.4	22
74	Genetic Diversity of Leafy Liverwort Species (Jungermanniidae, Marchantiophyta) in Poland: Diversity of Leafy Liverwort Species with Various Reproductive Modes. Biodiversity Research and Conservation, 2013, 27, 3-54.	0.2	4

ARTICLE IF CITATIONS Impact of distance to mature forest on the recolonisation of bryophytes in a regenerating Tasmanian 0.3 18 wet eucalypt forest. Australian Journal of Botany, 2013, 61, 633. Release thresholds for moss spores: the importance of turbulence and sporophyte length. Journal of Ecology, 2014, 102, 721-729. Bryophyte communities in the Amazon forest are regulated by height on the host tree and site 1.9 44 elevation. Journal of Ecology, 2015, 103, 441-450. So close and yet so far away: longâ $\in$ distance dispersal events govern bryophyte metacommunity reassembly. Journal of Ecology, 2016, 104, 1707-1719. Preemergence Control of Silvery Threadmoss (Bryum argenteum) Grown from Spores and Bulbils in 0.4 3 Axenic Culture. Weed Technology, 2016, 30, 198-206. Polytrichum commune spores nucleate ice and associated microorganisms increase the temperature of ice nucleation activity onset. Aerobiologia, 2016, 32, 353-361. The epiphytic bryophyte community of Atlantic oak woodlands shows clear signs of recovery 1.9 4 following the removal of invasive Rhododendron ponticum. Biological Conservation, 2017, 212, 96-104. Stand Structure and Substrate Diversity as Two Major Drivers for Bryophyte Distribution in a 1.7 16 Temperate Montane Ecosystem. Frontiers in Plant Šcience, 2017, 8, 874. Divergence in Life-History and Developmental Traits in Silvery-Thread Moss (Bryum argenteum Hedw.) 0.8 6 Genotypes between Golf Course Putting Greens and Native Habitats. Weed Science, 2018, 66, 642-650. Plant community development following reclamation of oil sands mine sites in the boreal forest: a 2.1 review. Environmental Reviews, 2018, 26, 286-298. Effects of Rate of Drying, Life History Phase, and Ecotype on the Ability of the Moss Bryum Argenteum to Survive Desiccation Events and the Influence on Conservation and Selection of Material for 17 1.1 Restoration. Frontiers in Ecology and Evolution, 2019, 7, . Terrestrial Bryophyte and Lichen Responses to Canopy Opening in Pine-Moss-Lichen Forests. Forests, 2019, 10, 233. Forest passerines as a novel dispersal vector of viable bryophyte propagules. Proceedings of the Royal 1.2 25 Society B: Biological Sciences, 2019, 286, 20182253. Floating ability, shoot length and abundance facilitate hydrochorous dispersal of moss and liverwort fragments. Journal of Vegetation Science, 2019, 30, 30-41. 1.1 Morphological characters and SNP markers suggest hybridization and introgression in sympatric populations of the pleurocarpous mosses Homalothecium lutescens and H. sericeum. Organisms 2 0.7 Diversity and Evolution, 2020, 20, 619-637. Tradeoffs between phenotypic plasticity and local adaptation influence the ecophysiology of the moss, Sphagnum magellanicum. Oecológia, 2020, 193, 867-877. The Fate of Bryophyte Sporophytesâ€"Phenology and Vectors of Buxbaumia viridis in the Kalkalpen 1.6 5 National Park, Austria. Plants, 2020, 9, 1320.

CITATION REPORT

92Comparing the effect of landscape context on vascular plant and bryophyte communities in a<br/>humanâ€dominated landscape. Journal of Vegetation Science, 2021, 32, .1.15

75

77

79

81

83

84

85

86

87

89

#	Article	IF	CITATIONS
93	Colonisation Processes. , 2006, , 35-54.		29
95	Epiphytic bryophytes on European aspen <i>Populus tremula</i> in old-growth forests in northeastern Finland and in adjacent sites in Russia. Canadian Journal of Botany, 2000, 78, 529-536.	1.2	30
96	The Fate of the Missing Spores — Patterns of Realized Dispersal beyond the Closest Vicinity of a Sporulating Moss. PLoS ONE, 2012, 7, e41987.	1.1	53
97	Spatial distribution patterns of Rhynchostegium megapolitanum at the landscape scale – an expanding species?. Applied Vegetation Science, 2007, 10, 111.	0.9	11
98	CHANGE OF MARCHANTIA SP SIZE AND COLOUR RELATED TO DISTANCE FROM WATER RESOURCE IN CUGHUP EMBUN, SOUTH SUMATERA. Biovalentia: Biological Research Journal, 2016, 2, .	0.1	0
100	Spore dispersal of Selaginella denticulata, S. helvetica, and S. selaginoides, and the significance of heterospory in Selaginellacae. American Fern Journal, 2020, 110, 58.	0.2	3
102	The danger and indeterminacy of forfeiting perching space of bryophytes from climate shift: a case study for 115 species in China. Environmental Monitoring and Assessment, 2022, 194, 233.	1.3	2
105	Sexual and asexual offspring germination in a tropical liverwort is mediated by phytochromes. Bryologist, 2022, 125, .	0.1	0
106	Disentangling spatial and island shape effects on bryophyte distribution in the Zhoushan Archipelago, China. Folia Geobotanica, 2022, 57, 83-101.	0.4	1
107	Taxonâ€dependent effects of dispersal limitation versus environmental filters on bryophyte assemblages―Multiple perspective studies in landâ€bridge islands. Ecology and Evolution, 2023, 13, .	0.8	1

110 Bryophytes. , 2024, , 475-603.