

CITATION REPORT

List of articles citing

Relations between hemostasis variables and cardiovascular risk factors in middle-aged adults. Atherosclerosis Risk in Communities (ARIC) Study Investigat

DOI: 10.1016/1047-2797(92)90098-b
Annals of Epidemiology, 1992, 2, 481-94.

Source: <https://exaly.com/paper-pdf/23264787/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
68	Leukocyte count correlates in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. <i>American Journal of Epidemiology</i> , 1992 , 136, 525-37	3.8	125
67	Association of coagulation factors and inhibitors with carotid artery atherosclerosis. Early results of the Atherosclerosis Risk in Communities (ARIC) Study. <i>Annals of Epidemiology</i> , 1992 , 2, 471-80	6.4	60
66	The Atherosclerosis Risk in Communities (ARIC) Study. Introduction and objectives of the hemostasis component. <i>Annals of Epidemiology</i> , 1992 , 2, 467-9	6.4	30
65	Alcohol consumption and ultrasonographically assessed carotid artery wall thickness and distensibility. The Atherosclerosis Risk in Communities (ARIC) Study Investigators. <i>Circulation</i> , 1993 , 88, 2787-93	16.7	70
64	Distribution and correlates of plasma fibrinogen in middle-aged women. Initial findings of the Postmenopausal Estrogen/Progestin Interventions (PEPI) study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1995 , 15, 2085-93	9.4	36
63	Postmenopausal hormone-replacement therapy and cardiovascular risk. <i>Annals of Medicine</i> , 1995 , 27, 149-56	1.5	2
62	Plasma fibrinogen levels and their correlates in 6457 coronary heart disease patients. The Bezafibrate Infarction Prevention (BIP) Study. <i>Journal of Clinical Epidemiology</i> , 1995 , 48, 757-65	5.7	49
61	Hemostatic factors as triggers of cardiovascular events. <i>Cardiology Clinics</i> , 1996 , 14, 239-50	2.5	15
60	Correlates of leukocyte counts in men. <i>Annals of Epidemiology</i> , 1996 , 6, 74-82	6.4	23
59	Thrombophilia in patients with hypercholesterolemia. <i>Metabolism: Clinical and Experimental</i> , 1996 , 45, 966-9	12.7	14
58	Late thromboembolic complication from a Palmaz stent in the common iliac artery. <i>CardioVascular and Interventional Radiology</i> , 1996 , 19, 190-2	2.7	4
57	Association of fibrinogen and coagulation factors VII and VIII with cardiovascular risk factors in the elderly: the Cardiovascular Health Study. Cardiovascular Health Study Investigators. <i>American Journal of Epidemiology</i> , 1996 , 143, 665-76	3.8	109
56	Long-term prognosis of women after myocardial infarction. SPRINT Study Group. Secondary Prevention Reinfarction Israeli Nifedipine Trial. <i>American Journal of Epidemiology</i> , 1997 , 146, 153-60	3.8	25
55	Recent advances in hemostatic risk factors and cardiovascular diseases. <i>Fibrinolysis and Proteolysis</i> , 1997 , 11, 31-34		1
54	Assays for thrombosis factors: relations to plasma lipids. <i>American Journal of Clinical Nutrition</i> , 1997 , 65, 1669S-1673S	7	5
53	Alcohol, hypertension and the cardiovascular system: a critical appraisal. <i>Addiction Biology</i> , 1997 , 2, 159-70	406	26
52	Thrombophilia in Patients with Hypertriglyceridemia. <i>Journal of Thrombosis and Thrombolysis</i> , 1997 , 4, 425-429	5.1	9

51	Fibrinogen and von Willebrand factor in IDDM: relationships to lipid vascular risk factors, blood pressure, glycaemic control and urinary albumin excretion rate: the EURODIAB IDDM Complications Study. <i>Diabetologia</i> , 1997 , 40, 698-705	10.3	57
50	Epidemiology of coagulation factors, inhibitors and activation markers: the Third Glasgow MONICA Survey. I. Illustrative reference ranges by age, sex and hormone use. <i>British Journal of Haematology</i> , 1997 , 97, 775-84	4.5	238
49	Epidemiology of coagulation factors, inhibitors and activation markers: The Third Glasgow MONICA Survey. II. Relationships to cardiovascular risk factors and prevalent cardiovascular disease. <i>British Journal of Haematology</i> , 1997 , 97, 785-97	4.5	127
48	Moderate beer consumption and the blood coagulation in patients with coronary artery disease. <i>Journal of Internal Medicine</i> , 1997 , 241, 47-51	10.8	67
47	Elevated plasma fibrinogen: cause or consequence of cardiovascular disease?. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1998 , 18, 621-5	9.4	72
46	Socioeconomic status and determinants of hemostatic function in healthy women. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999 , 19, 485-92	9.4	49
45	Plasma fibrinogen and its correlates in elderly Japanese men living in Japan and Hawaii. <i>Journal of Clinical Epidemiology</i> , 1999 , 52, 1201-6	5.7	9
44	Haemostasis and carotid artery wall thickness in non-insulin dependent diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2000 , 47, 25-35	7.4	12
43	The -323Ins10 polymorphism for factor VII is not associated with coronary atherosclerosis in symptomatic men. The REGRESS study group. <i>Thrombosis Research</i> , 2000 , 97, 275-80	8.2	8
42	Age-dependency in molecular markers of haemostasis, fibrinolysis and in soluble adhesion molecules. <i>Fibrinolysis and Proteolysis</i> , 2000 , 14, 281-287		1
41	Changes in hemostasis during treatment of hypertriglyceridemia with a diet rich in monounsaturated and n-3 polyunsaturated fatty acids in comparison with a low-fat diet. <i>Thrombosis Research</i> , 2001 , 101, 355-66	8.2	34
40	Association between physical activity and markers of inflammation in a healthy elderly population. <i>American Journal of Epidemiology</i> , 2001 , 153, 242-50	3.8	435
39	Association of hemostatic variables with MRI-detected cerebral abnormalities: the atherosclerosis risk in communities study. <i>Neuroepidemiology</i> , 2001 , 20, 96-104	5.4	25
38	Increased acute phase reactants are associated with levels of lipoproteins and increased carotid plaque volume. <i>European Journal of Vascular and Endovascular Surgery</i> , 2001 , 21, 227-34	2.3	32
37	Plasma fibrinogen as a predictor of total and cause-specific mortality in elderly Japanese-American men. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001 , 21, 1065-70	9.4	51
36	Relationship between plasma markers of endothelial cell integrity and the Framingham cardiovascular disease risk-factor scores in apparently healthy individuals. <i>Blood Coagulation and Fibrinolysis</i> , 2002 , 13, 513-8	1	29
35	Relationship between physical activity and inflammation among apparently healthy middle-aged and older US adults. <i>Archives of Internal Medicine</i> , 2002 , 162, 1286-92		327
34	Late complication of aortoiliac stent placement- atheroembolization of the lower extremities. <i>Journal of Surgical Research</i> , 2002 , 103, 153-9	2.5	26

33	Insulin resistance is accompanied by increased von Willebrand factor levels in nondiabetic women: a study of offspring of type 2 diabetic subjects compared to offspring of nondiabetic subjects. <i>Journal of Internal Medicine</i> , 2002 , 252, 155-63	10.8	6
32	Leisure-time physical activity and reduced plasma levels of obesity-related inflammatory markers. <i>Obesity</i> , 2003 , 11, 1055-64		162
31	Association between obesity and a prothrombotic state: the Framingham Offspring Study. <i>Thrombosis and Haemostasis</i> , 2004 , 91, 683-9	7	146
30	Effects of diabetes on the vascular system: current research evidence and best practice recommendations. <i>Journal of Vascular Nursing</i> , 2004 , 22, 2-11; quiz 12-3	1	9
29	Body fat is the main predictor of fibrinogen levels in healthy non-obese men. <i>Metabolism: Clinical and Experimental</i> , 2004 , 53, 984-8	12.7	18
28	Association of physical activity and body mass index with novel and traditional cardiovascular biomarkers in women. <i>JAMA - Journal of the American Medical Association</i> , 2006 , 295, 1412-9	27.4	280
27	Additive value of immunoassay-measured fibrinogen and high-sensitivity C-reactive protein levels for predicting incident cardiovascular events. <i>Circulation</i> , 2006 , 114, 381-7	16.7	65
26	Fibrinogen apheresis in the treatment of peripheral arterial disease. <i>Blood Purification</i> , 2007 , 25, 404-10	3.1	13
25	Haemorheological, platelet and endothelial indices in relation to global measures of cardiovascular risk in hypertensive patients: a substudy of the Anglo-Scandinavian Cardiac Outcomes Trial. <i>Journal of Internal Medicine</i> , 2007 , 261, 82-90	10.8	13
24	Obesity and thrombosis. <i>European Journal of Vascular and Endovascular Surgery</i> , 2007 , 33, 223-33	2.3	208
23	von Willebrand factor in cardiovascular disease: focus on acute coronary syndromes. <i>Circulation</i> , 2008 , 117, 1449-59	16.7	294
22	Hepatitis C infection is associated with increased coronary artery atherosclerosis defined by modified Reardon severity score system. <i>Circulation Journal</i> , 2008 , 72, 1960-5	2.9	82
21	Establishment of reference intervals for von Willebrand factor antigen and eight coagulation factors in a Korean population following the Clinical and Laboratory Standards Institute guidelines. <i>Blood Coagulation and Fibrinolysis</i> , 2010 , 21, 251-5	1	11
20	Fibrinogen promotes early atherosclerotic changes of the carotid artery in young, healthy adults. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 1003-8	4	18
19	Endothelial dysfunction in paroxysmal atrial fibrillation as a prothrombotic state. Comparison with permanent/persistent atrial fibrillation. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 298-304	4	20
18	Variability of fibrinogen measurements in post-myocardial infarction patients. Results from the AIRGENE study center Augsburg. <i>Thrombosis and Haemostasis</i> , 2012 , 107, 895-902	7	2
17	The relationship between body mass index, activated protein C resistance and risk of venous thrombosis. <i>Journal of Thrombosis and Haemostasis</i> , 2012 , 10, 1761-7	15.4	22
16	The role of flow-mediated dilatation and high sensitive C-reactive protein in paroxysmal atrial fibrillation. <i>Egyptian Heart Journal</i> , 2012 , 64, 211-217	1.3	

15	Fresh frozen plasma quality: relation to age and gender of blood donors. <i>Vox Sanguinis</i> , 2012 , 102, 116-241	3.1	11
14	Effects of pioglitazone and vildagliptin on coagulation cascade in diabetes mellitus--targeting thrombogenesis. <i>Expert Opinion on Therapeutic Targets</i> , 2013 , 17, 627-39	6.4	11
13	Macroeconomics and cardiovascular risk factors: the same view through a different lens?. <i>Circulation</i> , 2013 , 127, 1451-2	16.7	1
12	Thrombosis in central obesity and metabolic syndrome: mechanisms and epidemiology. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 669-80	7	81
11	Atherothrombotic risk in obesity. <i>Hamostaseologie</i> , 2013 , 33, 259-68	1.9	14
10	Changes in procoagulants track longitudinally with insulin resistance: findings from the coronary artery risk development in young adults (CARDIA) study. <i>Diabetic Medicine</i> , 2014 , 31, 462-5	3.5	8
9	Potential value of targeting von Willebrand factor in atherosclerotic cardiovascular disease. <i>Expert Opinion on Therapeutic Targets</i> , 2014 , 18, 43-53	6.4	18
8	Individual differences of plasma proteins and factors in fresh frozen plasma from Chinese regional blood donors. <i>Journal of Thrombosis and Thrombolysis</i> , 2019 , 47, 420-426	5.1	2
7	Lifestyle Intervention Improves Prothrombotic Coagulation Profile in Individuals at High Risk for Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e3198-e3207	5.6	5
6	Hemostatic risk factors for cardiovascular disease. <i>Developments in Cardiovascular Medicine</i> , 1996 , 135-151		1
5	Plasma fibrinogen and coronary heart disease in elderly Japanese-American men. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1996 , 16, 262-8	9.4	16
4	Fibrinogen and its relations to subclinical extracoronary and coronary atherosclerosis in hypercholesterolemic men. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1997 , 17, 45-50	9.4	22
3	Obesity and Hypertension. 2008 , 33-40		
2	Analysis of Coagulation Factor Activity of Normal Adults with APTT Limit Range. <i>Korean Journal of Clinical Laboratory Science</i> , 2015 , 47, 237-242	0.4	1
1	Sex-specific associations between serum lipids and hemostatic factors: the cross-sectional population-based KORA-fit study. 2022 , 21,		0