

A comparison of root and shoot competition between sp

Weed Research

32, 45-55

DOI: [10.1111/j.1365-3180.1992.tb01861.x](https://doi.org/10.1111/j.1365-3180.1992.tb01861.x)

Citation Report

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Effects of nitrogen fertilizer and emergence date on root and shoot competition between wheat and blackgrass. <i>Weed Research</i> , 1992, 32, 175-182.   | 0.8 | 20        |
| 2  | A study of root and shoot interactions between cereals and peas in mixtures. <i>Journal of Agricultural Science</i> , 1993, 120, 13-24.   | 0.6 | 47        |
| 3  | Comparison of Green Foxtail ( <i>Setaria viridis</i> ) and Wild Oat ( <i>Avena fatua</i> ) Growth, Development, and Competitiveness under Three Temperature Regimes. <i>Weed Science</i> , 1993, 41, 369-378. | 0.8 | 23        |
| 4  | Spring Wheat ( <i>Triticum aestivum</i> ) Growth and Yield as Influenced by Duration of Wild Oat ( <i>Avena</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 0.4 17   | 0.4 | 17        |
| 5  | Inter- and intraspecific competition of fat-hen ( <i>Chenopodium album</i> L.) and groundsel ( <i>Senecio</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 50   | 0.8 | 10        |
| 6  | The response of four spring-sown combinable arable crops to weed competition. <i>Weed Research</i> , 1994, 34, 137-146.   | 0.8 | 40        |
| 7  | Simulation of competition between barley and wild oats under different managements and climates. <i>Ecological Modelling</i> , 1994, 71, 269-287.   | 1.2 | 27        |
| 8  | Losses in grain yield of winter crops from <i>Lolium rigidum</i> competition depend on crop species, cultivar and season. <i>Weed Research</i> , 1995, 35, 503-509.   | 0.8 | 123       |
| 9  | A comparison of the dynamics of root growth and biomass partitioning in wild oat ( <i>Avena fatua</i> L.) and spring wheat. <i>Weed Research</i> , 1995, 35, 57-66.   | 0.8 | 3         |
| 10 | Comparative Analysis of Three Cruciferous Weeds: Growth, Development, and Competitiveness. <i>Weed Science</i> , 1995, 43, 75-80.   | 0.8 | 10        |
| 11 | Approaches for Improving Crop Competitiveness Through the Manipulation of Fertilization Strategies. <i>Weed Science</i> , 1995, 43, 491-497.  | 0.8 | 233       |
| 13 | Competition in a Global Change Environment: The Importance of Different Plant Traits for Competitive Success. <i>Journal of Biogeography</i> , 1995, 22, 297.   | 1.4 | 23        |
| 14 | Dog mustard ( <i>Erucastrum gallicum</i> ) response to crop competition. <i>Weed Science</i> , 1997, 45, 397-403.   | 0.8 | 3         |
| 15 | The effect of seed dormancy on percentage and rate of germination in <i>Polygonum persicaria</i> , and its relevance for crop-weed interaction. <i>Annals of Applied Biology</i> , 1998, 132, 289-299.        | 1.3 | 6         |
| 16 | Competition between safflower and weeds as influenced by crop genotype and sowing time. <i>Weed Research</i> , 1998, 38, 247-255.   | 0.8 | 20        |
| 17 | Competition between barley and <i>Lolium rigidum</i> for nitrate. <i>Weed Research</i> , 1998, 38, 453-460.   | 0.8 | 26        |
| 18 | Determination of allelochemicals in spring cereal cultivars of different competitiveness. <i>Weed Science</i> , 1999, 47, 498-504.  | 0.8 | 77        |
| 19 | Competitive attributes of <i>A. sativa</i> , <i>T. aestivum</i> , and <i>H. vulgare</i> are conserved in no-till cropping systems. <i>Weed Science</i> , 1999, 47, 712-719.                                   | 0.8 | 9         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 20 | The relative importance of root and shoot competition between water-seeded rice and <i>Echinochloa phyllopogon</i> . <i>Weed Research</i> , 1999, 39, 181-190.                         | 0.8 | 46        |
| 21 | Improving grasslands: the influence of soil moisture and nitrogen fertilization on the establishment of seedlings. <i>Journal of Applied Ecology</i> , 1999, 36, 263-270.              | 1.9 | 26        |
| 22 | Interference between <i>Avena sterilis</i> , <i>Phalaris minor</i> and five barley cultivars. <i>Weed Research</i> , 2000, 40, 549-559.  | 0.8 | 40        |
| 23 | Advances in weed management strategies. <i>Field Crops Research</i> , 2000, 67, 95-104.  | 2.3 | 49        |
| 24 | Enhancing the competitive ability of crops. , 2001, , 269-321.   |     | 88        |
| 25 | Competição por recursos do solo entre ervas daninhas e culturas. <i>Ciencia Rural</i> , 2001, 31, 707-714.   | 0.3 | 32        |
| 26 | Wheat- <i>Lolium multiflorum</i> competence: effect of nitrogen application onto Argentinean varieties aggressivity. <i>Cereal Research Communications</i> , 2001, 29, 451-458.        | 0.8 | 3         |
| 27 | Title is missing!. <i>Plant Ecology</i> , 2001, 152, 119-136.  | 0.7 | 83        |
| 28 | Influence of nitrogen on competition between winter cereals and sterile oat. <i>Weed Science</i> , 2001, 49, 77-82.  | 0.8 | 52        |
| 29 | Influence of nitrogen on competition between purple nutsedge, maize and soybean. <i>International Journal of Pest Management</i> , 2002, 48, 73-79.                                    | 0.9 | 11        |
| 30 | Residual effects of crop rotation and weed management on a wheat test crop and weeds. <i>Weed Science</i> , 2002, 50, 101-111.   | 0.8 | 8         |
| 31 | Estimating giant foxtail cohort productivity in soybean based on weed density, leaf area, or volume. <i>Weed Science</i> , 2002, 50, 72-78.  | 0.8 | 19        |
| 32 | Effect of nitrogen fertilization timing on the demographic processes of wild oat ( <i>Avena fatua</i> ) in barley ( <i>Hordeum vulgare</i> ). <i>Weed Science</i> , 2002, 50, 616-621. | 0.8 | 8         |
| 33 | Recruitment and competitive interaction between ramets and seedlings in a perennial medicinal herb, <i>Centella asiatica</i> . <i>Basic and Applied Ecology</i> , 2002, 3, 65-76.      | 1.2 | 16        |
| 34 | Variation Between Barley Cultivars in Early Response to Weed Competition. <i>Journal of Agronomy and Crop Science</i> , 2002, 188, 176-184.  | 1.7 | 54        |
| 35 | Nitrogen effect on competition between winter cereals and littleseed canarygrass. <i>Phytoparasitica</i> , 2003, 31, 252-264.  | 0.6 | 7         |
| 36 | Shoot and Root Competition in a <i>Lolium multiflorum</i> -Wheat Association. <i>Biological Agriculture and Horticulture</i> , 2003, 21, 15-33.  | 0.5 | 5         |
| 37 | Above- and belowground interference of purple and yellow nutsedge ( <i>Cyperus</i> spp.) with tomato. <i>Weed Science</i> , 2003, 51, 181-185.   | 0.8 | 38        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 38 | Influência de cultivares de arroz e épocas da adubação nitrogenada nas relações de interferência da cultura com cultivar simulador de infestação de arroz-vermelho. <i>Planta Daninha</i> , 2004, 22, 185-193. | 0.5 | 7         |
| 40 | Application method of nitrogen fertilizer affects weed growth and competition with winter wheat. <i>Weed Biology and Management</i> , 2004, 4, 103-113.  | 0.6 | 26        |
| 41 | Spring wheat seed size and seeding rate effects on yield loss due to wild oat ( <i>Avena fatua</i> ) interference. <i>Weed Science</i> , 2004, 52, 133-141.  | 0.8 | 53        |
| 42 | <i>Weed Biology and Management</i> , 2004, , .   |     | 4         |
| 43 | Competitiveness of Rice Cultivars as a Tool for Crop-Based Weed Management. , 2004, , 517-537.   |     | 18        |
| 44 | Variable crop plant establishment contributes to differences in competitiveness with wild oat among cereal varieties. <i>Canadian Journal of Plant Science</i> , 2005, 85, 771-776.                            | 0.3 | 26        |
| 45 | Nitrogen Fertilizer, Manure, and Compost Effects on Weed Growth and Competition with Spring Wheat. <i>Agronomy Journal</i> , 2005, 97, 1612-1621.  | 0.9 | 44        |
| 46 | Wild Mustard ( <i>Sinapis arvensis</i> L.) Competition with Three Winter Cereals as Affected by Nitrogen Supply. <i>Journal of Agronomy and Crop Science</i> , 2005, 191, 241-248.                             | 1.7 | 21        |
| 47 | Fertilizer, manure and compost effects on weed growth and competition with winter wheat in western Canada. <i>Crop Protection</i> , 2005, 24, 971-980.   | 1.0 | 84        |
| 48 | Barley ( <i>Hordeum vulgare</i> ) and Wild Oat ( <i>Avena fatua</i> ) Competition Is Affected by Crop and Weed Density1. <i>Weed Technology</i> , 2005, 19, 790-795.   | 0.4 | 35        |
| 49 | Competitive interactions between chick-pea genotypes and weeds. <i>Weed Research</i> , 2006, 46, 335-344.  | 0.8 | 31        |
| 50 | Competitive ability of wheat in conventional and organic management systems: A review of the literature. <i>Canadian Journal of Plant Science</i> , 2006, 86, 333-343.   | 0.3 | 105       |
| 51 | The ability of 29 barley cultivars to compete and withstand competition. <i>Weed Science</i> , 2006, 54, 783-792.  | 0.8 | 69        |
| 52 | Response of Corn ( <i>Zea mays</i> ) and Weeds to Planting Pattern and Herbicide Use. <i>Biological Agriculture and Horticulture</i> , 2006, 24, 117-134.  | 0.5 | 1         |
| 53 | Spatial pattern effect on corn ( <i>Zea mays</i> ) weeds competition in the humid Pampas of Argentina. <i>International Journal of Pest Management</i> , 2007, 53, 195-206.                                    | 0.9 | 16        |
| 54 | Cultivar and Seeding Rate Effects on the Competitive Ability of Spring Cereals Grown under Organic Production in Northern Canada. <i>Agronomy Journal</i> , 2007, 99, 1199-1207.                               | 0.9 | 53        |
| 55 | Integrated approaches to managing weeds in spring-sown crops in western Canada. <i>Crop Protection</i> , 2007, 26, 390-398.  | 1.0 | 57        |
| 56 | The competitive interactions between winter barley and <i>Avena sterilis</i> are site-specific. <i>Weed Research</i> , 2008, 48, 38-47.  | 0.8 | 6         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 57 | PHYTOTOXICITY OF 10 WINTER BARLEY VARIETIES AND THEIR COMPETITIVE ABILITY AGAINST COMMON POPPY AND IVY-LEAVED SPEEDWELL. <i>Experimental Agriculture</i> , 2008, 44, 385-397.  | 0.4 | 14        |
| 58 | Nitrogen Fertilizer Rate Effects on Weed Competitiveness is Species Dependent. <i>Weed Science</i> , 2008, 56, 743-747.  | 0.8 | 110       |
| 59 | Above- and below-ground competition between barley, wheat, lupin and vetch in a cereal and legume intercropping system. <i>Grass and Forage Science</i> , 2009, 64, 401-412.   | 1.2 | 79        |
| 60 | Weed suppression ability of three winter wheat varieties at different row spacing under organic farming conditions. <i>Weed Research</i> , 2009, 49, 526-533.  | 0.8 | 64        |
| 61 | Grain yield increase in cereal variety mixtures: A meta-analysis of field trials. <i>Field Crops Research</i> , 2009, 114, 361-373.  | 2.3 | 161       |
| 62 | Barley and Rigid Ryegrass ( <i>Lolium rigidum</i> ) Competition is Influenced by Crop Cultivar and Density. <i>Weed Technology</i> , 2009, 23, 40-48.  | 0.4 | 44        |
| 63 | Weed communities of transgenic glyphosate-tolerant soyabean crops in ex-pasture land in the southern Mesopotamic Pampas of Argentina. <i>Weed Research</i> , 2010, 50, 320-330.  | 0.8 | 17        |
| 64 | Nitrogen Use Efficiency and Nitrogen Fertilizer Recovery of Durum Wheat Genotypes as Affected by Interspecific Competition. <i>Agronomy Journal</i> , 2010, 102, 707-715.  | 0.9 | 58        |
| 65 | Wide Row Spacing and Rigid Ryegrass ( <i>Lolium rigidum</i> ) Competition Can Decrease Barley Yield. <i>Weed Technology</i> , 2010, 24, 310-318.   | 0.4 | 1         |
| 66 | Competitive interactions of fifty barley cultivars with <i>Avena sterilis</i> and <i>Asperugo procumbens</i> . <i>Field Crops Research</i> , 2010, 117, 90-100.  | 2.3 | 10        |
| 67 | Effects of Changes in Flax ( <i>Linum usitatissimum</i> L.) Density and Interseeding with Red Clover ( <i>Trifolium pratense</i> L.) on the Competitive Ability of Flax Against Brassica Weeds. <i>Agroecology and Sustainable Food Systems</i> , 2011, 35, 914-926. | 0.9 | 7         |
| 68 | Factores biológicos que determinan la competencia de <i>Commelina erecta</i> con otras malezas en sistemas de cultivo. <i>Planta Daninha</i> , 2011, 29, 97-106.   | 0.5 | 6         |
| 69 | Growth, gas exchange and competitive ability of <i>Sorghum halepense</i> populations under different soil water availability. <i>Canadian Journal of Plant Science</i> , 2011, 91, 1011-1025.  | 0.3 | 13        |
| 70 | Inter-plant competition for resources in maize crops grown under contrasting nitrogen supply and density: Variability in plant and ear growth. <i>Field Crops Research</i> , 2011, 121, 373-380.   | 2.3 | 122       |
| 71 | Leaf gas exchange and competitive ability of <i>Zea mays</i> and <i>Sorghum halepense</i> as affected by water competition. <i>Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science</i> , 2012, 60, 231-246.                | 0.2 | 1         |
| 72 | The Biology of Canadian Weeds. 27. <i>Avena fatua</i> L. (updated). <i>Canadian Journal of Plant Science</i> , 2012, 92, 1329-1357.  | 0.3 | 73        |
| 73 | Italian ryegrass ( <i>Lolium multiflorum</i> Lam.) density and N fertilization on wheat ( <i>Triticum aestivum</i> L.) yield in Argentina. <i>Crop Protection</i> , 2012, 32, 36-40.   | 1.0 | 23        |
| 74 | A mathematical model for the evaluation of cooperation and competition effects in intercrops. <i>European Journal of Agronomy</i> , 2013, 51, 9-17.  | 1.9 | 16        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 75 | Breeding Cereal Crops for Enhanced Weed Suppression: Optimizing Allelopathy and Competitive Ability. <i>Journal of Chemical Ecology</i> , 2013, 39, 213-231.  | 0.9 | 112       |
| 76 | Root and shoot competition: a meta-analysis. <i>Journal of Ecology</i> , 2013, 101, 1298-1312.  | 1.9 | 119       |
| 77 | Response of Corn and Redroot Pigweed to Nitrogen Fertilizer in Different Irrigation Regimes. <i>Agronomy Journal</i> , 2013, 105, 1107-1118.  | 0.9 | 4         |
| 78 | The Morphological Features and Canopy Parameters as Factors Affecting the Competition Between Winter Wheat Varieties and Weeds. <i>Journal of Plant Protection Research</i> , 2013, 53, 203-209.  | 1.0 | 2         |
| 79 | Weed suppression ability of six soybean [ <i>Glycine max</i> (L.) Merr.] varieties under natural weed development conditions. <i>Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science</i> , 2013, 61, 43-53. | 0.2 | 6         |
| 80 | The Suitability of Different Winter and Spring Wheat Varieties for Cultivation in Organic Farming. , 2014, , .  |     | 1         |
| 81 | Root discrimination of closely related crop and weed species using FT MIR-ATR spectroscopy. <i>Frontiers in Plant Science</i> , 2015, 6, 765.   | 1.7 | 24        |
| 82 | A review of the potential for competitive cereal cultivars as a tool in integrated weed management. <i>Weed Research</i> , 2015, 55, 239-248.   | 0.8 | 161       |
| 83 | Effects of nitrogen on the growth and competition between seedlings of two temperate forest tree species. <i>Scandinavian Journal of Forest Research</i> , 2015, , 1-7.   | 0.5 | 1         |
| 84 | Biology and management of <i>Avena fatua</i> and <i>Avena ludoviciana</i> : two noxious weed species of agro-ecosystems. <i>Environmental Science and Pollution Research</i> , 2017, 24, 19465-19479.   | 2.7 | 27        |
| 85 | Modelling of low input herbicide strategies for the control of wild oat in intensive winter wheat cropping systems. <i>Field Crops Research</i> , 2017, 201, 1-9.   | 2.3 | 4         |
| 86 | Competitive ability of five common weed species in competition with soybean. <i>International Journal of Pest Management</i> , 2017, 63, 30-36.   | 0.9 | 26        |
| 87 | Differences for traits associated with early N acquisition in a grain legume and early complementarity in grain legume-triticale mixtures. <i>AoB PLANTS</i> , 2018, 10, ply001.  | 1.2 | 10        |
| 88 | <i>Avena fatua</i> L. escapes and delayed emergence in wheat ( <i>Triticum aestivum</i> L.) crops of Argentina. <i>Crop Protection</i> , 2018, 103, 30-38.  | 1.0 | 7         |
| 89 | Weed Control Through Crop Plant Manipulations. , 2018, , 73-96.   |     | 11        |
| 90 | The combined effects of false seedbed technique, post-emergence chemical control and cultivar on weed management and yield of barley in Greece. <i>Phytoparasitica</i> , 2020, 48, 131-143.   | 0.6 | 23        |
| 91 | Estimations and projections of <i>Avena fatua</i> dynamics under multiple management scenarios in crop fields using simplified longitudinal monitoring. <i>PLoS ONE</i> , 2021, 16, e0245217.   | 1.1 | 6         |
| 93 | Interspecific interactions regulate plant reproductive allometry in cereal-legume intercropping systems. <i>Journal of Applied Ecology</i> , 2021, 58, 2579-2589.   | 1.9 | 6         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 94  | Crop Competitiveness. SpringerBriefs in Agriculture, 2014, , 9-20.   | 0.9 | 2         |
| 95  | Weed Suppression and Tolerance in Winter Oats. Weed Technology, 2017, 31, 740-751.   | 0.4 | 10        |
| 96  | Ecological risks of transgenic virus-resistant crops. , 2001, , 125-142.   |     | 3         |
| 97  | Características de plantas de arroz e a habilidade competitiva com plantas daninhas. Planta Daninha, 2003, 21, 165-174.  | 0.5 | 23        |
| 98  | Alocação de matéria seca e capacidade competitiva de cultivares de milho com plantas daninhas. Planta Daninha, 2011, 29, 373-382.  | 0.5 | 15        |
| 99  | Effects of different densities of wild oat ( <i>Avena fatua</i> L.) and nitrogen rates on oilseed rape ( <i>Brassica</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj ETQq0 0 0 rgBT /Overlock 10 Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj ETQq0 0 0 rgBT /Overlock 10     | 1.6 | 4         |
| 100 | Competition between <i>Avena sterilis</i> ssp. <i>sterilis</i> and wheat in South Western Spain. Spanish Journal of Agricultural Research, 2011, 9, 862.   | 0.3 | 8         |
| 101 | The Influence of interaction between spring wheat and spring barley on accumulation of nitrogen, phosphorus and potassium in plants. Annales Univesitatis Mariae Curie-Skłodowska Sectio E Agricultura, 2009, 64, .  | 0.1 | 0         |
| 102 | The intensity of competitive interactions between spring wheat ( <i>Triticum aestivum</i> L. emend. Fiori et.) Tj ETQq0 0 0 rgBT /Overlock 10 Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj ETQq0 0 0 rgBT /Overlock 10 Tj ETQq1 1 0.784314 rgBT /Overlock 10                 | 1.0 | 4         |
| 103 | Changes in spring wheat ( <i>Triticum aestivum</i> ssp. <i>vulgare</i> L.) and persian clover ( <i>Trifolium resupinatum</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj ETQq0 0 0 rgBT /Overlock 10 Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tj ETQq0 0 0 rgBT /Overlock 10 | 1.0 | 3         |
| 104 | <i>Avena fatua</i> L.: Wild oat. Acta Herbologica, 2017, 26, 75-86.  | 0.2 | 0         |
| 105 | Intercropping organic field peas with barley, oats and mustard improves weed control but has variable effects on grain yield and net returns. Canadian Journal of Plant Science, 0, , .  | 0.3 | 1         |
| 106 | Wpływ oddziaływań, pomiędzy pszenicą... jarzą... a zmianami jarym na akumulację w roślinach azotu, fosforu i potasu. Agronomy Science, 2009, 64, 93-106.   | 0.1 | 0         |
| 108 | Negative relationship between topsoil root production and grain yield in oat and barley. Agriculture, Ecosystems and Environment, 2023, 349, 108467.   | 2.5 | 0         |